



LETTER TO THE EDITOR

Ulcerate colitis and protein C system: is there a link of causality

Xuhong Lin¹ · Dandan Wei¹ · Huichao Wang² · Zhaoya Yu¹ · Yuxia Li¹ · Yanmin Li¹ · Yanan Wen¹ · Xuequn Ren³ · Guanchang Cheng⁴

Accepted: 19 November 2015 / Published online: 5 December 2015

© The Author(s) 2015. This article is published with open access at Springerlink.com

Dear Editor:

Ulcerate colitis (UC), the main type of inflammatory bowel disease (IBD), is an unexplained immune-mediated intestinal mucosal inflammation with higher cancer risk. Because of an interaction between inflammation and coagulation in the disease progression, hypercoagulative state is often seen in patients of UC. Endothelial cell damage plays an important role in the activation of coagulation system in patients with UC, while protein C (PC) system, a well-characterized anticoagulant system located on the vascular endothelial cell surface, which consists of protein S (PS), endothelial cell protein C receptor (EPCR), PC, protein C inhibitor, and thrombomodulin (TM), is a mediator of endothelial function. Recent studies indicated that PC system plays an important

role in regulating intestinal homeostasis both in human intestinal microvascular endothelial cells and animal models of colitis. Therefore, we speculate that the imbalance of PC system may be one of the pathogenesis of UC patients. Our *in vivo* study verified that the degree of inflammation was negatively correlated with the PC activity, and mouse colonic macrophages and mucosa microvascular endothelial cells are suggested as the major mediators involved. Until today, it is unclear why PC system changed, and how macrophages and mucosa microvascular endothelial cells interact with each other in UC pathology, so it is important to investigate the changes and mechanisms of PC system, and to find new drug target. Further research *in vitro* revealed that TNF- α and IL-6 levels were increased in the supernatant of macrophages from

This study is supported by The National Science Foundation of China (No.U1304802 to Dr. X.-H. Lin).

Xuhong Lin, Dandan Wei and Huichao Wang contributed equally to this work.

✉ Xuequn Ren
281773134@qq.com

✉ Guanchang Cheng
113072955@qq.com

Xuhong Lin
lxh80726@126.com

Dandan Wei
wei-dandan@163.com

Huichao Wang
uhchao@163.com

Zhaoya Yu
759789925@qq.com

Yuxia Li
1821941279@qq.com

Yanmin Li
471421134@qq.com

Yanan Wen
897890575@qq.com

- 1 Department of Clinical Laboratory, Translational Medicine Center, Huaihe Hospital Affiliated to Henan University, 115 Ximen Street, Kaifeng City, Henan Province, China
- 2 Department of Nephrology, First Affiliated Hospital of Henan University, 357 Ximen Street, Kaifeng City, Henan Province, China
- 3 Department of General Surgery, Huaihe Hospital Affiliated to Henan University, 115 Ximen Street, Kaifeng City, Henan Province, China
- 4 Department of Cardiovascular Medicine, Huaihe Hospital Affiliated to Henan University, 115 Ximen Street, Kaifeng City, Henan Province, China

dextran sulfate sodium (DSS) mice colonic tissue, and these conditions were only seen in CD14⁺CD64⁺ subtype macrophages. In addition, after incubation of TNF- α or IL-6 with colonic mucosal microvascular endothelial cells, the activity of PC system was inhibited, manifested as decreased APC activity, and downregulated expression of EPCR. Furthermore, β -arrestin-pJNK MAPK expressions were upregulated by real-time PCR and Western blot. Taken together into account, due to complex UC mechanism, its pathogenesis is unclear, and there is little research about colonic mucosal microvascular, especially the study for PC system changes in the pathogenesis of UC is rare. In our study, we found that in DSS-induced mouse colitis, colonic CD14⁺CD64⁺ macrophages produced more inflam-

matory cytokines, which result in inhibited PC pathway via mucosal microvascular endothelial cells, and β -arrestin-pJNK MAPK signal pathway may involved. Therefore, enhancing the activity of PC system provides a major theoretical and clinical significance to find new drug target for the treatment of UC.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.