

## MEETING ABSTRACT

## Open Access

# Emphysematous cholecystitis. Advantages of abdominal ultrasound in the ED

M Algaba-Montes<sup>1\*</sup>, A Oviedo-García<sup>1</sup>, D Nuñez-Hospital<sup>2</sup>, J Lopez-Libano<sup>3</sup>, JM Alvarez-Franco<sup>4</sup>, N Diaz-Rodriguez<sup>5</sup>, A Rodriguez-Lorenzo<sup>6</sup>*From* 9th WINFOCUS World Congress on Ultrasound in Emergency and Critical Care Hong Kong. 6-9 November 2013

## Background

Emphysematous cholecystitis (EC) is an entity with high morbidity and mortality, and therefore require a diagnosis agile and dynamic, allowing appropriate management to avoid complications. The emergency ultrasound (US) allows a versatile and comprehensive management, improving the prognosis of this disease in the majority of cases.

## Objective

we present a case of EC, diagnosed at ER, through the use of US scanning used by Emergency Physicians.

## Patients and methods

a patient with abdominal pain, with a final diagnosis of an EC assessing US, performed by EP.

## Results

We report the case of a 72 year old patient with prior stroke without sequelae and hypertensive, with abdominal pain of 7 days duration, high fever and bilious vomiting. Physical examination was marked hypotension (80/45 mmHg), distal coldness, pallor and sweating, 38.5 ° C, 145 spm. The distended abdomen with abolished peristalsis and positive Murphy right upper quadrant. Rest without findings of interest. Analytically glucose was 505 mg / dl, creatinine of 2.44 mg / dl, bilirubin 2.2 mg / dl, AST 350, LDH 407, amylase 125, 19500 leukos with neutrophilia and pH of 7.13, with lactic 12. Was performed in consultation abdominal US showed a thickened gallbladder wall (8 mm), well-circumscribed, oval, distended and gas in the same light, compatible with emphysematous cholecystitis. Support

measures were initiated, antibiotics, insulin therapy and emergency surgery was indicated. This allowed a favorable high after joining UCI in 7 days without further complications.

## Conclusion

EC is a rare entity that represents 1% of all cholecystitis, clinically indistinguishable, but with a worse prognosis (25% mortality) and more complications. Here debut comes as poorly controlled diabetes. The use of abdominal US in ER allows for both a rapid and versatile, with proper treatment start, this being vital to good patient outcomes.

## Authors' details

<sup>1</sup>Emergency Department, Valme Hospital, Seville, Members of the Working Group of Ultrasound SEMES\_Andalucía and SEMERGEN, Spain. <sup>2</sup>Emergency Department, Valme Hospital. Seville, Spain. <sup>3</sup>Critical Care Department, Miramar Hospital, Mallorca, Member of the Working Group of Ultrasound SEMERGEN, Spain. <sup>4</sup>Emergency Department, IB-Salut, Ibiza, Member of the Working Group of Ultrasound SEMERGEN, Spain. <sup>5</sup>Primary Care. Barbadás Primary Care Center. Ourense, Member of the Working Group of Ultrasound SEMERGEN, Spain. <sup>6</sup>Radiology Department. Perpetuo Socorro Hospital, Vigo, Member of the Working Group of Ultrasound SEMERGEN, Spain.

Published: 31 January 2014

## Reference

1. Fox JC, Solley M, Anderson CL, et al: Prospective evaluation of emergency physician performed bedside ultrasound to detect acute appendicitis. *Eur J Emerg Med* 2008, **15**(2):80-5.

doi:10.1186/2036-7902-6-S1-A7

Cite this article as: Algaba-Montes et al.: Emphysematous cholecystitis. Advantages of abdominal ultrasound in the ED. *Critical Ultrasound Journal* 2014 **6**(Suppl 1):A7.

<sup>1</sup>Emergency Department, Valme Hospital, Seville, Members of the Working Group of Ultrasound SEMES\_Andalucía and SEMERGEN, Spain  
Full list of author information is available at the end of the article