

RESEARCH

Open Access

Effect of a fever control protocol-based strategy on ventilator-associated pneumonia in severely brain-injured patients

Yoann Launey^{1,2,3*}, Nicolas Nessler^{1,2,3}, Audren Le Cousin^{1,3}, Fanny Feuillet⁴, Ronan Garlantezec⁵, Yannick Maldant^{1,2,3} and Philippe Seguin^{1,2,3}

Abstract

Introduction: Fever is associated with a poor outcome in severely brain-injured patients, and its control is one of the therapies used in this condition. But, fever suppression may promote infection, and severely brain-injured patients are frequently exposed to infectious diseases, particularly ventilator-associated pneumonia (VAP). Therefore, we designed a study to explore the role of a fever control protocol in VAP development during neuro-intensive care.

Methods: An observational study was performed on severely brain-injured patients hospitalized in a university ICU. The primary goal was to assess whether fever control was a risk factor for VAP in a prospective cohort in which a fever control protocol was applied and in a historical control group. Moreover, the density of VAP incidence was compared between the two groups. The statistical analysis was based on a competing risk model multivariate analysis.

Results: The study included 189 brain-injured patients (intervention group, $n = 98$, and historical control group, $n = 91$). The use of a fever control protocol was an independent risk factor for VAP (hazard ratio 2.73, 95% confidence interval (1.38, 5.38; $P = 0.005$)). There was a significant increase in the incidence of VAP in patients treated with a fever control protocol (26.1 versus 12.5 VAP cases per 1000 days of mechanical ventilation). In cases in which a fever control protocol was applied for >3 days, we observed a higher rate of VAP in comparison with the rate among patients treated for ≤ 3 days.

Conclusions: Fever control in brain-injured patients was a major risk factor for VAP occurrence, particularly when applied for >3 days.

Introduction

In severely brain-injured patients, the incidence of fever in the first week of hospitalization approaches 87% [1]. Numerous studies have noted that fever in these patients is associated with a prolonged length of stay in the intensive care unit (ICU) and hospital, severe physical dependency, cognitive impairment and increased mortality [2,3]. Thus, fever control is a common therapy used to treat patients with severe brain injuries [4]. Nevertheless, specific guidelines concerning the optimal delay, length, means and target of temperature control are lacking [4,5].

From another perspective, several experimental and clinical studies that have assessed fever control in other diseases have suggested that there are harmful effects associated with a fever control strategy [6-9]. These adverse effects of fever control, notably shivering, infectious diseases and/or hypotension, are negative factors affecting the prognoses of brain-injured patients [10,11]. In fact, fever, which is typically defined by an elevation of the core temperature above 38.2°C [12,13], may be an adaptive response to injury when an infectious disease is ongoing [14]. Nosocomial infections, particularly ventilator-associated pneumonia (VAP), are frequently observed in severely brain-injured patients, affecting from 8% to more than 70% of patients [15,16]. Therefore, we designed this study to define the role of a fever control protocol in VAP development during neuro-intensive care.

* Correspondence: yoann.launey@chu-rennes.fr

¹CHU de Rennes. Hpital Pontchaillou. Ple Anesthsie-SAMU-Urgences-Ranimations. 2, rue Henri Le Guilloux, 35033 Rennes, Cedex, France

²Inserm UMR991, quipe 2: Stress, Defense and Regeneration. 2, rue Henri Le Guilloux, 35000 Rennes, France

Full list of author information is available at the end of the article

Materials and methods

This observational study was conducted between January 2008 and December 2009 in the surgical ICU of a university hospital, which received, among others, severely brain-injured patients. The criteria for selection included a severe brain injury (traumatic brain injury (TBI), subarachnoid hemorrhage (SAH), or intracranial hemorrhage), a Glasgow Coma Scale score ≤ 8 and a need for mechanical ventilation (MV) for 48 hours or longer. Each selected patient underwent chart review. Patients who had aspiration pneumonia at admission or a brain computed tomography (CT) scan without any lesion or who received MV for less than 48 hours were excluded. This study was submitted to the local ethical committee of Rennes University Hospital (CHU Pontchaillou, Rennes, no. 14.45), which stated that no patient consent was needed.

In 2008, Greer *et al.* suggested that fever had a deleterious effect in brain-injured patients and that there was an evidence-based need to confirm the potential benefit of fever control in this population [2]. Consequently, from 1 January 2009 until 31 December 2009, a fever control protocol was applied in cases in which the core body temperature was $>38.2^{\circ}\text{C}$. Because the fever control protocol was the only therapeutic change, patients admitted from 1 January 2008 to 31 December 2008 served as historical controls. In the fever-control group, patients were prospectively and consecutively included, whereas in the historical control group, patients were identified and selected from the list of admissions to our ICU.

The primary endpoint was to assess whether fever control was a risk factor for VAP. The secondary endpoints were the density of incidence and the incidence of VAP, both nosocomial urinary tract infections (UTIs) and catheter-related bloodstream infections (CRBIs), the duration of MV, the length of stay in the ICU, and the mortality rate.

Patient management

During the two study periods, brain-injured patients were similarly managed according to identical local protocols based on international guidelines [17,18]. The two major objectives were as follows: 1) to infuse neurosedation combining midazolam and opioids; and 2) to maintain cerebral perfusion pressure ≥ 70 mmHg. All the patients were orotracheally intubated; VAP prevention measures included hand washing with alcohol based antiseptic, a 30 semi-recumbent patient position, MV with positive end-expiratory pressure >5 mmHg, and oral care based on local application of povidone-iodine solution six times daily [15]. Selective digestive decontamination or subglottic drainage was not performed, and ulcer prophylaxis was not systematically administered. The core body temperature was continuously monitored with a rectal

thermal probe (Mon-a-Therm, Covidien, Mansfield, MA, USA).

Fever control intervention was applied in cases in which the body core temperature was $>38.2^{\circ}\text{C}$ [12]. This protocol was established based on previously published studies [4,19,20] and successively included the following: external cooling, an infusion of 500 mL of cooled (4°C) saline serum and enteral administration of acetaminophen. A neuromuscular blocker (cisatracurium) was also used if shivering occurred. The objective was to reach a core body temperature of 36°C to 38.2°C .

Data collection and definitions

Gender, age, body mass index, the reason for ICU admission, a history of smoking, a positive alcohol test at admission and CT scan-identified lung contusions involving more than one lobe were recorded for each patient at the time of admission to the ICU. The severity of the brain injury was assessed at admission using the Glasgow Coma Scale score and the Simplified Acute Physiological Score II (SAPS II) [21].

We collected the daily minimum (T_{min}) and maximum (T_{max}) body core temperature and calculated the mean daily temperature as $(T_{\text{max}} + T_{\text{min}})/2$ during the first eight days of ICU hospitalization. We reported the delay between ICU admission and the initiation of the fever control protocol, the length of application of the fever control protocol and mortality at ICU discharge. Antibiotic prophylaxis, neuromuscular blocker agents, mannitol and thiopental use was reported for all patients. Additionally, acute respiratory distress syndrome (ARDS), defined as previously [22], and microbiological findings for VAP were reported.

VAP was defined as new and persistent pulmonary infiltrates on a chest radiograph occurring after 48 hours of MV, combined with at least two of the following criteria: 1) purulent tracheal secretions and/or body core temperature $>38^{\circ}\text{C}$ and/or leukocytosis $>10,000/\text{mm}^3$ or leukopenia $<3,000/\text{mm}^3$; and 2) microbiological confirmation via an endotracheal aspirate quantitative culture growing $\geq 10^6$ colony forming units (cfu)/mL [23]. The occurrence of VAP was reported and expressed as the density of incidence, defined as the number of VAP cases per 1,000 ventilator-days. We also defined early-onset VAP as VAP that occurred ≤ 5 days after admission and late-onset VAP as VAP diagnosed >5 days, according to the American Thoracic Society criteria [24].

UTIs and CRBIs were defined according to the CDC guidelines [25,26].

Statistical analysis

The categorical parameters are expressed as number and frequency. The quantitative parameters are expressed as the mean (standard deviation) if there was a normal

distribution or the median (interquartile range) otherwise. The categorical parameters were compared using the Chi-square test or, when necessary, Fisher's exact test. The quantitative parameters were compared using the two-tailed *t*-test (when normally distributed) or with the Mann Whitney *U* test. The temperature value was recorded on the date of the VAP diagnosis. To determine risk factors for VAP occurrence, we used a competing risk model, as proposed by Fine and Gray [27]. VAP before the 28th day of hospitalization in the ICU was an event of interest but could be precluded by death before the 28th day of hospitalization. Patients were therefore observed from the day of hospitalization to the occurrence of VAP before the 28th day of hospitalization in the ICU, to death before the 28th day of hospitalization, or to censoring (hospital discharge while living, without VAP occurrence at day 28 or before). We first compared patient characteristics according to whether the fever control protocol was used to identify potential confounding factors existing before VAP occurrence. A univariate analysis was first performed. Variables with *P*-values <0.20 were then included in the multivariate competing risk model, and a backward selection was applied. The model was adjusted for the risk factors selected according to use of the fever control protocol. By stratifying the duration of fever control, we analyzed the effect of the duration of fever control on VAP occurrence. The adjusted hazard ratios (HRs) and their corresponding 95% confidence intervals (CIs) were calculated, and the cumulative incidence curves for VAP were determined using the Nelson-Aalen non-parametric estimator. By stratifying the fever control duration, we could also describe the impact of the fever control duration on VAP based on the cumulative incidence curve with the Nelson-Aalen non-parametric estimator. A two-sided *P*-value <0.05 was considered statistically significant. The statistical models were built using SAS 9.3 (SAS Institute, Inc., Cary, NC, USA) and R Core Team (2012) software (R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria).

Results

During the study period, 331 brain-injured patients were screened, and 189 were included (intervention group, *n* = 98; historical control group, *n* = 91) (Figure 1). The primary characteristics of the patients are presented in Table 1. The proportion of subarachnoid hemorrhaging was significantly higher in the intervention group than in the historical control group (28% versus 14%, respectively; *P* = 0.006). The clinical characteristics are reported in Table 2.

Management of the fever control protocol

All patients in the intervention group were treated with the fever control protocol when their fever was >38.2°C.

The delay before its effective application was less than 24 hours, and its median duration was four days (range, two to six days). The daily mean temperature curves for each group are displayed in Figure 2.


The density of the incidence of VAP was significantly increased in the intervention group beginning on ICU day 5 (26.1 versus 12.5 VAP cases per 1,000 days of ventilation; *P* <0.001). The incidence of VAP was 38% (*n* = 37) versus 12% (*n* = 11) for the intervention group and historical control group, respectively. The mean delay for VAP occurrence was similar in the two groups (8.4 days). The mean length of MV was significantly increased in the intervention group (15.11 versus 10.7 days, *P* <0.001). Methicillin-sensitive *Staphylococcus aureus* (MSSA), *Streptococcus pneumoniae* and *Haemophilus influenzae* comprised 80% of the microbiological findings. The microbiological findings are detailed in Table 3. Considering the VAP-onset delay, we reported 10 early-onset VAP cases in the intervention group compared with 5 in the historical control group and 27 late-onset VAP cases in the intervention group compared with 6 in the historical control group.

The densities of incidence of UTIs and CRBIs were not significantly different between the intervention and the historical control groups (9.7 versus 14.7 UTIs per 1,000 urinary catheter-days, respectively; *P* = 0.9; 3.9 versus 1.6 CRBIs per 1,000 catheter-days, respectively; *P* = 0.19). The mortality rate also did not differ between the two groups (34% in the intervention group versus 23% in the historical control group; *P* = 0.107).

In the univariate analysis (Table 4), the patients who developed VAP were significantly younger, used tobacco more frequently and had more extensive lung injury at admission to the ICU. Application of the fever control protocol was a major risk factor for developing VAP (HR 3.06; 95% CI (1.58, 5.94)). In the multivariate analysis, the competing risk model was built after adjustment of the HR for a history of smoking, age, the SAPS II score, pentothal use and the use of neuromuscular blocker agents. The use of the fever control protocol was the only significant independent risk factor for VAP occurrence (HR 2.73, 95% CI (1.38, 5.38)) (Table 4). The cumulative incidence function curves for VAP are displayed in Figure 3. The risk of VAP appeared to be higher when the duration of the fever control protocol was >3 days (Figure 4).

Discussion

In this study, we found that the use of a fever control protocol during the early management of critically brain-injured patients was associated with an increased risk of VAP. During ICU hospitalization of brain-injured patients, sepsis and VAP are major causes of death [23]. In particular, VAP is estimated to occur in 25% to 70% of critically ill



patients [23,28,29]. The incidence of VAP in our severely brain-injured patients was 26%, and the microbiological results were relatively similar to those of previous studies [15,30,31], which predominantly found the following microorganisms: MSSA, *Haemophilus influenzae* and *Streptococcus pneumoniae*. It is noteworthy that VAP incidence could be underestimated in the intervention group because fever is a clinical sign of VAP; consequently, the difference in VAP occurrence between the two groups could be minimized. The deleterious effect of such infections has recently been reported in two meta-

analyses, which showed that VAP was associated with increased mortality [32,33]. However, in a subgroup of trauma patients, no significant increase in mortality was observed [33].

VAP in severely brain-injured patients has a different pattern than in other critically ill patients. Several factors favor the development of VAP. In brain-injured patients, VAP typically occurs early in the ICU stay, between day 5 and day 11 after hospitalization. The presence of abnormal protection of the glottal area and consecutive early micro-inhalation is typically suggested to explain the occurrence of VAP. These aspirations predominantly contain oropharyngeal bacteria from the commensal flora of the patient, as shown by Ewig *et al.* [34]. In this context, pneumonia results from extensive tracheal and bronchial inocula in patients with altered immune responses and ventilation-related impairment (including due to an endotracheal tube, sedation, or neuromuscular blockers) [35,36].

Fever may be an adaptive response to injury, particularly in cases of ongoing infectious disease. When this response is suppressed, several studies have indicated an increased risk of infectious disease and, especially, VAP. In particular, randomized studies have shown an increased rate of VAP in groups treated with therapeutic hypothermia [37,38]. For instance, although the methodology is debatable, a meta-analysis including eight studies of patients with TBIs reported an increased rate of VAP in the group receiving therapeutic hypothermia (OR 2.37, 95% CI (1.37, 4.10)) [37]. Similarly, Geurts *et al.* found an increased risk of VAP (risk ratio 1.44, 95% CI (1.10, 1.90)) and sepsis (risk ratio 1.80, 95% CI

Table 1 Baseline characteristics

Baseline characteristics	Intervention group	Control group	P-value
	Number = 98	Number = 91	
Age	45 (25 to 57)	53 (27 to 66)	0.022
Gender (male)	72 (73%)	64 (70%)	0.631
BMI	25 4	24 5	0.123
Reason for admission			0.012
TBI	64 (65%)	61 (67%)	
SAH	27 (28%)	13 (14%)	
ICH	7 (7%)	17 (19%)	
Smoking	38 (39%)	23 (25%)	0.048
Positive alcohol test	11 (11%)	14 (15%)	0.399
Lung contusion	28 (29%)	17 (19%)	0.111
GCS	6 (3 to 8)	6 (3 to 7)	0.338
SAPS II	49 11	53 14	0.013

BMI: body mass index; GCS: Glasgow Coma Scale; ICH: intracranial hemorrhage; SAH: subarachnoid hemorrhage; SAPS II: Simplified Acute Physiology Score II; TBI: traumatic brain injury.

Table 2 Clinical characteristics during ICU management

Clinical characteristics	Intervention group	Control group	P-value
	Number = 98	Number = 91	
ARDS, yes	4 (4%)	2 (2%)	0.684
Antibiotic prophylaxis	43 (44%)	43 (47%)	0.645
NMB agents	13 (13%)	5 (5%)	0.116
Pentothal use, yes	39 (40%)	19 (21%)	0.005
Mannitol use, yes	49 (50%)	21 (23%)	<0.001
ICU length of stay, days	17 13	11 8	<0.001
Mortality, number (%)	23 (23%)	31 (34%)	0.107

ARDS: acute respiratory distress syndrome; ICU: intensive care unit; NMB: neuromuscular blocker.

(1.04, 3.10)) in patients treated with therapeutic hypothermia, regardless of what surface or endovascular cooling devices were used [38].

To overcome this adverse effect, normothermia-targeted temperature management has been suggested. However, data on the effects of normothermia in neurologically injured patients are limited. Several studies have investigated and compared the efficacy of different devices in maintaining normothermia but, despite their conclusions, have reported no difference in adverse effects. These studies were not designed to evaluate the rate of infection [39-41]. Additionally, Puccio *et al.* retrospectively compared the effects of a normothermia protocol on TBI management

[42]. They found a beneficial effect on intracranial pressure and on the duration of intracranial hypertension episodes in the normothermia group. The systematic initiation of a normothermia protocol, even prophylactically, allowed fever control and possibly attenuation of secondary brain injuries. Similarly, Badjatia *et al.* assessed the effect of normothermia in SAH in a case control study [11]. They found better fever control, without a significant increase in the rate of VAP (a 58% rate of VAP in the therapeutic normothermia group versus 51% in the historical control group; $P = 0.08$).

In our study, we observed that a sustained intervention for >3 days was more likely to aggravate the incidence of VAP compared with an intervention with a duration of ≤ 3 days. To the best of our knowledge, no previous study has investigated the effect of the duration of normothermia. It appears that the targeted body temperature does not affect VAP occurrence; however, counteracting the physiological fever response to stress and its duration may be a key factor in the increased occurrence of VAP. More specifically, although fever is deleterious in the short and medium term in brain-injured patients, fever control, whether by hypo- or normothermia, allows reduction of the intracranial pressure at the expense of increased susceptibility to infections. Moreover, in therapeutic hypothermia, the hypothesized anti-inflammatory effect has been associated with impaired cellular and humoral immunity [43]. Unfortunately,




Figure 2 Mean daily body core temperature curves for the first eight days of ICU stay. Upper (dashed line) and lower (dark line) limits of standard deviation are displayed for the control group and the intervention group, respectively.

Table 3 Microbiological findings for VAP

Bacterium	Total	Intervention group	Control group
	Number = 62	Number = 48	Number = 14
Gram-positive cocci			
MSSA	22	17	5
MRSA	1	0	1
<i>Streptococcus pneumoniae</i>	11	8	3
<i>Streptococcus anginosus</i>	1	0	1
Gram-negative bacilli			
<i>Haemophilus influenzae</i>	17	14	3
<i>Escherichia coli</i>	3	3	0
<i>Moraxella catarrhalis</i>	1	0	1
<i>Klebsiella oxytoca</i>	1	1	0
<i>Proteus mirabilis</i>	1	1	0
<i>Citrobacter freundii</i>	1	1	0
<i>Enterobacter cloacae</i>	1	1	0
<i>Acinetobacter baumannii</i>	1	1	0

MSSA: methicillin-sensitive *Staphylococcus aureus*; MRSA: methicillin-resistant *Staphylococcus aureus*; VAP: ventilator-associated pneumonia.

to our knowledge, no study has assessed the underlying brain physiology in fever control targeting normothermia.

Concerning the strengths and limitations of our study, the survey of major nosocomial infections in our ICU was prospective starting in 2002, and the diagnostic criteria did not change between the two study periods. Although the use of competing risk models in the multivariate analysis ensured the strength of our study, the major limitation is the design of the study, which used a historical control group, consequently implying limited matching between the two study groups. The interpretation of the

results should consider the difference in the mean age and the reason for admission between each group, particularly the higher number of SAH in the control group. Additionally, we have accounted for the confounding factors that may impact the occurrence of VAP, but red blood cell transfusion, level of sedation and antibiotic use during ICU stay were not reported. Head of bed elevation was not reported but standard care in our ICU uses a 30 semi-recumbent patient position. Finally, the fever threshold chosen to start the fever control protocol should be discussed; certain studies have used a temperature threshold of 38C to start fever control interventions, but solid evidence to clearly support this threshold is lacking.

Conclusions

Protocol-based control of fever in patients with severe head injuries is associated with an increased risk of VAP during the patients ICU stay. Our results suggest that when targeted normothermia is used, the probability of developing an infectious fever within 72 hours is low. However, if fever occurs later, the probability of an infectious fever is higher and should alert intensivists to an ongoing infection. Because data on the impact of fever control and its side effects on the outcome of brain-injured patients are scarce, a large randomized study is required.

Key messages

The fever control in severely brain-injured patients is a major risk factor for VAP. Moreover, the VAP rate appears to be higher if fever control lasts longer than 3 days.

Table 4 Risk factors for VAP occurrence before day 28 (univariate and multivariate analyses)

Parameters	Univariate analysis			Multivariate analysis		
	HR	95% CI (HR)	P-value	HR	95% CI (HR)	P-value
Age	0.97	(0.95, 0.98)	<0.001	0.97	(0.95, 0.98)	<0.001
Gender (male)	1.82	(0.88, 3.78)	0.110			
BMI	0.99	(0.93, 1.05)	0.730			
Smoking	1.89	(1.09, 3.30)	0.024	1.67	(0.95, 2.93)	0.078
Alcohol intake	0.85	(0.35, 2.09)	0.730			
Lung contusion	2.41	(1.37, 4.24)	0.002			
GCS	0.97	(0.85, 1.12)	0.710			
SAPS II	0.98	(0.95, 1.00)	0.068	1.01	(0.98, 1.04)	0.590
NMB agents	2.11	(0.97, 4.61)	0.061			
Mannitol	1.25	(0.72, 2.17)	0.430			
Thiopental	1.36	(0.78, 2.38)	0.270	0.95	(0.53, 1.69)	0.950
Fever control	3.06	(1.58, 5.94)	<0.001	2.73	(1.38, 5.38)	0.005

BMI: body mass index; GCS: Glasgow Coma Scale; NMB: neuromuscular blocker; SAPS II: Simplified Acute Physiology Score II; VAP: ventilator-associated pneumonia.




Figure 3 Cumulative incidence function curve for VAP occurrence according to fever control protocol management. The non-parametric estimator according to Nelson-Aalen. VAP: ventilator-associated pneumonia.




Figure 4 Cumulative incidence function curve for VAP occurrence according to the duration of the fever control protocol when applied. VAP: ventilator-associated pneumonia.

Abbreviations

ARDS: acute respiratory distress syndrome; CI: confidence interval; CRBI: catheter-related bloodstream infection; CT: computed tomography; HR: hazard ratio; ICU: intensive care unit; MSSA: methicillin-sensitive *Staphylococcus aureus*; MV: mechanical ventilation; SAH: subarachnoid hemorrhage; SAPS II: Simplified Acute Physiological Score II; TBI: traumatic brain injury; UTI: urinary tract infection; VAP: ventilator-associated pneumonia.

Competing interests

The authors declare that they have no competing interests.

Authors contributions

YL, AL and PS designed and initiated the study. AL collected data and created the database of brain-injured patients. RG and FF participated in the statistical analysis of the data. YL, AL, PS and FF created the tables and figures. YL, AL, NN, PS and YM interpreted the data. YL, NN, PS and YM contributed in the writing of the manuscript. All authors read and approved the manuscript.

Author details

¹CHU de Rennes. Hpital Pontchaillou. Ple Anesthésie-SAMU-Urgences-Ranimations. 2, rue Henri Le Guilloux, 35033 Rennes, Cedex, France. ²Inserm UMR991, quipe 2: Stress, Defense and Regeneration. 2, rue Henri Le Guilloux, 35000 Rennes, France. ³Université Rennes 1, 35033 Rennes, Cedex, France. ⁴Plateforme de Biométrie Cellule de promotion la recherche clinique, CHU Nantes, EA 4275 SPHERE Biostatistics, Pharmacoepidemiology & Human Sciences Research, UFR de Pharmacie, Université de Nantes. 1, rue Gaston Veil, 44000 Nantes, France. ⁵Ecole Nationale Hautes Etudes de Santé Publique. 2, Avenue du Professeur Lon Bernard, 35033 Rennes, Cedex, France.

Received: 18 July 2014 Accepted: 25 November 2014

Published online: 15 December 2014

References

1. Stocchetti N, Rossi S, Zanier ER, Colombo A, Beretta L, Citerio G: **Pyrexia in head-injured patients admitted to intensive care.** *Intensive Care Med* 2002, **28**:1555-1562.
2. Greer DM, Funk SE, Reaven NL, Ouzounelli M, Uman GC: **Impact of fever on outcome in patients with stroke and neurologic injury: a comprehensive meta-analysis.** *Stroke* 2008, **39**:3029-3035.
3. Diringner MN, Reaven NL, Funk SE, Uman GC: **Elevated body temperature independently contributes to increased length of stay in neurologic intensive care unit patients.** *Crit Care Med* 2004, **32**:1489-1495.
4. Badjatia N: **Hyperthermia and fever control in brain injury.** *Crit Care Med* 2009, **37**:S250-S257.
5. McIlvoy L: **Fever management in patients with brain injury.** *AACN Adv Crit Care* 2012, **23**:204-211.
6. Jiang Q, Cross AS, Singh IS, Chen TT, Viscardi RM, Hasday JD: **Febrile core temperature is essential for optimal host defense in bacterial peritonitis.** *Infect Immun* 2000, **68**:1265-1270.
7. Su F, Nguyen ND, Wang Z, Cai Y, Rogiers P, Vincent JL: **Fever control in septic shock: beneficial or harmful?** *Shock* 2005, **23**:516-520.
8. Hersch M, Raveh D, Izbicki G: **Effect of intravenous propacetamol on blood pressure in febrile critically ill patients.** *Pharmacotherapy* 2008, **28**:1205-1210.
9. Schulman CI, Namias N, Doherty J, Manning RJ, Li P, Alhaddad A, Lasko D, Amortegui J, Dy CJ, Dlugasch L, Baracco G, Cohn SM: **The effect of antipyretic therapy upon outcomes in critically ill patients: a randomized, prospective study.** *Surg Infect* 2005, **6**:369-375.
10. Oddo M, Frangos S, Maloney-Wilensky E, Andrew Kofke W, Le Roux PD, Levine JM: **Effect of shivering on brain tissue oxygenation during induced normothermia in patients with severe brain injury.** *Neurocrit Care* 2010, **12**:10-16.
11. Badjatia N, Fernandez L, Schmidt JM, Lee K, Claassen J, Connolly ES, Mayer SA: **Impact of induced normothermia on outcome after subarachnoid hemorrhage: a case control study.** *Neurosurgery* 2010, **66**:696-700. discussion 700-701.
12. O'Grady NP, Barie PS, Bartlett JG, Bleck T, Carroll K, Kalil AC, Linden P, Maki DG, Nierman D, Pasculle W, Masur H: **Guidelines for evaluation of new fever in critically ill adult patients: 2008 update from the American College of Critical Care Medicine and the Infectious Diseases Society of America.** *Crit Care Med* 2008, **36**:1330-1349.
13. Dellinger RP, Levy MM, Carlet JM, Bion J, Parker MM, Jaeschke R, Reinhart K, Angus DC, Brun-Buisson C, Beale R, Calandra T, Dhainaut JF, Gerlach H, Harvey M, Marini JJ, Marshall J, Ranieri M, Ramsay G, Sevransky J, Thompson BT, Townsend S, Vender JS, Zimmerman JL, Vincent JL: **Surviving Sepsis Campaign: international guidelines for management of severe sepsis and septic shock: 2008.** *Crit Care Med* 2008, **36**:296-327.
14. Launey Y, Nesselner N, Malldant Y, Seguin P: **Clinical review: fever in septic ICU patients: friend or foe?** *Crit Care* 2011, **15**:222.
15. Seguin P, Tanguy M, Laviolle B, Tirel O, Malldant Y: **Effect of oropharyngeal decontamination by povidone-iodine on ventilator-associated pneumonia in patients with head trauma.** *Crit Care Med* 2006, **34**:1514-1519.
16. Roquilly A, Mahe PJ, Seguin P, Guitton C, Floch H, Tellier AC, Merson L, Renard B, Malledant Y, Flet L, Sebille V, Volteau C, Masson D, Nguyen JM, Lejus C, Asehnoune K: **Hydrocortisone therapy for patients with multiple trauma: the randomized controlled HYPOLYTE study.** *JAMA* 2011, **305**:1201-1209.
17. Bratton SL, Chestnut RM, Ghajar J, McConnell Hammond FF, Harris OA, Hartl R, Manley GT, Nemecek A, Newell DW, Rosenthal G, Schouten J, Shutter L, Timmons SD, Ullman JS, Videtta W, Wilberger JE, Wright DW: **Guidelines for the management of severe traumatic brain injury. I. Blood pressure and oxygenation.** *J Neurotrauma* 2007, **24** Suppl 1:S7-S13.
18. Connolly ES, Rabinstein AA, Carhuapoma JR, Derdeyn CP, Dion J, Higashida RT, Hoh BL, Kirkness CJ, Naidech AM, Ogilvy CS, Patel AB, Thompson BG, Vespa P: **Guidelines for the management of aneurysmal subarachnoid hemorrhage: a guideline for healthcare professionals from the American Heart Association/American Stroke Association.** *Stroke* 2012, **43**:1711-1737.
19. Badjatia N, Boddock M, Guanci M, Rordorf GA: **Rapid infusion of cold saline (4 degrees C) as adjunctive treatment of fever in patients with brain injury.** *Neurology* 2006, **66**:1739-1741.
20. Aiyagari V, Diringner MN: **Fever control and its impact on outcomes: what is the evidence?** *J Neurol Sci* 2007, **261**:39-46.
21. Le Gall JR, Lemeshow S, Saulnier F: **A new Simplified Acute Physiology Score (SAPS II) based on a European/North American multicenter study.** *JAMA* 1993, **270**:2957-2963.
22. Bernard GR, Artigas A, Brigham KL, Carlet J, Falke K, Hudson L, Lamy M, Legall JR, Morris A, Spragg R: **The American-European Consensus Conference on ARDS. Definitions, mechanisms, relevant outcomes, and clinical trial coordination.** *Am J Respir Crit Care Med* 1994, **149**:818-824.
23. Chastre J, Fagon JY: **Ventilator-associated pneumonia.** *Am J Respir Crit Care Med* 2002, **165**:867-903.
24. American Thoracic Society, Infectious Diseases Society of America: **Guidelines for the management of adults with hospital-acquired, ventilator-associated, and healthcare-associated pneumonia.** *Am J Respir Crit Care Med* 2005, **171**:388-416.
25. Centers for Disease Control and Prevention: **Catheter-associated urinary tract infection (CAUTI) event.** In *CDC/NHSN Surveill Defin Specif Types Infect*, Volume 7. 2014:1-15.
26. Centers for Diseases Control and Prevention: **Central line-associated bloodstream infection (CLABSI) event.** In *CDC/NHSN Surveill Defin Specif Types Infect*, Volume 7. 2014. 4:1-19.
27. Fine J, Gray R: **A Proportional hazards model for the subdistribution of a competing risk.** *J Am Stat Assoc* 1999, **94**:496-509.
28. Royon V, Guitard PG, Abriou C, Frebourg N, Menard JF, Clavier T, Dureuil B, Veber B: **Hypothermia at admission increases the risk of pulmonary contusion's infection in intubated trauma patients.** *Ann Fr Anesthésie Réanimation* 2012, **31**:870-875.
29. Tejada Artigas A, Bello Dronza S, Chacn Valls E, Muñoz Marco J, Villuendas Usn MC, Figueras P, Suarez FJ, Hernandez A: **Risk factors for nosocomial pneumonia in critically ill trauma patients.** *Crit Care Med* 2001, **29**:304-309.
30. Bronchard R, Albaladejo P, Brezac G, Geoffroy A, Seince PF, Morris W, Branger C, Marty J: **Early onset pneumonia: risk factors and consequences in head trauma patients.** *Anesthesiology* 2004, **100**:234-239.
31. Sirvent JM, Torres A, Vidaur L, Armengol J, de Batlle J, Bonet A: **Tracheal colonisation within 24 h of intubation in patients with head trauma: risk factor for developing early-onset ventilator-associated pneumonia.** *Intensive Care Med* 2000, **26**:1369-1372.

32. Melsen WG, Rovers MM, Koeman M, Bonten MJM: **Estimating the attributable mortality of ventilator-associated pneumonia from randomized prevention studies.** *Crit Care Med* 2011, **39**:2736-2742.
33. Melsen WG, Rovers MM, Groenwold RH, Bergmans DC, Camus C, Bauer TT, Hanisch EW, Klarin B, Koeman M, Krueger WA, Lacherade JC, Lorente L, Memish ZA, Morrow LE, Nardi G, van Nieuwenhoven CA, O'Keefe GE, Nakos G, Scannapieco FA, Seguin P, Staudinger T, Topeli A, Ferrer M, Bonten MJ: **Attributable mortality of ventilator-associated pneumonia: a meta-analysis of individual patient data from randomised prevention studies.** *Lancet Infect Dis* 2013, **13**:665-671.
34. Ewig S, Torres A, El-Ebiary M, Fbregas N, Hernandez C, Gonzalez J, Nicols JM, Soto L: **Bacterial colonization patterns in mechanically ventilated patients with traumatic and medical head injury. Incidence, risk factors, and association with ventilator-associated pneumonia.** *Am J Respir Crit Care Med* 1999, **159**:188-198.
35. George DL, Falk PS, Wunderink RG, Leeper KV Jr, Meduri GU, Steere EL, Corbett CE, Mayhall CG: **Epidemiology of ventilator-acquired pneumonia based on protected bronchoscopic sampling.** *Am J Respir Crit Care Med* 1998, **158**:1839-1847.
36. Hubmayr RD, Burchardi H, Elliot M, Fessler H, Georgopoulos D, Jubran A, Limper A, Pesenti A, Rubenfeld G, Stewart T, Villar J, American Thoracic Society Assembly on Critical Care, European Respiratory Society, European Society of Intensive Care Medicine, Societe de Reanimation de Langue Francaise: **Statement of the 4th International Consensus Conference in Critical Care on ICU-Acquired Pneumonia**. Chicago, Illinois, May 20. *Intensive Care Med* 2002, **28**:1521-1536.
37. Peterson K, Carson S, Carney N: **Hypothermia treatment for traumatic brain injury: a systematic review and meta-analysis.** *J Neurotrauma* 2008, **25**:62-71.
38. Geurts M, Macleod MR, Kollmar R, Kremer PHC, van der Worp HB: **Therapeutic hypothermia and the risk of infection: a systematic review and meta-analysis.** *Crit Care Med* 2014, **42**:231-242.
39. Badjatia N, O'Donnell J, Baker JR, Huang D, Ayata C, Greer DM, Carter BS, Ogilvy CS, McDonald CT: **Achieving normothermia in patients with febrile subarachnoid hemorrhage: feasibility and safety of a novel intravascular cooling catheter.** *Neurocrit Care* 2004, **1**:145-156.
40. Diringner MN, Neurocritical Care Fever Reduction Trial Group: **Treatment of fever in the neurologic intensive care unit with a catheter-based heat exchange system.** *Crit Care Med* 2004, **32**:559-564.
41. Broessner G, Beer R, Lackner P, Helbok R, Fischer M, Pfausler B, Rhorer J, Kppers-Tiedt L, Schneider D, Schmutzhard E: **Prophylactic, endovascularly based, long-term normothermia in ICU patients with severe cerebrovascular disease: bicenter prospective, randomized trial.** *Stroke J Cereb Circ* 2009, **40**:e657-e665.
42. Puccio AM, Fischer MR, Jankowitz BT, Yonas H, Darby JM, Okonkwo DO: **Induced normothermia attenuates intracranial hypertension and reduces fever burden after severe traumatic brain injury.** *Neurocrit Care* 2009, **11**:82-87.
43. Sahuquillo J, Vilalta A: **Cooling the injured brain: how does moderate hypothermia influence the pathophysiology of traumatic brain injury.** *Curr Pharm Des* 2007, **13**:2310-2322.

doi:10.1186/s13054-014-0689-4

Cite this article as: Launey *et al.*: Effect of a fever control protocol-based strategy on ventilator-associated pneumonia in severely brain-injured patients. *Critical Care* 2014 **18**:689.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

