

RESEARCH ARTICLE

Open Access

Cytotoxicity of *Elaeophorbium drupifera* and other Cameroonian medicinal plants against drug sensitive and multidrug resistant cancer cells

Victor Kuete^{1,2}, Igor K Voukeng², Roger Tsobou³, Armelle T Mbaveng², Benjamin Wiench¹, Veronique P Beng⁴ and Thomas Efferth^{1*}

Abstract

Background: Multidrug resistance (MDR) is a major hurdle for cancer treatment worldwide and accounts for chemotherapy failure in over 90% of patients with metastatic cancer. Evidence of the cytotoxicity of Cameroonian plants against cancer cell lines including MDR phenotypes is being intensively and progressively provided. The present work was therefore designed to evaluate the cytotoxicity of the methanol extracts of twenty-two Cameroonian medicinal plants against sensitive and MDR cancer cell lines.

Methods: The methanol maceration was used to obtain the crude plant extracts whilst the cytotoxicity of the studied extracts was determined using a resazurin reduction assay.

Results: A preliminary assay on leukemia CCRF-CEM cells at 40 µg/mL shows that six of the twenty plant extracts were able to enhance less than 50% of the growth proliferation of CCRF-CEM cells. These include *Crinum zeylanicum* (32.22%), *Entada abyssinica* (34.67%), *Elaeophorbium drupifera* (35.05%), *Dioscorea bulbifera* (45.88%), *Eremomastax speciosa* (46.07%) and *Polistigma thonningii* (45.11%). Among these six plants, *E. drupifera* showed the best activity with IC₅₀ values below or around 30 µg/mL against the nine tested cancer cell lines. The lowest IC₅₀ value of 8.40 µg/mL was recorded with the extract of *E. drupifera* against MDA-MB231 breast cancer cell line. The IC₅₀ values below 10 µg/mL were recorded with the extracts of *E. drupifera* against MDA-MB231 breast cancer cells, *C. zeylanicum* against HCT116 p53^{+/+} and HCT116p53^{-/-} colon cancer cells and *E. abyssinica* against HCT116 p53^{+/+} cells.

Conclusion: The results of the present study provide evidence of the cytotoxic potential of some Cameroonian medicinal plants and a baseline information for the potential use of *Elaeophorbium drupifera* in the treatment of sensitive and drug-resistant cancer cell lines.

Keywords: Cameroon, Cytotoxicity, *Elaeophorbium drupifera*, Medicinal plants

Background

The escape of cancer cells from chemotherapy by multidrug resistance (MDR) mechanisms is until now a major reason for systemic cancer treatment failure. So far, limited progress has been made in the fight against MDR cancer, and even the use of combination chemotherapy cannot solve the problem [1,2]. Medicinal plants and alternative medicine are undeniable sources of new exploitable active principles to manage infectious and degenerative

diseases. The structural diversity of chemicals from the medicinal plants makes them valuable tools in the search for potentially active drugs on sensitive and resistant phenotypes. It is estimated that more than 60% of the approved anticancer drugs in the United States of America (from 1983 to 1994) were from natural origin [3,4]. In Cameroon, medicinal plants are traditionally used to manage infectious diseases and different types of cancers [5]. Evidence of the cytotoxicity of these plants against cancer cell lines has been provided [6-11]. In a recent research program, we started to investigate the cytotoxicity of Cameroonian plants against drug-resistant cancer cell lines. The idea is to identify plants able to kill drug-resistant

* Correspondence: efferth@uni-mainz.de

¹Department of Pharmaceutical Biology, Institute of Pharmacy and Biochemistry, University of Mainz, Staudinger Weg 5, Mainz 55128, Germany
Full list of author information is available at the end of the article

cancer cells with similar efficacy as their drug-sensitive counterparts. Some of the plants identified so far include *Echinops giganteus*, *Imperata cylindrica*, *Piper capense* and *Xylopia aethiopica* which displayed considerable activities against the P-glycoprotein-expressing adriamycin-resistant cell line, CEM/ADR5000 [7,12]. This encourage us to move forward to search for new cytotoxic agents from Cameroonian medicinal plants, with emphasis on MDR phenotypes with different mechanism of action. The present work was therefore designed to evaluate the cytotoxicity of twenty-two Cameroonian plants against both sensitive and drug-resistant cancer cell lines.

Methods

Plant material

All medicinal plants used in the present work were collected at various locations of Dschang, West-Region of Cameroon, between January and April 2012. The plants were identified at the National Herbarium (Yaounde, Cameroon), where voucher specimens were deposited under the reference numbers indicated in Table 1. The air-dried and powdered plant material was soaked in methanol for 48 h, at room temperature. The methanol extract was concentrated under reduced pressure to give the crude extract. This extract was then conserved at 4°C until further use.

Extraction

The air-dried and powdered plant samples (1 kg) were soaked in methanol (3 L) for 48 h, at room temperature. The methanol extract was concentrated under *vacuum* to give the crude extract. This extract was then conserved at 4°C until use.

Chemicals

Doxorubicin, vinblastine and daunorubicin were provided by the University Medical Center of the Johannes Gutenberg University (Mainz, Germany) and dissolved in PBS (Invitrogen, Eggenstein, Germany) at a concentration of 10 mM. Geneticin was purchased from Sigma-Aldrich (Munich, Germany) at a concentration of 50 mg/mL in sterile-filtered H₂O.

Cell cultures

Drug-sensitive CCRF-CEM and multidrug-resistant CEM/ADR5000 leukemia cells were maintained in RPMI 1640 medium (Invitrogen) supplemented with 10% fetal calf serum in a humidified 5% CO₂ atmosphere at 37°C. Sensitive and resistant cells were kindly provided by Dr. Axel Sauerbrey (Department of Pediatrics, University of Jena, Jena, Germany). The generation of the resistant subline was previously described [52]. The specific overexpression of P-glycoprotein, but not other ABC transporters has been reported [53,54]. Breast cancer

cells, transduced with control vector (MDA-MB-231-pcDNA3) or with cDNA for the breast cancer resistance protein *BCRP* (MDA-MB-231-*BCRP* clone 23), were maintained under standard conditions as described above for CCRF-CEM cells. Human wild-type HCT116 (*p53*^{+/+}) colon cancer cells as well as knockout clones HCT116 (*p53*^{-/-}) derived by homologous recombination were a generous gift from Dr. B. Vogelstein and H. Hermeking (Howard Hughes Medical Institute, Baltimore, MD). Human glioblastoma multiforme U87MG cells (non-transduced) and U87MG cell line transduced with an expression vector harboring an epidermal growth factor receptor (*EGFR*) gene with a genomic deletion of exons 2 through 7 (U87MG.Δ*EGFR*) were kindly provided by Dr. W. K. Cavenee (Ludwig Institute for Cancer Research, San Diego, CA) [55]. MDA-MB-231-*BCRP*, U87MG.Δ*EGFR* and HCT116 (*p53*^{-/-}) were maintained in DMEM medium containing 10% FBS (Invitrogen) and 1% penicillin (100 U/mL)-streptomycin (100 µg/mL) (Invitrogen) and were continuously treated with 800 ng/mL and 400 µg/mL geneticin, respectively. Human HepG2 hepatocellular carcinoma cells and normal AML12 hepatocytes were obtained from the American Type Culture Collection (ATCC, USA). The above medium without geneticin was used to maintain MDA-MB-231, U87MG, HCT116 (*p53*^{+/+}), HepG2 and AML12 cell lines. The cells were passaged twice weekly. All experiments were performed with cells in the logarithmic growth phase.

Resazurin reduction assay

Resazurin reduction assay [56] was performed to assess the cytotoxicity of the studied samples toward various sensitive and resistant cancer cell lines. The assay is based on the reduction of resazurin, to the highly fluorescent resorufin by viable cells. Non-viable cells rapidly lose the metabolic capacity to reduce resazurin and thus produce no fluorescent signal. Briefly, adherent cells were detached by treatment with 0.25% trypsin/EDTA (Invitrogen, Darmstadt, Germany) and an aliquot of 1×10^4 cells was placed in each well of a 96-well cell culture plate (Thermo Scientific, Langenselbold, Germany) in a total volume of 200 µL. Cells were allowed to attach overnight and then treated with different concentrations of the studied sample. For suspension cells, aliquots of 2×10^4 cells per well were seeded in 96-well-plates in a total volume of 100 µL. The studied sample was immediately added in varying concentrations in an additional 100 µL of culture medium to obtain a total volume of 200 µL/well. After 24 h or 48 h, 20 µL resazurin (Sigma-Aldrich, Schnellendorf, Germany) 0.01% w/v in double-distilled water (ddH₂O) were added to each well and the plates incubated at 37°C for 4 h. Fluorescence was measured on an Infinite M2000 Pro™ plate reader (Tecan, Crailsheim, Germany) using an

Table 1 Pharmacognosy of twenty-two studied Cameroonian medicinal plants

Plant species, family/(Voucher specimen) ^a	Traditional use	Part used traditionally	Part used in this study (extraction yield in %) ^b	Potential active constituents	Previously screened activity
<i>Ageratum conyzoides</i> Linn. (Asteraceae)/(19050/SFR-Cam)	Purgative, fever, ulcers and wound, mental and infectious diseases, headaches, crawl-crawl, diarrhea [13].	Leaves, whole plant [13]	Leaves (8.52%)	β -caryophyllene, precocene I, friedelin, Lycopsamine, echinatine, β -sitosterol, stigmasterol, 5-methoxynobiletin, linderoflavone B, eupalestin, sabinene, α and β pinene, 1.6%, β -phellandrene, 1,8-cineole and limonene, ocimene, eugenol [13]	Antimicrobial, anticonvulsant, analgesic, anti-inflammatory, antipyretic, insecticidal [13]
<i>Albizia gummifera</i> (Mimosaceae)/(41196/HNC)	Bacterial infections, skin diseases, malaria and stomach pain [14]	Bark	Bark (11.51%)	Vitalboside A, vitalboside-A-2-methylglucuronate, lupeol, lupenone [14]	Antiplasmodial [15]
<i>Aloe barbadensis</i> Mill. (Liliaceae), ICNA	Abrasions and burns, emollient and moisturizer [16]	Leaves [16]	Leaves (8.15%)	Aloin A, B, aloesin, aloesin A, aloe-emodin, rhein, aloe-emodin-9-anthrone [16]	Antidiabetic, antiviral, angiogenic, toxicity, immunomodulator [16]
<i>Cissus quadrangularis</i> Linn. (Vitaceae)/(18668/SRF-Cam)	Fracture healing, eye diseases, chronic ulcer, tumors, asthma [17]	Stem, pulps [17]	Stem (6.83%)	Alpha and β -amyrin, β -sitosterol, ketosteroid, phenols, tannins, carotene [17]	Antiosteoporotic, analgesic, hypotensive, antibacterial, antifungal [18]
<i>Crinum zeylanicum</i> Linn. (Amaryllidaceae)/(18263/SRF-Cam)	Rheumatism, earache, malaria, poison [19]	Bulbs [19]	Whole plant (6.85%)	Flexinine, 6-hydroxypowelline, zeylamine, hamayne, 3-acetylhamayne, crinamine, 6-hydroxycrinamine, 6-methoxycrinamine, crinine, ambelline, 6-hydroxybuphandrine, 6-ethoxybuphandrine, 6-ethoxybuphanidrine, lycorine, 11-O-acetoxymbelline, galanthamine, sanguinine, 3-O-acetylsanguinine [19]	Antiproliferative compounds [19]
<i>Croton macrostachys</i> Hochst. (Euphorbiaceae)/(40501/HNC)	Antidiabetic [20]	Roots, bark [20]	Bark (12.72%)	Taraxer-14-en-28-oic acid, trachyloban-19-oic acid, trachyloban-18-oic acid, neoclerodan-5,10-en-19,6 β ;20,12-diolide, 3 α , 19-dihydroxytrachylobane, 3 α ,18,19-trihydroxytrachylobane [20]	Not reported
<i>Dioscorea bulbifera</i> L. (Dioscoreaceae)/(14274/HNC)	Sore throat and struma, leprosy and tumors, diabetes, microbial infections [21,22]	Rhizome [21]	Rhizome (15.8%)	Kaempferol-3,5-dimethyl ether, caryatin, (+)-catechin, myricetin, quercetin-3-O-galactopyranoside, myricetin-3-O-galactopyranoside, myricetin-3-O-glucopyranoside, diosbulbin B [21]	Analgesic, anti-inflammatory [23], antimicrobial [22]

Table 1 Pharmacognosy of twenty-two studied Cameroonian medicinal plants (Continued)

<i>Dioscorea dumetorum</i> (Kunth) Pax Trusted (Dioscoreaceae)/ (24431/ SRF-Cam)	Diabetes, topical anesthetic, poison [24]	Leaves	Leaves (6.95%)	Dumetorine, dihydrodioscorine, demethylbatatasin IV, dihydroresveratrol [24,25]	Not reported
<i>Dissotis perkinsiae</i> Gilg. (Melastomataceae)/ (6991/ SRF-Cam)	Typhoid fever (Personal information)	Leaves, stem	Stem with leaves (10.35%)	Not reported	Not reported
<i>Elaeophorbia drupifera</i> (Thonn.) Stapf. (Euphorbiaceae)/ (57644/HNC)	Hypertension, diabetes [26]	Leaves [26]	Leaves	Euphol, tirucallo, euphorbol, ingenol elaeophorbate, epitaraxerol, taraxerone, friedelin, lup-20(29)-en-3-one or lupenone, lupeol, olean-12-ene-3-one, olean-12-ene-3-ol, elaeophorbate [27,28]	Leaves extract moderately inhibit HIV-1 and HIV-2 proviral DNA copying [29], relaxant effect on vascular smooth muscles on rats [30]
<i>Entada abyssinica</i> Steud. ex A. Rich. (Mimosaceae)/ (26967/SRF-Cam)	Bronchitis, coughs, arthritic pain, miscarriage, fever, abdominal pain [31]	Bark, Juice [31]	Bark (13.95%)	Not reported	Not reported
<i>Eremomastax speciosa</i> (Hochst) Cufod (Acanthaceae)/ (16371/SRF-Cam)	Dysentery, anemia, irregular menstruation, hemorrhoids, urinary tract infection [32]	Stem, leaves [32]	Stem with leaves (8.15%)	Not reported	Anti-diarrhoeal, anti-ulcerogenic [32]
<i>Gossypium barbadense</i> L. (Malvaceae)/(25771/HNC)	Cold, bronchitis, palpitations, wounds, systematic diarrhea [33]	leaves, young shoots [33]	Leaves (8.15%)	Gossypol, hemigossypol, 6-methoxyhemigossypol, 6-deoxyhemigossypol, 6-methoxygossypol, 6,6'-dimethoxygossypol [34,35]	Antimicrobial [33]
<i>Kigelia Africana</i> (Lam.) Benth (Bignoniaceae)/ (23220/SRF-Cam)	Dysentery, ringworm, tape-worm, malaria, diabetes, post-partum haemorrhage, pneumonia, toothache, syphilis, gonorrhoea [36]	Roots, fruits, leaves, bark [36]	Bark (15.85%)	Pinnatal, norvibutinal, β -sitosterol, 7-hydroxy viteoid II, 7-hydroxy eucommic acid, jiofuran, jioglulotide, kigelin, specioside, verminoside, stigmasterol, lapachol [36]	Antibacterial and antifungal, cytotoxic, analgesic, anti-inflammatory, antimalarial, antiprotozoal, central nervous system stimulant, antidiabetic [36]
(Bignoniaceae)/ <i>Markhamia tomentosa</i> K.Schum. (1974/SRF-Cam)	Anti-snake venom, sore eyes, heart pain, scrotal elephantiasis [37]	Leaves	Bark (5.31%)	2-acetylnaphtho[2,3-b]furan-4,9-dione, 2-acetyl-6-methoxynaphtho[2,3-b]furan-4,9-dione, oleanolic acid, pomolic acid, 3-acetylpomolic acid, tormentic acid, β -sitosterol, β -sitosterol-3-O- β -D-glucopyranoside [38]	Antimicrobial, antiprotozoal [37,38]
Stem and leaves	Malaria [39]	Leaves, roots, stems [39]	Stem with leaves (8.13%)	Paullinoside A, paullinonides A and B, β -amyrin, 13 β , 17 β -dihydroxy-28-norolean-12-ene, β -sitosterol, β -sitosterol glucopyranoside [40]	Antiparasitic [39]
<i>Polio stigma thonningii</i> (Schum.) Milne-Readhead (Caesalpiniaceae)/ (32129/HNC)	Leprosy, smallpox, coughs, wounds, ulcers [41]	bark, root, leaves [41]	Bark (13.95%)	Pliostigmin, quercetin, quercitrin, 6-C-methylquercetin 3-methyl ether, 6-C-methylquercetin 3,7,3'-trimethyl ether, 6,8-di-C-methylkaempferol 3-methyl ether,	Antihelmintic, antitussive, bronchodilator, antibacterial [41]

Table 1 Pharmacognosy of twenty-two studied Cameroonian medicinal plants (Continued)

<i>Pseudarthria confertiflora</i> (A. Rich.) Bak. Fabaceae/ (17465/SRF-Cam)	Typhoid fever (Personal information)	Leaves	Leaves (10.75%)	6,8-di-C-methylkaempferol 3,7-dimethyl ether [42] Not reported	Not reported
<i>Spathodea campanulata</i> P. Beauv. (Bignoniaceae)/ (29470/SRF-Cam)	Kidney diseases, urethra inflammation, poison, enemas, fungus skin diseases, herpes, stomachache, diarrhea [43]	Flowers, leaves, bark [43]	Bark (15.81%)	Ajugol, <i>p</i> -hydroxy- benzoic acid, methyl <i>p</i> -hydroxy- benzoate [43]	Molluscicidal, hypoglycemic, anti-HIV, anti- malarial [43]
<i>Spilanthes filicaulis</i> (Schum et Thonn.) C.D. (Asteraceae)/ (42040/HNC)	Toothache, stomach ache, gastritis [44]	Whole plant	Whole plant (5.62%)	Not reported	Analgesic, antimicrobial [45], antiulcerogenic [46]
<i>Stereospermum acuminatissimum</i> K. Schum. (Bignoniaceae)/ (45705/HNC)	Haemostatic, cicatrizing [47]	Leaves, barks [48]	Bark (13.15%)	1,3,7-trimethylguanin-1/3-ium, 3,7-dimethylguanin-1/3-ium, 2-(4-hydroxyphenyl)ethyl hentriacontanoate, sterequinones A, B, C, E, F, H, zenkequinones A, B, <i>p</i> -coumaric acid, methyl caffeate, caffeic acid, psilalic acid, syringaldehyde, norviburtinal, specioside, verminoside, tyrosol, eutigoside A, ellagic acid, atranorin, ursolic acid, pomolic acid, quinovic acid, oleanolic acid, triacontan-1,30-dioldiferulate, 2-(4-hydroxyphenyl)ethyl dotriacontanoate [47,48]	Antiprotozoal [49]
<i>Terminalia glaucescens</i> Planch. (Combretaceae)/ (9468/SRFCAM)	Dysentery, AIDS [50]	Bark	Bark (13.84%)	glaucinoic acid, arjunic acid, arjungenin, sericoside, friedelin [50]	Antiplasmodial, cytotoxic, antimicrobial, aldose reductase inhibition [50,51]


^aPlants were identified at the Cameroon National Herbarium (HNC); ICNA: Voucher with no identification code at the HNC; ^bThe percentage of the methanol extract.

excitation wavelength of 544 nm and an emission wavelength of 590 nm. Each assay was done at least two times, with six replicate each. The viability was evaluated based on a comparison with untreated cells. IC₅₀ values represent the sample's concentrations required to inhibit 50% of cell proliferation and were calculated from a calibration curve by linear regression using Microsoft Excel.

Results and discussion

In a prescreening of twenty-two plants, we tested a single concentration of 40 µg/mL for each sample against the sensitive CCRF-CEM leukemia cell line. The results depicted in Figure 1 indicate that six of the twenty-two plant extracts were able to display less than 50% growth

proliferation of CCRF-CEM cells. These include *Crinum zeylanicum* (32.22%), *Entada abyssinica* (34.67%), *Elaeophorbium drupifera* (35.05%), *Dioscorea bulbifera* (45.88%), *Eremomastax speciosa* (46.07%) and *Polistigma thonningii* (45.11%). The IC₅₀ values of these samples were then determined on a panel of cancer cell lines, including both sensitive and MDR phenotypes. The results are shown in Table 2. Only the *Elaeophorbium drupifera* extract as well as the control drug doxorubicin inhibited the proliferation of the nine studied cancer cell lines, with IC₅₀ values below 40 µg/mL. Other extracts showed selective activities, the IC₅₀ values being obtained on 6/10 tested cells lines for *Crinum zeylanicum*, 4/10 for *Dioscorea bulbifera* and *Entada abyssinica*, 3/10 for *Eremomastax*



speciosa and *Polistigma thonningii* (Table 2). According to the criteria of the American National Cancer Institute, 30 µg/mL is the upper IC₅₀ limit considered promising for purification of a crude extract [57]. Consequently, the highest concentration tested (40 µg/mL) in our screening was slightly above this limit. Considering this cutoff point, the IC₅₀ values below or around 30 µg/mL were recorded with only the *E. drupifera* extract against the nine tested cancer cell lines (Table 2). However, other extract also

displayed activities with IC₅₀ values below 30 µg/mL on at least one of the cancer cell line tested.

MDR is a major hurdle for cancer treatment worldwide and accounts for chemotherapy failure in over 90% of patients with metastatic cancer [1,58]. In the present work, we investigated both sensitive and MDR cell lines. The degrees of resistance were calculated by dividing the IC₅₀ value of the resistant cell line by the corresponding parental sensitive cell line. We tested cell models overexpressing

Table 2 Cytotoxicity of the studied extracts towards sensitive and drug-resistant cancer cell lines and normal cells as determined by the resazurin assay

Cell lines	Samples, IC ₅₀ values (µg/mL) and degree of resistance ^a (in brackets)						
	<i>Crinum zeylanicum</i>	<i>Dioscorea bulbifera</i>	<i>Elaeophoria drupifera</i>	<i>Entada abyssinica</i>	<i>Eremomastax speciosa</i>	<i>Polistigma thonningii</i>	Doxorubicin
CCRF-CEM	17.22 ± 2.19	19.77 ± 2.22	11.86 ± 1.19	15.81 ± 1.47	23.65 ± 2.51	26.44 ± 1.18	0.11 ± 0.01
CEM/ADR5000	23.67 ± 1.97 (1.38)	- (>2.02)	13.72 ± 0.97(1.12)	- (>2.53)	38.71 ± 4.12 (1.64)	- (>1.51)	195.12 ± 14.30 (1772)
MDA-MB231	18.01 ± 1.61	33.17 ± 2.91	8.40 ± 0.55	29.14 ± 1.83	35.13 ± 2.49	34.19 ± 2.24	1.10 ± 0.01
MDA-MB231/ <i>BCRP</i>	11.18 ± 1.11 (0.62)	- (>1.21)	30.96 ± 2.74(3.69)	- (>1.37)	- (>1.14)	- (>1.17)	7.83 ± 0.01 (7.11)
HCT116 (<i>p53</i> ^{+/+})	4.32 ± 0.52	36.14 ± 2.37	25.36 ± 2.13	9.55 ± 1.11	-	-	1.43 ± 0.02
HCT116 (<i>p53</i> ^{-/-})	7.45 ± 0.64 (1.73)	- (<0.69)	28.61 ± 3.08(1.13)	14.38 ± 1.25 (1.51)	-	-	4.06 ± 0.04 (2.84)
U87MG	-	-	23.58 ± 2.20	-	-	34.22 ± 2.74	1.06 ± 0.03
U87MG.Δ <i>EGFR</i>	-	27.76 ± 1.86 (<0.69)	16.03 ± 0.88(0.68)	-	-	- (>1.17)	6.11 ± 0.04 (5.76)
Hep-G2	-	-	23.23 ± 1.67 (1.72)	-	-	-	1.41 ± 0.12 (<0.04)
AML12	-	-	-	-	-	-	-

^aThe degree of resistance (in brackets) was determined as the ratio of IC₅₀ value of the resistant/IC₅₀ sensitive cell line; (-): >40 µg/mL.

two ATP-binding cassette transporters, *i.e.* P-glycoprotein (ABCB1/MDR1) or breast cancer resistance protein (ABCG2/BCRP). Furthermore, we tested a p53 knockout cell line and a transfectant cell line harboring a mutation-activated *EGFR* gene ($\Delta EGFR$) as examples for resistance-inducing tumor suppressors and oncogenes. Finally, we investigated HepG2 liver cancer cells and AML12 normal hepatocytes to compare carcinoma cells with normal cells. The degree of resistance on the tested cell line toward the control drug doxorubicin was generally high, showing that the studied cell lines can obviously be considered as suitable cell models to study drug resistance. For the most active extract *E. drupifera*, it can be observed that the degrees of resistance were in all cases lower than those of doxorubicin, suggesting that this sample can be exploited in a possible fight against cancer diseases involving MDR phenotypes. In addition, collateral sensitivity (sample more active on resistant cells than on sensitive cells) was observed with the extract of *E. drupifera* against U87MG. $\Delta EGFR$, highlighting its good antiproliferative activity.

To the best of our knowledge, the cytotoxicity of the six most active extracts (*C. zeylanicum*, *D. bulbifera*, *E. drupifera*, *E. abyssinica*, *E. speciosa* and *P. thoningii*) is being reported for the first time. Nevertheless, compounds with activities against malignant cells such as crinine, 6-hydroxybuphanidrine and 6-ethoxybuphanidrine were isolated from *C. zeylanicum* [19]. Also, lupeol [27,28] a moderately active cytotoxic compound [59] was identified in *E. drupifera*, the plant that displayed the best activity as observed in this study. The presence of such compounds could probably explain their antiproliferative activity.

Conclusion

In conclusion, the results of the present study provide evidence of the cytotoxic potential of some Cameroonian medicinal plants and highlight the good activity of *Elaeophorbia drupifera* on sensitive and drug-resistant cancer cell lines. This plant is a potential cytotoxic source, that could be explored in more details in the future to develop novel anticancer drugs against sensitive and resistant phenotypes.

Competing interest

The authors declare that they have no competing interests.

Authors' contributions

VK, IKV, ATM, RT, BW, and VPB carried out the study; VK and TE designed the experiments. VK wrote the manuscript; VK and TE supervised the work. All authors read and approved the final manuscript.

Acknowledgments

VK is very grateful to the Alexander von Humboldt foundation for an 18 months' fellowship in Germany through the "Georg Foster Research Fellowship for Experienced Researcher" program.

Author details

¹Department of Pharmaceutical Biology, Institute of Pharmacy and Biochemistry, University of Mainz, Staudinger Weg 5, Mainz 55128, Germany. ²Department of Biochemistry, Faculty of Science, University of Dschang, Dschang, Cameroon. ³Department of Plant Biology, Faculty of Science, University of Dschang, Dschang, Cameroon. ⁴Department of Organic Biochemistry, Faculty of Science, University of Yaoundé I, Yaoundé, Cameroon.

Received: 24 June 2013 Accepted: 30 September 2013

Published: 2 October 2013

References

1. Liu J, Zhao Y, Guo Q, Wang Z, Wang H, Yang Y, Huang Y: TAT-modified nanosilver for combating multidrug-resistant cancer. *Biomaterials* 2012, **33**(26):6155–6161.
2. Broxterman H, Gotink K, Verheul H: Understanding the causes of multidrug resistance in cancer: a comparison of doxorubicin and sunitinib. *Drug Resist Update* 2009, **12**:114–126.
3. Newman DJ, Cragg GM: Natural products as sources of new drugs over the last 25 years. *J Nat Prod* 2007, **70**(3):461–477.
4. Stevigny C, Bailly C, Quetin-Leclercq J: Cytotoxic and antitumor potentialities of aporphinoid alkaloids. *Curr Med Chem Anticancer Agents* 2005, **5**(2):173–182.
5. Adjanohoun J, Aboubakar N, Dramane K, Ebot M, Ekpre J, Enow-Orock E, Focho D, Gbile Z, Kamanyi A, Kamsu-Kom J: *Traditional medicine and pharmacopoeia: contribution to ethnobotanical and floristic studies in Cameroon*. Lagos-Nigeria: Technical and Research Commission of Organisation of African Unity (OAU/STRC); 1996.
6. Kuete V, Mbaveng AT, Tsaffack M, Beng VP, Etoa FX, Nkengfack AE, Meyer JJM, Lall N: Antitumor, antioxidant and antimicrobial activities of *Bersama engleriana* (Melianthaceae). *J Ethnopharmacol* 2007, **115**(3):494–501.
7. Kuete V, Krusche B, Youns M, Voukeng I, Fankam AG, Tankeo S, Lacmata S, Effertth T: Cytotoxicity of some Cameroonian spices and selected medicinal plant extracts. *J Ethnopharmacol* 2011, **134**(3):803–812.
8. Dzoym J, Nkuete AH, Kuete V, Tala MF, Wabo HK, Guru SK, Rajput VS, Sharma A, Tane P, Khan IA, et al: Cytotoxicity and antimicrobial activity of the methanol extract and compounds from *Polygonum limbatum*. *Planta Med* 2012, **78**(8):787–792.
9. Choumessi AT, Danel M, Chassaing S, Truchet I, Penlap VB, Pieme AC, Asonganyi T, Ducommun B, Valette A: Characterization of the antiproliferative activity of *Xylopi aethiopica*. *Cell Div* 2012, **7**(1):8.
10. Dzoym J, Guru S, Pieme C, Kuete V, Sharma A, Khan I, Saxena A, Vishwakarma R: Cytotoxic and antimicrobial activity of selected Cameroonian edible plants. *BMC Complement Altern Med* 2013, **13**(1):78.
11. Tamokou Jde D, Chouna J, Fischer-Fodor E, Chereches G, Barbos O, Damian G, Benedec D, Duma M, Efouet A, Wabo H, et al: Anticancer and antimicrobial activities of some antioxidant-rich cameroonian medicinal plants. *PLoS One* 2013, **8**(2):e55880.
12. Kuete V, Sandjo L, Wiench B, Effertth T: Cytotoxicity and modes of action of four Cameroonian dietary spices ethno-medically used to treat Cancers: *Echinops giganteus*, *Xylopi aethiopica*, *Imperata cylindrica* and *Piper capense*. *J Ethnopharmacol* 2013. doi: 10.1016/j.jep.2013.06.029.
13. Okunade AL: *Ageratum conyzoides* L. (Asteraceae). *Fitoterapia* 2002, **73**(1):1–16.
14. Rukunga GM, Waterman PG: A new oleanane glycoside from the stem bark of *Albizia gummifera*. *Fitoterapia* 2001, **72**(2):140–145.
15. Rukunga GM, Muregi FW, Tolo FM, Omar SA, Mwitari P, Muthaura CN, Omlin F, Lwande W, Hassanali A, Githure J, Iraqi FW, Mungai GM, Kraus W, Kofi-Tsekpo: The antiplasmodial activity of spermine alkaloids isolated from *Albizia gummifera*. *Fitoterapia* 2007, **78**(7–8):455–459.
16. Boudreau MD, Beland FA: An evaluation of the biological and toxicological properties of *Aloe barbadensis* (miller), *Aloe vera*. *J Environ Sci Health C Environ Carcinog Ecotoxicol Rev* 2006, **24**(1):103–154.
17. Jainu M, Devi S: *In vitro* and *in vivo* evaluation of free radical scavenging potential of *Cissus quadrangularis*. *Afr J Biomed Res* 2005, **8**:95–99.
18. Jainu M, Devi CS: Gastroprotective action of *Cissus quadrangularis* extract against NSAID induced gastric ulcer: role of proinflammatory cytokines and oxidative damage. *Chem Biol Interact* 2006, **161**(3):262–270.

19. Berkov S, Romani S, Herrera M, Viladomat F, Codina C, Momekov G, Ionkova I, Bastida J: **Antiproliferative alkaloids from *Crinum zeylanicum***. *Phytother Res* 2011, **25**(11):1686–1692.
20. Kapingu C, Guillaume D, Mbawambo H, Moshi J, Uliso C, Mahunnah A: **Diterpenoids from the roots of *Croton macrostachys***. *Phytochemistry* 2000, **54**:767–770.
21. Gao H, Kuroyanagi M, Wu L, Kawahara N, Yasuno T, Nakamura Y: **Antitumor-promoting constituents from *Dioscorea bulbifera* L. in JB6 mouse epidermal cells**. *Biol Pharm Bull* 2002, **25**(9):1241–1243.
22. Kuete V, Bertrandponno R, Mbaveng AT, Tapondjou LA, Meyer JJ, Barboni L, Lall N: **Antibacterial activities of the extracts, fractions and compounds from *Dioscorea bulbifera***. *BMC Complement Altern Med* 2012, **12**:228.
23. Mbiantcha M, Kamanyi A, Teponno RB, Tapondjou AL, Watcho P, Nguielefack TB: **Analgesic and anti-inflammatory properties of extracts from the bulbs of *Dioscorea bulbifera* L. var sativa (Dioscoreaceae) in mice and rats**. *Evid Based Complement Alternat Med* 2011, **2011**:912935.
24. Corley G, Tempesta S: **Convulsant alkaloids from *Dioscorea dumetorum***. *Tetrahedron Lett* 1985, **26**(13):1615–1618.
25. Adesanya S, Ogundana S, Roberts M: **Dihydrostilbene phytoalexins from *Dioscorea bulbifera* and *D. dumetorum***. *Phytochemistry* 1989, **28**(3):773–774.
26. Eno A, Azah N: **Effect of ethanolic extract from *Elaeophorbium drupifera* leaves on the gastrointestinal smooth muscle of the rabbit**. *Niger J Physiol Sci* 2004, **19**(1–2):60–68.
27. Kinghorn A, Evans F: **Occurrence of ingenol in *Elaeophorbium* species**. *Planta Med* 1974, **26**:150–154.
28. Ahiahonu PW, Goodenowe DB: **Triterpenoids from leaves of *Elaeophorbium drupifera***. *Fitoterapia* 2007, **78**(5):337–341.
29. Ayisi N, Nyadedzor C: **Comparative *in vitro* effects of AZT and extract of *Ocimum gratissimum*, *Ficus polita*, *Clauseria anisata*, *Alchornea cordifolia*, and *Elaeophorbium drupifera* against HIV-1 and HIV-2 infections**. *Antivir Res* 2003, **58**:25–33.
30. Eno AE, Owo OI: **Cardiovascular effects of an extract from the roots of a shrub *Elaeophorbium drupifera***. *Phytother Res* 1999, **13**(7):549–554.
31. Olajide OA, Alada AR: **Studies on the anti-inflammatory properties of *Entada abyssinica***. *Fitoterapia* 2001, **72**(5):492–496.
32. Oben J, Assi S, Agbor G, Musoro D: **Effect of *Eremomastax speciosa* on experimental diarrhoea**. *Afr J Trad Complement Altern Med* 2006, **3**(1):95–100.
33. Essien E, Aboaba O, Ogunwande A: **Constituents and antimicrobial properties of the leaf essential oil of *Gossypium barbadense* (Linn.)**. *J Med Plant Res* 2011, **5**:702–705.
34. Stipanovic D, Bell A, Mace E, Howell R: **Antimicrobial terpenoids of *Gossypium*: 6-methoxygossypol and 6,6'-dimethoxygossypol**. *Phytochemistry* 1975, **14**:1077–1081.
35. Bell A, Stipanovic D, Howell R, Fryxell A: **Antimicrobial terpenoids of *Gossypium*: Hemigossypol, 6-methoxyhemigossypol and 6-deoxyhemigossypol**. *Phytochemistry* 1975, **14**:225–231.
36. Saini S, Kaur H, Verma B, Ripudaman, Singh K: ***Kigelia africana* (Lam.) Benth: - an overview**. *Nat Prod Rad* 2009, **8**(2):190–197.
37. Aladesanmi A, Iwalewa E, Adebajo A, Akinkunmi E, Taiwo B, Olorunmola F, Lamikanra A: **Antimicrobial and antioxidant activities of some Nigerian medicinal plants**. *Afr J Trad Complement Altern Med* 2007, **4**(2):173–184.
38. Tantangmo F, Lenta BN, Boyom FF, Ngouela S, Kaiser M, Tsamo E, Weniger B, Rosenthal PJ, Vonthron-Sénécheau C: **Antiprotozoal activities of some constituents of *Markhamia tomentosa* (Bignoniaceae)**. *Ann Trop Med Parasitol* 2010, **104**(5):391–398.
39. Okpekon T, Yolou S, Gleye C, Roblot F, Loiseau P, Bories C, Grellier P, Frappier F, Laurens A, Hocquemiller R: **Antiparasitic activities of medicinal plants used in Ivory Coast**. *J Ethnopharmacol* 2004, **90**(1):91–97.
40. Miemanang R, Krohn K, Hussain H, Dongo E: **Paullinoside a and paullinamide a: a new cerebroside and a new ceramide from leaves of *Paullinia pinnata***. *Z Naturforsch* 2006, **61b**:1123–1127.
41. Akinpelu DA, Obuotor EM: **Antibacterial activity of *Piliostigma thonningii* stem bark**. *Fitoterapia* 2000, **71**(4):442–443.
42. Ibewuik JC, Ogundaini AO, Ogungbamila FO, Martin M-T, Gallard J-F, Bohlin L, Pais M: **Piliostigmin, a 2-phenoxychromone, and C-methylflavonols from *Piliostigma thonningii***. *Phytochemistry* 1996, **43**(3):687–690.
43. Pianaro A, Pinto P, Ferreira T, Ishikawa K, Braz-Filho R: **Iridoid glucoside and antifungal phenolic compounds from *Spathodea campanulata* roots**. *Semina: Ciências Agrárias, Londrina* 2007, **28**(2):251–256.
44. Simbo D: **An ethnobotanical survey of medicinal plants in Babungo, Northwest Region, Cameroon**. *Ethnobiol Ethnomed* 2010, **6**:8.
45. Ekor M, Ashorobi R, Ibitoye SF, Kasimi L: **Acute toxicity, analgesic potential and preliminary antimicrobial studies of the aqueous plant extract of *Spilanthes filicaulis***. *Niger J Health Biomed Sci* 2005, **4**(1):30–34.
46. Tan V, Njimi K, Ayafor F: **Screening of some African Medicinal plants for anticarcinogenic activity: part 1**. *Phytother res* 1997, **11**:45–47.
47. Ramsay KS, Wafo P, Ali Z, Khan A, Oluyemisi OO, Marasini BP, Khan IA, Bonaventure NT, Choudhary MI, Atta ur R: **Chemical constituents of *Stereospermum acuminatissimum* and their urease and alpha-chymotrypsin inhibitions**. *Fitoterapia* 2012, **83**(1):204–208.
48. Sob SV, Wabo HK, Tang CP, Tane P, Ngadjui BT, Ye Y: **Phenol esters and other constituents from the stem barks of *Stereospermum acuminatissimum***. *J Asian Nat Prod Res* 2011, **13**(12):1128–1134.
49. Ndjakou LB, Vonthron-Sénécheau C, Fongang Soh R, Tantangmo F, Ngouela KM, Tsamo E, Anton R, Weniger B: ***In vitro* antiprotozoal activities and cytotoxicity of some selected Cameroonian medicinal plants**. *J Ethnopharmacol* 2007, **111**(1):8–12.
50. Atta-ur-Rahman, Zareen S, Choudhary I, Akhtar N, Shujaat S, Ngounou N: **Some chemical constituents of *Terminalia glaucescens* and their enzymes inhibition activity**. *Z Naturforsch* 2005, **60b**:347–350.
51. Adebayo E, Ishola O: **Phytochemical and antimicrobial screening of crude extracts from the root, stem bark, and leaves of *Terminalia glaucescens***. *Afr J Pharm Pharmacol* 2009, **3**(5):217–221.
52. Kimmig A, Gekeler V, Neumann M, Frese G, Handgretinger R, Kardos G, Diddens H, Niethammer D: **Susceptibility of multidrug-resistant human leukemia cell lines to human interleukin 2-activated killer cells**. *Cancer Res* 1990, **50**(21):6793–6799.
53. Efferth T, Sauerbrey A, Olbrich A, Gebhart E, Rauch P, Weber HO, Hengstler JG, Halatsch ME, Volm M, Tew KD, Ross DD, Funk JO: **Molecular modes of action of artesunate in tumor cell lines**. *Mol Pharmacol* 2003, **64**(2):382–394.
54. Gillet J, Efferth T, Steinbach D, Hamels J, de Longueville F, Bertholet V, Remacle J: **Microarray-based detection of multidrug resistance in human tumor cells by expression profiling of ATP-binding cassette transporter genes**. *Cancer Res* 2004, **64**(24):8987–8993.
55. Huang HS, Nagane M, Klingbeil CK, Lin H, Nishikawa R, Ji XD, Huang CM, Gill GN, Wiley HS, Cavenee WK: **The enhanced tumorigenic activity of a mutant epidermal growth factor receptor common in human cancers is mediated by threshold levels of constitutive tyrosine phosphorylation and unattenuated signaling**. *J Biol Chem* 1997, **272**(5):2927–2935.
56. O'Brien J, Wilson I, Orton T, Pognan F: **Investigation of the Alamar Blue (resazurin) fluorescent dye for the assessment of mammalian cell cytotoxicity**. *Eur J Biochem* 2000, **267**(17):5421–5426.
57. Suffness M, Pezzuto JM: **Assays related to cancer drug discovery**. In *Methods in Plant Biochemistry: Assays for Bioactivity*. Edited by Hostettmann K. London: Academic Press; 1990:6.
58. Longley D, Johnston P: **Molecular mechanisms of drug resistance**. *J Pathol* 2005, **205**:275–292.
59. Kuete V, Efferth T: **Pharmacogenomics of Cameroonian traditional herbal medicine for cancer therapy**. *J Ethnopharmacol* 2011, **137**(1):752–766.

doi:10.1186/1472-6882-13-250

Cite this article as: Kuete et al.: Cytotoxicity of *Elaeophorbium drupifera* and other Cameroonian medicinal plants against drug sensitive and multidrug resistant cancer cells. *BMC Complementary and Alternative Medicine* 2013 **13**:250.