

RESEARCH ARTICLE

Open Access

Association of weight misperception with weight loss in a diabetes prevention program

Andrea L Hernan^{1*}, Vincent L Versace¹, Tiina Laatikainen^{1,2,3,4}, Erkki Vartiainen^{1,4}, Edward D Janus^{1,5} and James A Dunbar¹

Abstract

Background: Weight misperception may have an impact on perceived risk and susceptibility for chronic diseases. Little has been reported on the long term effects of this misperception in chronic disease interventions, particularly in field of diabetes prevention. The aim of this study was to investigate the relationship between weight misperception and weight loss during a diabetes prevention project conducted in south-east Australia with individuals at moderate to high risk of developing diabetes.

Methods: A total of n=251 at risk individuals provided self-reported weight during recruitment from 2004-2006. Objectively measured weight was assessed at baseline (0-21 days after recruitment), and subsequently at three months and 12 months after the intervention. Differences between self-reported and actual weight status are presented as percentages. Linear regression was used to investigate the relationship between weight misperception and weight loss, adjusting for baseline weight and BMI.

Results: Those who had high levels of under-reporting at baseline had greater weight loss at three and 12 months compared with those who under-reported to some degree, and those over-reporting their weight. A significant association was found between weight misperception and weight loss at the three and the 12 month time points. Baseline weight was not associated with weight loss.

Conclusions: Weight misperception should be acknowledged as a factor to be addressed when screening and identifying individuals at risk for diabetes. Screening and giving feedback is important in terms of awareness of participants' actual weight status and may have an effect on program outcomes.

Keywords: Weight misperception, Weight loss, Diabetes prevention, Risk screening, Risk perception

Background

Waist circumference and weight are the main predictors of type 2 diabetes, and therefore weight loss and accurate weight perception are key parts in prevention.

It is widely acknowledged in the weight misperception literature that individuals tend to over-report their height and under-report their weight [1]. Various factors have been attributed to weight misperception such as age, gender, ethnicity, socio-economic status, being overweight or obese and a social desirability for tallness and thinness [1-3].

Under-reporting weight in overweight or obese individuals has been linked with reduced perception of health risks [4] and of the advantages of weight loss [5], and also less physical activity and attempts at weight loss [6]. However, when overweight perception is corrected positive associations between weight perception and weight loss have been identified [7-10].

Weight misperception may have an impact on perceived risk and susceptibility for chronic diseases. Little has been reported on the long term effects of this misperception in chronic disease interventions, particularly in field of diabetes prevention. If individuals misperceive their weight status and therefore disease risk, the effectiveness of intervention screening and program uptake may be reduced.

The aim of this paper was to investigate the association between weight misperception and weight loss during an

* Correspondence: andrea.hernan@greaterhealth.org

¹Greater Green Triangle University Department of Rural Health, Flinders University and Deakin University, PO Box 423, Warrnambool, VIC 3280, Australia

Full list of author information is available at the end of the article

Australian diabetes prevention program conducted with moderate to high risk individuals [11]. We hypothesise that there would be a positive relationship between weight misperception and weight loss at three and 12 months post intervention. The underlying assumption for this hypothesis was that individuals who have high levels of weight misperception are likely to be alarmed and more motivated into taking action upon realisation of actual weight status.

Methods

The methods, recruitment process and results for the Greater Green Triangle Diabetes Prevention Project (GGT DPP) conducted in south east Australia between 2004-2006 have been previously described [11]. The GGT DPP was a diet and lifestyle intervention in obese and overweight subjects at moderate to high risk of developing type 2 diabetes. Briefly, recruitment occurred through opportunistic screening by study nurses of rural individuals aged 40-75 years presenting at local general practices. Individuals who appeared to be over the age of 40 and who were considered overweight were approached during recruitment. A self-administered risk screening tool (Finnish Diabetes Risk Score, FINDRISC) was used to identify individuals at moderate to high risk of developing type 2 diabetes [12].

During recruitment individuals were asked to complete self-reported anthropometric information on the FINDRISC form which included a height, weight and waist circumference category [13]. Participating individuals attended a baseline testing session within three weeks of being screened, although most were tested on the same day as providing self-reported anthropometric information. Objectively measured weight was among the anthropometric measures taken by specially trained study nurses using the European Health Risk Monitoring protocol [14]. Participants attended a further three and 12 month testing session post-intervention where weight was measured amongst other anthropometric, biochemical and behavioural measures. Weight was measured using the same standardised mechanical column beam balance scales (Seca) at the three time points of testing.

Analysis

A retrospective post-hoc analysis was undertaken with data collected during the implementation of the GGT DPP. Weight misperception was calculated as the difference between participants' self-reported weight before baseline testing and the actual weight recorded at baseline testing. This variable was converted to per cent values. The weight recorded at baseline was measured in kg to one decimal place and was used to calculate BMI (kg/m^2). The weight loss variables (i.e. three months and 12 months) were calculated as the difference between

the clinically recorded weights relative to the weight at baseline and were converted to per cent values.

Linear regression ('Enter' method) was used to investigate the relationships between the dependent variables of percentage weight change at three and 12 months, and the explanatory variables of weight misperception, baseline weight and BMI. Weight misperception was included in all models with baseline weight and BMI included separately. These analyses were carried out on the pooled sample of males and females, and also stratified by gender. Linear regression was also used to assess the influence of age and years of formal education on weight misperception. Out of $n = 311$ eligible participants, $n = 251$ are included for these analyses at baseline because participants from one site ($n = 60$) were excluded as only objective measurements were recorded. At three months there were $n = 201$ participants, and at 12 months there were $n = 208$ participants included in the analyses.

Ethics approval

Written informed consent was obtained from the GGT DPP participants for the publication of this report. This study was approved by the Flinders University Clinical Research Committee (reference number 105/034).

Results

There were a total of $n = 62$ males and $n = 189$ females, the mean age for males was 57 years and the mean age for females was 56 years, the majority were married (males 74.2%, females 70.4%), and 63.3% of males and 49.2% of females were employed. Males had a mean 11.7 years of formal education compared to females 11.9 years. Mean weight for the pooled sample at baseline was 92.6 kg and mean weight loss at 3 months was 2.53 kg and mean weight loss at 12 months was 2.45 kg.

Weight misperception by categories of under or over-reporting and associated weight loss at three and 12 month follow up are presented in Table 1. Those who had a high level of under-reporting at baseline ($>5\%$) had greater weight loss at three and 12 months (-3.52% and -3.70% respectively) compared with those who under-reported to some degree or over-reported their weight.

There was a significant relationship between weight misperception and weight loss both at three and 12 months ($B = 0.205$ and $B = 0.220$ respectively) for both genders combined (Table 2). The same positive relationship was found for males and females, but was only significant in the latter. This was possibly because of the relatively low number of males in the sample (data not shown).

The variables baseline BMI and baseline weight were not found to be statistically significant, indicating that the relationship between weight misperception and weight

Table 1 Weight loss at 3 and 12 months by weight misperception categories

	n	3 month follow up		12 month follow up			
		n	Weight % Δ 3 months (SE)	Weight kg Δ 3 months (SE)	n	Weight % Δ 12 months (SE)	Weight kg Δ 12 months (SE)
Weight misperception category							
Under-report >5%	70	58	-3.5 (0.5)	-3.5 (0.5)	59	-3.7 (0.6)	-3.6 (0.6)
Under-report 2%-4.99%	72	59	-2.6 (0.5)	-2.8 (0.4)	60	-3.0 (0.8)	-2.9 (0.8)
Under or over report -1.99% - 1.99%	87	67	-1.2 (0.4)	-1.6 (0.3)	71	-1.7 (0.65)	-1.7 (0.6)
Over-report >2%	22	18	-2.3 (0.7)	-2.3 (0.5)	18	-0.5 (1.24)	-0.04 (1.28)
Total	251	201	-2.4 (0.3)	-2.5 (0.2)	208	-2.6 (0.4)	-2.5 (0.4)

Δ = change, kg = kilograms, SE = standard error.

loss at the two time periods were independent of baseline weight (Table 2). That is neither baseline weight nor BMI were significant predictors of weight loss. Approximately 7% of the variation of the weight loss at three months and 3% at 12 months were explained by weight misperception.

Age and years of formal education were non-significant explanatory variables of weight loss at three months (age: $B = 0.038$, $p = 0.162$; formal education: $B = 0.007$, $p = 0.922$) and 12 months (age, $B = 0.034$, $p = 0.146$; formal education: $B = 0.006$, $p = 0.922$) when included in multiple regression models with weight misperception. Weight misperception was significantly associated with weight loss at three months ($B = 0.190$, $p < 0.001$ in the model including age and $B = 0.205$, $p < 0.001$ in the model including formal education) and 12 months ($B = 0.158$, $p < 0.001$ in the model including age and $B = 0.173$, $p < 0.001$ in the model including formal education).

Discussion

This study identified weight misperception as a variable associated with weight loss in a diabetes prevention program over time. This association existed independently of obesity level at baseline, indicating that misperception has the same effect in overweight and obese individuals. Furthermore, those who had high levels of under-reporting weight at baseline had greater weight loss compared with those who either under-reported to some degree or over-reported their weight.

We believe that individuals who have high levels of weight misperception (under-reporters) are likely to be

alarmed into taking action and more motivated to lose weight compared with those who over-report or have accurate weight assessments. This finding reflects the social cognitive and self-regulation theories of health behaviour change [15-18], where motivation is formed through the development of outcome expectations, self-efficacy and risk perception [17,19,20]. According to these theories correcting weight misperception may be considered as a precursor or additional element which contributes to building risk perception, like the psychological and behavioural changes that have been shown to predict bio physiological outcomes in this diabetes prevention program [21].

There is a paucity of literature investigating the association between weight misperception and observed weight loss in prevention studies. It is possible that other predictors for diabetes, such as family history and experience of chronic disease, are stronger contributors for building risk perception than weight perception. Harwell et al. (2001) found that rural survey participants who were obese and had a family history of diabetes were more likely to consider themselves at risk for diabetes [22]. Dorman et al. (2012) found that those with increased familial risk for diabetes, CHD and stroke had high levels of perceived risk and worry about diabetes [23]. In contrast, Adriannse et al (2003) found that participants in a diabetes screening program with diabetes risk factors such as having a family history, obesity and hypertension, did not perceive themselves to be at increased risk for the disease [24]. The findings from this

Table 2 Regression models examining weight misperception and weight loss at 3 and 12 months

		Weight loss after 3 months				Weight loss after 12 months			
		Constant	B	p-value	Adj r ²	Constant	B	p-value	Adj r ²
Model 1	Weight Misperception (%)	-1.790	0.205	<0.001	0.068	-1.94	0.220	0.006	0.032
Model 2	Weight Misperception (%)	-2.35	0.210	<0.001	0.065	0.378	0.208	0.01	0.033
	Baseline BMI (kg/m ²)		0.017	0.679			-0.07	0.247	
Model 3	Weight Misperception (%)	-2.90	0.213	<0.001	0.068	-0.21	0.211	0.009	0.031
	Baseline Weight (kg)		0.012	0.368			-0.019	0.365	

Δ = change, kg = kilograms, BMI = Body Mass Index, Adj = Adjusted.

study emphasises the importance of population recognition and awareness of weight as one of the main contributors of risk for developing diabetes.

Weight misperception was an important explanatory variable indicating that the risk screening and giving feedback are essential components of the program. However weight misperception by itself has limited value for predicting percentage weight loss after three months and 12 months, as the intervention was the main contributor to weight loss. Further research is needed to confirm the predictive value of weight misperception towards building up risk perception and outcome expectations and their relationship to weight loss.

Strengths and limitations

The lower proportion of males in the sample could be viewed as a limitation with potential implications for internal validity through selection bias. By carrying out an analysis stratified by gender it was found that the direction of the regression coefficient was the same for both genders despite being non-significant for males. This study was a post-hoc retrospective analysis of a prospective cohort so there are limitations when undertaking analysis of this kind. This includes being unable to assess other potential confounding factors which could change the associations found.

These results can only be generalized to intervention populations recruited from general practitioner clinics for diabetes prevention programs given the non-random selection of participants from the clinic waiting rooms. It is known that those recruited in interventions already have some motivation and intention for behaviour change [25]. These results, however, do provide important information for planning and implementation of lifestyle interventions.

Conclusions

The results from this study suggest the importance of population screening and creating obesity awareness for diabetes risk. We suggest that weight misperception should be addressed during the screening process for diabetes prevention programs in order to better develop risk perception, build perceived susceptibility, and understand the seriousness of the disease risk.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

AH conceived and designed the study. TL, EDJ and JAD were involved in acquisition of data. VV with AH, TL, EV and EDJ analysed and interpreted the data. AH drafted the manuscript and was responsible for its revisions. VV, TL, EV, EDJ and JAD contributed to specific sections in the manuscript. JAD and was the chief investigator for the GGT DPP. All authors read and approved the final manuscript.

Acknowledgements

We would like to thank the participating general practices in Hamilton, Horsham and Mt Gambier, and the study nurses. We would also like to acknowledge Benjamin Philpot for his statistical and conceptual advice. The study was funded by the Australian Government Department of Health and Ageing, Canberra, Australia.

Author details

¹Greater Green Triangle University Department of Rural Health, Flinders University and Deakin University, PO Box 423, Warrnambool, VIC 3280, Australia. ²Institute of Public Health and Clinical Nutrition, University of Eastern Finland, PO Box 1627, Kuopio 70211, Finland. ³Hospital District of North Karelia, Tikkamäentie 16, Joensuu 80210, Finland. ⁴National Institute for Health and Welfare, Mannerheimintie 166, Helsinki 00300, Finland. ⁵Department of Medicine, North West Academic Centre, The University of Melbourne, Western Hospital, Footscray, VIC 3011, Australia.

Received: 26 March 2013 Accepted: 29 January 2014

Published: 30 January 2014

References

1. Gorber SC, Tremblay M, Moher D, Gorber B: A comparison of direct vs. self-report measures for assessing height, weight and body mass index: a systematic review. *Obes Rev* 2007, **8**(4):307–326.
2. Park JY, Mitrou PN, Keogh RH, Luben RN, Wareham NJ, Khaw KT: Effects of body size and sociodemographic characteristics on differences between self-reported and measured anthropometric data in middle-aged men and women: the EPIC-Norfolk study. *Eur J Clin Nutr* 2011, **65**(3):357–367.
3. Stommel M, Schoenborn C: Accuracy and usefulness of BMI measures based on self-reported weight and height: findings from the NHANES & NHIS 2001–2006. *BMC Public Health* 2009, **9**(1):421.
4. Gregory CO, Blanck HM, Gillespie C, Maynard LM, Serdula MK: Perceived health risk of excess body weight among overweight and obese men and women: differences by sex. *Prev Med* 2008, **47**(1):46–52.
5. Riddell LJ, Inman V: Body weight and weight loss: are health messages reaching their target? *Asia Pac J Clin Nutr* 2007, **16**(4):683.
6. Duncan DT, Wolin KY, Scharoun-Lee M, Ding EL, Warner ET, Bennett GG: Does perception equal reality? Weight misperception in relation to weight-related attitudes and behaviors among overweight and obese US adults. *Int J Behav Nutr Phys Act* 2011, **8**(1):20.
7. DeVille-Almond J, Tahrani AA, Grant J, Gray M, Thomas GN, Taheri S: Awareness of obesity and diabetes: a survey of a subset of british male drivers. *Am J Mens Health* 2011, **5**(1):30–37.
8. Inoue M, Toyokawa S, Inoue K, Suyama Y, Miyano Y, Suzuki T, Miyoshi Y, Kobayashi Y: Lifestyle, weight perception and change in body mass index of Japanese workers: MY Health Up Study. *Pub Health* 2010, **124**(9):530–537.
9. Lemon SC, Rosal MC, Zapka J, Borg A, Andersen V: Contributions of weight perceptions to weight loss attempts: differences by body mass index and gender. *Body Image* 2009, **6**(2):90–96.
10. Yaemsiri S, Slining M, Agarwal S: Perceived weight status, overweight diagnosis, and weight control among US adults: the NHANES 2003–2008 Study. *Int J Obes* 2010, **35**(8):1063–1070.
11. Laatikainen T, Dunbar J, Chapman A, Kilkkinen A, Vartiainen E, Heistaro S, Philpot B, Absetz P, Bunker S, O'Neil A, et al: Prevention of type 2 diabetes by lifestyle intervention in an Australian primary health care setting: Greater Green Triangle (GGT) Diabetes Prevention Project. *BMC Public Health* 2007, **7**(1):249.
12. Lindstrom J, Tuomilehto J: The diabetes risk score: a practical tool to predict type 2 diabetes risk. *Diabetes Care* 2003, **26**(3):725–731.
13. Hernan A, Philpot B, Janus E, Dunbar J: Recruitment into diabetes prevention programs: what is the impact of errors in self-reported measures of obesity? *BMC Public Health* 2012, **12**:510.
14. Tolonen H, Kuulasmaa K, Laatikainen T, Wolf H: *European Health Risk Monitoring Project. Recommendation for indicators, international collaboration, protocol and manual of operations for chronic disease risk factor surveys.* Helsinki: Finnish National Public Health Institute; 2002.
15. Gollwitzer PM: Implementation intentions: strong effects of simple plans. *Am Psychol* 1999, **54**(7):493.
16. Schwarzer R: Modeling Health Behavior Change: How to Predict and Modify the Adoption and Maintenance of Health Behaviors. *Appl Psychol* 2008, **57**(1):1–29.

17. Schwarzer R, Fuchs R: **Self efficacy and health behaviors**. In *Predicting health behavior: Research and practice with social cognition models*. Edited by Connor M, Norman P. Buckingham, UK: Open University Press; 1996:163–196.
18. Uutela A, Absetz P, Nissinen A, Valve R, Talja M, Fogelholm M: **Health Psychological Theory in Promoting Population Health in Pajjat-Hame, Finland: First Steps toward a Type 2 Diabetes Prevention Study**. *J Health Psychol* 2004, **9**(1):73–84.
19. Bandura A: *Self-efficacy: The exercise of control*. New York: Freeman; 1997.
20. Schwarzer R: **Self-efficacy in the adoption and maintenance of health behaviors: theoretical approaches and a new model**. In *Self-efficacy: thought control of action*. Edited by Schwarzer R. Washington, DC: Hemisphere; 1992:217–242.
21. Laatikainen T, Philpot B, Hankonen N, Sippola R, Dunbar JA, Absetz P, Reddy P, Davis-Lameloise N, Vartiainen E: **Predicting changes in lifestyle and clinical outcomes in preventing diabetes: the Greater Green Triangle Diabetes Prevention Project**. *Prev Med* 2012, **54**:157–161.
22. Harwell TS, Dettori N, Flook BN, Priest L, Williamson DF, Helgeson SD, Gohdes D: **Preventing Type 2 Diabetes Perceptions about risk and prevention in a population-based sample of adults ≥ 45 years of age**. *Diabetes Care* 2001, **24**(11):2007–2008.
23. Dorman JS, Valdez R, Liu T, Wang C, Rubinstein WS, O'Neill SM, Acheson LS, Ruffin MT IV, Khoury MJ: **Health beliefs among individuals at increased familial risk for type 2 diabetes: Implications for prevention**. *Diabetes Res Clin Pract* 2012, **96**(2):156–162.
24. Adriaanse M, Snoek F, Dekker J, Spijkerman A, Nijpels G, Van der Ploeg H, Heine R: **Perceived risk for Type 2 diabetes in participants in a stepwise population-screening programme**. *Diabet Med* 2003, **20**(3):210–215.
25. Hankonen N, Vollmann M, Renner B, Absetz P: **What is setting the stage for abdominal obesity reduction? A comparison between personality and health-related social cognitions**. *J Behav Med* 2010, **33**(5):415–422.

doi:10.1186/1471-2458-14-93

Cite this article as: Hernan *et al.*: Association of weight misperception with weight loss in a diabetes prevention program. *BMC Public Health* 2014 **14**:93.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

