

RESEARCH ARTICLE

Open Access

SB 206553, a putative 5-HT_{2C} inverse agonist, attenuates methamphetamine-seeking in rats

Steven M Graves^{1,2*} and T Celeste Napier^{1,2}

Abstract

Background: Methamphetamine (meth) dependence presents a substantial socioeconomic burden. Despite the need, there is no FDA-approved pharmacotherapy for psychostimulant dependence. We consider 5-HT_{2C} receptors as viable therapeutic targets. We recently revealed that the atypical antidepressant, mirtazapine, attenuates meth-seeking in a rodent model of human substance abuse. Mirtazapine historically has been considered to be an antagonist at 5-HT_{2C} receptors, but more recently shown to exhibit inverse agonism at constitutively active 5-HT_{2C} receptors. To help distinguish the roles for antagonism vs. inverse agonism, here we explored the ability of a more selective 5-HT_{2C} inverse agonist, SB 206553 to attenuate meth-seeking behavior, and compared its effects to those obtained with 5-HT_{2C} antagonists, SDZ Ser 082 and SB 242084. To do so, rats were trained to self-administer meth and tested for seeking-like behavior in cue reactivity sessions consisting of contingently presenting meth-associated cues without meth reinforcement. We also explored motor function to determine the influence of SB 206553 and SDZ Ser 082 on motor activity in the presence and absence of meth.

Results: Like mirtazapine, pretreatment with SB 206553 (1.0, 5.0, and 10.0 mg/kg), attenuated meth-seeking. In contrast, the antagonists, SDZ Ser 082 (0.1, 0.3, and 1.0 mg/kg) and SB 242084 (3.0 mg/kg) had no effect on cue reactivity (CR). SB 242084 (3.0 mg/kg) failed to attenuate the effects of 5.0 and 10 mg/kg SB 206553 on CR. Motor function was largely unaltered by the 5-HT_{2C} ligands; however, SB 206553, at the highest dose tested (10.0 mg/kg), attenuated meth-induced rearing behavior.

Conclusions: The lack of effect by 5-HT_{2C} antagonists suggests that meth-seeking and meth-evoked motor activity are independent of endogenous 5-HT acting at 5-HT_{2C} receptors. While SB 206553 dramatically impacted meth-evoked behaviors it is unclear whether the observed effects were 5-HT_{2C} receptor mediated. Thus, SB 206553 deserves further attention in the study of psychostimulant abuse disorders.

Keywords: Methamphetamine, Serotonin, Self-administration, Motor, Inverse agonist, Seeking

Background

Psychostimulants such as cocaine and methamphetamine (meth) dramatically enhance transmission of monoamines, including serotonin (5-HT). It is clear that the 5-HT_{2C} receptor subtype is involved in stimulant-mediated behaviors [1,2]. For example, 5-HT_{2C} receptor agonists decrease cue-induced reinstatement of cocaine-seeking [3-6] (5-HT_{2C} antagonists have no effect [5,7]). These latter findings suggest that cue-associated seeking behavior

is not dependent on 5-HT levels at synapses expressing 5-HT_{2C} receptors (illustrated by the lack of effect of antagonists on cocaine cue-induced reinstatement); nonetheless, the systems regulating seeking behavior are under negative control by activated 5-HT_{2C} receptors (illustrated by the ability of 5-HT_{2C} agonists to decrease cue-induced reinstatement of cocaine-seeking). The majority of behavioral studies focusing on 5-HT_{2C} receptors have relied on agonist/antagonist relationships. However, evidence indicates constitutive activity (i.e. agonist-independent activation) of 5-HT_{2C} receptors *in vivo* [8-10] and it currently is unknown how constitutively active 5-HT_{2C} receptors impact stimulant-induced behaviors.

5-HT_{2C} receptors are the only known G protein coupled receptor to undergo mRNA editing by adenosine

* Correspondence: Steven_Graves@rush.edu

¹Department of Pharmacology Rush, University Medical Center, 1735 W Harrison Street, Cohn Research Building, Chicago, IL 60612, USA

²Center for Compulsive Behavior and Addiction, Rush University Medical Center, 1735 W Harrison Street, Cohn Research Building, Chicago, IL 60612, USA

deaminases [11,12]; a process that has important implications for the level of constitutive activity. Adenosine to inosine switches result in amino acid substitutions in the second intracellular loop of the receptor protein [13-15]. The unedited 5-HT_{2C} receptor displays high levels of constitutive activity [16-20], and these receptor protein substitutions decrease constitutive activity. Constitutive activity can be attenuated pharmacologically by inverse agonists, these drugs stabilize the receptor in its inactive conformation [21,22]. For example, systemic administration of the 5-HT_{2C} inverse agonist SB 206553 (SB206) enhances dopamine (DA) efflux in the nucleus accumbens and striatum in rats [8], whereas an agonist either decreases or has no effect on accumbal and striatal DA [8,23-27]. Supporting a role of constitutive activity in this effect, the SB206-induced effects were verified to occur independent of endogenous 5-HT [8]. We previously revealed that the atypical antidepressant, mirtazapine, attenuates methamphetamine (meth)-induced sensitization [28], place conditioning [29-31], and seeking [31] in rodent models of human substance abuse. Mirtazapine historically has been considered to be an antagonist at 5-HT_{2C} and other receptors [32-34]; however, more recent studies indicate its action as an inverse agonist at constitutively active 5-HT_{2C} receptors [35,36]. These new observations raised the following questions: Would a more selective 5-HT_{2C} receptor inverse agonist, like mirtazapine, also reduce meth-seeking? A related question is whether seeking for meth is similar to, or differs from, that for cocaine with regard to sensitivity to selective 5-HT_{2C} inverse agonists and/or antagonists?

To answer the above questions, we used two 5-HT_{2C} antagonists of different chemical structures, SB 242084 (SB242) and SDZ Ser 082 (SDZ) as well as the putative 5-HT_{2C} receptor inverse agonist SB206. Dose-response evaluations on meth-seeking behaviors were made for these drugs in using cue reactivity (CR) to meth-associated cues established during a self-administration protocol. We have previously demonstrated that CR offers an effective means to study drug-seeking that, as is more typical in the human scenario, does not depend on prior extinction training [31]. We also revealed that the ability of mirtazapine to alter seeking behavior is similar for CR and cue-induced reinstatement (following extinction) protocols [37]. Finally, to ascertain whether 5-HT_{2C} receptor compounds influenced motor activity, rats tested for CR also underwent motor assessments with acute 5-HT_{2C} ligand pretreatment.

Methods

Subjects

Fifty male Sprague-Dawley rats were purchased from Harlan (Indianapolis, IN), acclimated to housing in our

local vivarium for 5 days, and handled a minimum of 3 times prior to surgery. Food and water were provided *ad libitum* throughout the study. Rats were maintained in accordance with the Guide for Care and Use of Laboratory Animals (National Research Council, Washington DC) and with the guidelines and approval of the Rush University Institutional Animal Care and Use Committee.

Drugs

(+)-Methamphetamine HCl (Sigma, St. Louis, MO) was dissolved in sterile saline. The stimulant was self-administered at 0.1 mg/kg/0.1 mL infusion intravenously (iv) and non-contingently administered intraperitoneally (ip) for motor assessments as 1.0 mg/kg. SDZ Ser 082 (SDZ; Tocris, Ellisville, MO), a 5-HT_{2C} antagonist, was dissolved in saline and injected ip (0.1, 0.3, and 1.0 mg/kg). SB 242084 (SB242; Tocris), also a 5-HT_{2C} antagonist, was dissolved in 8 % β -cyclodextran and 1 % citric acid in deionized water and injected ip (3.0 mg/kg). SB 206553 (SB206) (Tocris), a 5-HT_{2C} inverse agonist, was dissolved in 1 % lactic acid in deionized water and injected ip (1.0, 5.0, and 10.0 mg/kg). Serotonin_{2C} ligands were administered in volumes of 1.0 ml/kg with the exception of 10.0 mg/kg SB206, which was administered at 2 ml/kg (from a 5.0 mg/ml stock solution). All drugs were administered as the base. pK_i values of SDZ, SB242 and SB206 for 5-HT₂ receptor subtypes are provided in Table 1.

Surgical procedures

Rats were instrumented with a jugular vein catheter under continuous isoflurane anesthesia. Custom built catheters were constructed with silastic tubing (0.3 mm i.d. x 0.64 mm o.d.; Dow Corning Co., Midland, MI) and implanted into the right jugular vein. The distal end of the catheter extended to the mid-scapular region with a metal guide canulae (22 gauge; Plastics One Inc., Roanoke, VA) and anchored to a plastic mesh. Rats were allowed to recover for a minimum of 5 days prior to beginning self-administration procedures. During this time, rats were handled, inspected and weighed daily to ensure that surgical wounds were healing properly (without infection), that normal behavioral grooming/eating patterns were retained, and that preoperative weight was regained.

Table 1 5-HT_{2C} receptors

| | 5-HT _{2A} | 5-HT _{2B} | 5-HT _{2C} |
|-------------|--------------------|--------------------|--------------------|
| SDZ Ser 082 | 6.3 | 6.7 | 8.1 |
| SB 242084 | 6.1 | 6.8 | 8.2 |
| SB 206553 | 5.6 | 7.7 | 7.8 |

pK_i values (nM) for 5-HT₂ receptor ligands in HEK-293 or CHO-K1 cells expressing human recombinant 5-HT₂ receptors [45].

Self-Administration acquisition

Rats were trained to self-administer meth 3 hr/day for 14 consecutive days in standard operant chambers enclosed in ventilated, sound attenuating cabinets (Med-Associates, St. Albans, VT). Each operant chamber contained two levers; the left lever assigned as the “active” lever and the right the “inactive” lever. Above each lever was a cue light and located on the opposite wall was an in-house light. A cue light above the active lever was activated in association with the infusion pump and the in-house light was subsequently activated for 20s, indicating a time-out period during which responses had no programmed consequences. Responding on the inactive lever also had no programmed consequences. On days 1–7, rats self-administered meth for 3 hr/day on a fixed ratio (FR) 1 schedule of reinforcement. On days 8–14, rats self-administered on a FR5 for 3 hr/day (refer to Figure 1) to enhance lever pressing behaviors and resistance to extinction [38]. Starting on day 8, rats were acclimated to injection procedures *via* daily ip vehicle pretreatments (30 min). The number of active lever presses, inactive lever presses, and infusions were recorded for all sessions. Stable self-administration behavior was operationally defined as <15 % infusion variability between days 13 and 14; failure to reach this criterion resulted in subject exclusion from the study.

Cue reactivity testing

Meth-seeking behavior was assessed *via* CR tests. CR testing consisted of single, 1 hr extinction sessions during which cues (i.e., cue light, time-out house light, and activation of infusion pump) were contingently presented on an FR1 schedule. During CR tests, infusion lines were filled with sterile saline and remained connected to saline-filled syringes (syringes were disconnected from

pumps to prevent excessive fluid intake). Meth was not accessible during CR testing. The number of active and inactive lever presses was measured continuously and tallied in 15 min intervals. On day 15, rats were given a 30 min pretreatment of vehicle and tested for baseline meth-seeking (CR1); baseline was used to equally distribute rats into treatment groups. Rats were tested for an additional four CR tests (CR2-5); between each CR test, rats were allowed to self-administer meth for 3 hr/day on an FR5 schedule (referred to as *intermittent self-administration*; see Figure 1) to prevent extinction training. Rats failing to administer at least 50 % of the number of infusions administered on day 14, for two consecutive days during intermittent self-administration sessions, were removed from the study. Rats were tested for CR after a 30 min pretreatment of 0.1, 0.3, 1.0 mg/kg SDZ and vehicle (treatment group 1) or 1.0, 5.0, 10.0 mg/kg SB206, and vehicle (group 2). The 5-HT_{2C} receptor antagonist, SB242, was tested against the 5-HT_{2C} receptor inverse agonist SB206 in the same rats, wherein SB242 (3.0 mg/kg ip) was administered 45 min, and SB206 (5.0 or 10.0 mg/kg ip) was administered 30 min, prior to the onset of CR testing. Dose order was randomized for all treatment groups. Doses of SB206 and SB242 were guided by literature demonstrating significant neurochemical effects in the nucleus accumbens [8]. Doses of SDZ were selected based on reports revealing an enhancement of the interoceptive cues of cocaine [39] and cocaine-evoked motor activity in naïve rats [40]. We have previously shown that the described CR paradigm is comparable to cue-induced reinstatement and that pharmacological intervention paradigms similar to those tested in the current report similarly reduce CR and cue-induced reinstatement [37]. Assessments were made at an early stage of withdrawal (24 hr) based on our aforementioned work [37] as well as evidence indicating that psychostimulant administration dysregulates neuronal function in the nucleus accumbens shell (a region with constitutively active 5-HT_{2C} receptors [10]) as early as 1–3 days of withdrawal [41] as well as biochemical indices indicating psychostimulant-induced plasticity in multiple brain regions during early phase withdrawal (1–3 days) using place conditioning paradigms [42,43] and motor sensitization [44].




Figure 1 Timeline of experimental protocol. Rats self-administered methamphetamine (meth) for 14 consecutive days (3 hrs/day). Days (D) 1–7 a fixed ratio (FR) 1 schedule of reinforcement was used for training. To increase responding rates, an FR5 was employed for days 8–14. On day 15, rats were acclimated to the cue reactivity (CR) protocol. Assessments of 5-HT_{2C} ligands on meth-seeking were made days 18, 21, 24, and 27 (i.e., CR2-CR5); treatments for CR were randomized. Between each CR test, rats were allowed to self-administer meth for two consecutive days on an FR5 schedule (3 hrs/day). Five days after CR5 (on day 33), motor assessments were initiated.

Motor assessments

A subset of rats tested for CR dose–response assessments was used to determine the motor effects of SDZ (1.0 mg/kg ip) and SB206 (5.0 and 10.0 mg/kg ip) in the presence and absence of meth (1.0 mg/kg ip). For this study, rats were withdrawn from meth and remained in their home cages during protocol days 28–32 and received no treatment (refer to Figure 1). After this 5 day period, rats were tested for motor activity for 3

consecutive days (days 33–35). All motor assessments were conducted using automated small animal activity boxes (Accuscan Instruments, Columbus, OH) equipped with two banks of photobeams positioned at different heights to characterize motor activity in three dimensional space. Rats were habituated to activity chambers for 1 hr prior to each motor test. On day 33, rats were administered 1 ml/kg of the respective vehicle for each test drug (rats for effects of SDZ were administered saline and rats tested for SB206-induced effects were administered 1 % lactic acid in deionized water). The injected rats were immediately returned to motor boxes for 1 hr after which rats were injected with either SDZ (1.0 mg/kg), or SB206 (5.0 or 10.0 mg/kg) and behavior was recorded for an additional 1 hr. Motor data collected 30 min post-injection were subsequently analyzed; this time frame reflected the one that was relevant to CR behaviors. On day 34, rats were administered a 30 min pretreatment of vehicle (saline or 1 % lactic acid), then administered 1 mg/kg meth (ip) and behavior recorded for 1 hr. On day 35, the procedure from day 34 was repeated using SDZ (1.0 mg/kg), or SB206 (5.0 or 10.0 mg/kg) (ip) instead of respective vehicles. Peak meth effects occurred 15 min post meth injection; meth-evoked motor activity was therefore analyzed for the last 45 min of testing (i.e., 15 min post meth injection). Horizontal activity (number of beam breaks in the horizontal plane), vertical activity (number of beam breaks in the vertical plane indicating rearing-like behavior), and total distance (cm traversed within the chamber) were recorded. These assessments provide a reliable index of overall motor patterns evoked by this dose of meth [28]. Stereotypy (rapid, repetitive behaviors) is a prominent component of meth-induced motor activity [28]; therefore, stereotypy number (the number of beam breaks repetitively disrupted) also was analyzed for meth-evoked motor activity. Rats tested for effects of SDZ on motor function were also tested for SDZ effects on CR; similarly, rats tested for effects of SB206 on motor function had prior exposure to SB206 during CR assessments.

Statistical analysis

Active lever presses, inactive lever presses, and number of infusions achieved during self-administration sessions were analyzed using a one-way rmANOVA with Newman-Keuls *post-hoc* analysis. The differences in lever pressing during CR tests were detected using two-way rmANOVA with Newman-Keuls *post-hoc* analysis. Lever pressing behavior for days 16 and 26 of intermittent self-administration, and motor activity were analyzed using a paired *t*-test. For all tests, $\alpha = 0.05$. Data are presented as mean \pm SEM. Data greater than two standard deviations from the mean were considered to

be statistical outliers and were excluded from the analyses; for CR, outliers were determined from behavior collapsed across the first 30 min of the tests.

Results


Stable meth self-administration

Fifty rats acquired the self-administration task; four rats were removed from the study as they displayed >15 % infusion variability between the last two training sessions (i.e., days 13 and 14); three rats died after a self-administration session. An additional 3 rats were removed due to failure to maintain stable self-administration behavior during periods of intermittent self-administration (i.e., responding for two consecutive days fell below 50 % of the number of reinforcements received on day 14 of self-administration). Of the 40 rats that completed the study, there were no differences for active lever pressing ($p = 0.89$; one statistical outlier removed), inactive lever pressing ($p = 0.23$; three statistical outliers removed), or infusions ($p = 0.76$; one statistical outlier removed) for days 11–14 (paired *t*-test; Figure 2A).

Meth-seeking behavior: cue reactivity


A within-subjects design was used wherein each rat was tested with three doses (randomized) of either the antagonist (SDZ; $n = 11$), inverse agonist (SB206; $n = 8$) and the respective vehicles. Similarly, a within-subjects design was used to test for antagonist/inverse agonist interactions with 3.0 mg/kg SB242 *vs.* 10.0 ($n = 11$) and 5.0 mg/kg ($n = 10$) SB206, and respective vehicles. There were no significant differences among rats assigned to SDZ (treatment group 1), SB206 (treatment group 2), or interaction studies (treatment groups 3 and 4) for active or inactive lever pressing during CR1 (ANOVA, $p > 0.05$; data not shown). Additionally, active lever pressing comparing CR after vehicle pretreatment was not different among the four treatment groups (ANOVA, $p > 0.05$; data not shown). Self-administration behavior was not altered by the interposed CR testing sessions; the first and last day of intermittent self-administration were not significantly different for active lever presses, inactive lever presses, or infusions comparing day 16 and day 26 (paired *t*-test, $p > 0.05$; Figure 2B). In agreement with our prior report validating consistent self-administration and seeking [37], these data verify that behavior is persistent throughout the paradigm.

Independent of the CR test number or pretreatment, meth-seeking behavior occurred most intensely within the first 15 min of CR testing (Figures 3 and 4) as previously reported [37]; by 30 min, levels emulated those expressed on the inactive lever (data not shown). It is also important to note that the number of lever presses achieved during self-administration sessions are far greater than the number during CR tests with a vehicle



pretreatment. This is a consequence of i) lever presses during CR tests are not reinforced with meth and ii) self-administration sessions are 3 hrs long whereas CR tests are shown in 15 min intervals.

SDZ, the 5-HT_{2C} antagonist, had no effect on number of active (Figure 3A) or inactive lever pressing (data not shown) at any dose tested. Two-way rmANOVA comparing active lever pressing revealed no significant Treatment effect ($F_{(3,36)} = 0.81$), a significant Time effect ($F_{(1,36)} = 97.57$), and no Treatment x Time interaction ($F_{(3,36)} = 0.49$). In contrast, administration of the 5-HT_{2C} inverse agonist SB206 attenuated active lever pressing in the first 15 min at all doses tested (Figure 3B) resulting



in an approximately 25 %, 50 %, and 95 % reduction in active lever pressing by 1.0, 5.0, and 10.0 mg/kg, respectively. Two-way rmANOVA revealed a significant effect of Treatment ($F_{(3,24)} = 5.97$) and Time ($F_{(1,24)} = 34.70$), and a Treatment x Time interaction ($F_{(3,24)} = 5.80$). Inactive lever pressing was attenuated by approximately 60, 80, and 100 % for 1.0, 5.0, and 10.0 mg/kg SB206, respectively, in the first 15 min of testing (data not shown). While the percentage change in inactive lever pressing appears dramatic, this only corresponds to approximately 2, 3, and 4 fewer inactive lever presses by 1.0, 5.0, and 10.0 mg/kg, respectively.

Similar to the effects seen with SDZ, 3.0 mg/kg of SB242 (also a 5-HT_{2C} receptor antagonist) had no effect on active (Figure 4) or inactive (data not shown) lever pressing when administered alone. Pretreatment with 3.0 mg/kg SB242 15 min prior to administration of 10.0




Figure 4 Attenuation of active lever pressing by SB206 was not reversed by the 5-HT_{2C} antagonist SB242. Shown are active lever presses in 15 min intervals for the first 30 min of cue reactivity (CR) testing from rats that completed interaction studies with 3.0 mg/kg SB 242084 (SB242) vs. 10.0 mg/kg SB 206553 (SB206; n = 11) or 5.0 mg/kg SB206 (n = 9; one statistical outlier removed). Rats were tested after a pretreatment of vehicle or 3.0 mg/kg SB242, and vehicle, 5.0, or 10.0 mg/kg SB206. The x-axis legend refers to time elapsed after treatment and immediately placing the subject into the operant chamber; data represent total active lever presses between 30–45 min and 45–60 min after treatment. **(A)** 5-HT_{2C} antagonism with 3.0 mg/kg SB242 had no effect on CR when administered alone and did not block the effects of 10.0 mg/kg SB206. SB206 significantly decreased active lever presses in the first 15 min of testing (**p < 0.001; Newman-Keuls comparing veh/veh vs. 10.0 mg/kg SB206) and significance was retained even following pretreatment with SB242 (***p < 0.001; comparing veh/veh vs. SB242/SB206). **(B)** 5-HT_{2C} antagonism with 3.0 mg/kg SB242 had no effect on active lever pressing when administered alone and did not block the effects of 5.0 mg/kg SB206. SB206 significantly decreased active lever presses in the first 15 min of testing (**p < 0.01; Newman-Keuls comparing veh/veh vs. 5.0 mg/kg SB206) and significance was retained following pretreatment with SB242 (***p < 0.001 comparing veh/veh vs. SB242/SB206).

induced reinstatement of cocaine-seeking [5,7]. However, because the SB242 did not prevent SB206-induced decreases in lever pressing, it is unclear whether SB206 effects are 5-HT_{2C} dependent.

Normal exploratory and meth-evoked motor activity

To assess the ability for the 5-HT_{2C} ligands to impact non-operant-related motor activity, SDZ and SB206 were tested for motor effects in the absence and presence of meth. In the absence of meth, neither the SDZ (1.0 mg/kg) nor SB206 (5.0 or 10.0 mg/kg) altered motor activity (Table 2). SDZ (1.0 mg/kg) had no effect on meth-evoked motor activity, (Table 3). Likewise, the lower dose of 5.0 mg/kg SB206, which was sufficient to attenuate meth-seeking, had no effect on meth-evoked

or 5.0 mg/kg of SB206 however, did not attenuate the SB206-induced suppression of active (Figure 4) and inactive (data not shown) lever pressing. Two-way rmANOVA revealed a significant Treatment effect ($F_{(3,40)} = 6.04$), Time effect ($F_{(1,40)} = 21.97$), and Treatment x Time interaction ($F_{(3,40)} = 7.36$) comparing the effects of 3.0 mg/kg SB242 and 10.0 mg/kg SB206 as well as a significant Treatment effect ($F_{(3,32)} = 4.90$), Time effect ($F_{(1,32)} = 36.71$), and Treatment x Time interaction ($F_{(3,32)} = 5.151$) comparing the effects of 3.0 mg/kg SB242 and 5.0 mg/kg SB206. Our antagonist data are consistent with the inability of 5-HT_{2C} antagonism to alter cue-

Table 2 Effects of SDZ Ser 082 and SB 206553 on motor activity

| | Horizontal Activity | Total Distance | Vertical Activity |
|------------------|---------------------|----------------|-------------------|
| Vehicle | 727 ± 85 | 189 ± 66 | 46 ± 10 |
| 1.0 mg/kg SDZ | 572 ± 164 | 182 ± 65 | 88 ± 32 |
| Vehicle | 634 ± 66 | 186 ± 63 | 37 ± 14 |
| 5.0 mg/kg SB206 | 564 ± 132 | 107 ± 34 | 9 ± 3 |
| Vehicle | 379 ± 97 | 82 ± 42 | 32 ± 11 |
| 10.0 mg/kg SB206 | 453 ± 41 | 132 ± 30 | 11 ± 5 |

Rats that had undergone repeated cue reactivity testing were withdrawn from methamphetamine and assessed for motor effects after pretreatment with 1.0 mg/kg SDZ Ser 082 (SDZ; 5-HT_{2C} receptor antagonist), 5.0 mg/kg SB 206553 (SB206; putative 5-HT_{2C} receptor inverse agonist), 10.0 mg/kg SB206 and respective vehicles. Serotonin_{2C} receptor antagonism and inverse agonism had no effect on motor activity.

Table 3 Meth-evoked motor activity: SDZ Ser 082 and SB 206553 effects

| | Horizontal Activity | Total Distance | Vertical Activity | Stereotypy Count |
|------------------|---------------------|----------------|-------------------|------------------|
| Vehicle | 8797 ± 738 | 3832 ± 336 | 1142 ± 122 | 477 ± 33 |
| 1.0 mg/kg SDZ | 8598 ± 858 | 3723 ± 392 | 996 ± 190 | 460 ± 29 |
| Vehicle | 7448 ± 1025 | 3113 ± 444 | 797 ± 141 | 446 ± 29 |
| 5.0 mg/kg SB206 | 9067 ± 711 | 4326 ± 572 | 594 ± 120 | 471 ± 18 |
| Vehicle | 8429 ± 1133 | 3705 ± 692 | 1124 ± 201 | 471 ± 26 |
| 10.0 mg/kg SB206 | 6448 ± 1276 | 2245 ± 499 | 398 ± 91** | 394 ± 58 |

Rats that had undergone repeated cue reactivity testing were withdrawn from methamphetamine and assessed for methamphetamine-evoked (1.0 mg/kg) motor effects after a 30 min pretreatment with 1.0 mg/kg SDZ Ser 082 (SDZ; 5-HT_{2C} receptor antagonist), 5.0 mg/kg SB 206553 (SB206; putative 5-HT_{2C} receptor inverse agonist), 10.0 mg/kg SB206 and respective vehicles. Serotonin_{2C} receptor antagonism had no effect on methamphetamine-evoked motor activity whereas inverse agonism with 10.0 mg/kg SB206 significantly attenuated vertical activity (paired *t*-test; ***p* < 0.01).

motor activity (Table 3). The higher dose (10.0 mg/kg) only significantly attenuated meth-evoked vertical activity.

Discussion

SB 206553 attenuates methamphetamine-seeking

Results from the current study are the first to investigate the role of 5-HT_{2C} receptor activity in meth-mediated behaviors and the first to test SB206 in a rodent model of addiction. We revealed that the putative inverse agonist robustly and dose-dependently attenuated meth-seeking without impacting exploratory motor activity at 5.0 mg/kg and 10.0 mg/kg. The 5-HT_{2C} antagonists had no effect on either meth-seeking or meth-induced motor activity. These results suggest that our prior work with mirtazapine may reflect the ability of mirtazapine to act as an inverse agonist, not as a 5-HT_{2C} antagonist.

The finding that the antagonist SB242 did not attenuate the effects of the inverse agonist SB206 was unexpected, and raised the question regarding the receptor selectivity of SB206. SB206 has high affinity for both the 5-HT_{2C} and 5-HT_{2B} receptors (*p*K_i = 8.5 and *p*K_i = 8.26, respectively in HEK-293 or CHO-K1 cells expressing human recombinant 5-HT_{2C} or 5-HT_{2B} receptors; Table 1) [45] with 100 fold or greater selectivity over other receptor targets including 5-HT_{2A} receptors [46]. In the current study, SB206 may antagonize 5-HT_{2B} receptors, which are known to regulate behavioral and DA-enhancing effects of amphetamine [47]. However, the increase in accumbal DA seen with 5.0 mg/kg SB206 is not altered with coadministration of the 5-HT_{2B} antagonist LY 266097 indicating that 5.0 mg/kg SB206 does not substantially antagonize 5-HT_{2B} receptors [47].

Antagonism of 5-HT_{2A} receptors are a third 5-HT receptor that may be contributing to our findings. 5-HT_{2A} antagonism with M100907 attenuates cue-induced reinstatement of cocaine-seeking [48] and meth-seeking (unpublished data). As mentioned above, SB206 demonstrates 100 fold selectivity over the 5-HT_{2A} receptor. 5-HT_{2A} antagonism by SB206, particularly at 10.0 mg/kg, is plausible but unlikely at 1.0 and 5.0 mg/kg, for

amphetamine-evoked DA concentrations in the nucleus accumbens and striatum are attenuated by 5-HT_{2A} antagonism but 5.0 mg/kg SB206 has no effect [49] suggesting that at this dose, SB206 does not act on 5-HT_{2A} receptors. Additionally, 5.0 mg/kg SB206 and 5-HT_{2A} antagonism with SR 46349B have oppositional effects demonstrated by striatal [¹¹C]raclopride binding lending further evidence to the selectivity of SB206 for 5-HT_{2C} over 5-HT_{2A} receptors [50]. Taken together, the available *in vivo* data suggests that SB206 does not antagonize 5-HT_{2A} receptors at 5.0 mg/kg.

SB206 may also act as a positive allosteric modulator of the α7 nicotinic acetylcholine receptor with an EC₅₀ of 1.5 μM for potentiation of responses evoked in GH4C1 cells by EC₂₀ nicotine [51]. Dunlop et al., find that 3, 10, and 30 mg/kg SB206 reverses MK-801-induced deficits in prepulse inhibition, a classical model of schizophrenia, but only confirm actions as a positive allosteric modulator using a nicotinic antagonist against 10.0 mg/kg SB206. It is unclear whether lower doses, such as 1.0 and 5.0 mg/kg SB206, would also act as a positive allosteric modulator at α7 nicotinic receptors. Consistent with our findings, antagonism of α7 nicotinic receptors in the ventral tegmental area attenuates the reward-facilitating effect of cocaine [52] whereas nicotine administration attenuates reinstatement of meth-seeking [53]. The role of α7 nicotinic receptors and in particular, positive allosteric modulation of this receptor, in meth abuse is understudied. Future investigation are needed to determine if this receptor is engaged by SB206 in the dose range tested before actions at the α7 receptor can be ruled out.

Debating receptor specificity based on our findings is difficult; however, it is important to note that a prior report showed that 1.0 mg/kg SB242 attenuates 5.0 mg/kg SB206-induced increases in striatal and accumbal DA [8]. Reasons why Spampinato and colleagues were able to antagonize SB206 effects and we were not may relate to the treatment history of rodent subjects. In the aforementioned neurochemical study [8], subjects were stimulant naïve, halothane anesthetized rats. In our

study, rats had an extensive history of meth self-administration. This chronic meth history may have modified the 5-HT_{2C} system, for we have observed a functionally upregulated response to systemic 5-HT_{2A/C} receptor agonism in the ventral pallidum of rats chronically administered meth [54]. Repeated meth administration may result in changes in mRNA editing profiles (and thus, levels of constitutive activity). Proof of concept for this hypothesis is seen with studies on stress, antidepressant, and antipsychotic administration where mRNA editing of the 5-HT_{2C} receptor is altered [55-58]. In addition to changing constitutive activity, mRNA editing of the 5-HT_{2C} receptor also alters the binding affinity of ligands whereby increased editing most often results in decreased agonist affinity for the receptor with enhanced or no effect on antagonists and inverse agonist binding affinity [16-18,20]. It is unknown how editing patterns affect the affinities of SB206 and SB242. Thus, the inability of SB242 to augment the effects of SB206 on CR may reflect changes in affinity for constitutively active isoforms and/or in the allosteric constant (i.e., an index of the concentration of constitutively active *vs.* non-constitutively active receptors). Unfortunately *ex vivo* binding studies, which would greatly facilitate our studies, are hampered by a lack of available tritiated ligands with 5-HT_{2C} specificity. While future studies are needed to explore the mechanism of action for SB206 and determine receptor specificity *in vivo*, we speculate that despite a lack of antagonism by SB242, 5-HT_{2C} receptors may still be involved. Moreover, continued study is necessary to determine the effects of 5-HT_{2C} receptor ligands on neuron function; to provide further insights our laboratory is conducting patch clamp studies in *ex vivo* slice preparations to determine the effects of 5-HT_{2C} ligands, including SB206, on neurophysiology.

SB 206553, but not SDZ Ser 082, attenuates methamphetamine-evoked motor activity

Acute antagonism of 5-HT_{2C} receptors enhances the motor effects of acute cocaine administration; however, this enhancement is lost in rats that are motorically sensitized by repeated injections of cocaine [40]. This is comparable to our current findings wherein rats with a history of meth self-administration, 5-HT_{2C} antagonism did not alter meth-evoked motor activity in rats. A single study testing SB206 on cocaine-evoked motor activity finds an attenuation of cocaine-evoked motor activity by 1.0 mg/kg SB206 but a potentiation with 4.0 mg/kg [59]; however, this was an acute study with a single injection of cocaine making it difficult to compare with the current investigations using meth self-administration. In rats with a history of meth self-administration, only meth-evoked vertical activity was attenuated by 10.0 mg/kg SB206. Both our findings and [59] are in

agreement in that SB206 does not alter normal exploratory motor behavior suggesting that SB206 is not sedating; differences in our study and [59] regarding the ability to augment psychostimulant-evoked motor activity may reflect behavioral differences consequent to meth-induced plasticity. This hypothesis is supported by [40] and our current findings wherein 5-HT_{2C} antagonism fails to augment cocaine-evoked and meth-evoked motor activity in rats with a history of repeated cocaine [40] or meth administration.

SB 206553 decreases inactive lever pressing

Antagonists had no effect, whereas SB206 attenuated inactive lever pressing. Mirtazapine, also an inverse agonist at the 5-HT_{2C} receptor [35,36], similarly attenuates inactive lever pressing in a cue-induced reinstatement paradigm without effecting non-operant related motor activity in a small animal activity chamber in rats with a history of meth self-administration or rotorod performance in naïve rats [37]. Moreover, in naïve rats, administration of 1, 3, and 4 mg/kg SB206 has no effect on motor activity (measured in activity chambers) [59]. A similar pattern is found for GABA_B receptor activation. The GABA_B agonist (CGP44542) decreases inactive lever pressing in an operant task [60], yet has no effect on performance in an intracranial self-stimulation procedure [61]. In summary, as SB206 had no effect on motor activity in rats with a meth history (current report) or in naïve rats [59], and that mirtazapine, also an inverse agonist at 5-HT_{2C} receptors [35,36], does not impair motor activity in naïve rats [37], we considered that decreased lever pressing of both active and inactive levers seen in the current study was not a consequence of decreased motor activity. Accordingly, we propose that SB206 blunted the *salience* of the levers, both active and inactive, rather than impairing the ability of the rats to perform the operant task. This interpretation is supported by our prior demonstration that significance of meth-associated context cues are reduced by mirtazapine [31]. Additional studies aimed at exploring the effect of SB206 on non-drug reinforced behaviors may advance this hypothesis; however, as 5-HT_{2C} receptors regulate food intake [6,62,63] as well as intracranial self-stimulation behavior [64,65], such evaluations may be challenging to conduct and difficult to interpret.

Conclusions

The presented findings are the first to explore the effects of 5-HT_{2C} antagonists and putative inverse agonist, SB206, on meth-seeking and meth-induced motor activity. Serotonin_{2C} antagonism had no effect on meth-seeking or motor activity (in the presence or absence of meth); in contrast, SB206, attenuated meth-seeking. This effect was selective, as SB206 agonist did

not alter exploratory behavior, and only the higher dose reduced one motor parameter evoked by meth (i.e., vertical activity). There is a growing interest in assessing the role of constitutively active 5-HT_{2C} receptors *in vivo*. It is currently unclear whether SB206 elicits the observed effects *via* 5-HT_{2C} receptor inverse agonism; nonetheless, based on the dramatic reduction in seeking behavior, further investigation of this compound is merited as well as continued study of 5-HT_{2C} regulation of psychostimulant-mediated behaviors.

Abbreviations

meth: Methamphetamine; DA: Dopamine; SDZ: SDZ Ser 082; SB206: SB 206553; SB242: SB 242084; 5-HT: Serotonin; CR: Cue reactivity.

Competing interests

The authors have no competing interests or financial disclosures to report.

Authors' contributions

Experiments and data analysis were conducted by SMG. Experimental design and manuscript writing were done by SMG and TCN. All authors read and approved the final manuscript.

Acknowledgements

Work supported by USPHSG DA015760 to TCN and DA024923 to SMG and TCN and the Daniel F. and Ada L. Rice Foundation.

Received: 10 January 2012 Accepted: 1 June 2012

Published: 14 June 2012

References

- Higgins GA, Fletcher PJ: Serotonin and drug reward: focus on 5-HT_{2C} receptors. *Eur J Pharmacol* 2003, **480**:151–162.
- Hayes DJ, Greenshaw AJ: 5-HT receptors and reward-related behaviour: a review. *Neurosci Biobehav Rev* 2011, **35**:1419–1449.
- Fletcher PJ, Grottick AJ, Higgins GA: Differential effects of the 5-HT(2A) receptor antagonist M100,907 and the 5-HT(2 C) receptor antagonist SB242,084 on cocaine-induced locomotor activity, cocaine self-administration and cocaine-induced reinstatement of responding. *Neuropsychopharmacology* 2002, **27**:576–586.
- Burbassi S, Cervo L: Stimulation of serotonin_{2C} receptors influences cocaine-seeking behavior in response to drug-associated stimuli in rats. *Psychopharmacology (Berl)* 2008, **196**:15–27.
- Neisewander JL, Acosta JL: Stimulation of 5-HT_{2C} receptors attenuates cue and cocaine-primed reinstatement of cocaine-seeking behavior in rats. *Behav Pharmacol* 2007, **18**:791–800.
- Cunningham KA, Fox RG, Anastasio NC, Bubar MJ, Stutz SJ, Moeller FG, Gilbertson SR, Rosenzweig-Lipson S: Selective serotonin 5-HT(2C) receptor activation suppresses the reinforcing efficacy of cocaine and sucrose but differentially affects the incentive-salience value of cocaine- vs. sucrose-associated cues. *Neuropharmacology* 2011, **61**:513–523.
- Filip M: Role of serotonin (5-HT)₂ receptors in cocaine self-administration and seeking behavior in rats. *Pharmacol Rep* 2005, **57**:35–46.
- De Deurwaerdere P, Navailles S, Berg KA, Clarke WP, Spampinato U: Constitutive activity of the serotonin_{2C} receptor inhibits *in vivo* dopamine release in the rat striatum and nucleus accumbens. *J Neurosci* 2004, **24**:3235–3241.
- Leggio GM, Cathala A, Neny M, Rouge-Pont F, Drago F, Piazza PV, Spampinato U: *In vivo* evidence that constitutive activity of serotonin(2C) receptors in the medial prefrontal cortex participates in the control of dopamine release in the rat nucleus accumbens: differential effects of inverse agonist versus antagonist. *J Neurochem* 2009.
- Navailles S, Moison D, Ryczko D, Spampinato U: Region-dependent regulation of mesoaccumbens dopamine neurons *in vivo* by the constitutive activity of central serotonin_{2C} receptors. *J Neurochem* 2006, **99**:1311–1319.
- Burns CM, Chu H, Rueter SM, Hutchinson LK, Canton H, Sanders-Bush E, Emeson RB: Regulation of serotonin-2 C receptor G-protein coupling by RNA editing. *Nature* 1997, **387**:303–308.
- Niswender CM, Sanders-Bush E, Emeson RB: Identification and characterization of RNA editing events within the 5-HT_{2C} receptor. *Ann N Y Acad Sci* 1998, **861**:38–48.
- Sanders-Bush E, Fentress H, Hazelwood L: Serotonin 5-HT₂ receptors: molecular and genomic diversity. *Mol Interv* 2003, **3**:319–330.
- Werry TD, Loiacono R, Sexton PM, Christopoulos A: RNA editing of the serotonin 5HT_{2C} receptor and its effects on cell signalling, pharmacology and brain function. *Pharmacol Ther* 2008, **119**:7–23.
- Gardiner K, Du Y: A-to-I editing of the 5HT_{2C} receptor and behaviour. *Brief Funct Genomic Proteomic* 2006, **5**:37–42.
- Fitzgerald LW, Iyer G, Conklin DS, Krause CM, Marshall A, Patterson JP, Tran DP, Jonak GJ, Hartig PR: Messenger RNA editing of the human serotonin 5-HT_{2C} receptor. *Neuropsychopharmacology* 1999, **21**:825–905.
- Herrick-Davis K, Grinde E, Niswender CM: Serotonin 5-HT_{2C} receptor RNA editing alters receptor basal activity: implications for serotonergic signal transduction. *J Neurochem* 1999, **73**:1711–1717.
- Niswender CM, Copeland SC, Herrick-Davis K, Emeson RB, Sanders-Bush E: RNA editing of the human serotonin 5-hydroxytryptamine 2 C receptor silences constitutive activity. *J Biol Chem* 1999, **274**:9472–9478.
- Berg KA, Cropper JD, Niswender CM, Sanders-Bush E, Emeson RB, Clarke WP: RNA-editing of the 5-HT(2 C) receptor alters agonist-receptor-effector coupling specificity. *Br J Pharmacol* 2001, **134**:386–392.
- Berg KA, Dunlop J, Sanchez T, Silva M, Clarke WP: A conservative, single-amino acid substitution in the second cytoplasmic domain of the human Serotonin_{2C} receptor alters both ligand-dependent and -independent receptor signaling. *J Pharmacol Exp Ther* 2008, **324**:1084–1092.
- Kenakin T: Drug efficacy at G protein-coupled receptors. *Annu Rev Pharmacol Toxicol* 2002, **42**:349–379.
- Kenakin T: Principles: receptor theory in pharmacology. *Trends Pharmacol Sci* 2004, **25**:186–192.
- Willins DL, Meltzer HY: Serotonin 5-HT_{2C} agonists selectively inhibit morphine-induced dopamine efflux in the nucleus accumbens. *Brain Research* 1998, **781**:291–299.
- Lucas G, Spampinato U: Role of striatal serotonin_{2A} and serotonin_{2C} receptor subtypes in the control of *in vivo* dopamine outflow in the rat striatum. *J Neurochem* 2000, **74**:693–701.
- Gobert A, Rivet JM, Lejeune F, Newman-Tancredi A, Dhumeau-Auclair A, Nicolas JP, Cistarelli L, Melon C, Millan MJ: Serotonin(2C) receptors tonically suppress the activity of mesocortical dopaminergic and adrenergic, but not serotonergic, pathways: a combined dialysis and electrophysiological analysis in the rat. *Synapse* 2000, **36**:205–221.
- Di Matteo V, Di Giovanni G, Di Mascio M, Esposito E: Selective blockade of serotonin_{2C/2B} receptors enhances dopamine release in the rat nucleus accumbens. *Neuropharmacology* 1998, **37**:265–272.
- Di Matteo V, Di Mascio M, Di Giovanni G, Esposito E: Acute administration of amitriptyline and mianserin increases dopamine release in the rat nucleus accumbens: possible involvement of serotonin_{2C} receptors. *Psychopharmacology* 2000, **150**:45–51.
- McDaid J, Tedford CE, Mackie AR, Dallimore JE, Mickiewicz AL, Shen F, Angle JM, Napier TC: Nullifying drug-induced sensitization: behavioral and electrophysiological evaluations of dopaminergic and serotonergic ligands in methamphetamine-sensitized rats. *Drug Alcohol Depend* 2007, **86**:55–66.
- Voigt RM, Napier TC: Context-dependent effects of a single administration of mirtazapine on the expression of methamphetamine-induced conditioned place preference. *Front Behav Neurosci* 2011, **5**:92.
- Voigt RM, Mickiewicz AL, Napier TC: Repeated mirtazapine nullifies the maintenance of previously established methamphetamine-induced conditioned place preference in rats. *Behav Brain Res* 2011, **225**:91–96.
- Herrold AA, Shen F, Graham MP, Harper LK, Specio SE, Tedford CE, Napier TC: Mirtazapine treatment after conditioning with methamphetamine alters subsequent expression of place preference. *Drug Alcohol Depend* 2009, **99**:231–239.
- Kooyman AR, Zwart R, Vanderheijden PM, Van Hooff JA, Vijverberg HP: Interaction between enantiomers of mianserin and ORG3770 at 5-HT₃ receptors in cultured mouse neuroblastoma cells. *Neuropharmacology* 1994, **33**:501–507.

33. de Boer TH, Maura G, Raiteri M, de Vos CJ, Wieringa J, Pinder RM: **Neurochemical and autonomic pharmacological profiles of the 6-aza-analogue of mianserin, Org 3770 and its enantiomers.** *Neuropharmacology* 1988, **27**:399–408.
34. Millan MJ, Gobert A, Rivet JM, Adhumeau-Auclair A, Cussac D, Newman-Tancredi A, Dekeyne A, Nicolas JP, Lejeune F: **Mirtazapine enhances frontocortical dopaminergic and corticolimbic adrenergic, but not serotonergic, transmission by blockade of alpha2- adrenergic and serotonin2C receptors: a comparison with citalopram.** *Eur J Neurosci* 2000, **12**:1079–1095.
35. Chanrion B, la Mannoury CC, Gavarini S, Seimandi M, Vincent L, Pujol JF, Bockaert J, Marin P, Millan MJ: **Inverse agonist and neutral antagonist actions of antidepressants at recombinant and native 5-hydroxytryptamine2C receptors: differential modulation of cell surface expression and signal transduction.** *Mol Pharmacol* 2008, **73**:748–757.
36. Labasque M, Meffre J, Carrat G, Becamel C, Bockaert J, Marin P: **Constitutive Activity of Serotonin2C Receptors at G Protein-Independent Signaling: Modulation by RNA Editing and Antidepressants.** *Mol Pharmacol* 2010.
37. Graves SM, Napier TC: **Mirtazapine alters cue-associated methamphetamine seeking in rats.** *Biol Psychiatry* 2011, **69**:275–281.
38. Acosta JJ, Thiel KJ, Sanabria F, Browning JR, Neisewander JL: **Effect of schedule of reinforcement on cue-elicited reinstatement of cocaine-seeking behavior.** *Behav Pharmacol* 2008, **19**:129–136.
39. Filip M, Bubar MJ, Cunningham KA: **Contribution of serotonin (5-HT) 5-HT2 receptor subtypes to the discriminative stimulus effects of cocaine in rats.** *Psychopharmacology (Berl)* 2006, **183**:482–489.
40. Filip M, Bubar MJ, Cunningham KA: **Contribution of serotonin (5-hydroxytryptamine; 5-HT) 5-HT2 receptor subtypes to the hyperlocomotor effects of cocaine: acute and chronic pharmacological analyses.** *J Pharmacol Exp Ther* 2004, **310**:1246–1254.
41. Kourrich S, Thomas MJ: **Similar neurons, opposite adaptations: psychostimulant experience differentially alters firing properties in accumbens core versus shell.** *J Neurosci* 2009, **29**:12275–12283.
42. Rademacher DJ, Napier TC, Meredith GE: **Context modulates the expression of conditioned motor sensitization, cellular activation and synaptophysin immunoreactivity.** *Eur J Neurosci* 2007, **26**:2661–2668.
43. Rademacher DJ, Kovacs B, Shen F, Napier TC, Meredith GE: **The neural substrates of amphetamine conditioned place preference: implications for the formation of conditioned stimulus-reward associations.** *Eur J Neurosci* 2006, **24**:2089–2097.
44. McDaid J, Graham MP, Napier TC: **Methamphetamine-induced sensitization differentially alters pCREB and ΔFosB throughout the limbic circuit of the mammalian brain.** *Mol Pharmacol* 2006, **70**:2064–2074.
45. Knight AR, Misra A, Quirk K, Benwell K, Revell D, Kennett G, Bickerdike M: **Pharmacological characterisation of the agonist radioligand binding site of 5-HT(2A), 5-HT(2B) and 5-HT(2 C) receptors.** *Naunyn Schmiedebergs Arch Pharmacol* 2004, **370**:114–123.
46. Kennett GA, Wood MD, Bright F, Cilia J, Piper DC, Gager T, Thomas D, Baxter GS, Forbes IT, Ham P, et al: **In vitro and in vivo profile of SB 206553, a potent 5-HT2C/5-HT2B receptor antagonist with anxiolytic-like properties.** *Br J Pharmacol* 1996, **117**:427–434.
47. Auclair AL, Cathala A, Sarrazin F, Depoortere R, Piazza PV, Newman-Tancredi A, Spampinato U: **The central serotonin receptor: a new pharmacological target to modulate the mesoaccumbens dopaminergic pathway activity.** *J Neurochem* 2010, **114**:1323–1332.
48. Nic Dhonnchadha BA, Fox RG, Stutz SJ, Rice KC, Cunningham KA: **Blockade of the serotonin 5-HT2a receptor suppresses cue-evoked reinstatement of cocaine-seeking behavior in a rat self-administration model.** *Behav Neurosci* 2009, **123**:382–396.
49. Porras G, Di MV, Fracasso C, Lucas G, De DP, Caccia S, Esposito E, Spampinato U: **5-HT2A and 5-HT2C/2B receptor subtypes modulate dopamine release induced in vivo by amphetamine and morphine in both the rat nucleus accumbens and striatum.** *Neuropsychopharmacology* 2002, **26**:311–324.
50. Egerton A, Ahmad R, Hirani E, Grasby PM: **Modulation of striatal dopamine release by 5-HT2A and 5-HT2C receptor antagonists: [11 C]raclopride PET studies in the rat.** *Psychopharmacology (Berl)* 2008, **200**:487–496.
51. Dunlop J, Lock T, Jow B, Sztzia F, Grauer S, Jow F, Kramer A, Bowlby MR, Randall A, Kowal D, et al: **Old and new pharmacology: positive allosteric modulation of the alpha7 nicotinic acetylcholine receptor by the 5-hydroxytryptamine(2B/C) receptor antagonist SB-206553 (3,5-dihydro-5-methyl-N-3-pyridinylbenzo[1,2-b:4,5-b']di pyrrole-1(2 H)-carboxamide).** *J Pharmacol Exp Ther* 2009, **328**:766–776.
52. Panagis G, Kastellakis A, Spyraiki C, Nomikos G: **Effects of methyllycaconitine (MLA), an alpha 7 nicotinic receptor antagonist, on nicotine- and cocaine-induced potentiation of brain stimulation reward.** *Psychopharmacology (Berl)* 2000, **149**:388–396.
53. Hiranita T, Nawata Y, Sakimura K, Anggadiredja K, Yamamoto T: **Suppression of methamphetamine-seeking behavior by nicotinic agonists.** *Proc Natl Acad Sci U S A* 2006, **103**:8523–8527.
54. Napier TC, Istre ED: **Methamphetamine-induced sensitization includes a functional upregulation of ventral pallidal 5-HT2A/2 C receptors.** *Synapse* 2008, **62**:14–21.
55. Huang XF, Tan YY, Huang X, Wang Q: **Effect of chronic treatment with clozapine and haloperidol on 5-HT(2A and 2 C) receptor mRNA expression in the rat brain.** *Neurosci Res* 2007, **59**:314–321.
56. Du Y, Stasko M, Costa AC, Davisson MT, Gardiner KJ: **Editing of the serotonin 2 C receptor pre-mRNA: Effects of the Morris Water Maze.** *Gene* 2007, **391**:186–197.
57. Iwamoto K, Nakatani N, Bundo M, Yoshikawa T, Kato T: **Altered RNA editing of serotonin 2 C receptor in a rat model of depression.** *Neurosci Res* 2005, **53**:69–76.
58. Englander MT, Dulawa SC, Bhansali P, Schmauss C: **How stress and fluoxetine modulate serotonin 2 C receptor pre-mRNA editing.** *J Neurosci* 2005, **25**:648–651.
59. McCreary AC, Cunningham KA: **Effects of the 5-HT2C/2B antagonist SB 206553 on hyperactivity induced by cocaine.** *Neuropsychopharmacology* 1999, **20**:556–564.
60. Paterson NE, Froestl W, Markou A: **The GABAB receptor agonists baclofen and CGP44532 decreased nicotine self-administration in the rat.** *Psychopharmacology (Berl)* 2004, **172**:179–186.
61. Paterson NE, Markou A: **The metabotropic glutamate receptor 5 antagonist MPEP decreased break points for nicotine, cocaine and food in rats.** *Psychopharmacology* 2004.
62. Fletcher PJ, Sinyard J, Higgins GA: **Genetic and pharmacological evidence that 5-HT2C receptor activation, but not inhibition, affects motivation to feed under a progressive ratio schedule of reinforcement.** *Pharmacol Biochem Behav* 2010, **97**:170–178.
63. Fletcher PJ, Tampakeras M, Sinyard J, Slassi A, Isaac M, Higgins GA: **Characterizing the effects of 5-HT(2 C) receptor ligands on motor activity and feeding behaviour in 5-HT(2 C) receptor knockout mice.** *Neuropharmacology* 2009, **57**:259–267.
64. Hayes DJ, Clements R, Greenshaw AJ: **Effects of systemic and intra-nucleus accumbens 5-HT2C receptor compounds on ventral tegmental area self-stimulation thresholds in rats.** *Psychopharmacology (Berl)* 2009, **203**:579–588.
65. Katsidoni V, Apazoglou K, Panagis G: **Role of serotonin 5-HT2A and 5-HT2C receptors on brain stimulation reward and the reward-facilitating effect of cocaine.** *Psychopharmacology (Berl)* 2011, **213**:337–354.

doi:10.1186/1471-2202-13-65

Cite this article as: Graves and Napier: SB 206553, a putative 5-HT_{2C} inverse agonist, attenuates methamphetamine-seeking in rats. *BMC Neuroscience* 2012 **13**:65.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

