

Epigenetic regulation of prostate cancer

Suyin P. Chin · Joanne L. Dickinson · Adele F. Holloway

Received: 1 February 2011 / Accepted: 23 May 2011 / Published online: 14 June 2011
© Springer-Verlag 2011

Abstract Prostate cancer is a commonly diagnosed cancer in men and a leading cause of cancer deaths. Whilst the underlying mechanisms leading to prostate cancer are still to be determined, it is evident that both genetic and epigenetic changes contribute to the development and progression of this disease. Epigenetic changes involving DNA hypo- and hypermethylation, altered histone modifications and more recently changes in microRNA expression have been detected at a range of genes associated with prostate cancer. Furthermore, there is evidence that particular epigenetic changes are associated with different stages of the disease. Whilst early detection can lead to effective treatment, and androgen deprivation therapy has a high response rate, many tumours develop towards hormone-refractory prostate cancer, for which there is no successful treatment. Reliable markers for early detection and more effective treatment strategies are, therefore, needed. Consequently, there is a considerable interest in the potential of epigenetic changes as markers or targets for therapy in prostate cancer. Epigenetic modifiers that demethylate DNA and inhibit histone deacetylases have recently been explored to reactivate silenced gene expression in cancer. However, further understanding of the mechanisms and the effects of chromatin modulation in prostate cancer are required. In this review, we examine the current literature on epigenetic changes associated with prostate cancer and discuss the potential use of epigenetic modifiers for treatment of this disease.

Keywords Prostate cancer · Epigenetics · DNA methylation · Histone acetylation · MicroRNA

S. P. Chin · J. L. Dickinson · A. F. Holloway (✉)
Menzies Research Institute Tasmania, University of Tasmania,
Private Bag 23,
Hobart, Tasmania 7000, Australia
e-mail: A.F.Holloway@utas.edu.au

Introduction

Prostate cancer is the most commonly diagnosed cancer for men living in developed countries (other than non-melanoma skin cancer). According to Cancer Research UK, an estimated 913,000 men worldwide were diagnosed in developed countries in 2008. According to the US National Cancer Institute, it was estimated that almost 217,730 men would be diagnosed in the USA alone in 2010 and more than 32,050 would die as a direct result of the disease. The use of prostate-specific antigen (PSA) as a screening tool has allowed the detection of prostate cancer in the early stages whilst it is still locally confined. Whilst more than 70% of diagnosed cases now survive beyond 5 years, this cancer is still associated with significant mortality and morbidity. Metastatic prostate tumours are responsible for the majority of deaths associated with this cancer. The most frequent site of prostate cancer metastasis is to bone; over 80% of men who die of prostate cancer have metastatic bony lesions (Bubendorf et al. 2000). In terms of current treatments for prostate cancer, we are still unable to identify with certainty those tumours requiring aggressive and immediate intervention (associated with considerable morbidity) and those where a “watchful-waiting” approach may be more appropriate. Thus, identification of markers predicting tumour behaviour has become of intense interest to researchers working to discover new prognostic and diagnostic markers and new targets for treatment.

Prostatic intraepithelial neoplasia (PIN) and, in particular, high-grade PIN has been identified as precancerous lesions most likely leading to prostatic carcinoma. On the prostate morphological spectrum, PIN refers to precancerous lesions involving cell proliferation within prostatic ducts, ductules and acini (De Marzo et al. 2004). PIN is believed to pre-date carcinoma by 10 or more years. High-grade PIN is considered clinically significant as men with high-grade

PIN have up to 50% chance of subsequently developing prostate cancer (Lee et al. 2011). A second morphological abnormality termed atypical small acinar proliferation has also been associated with increased risk of diagnosis with prostate cancer in subsequent biopsies. Prostate tumours are most commonly graded using the Gleason score, determined by the histological characteristics of the glandular architecture within the tumour.

It is clear that epigenetic changes within a cell play a significant role in the development and progression of cancer (Esteller 2008; Jones and Baylin 2007) and, as in most other human cancers, prostate cancer development and progression appears to involve an interplay between both genetic and epigenetic changes. There is now considerable evidence that changes in gene expression which involve epigenetic alterations may be an important factor in prostate cancer progression, and development of panels of epigenetically modified genes as markers of disease progression is of considerable topical interest.

Epigenetic mechanisms

Epigenetic alterations are heritable changes in gene expression that occur without changes in DNA sequence, with the broadest definition including all factors other than DNA sequence changes that heritably influence gene expression (Berger et al. 2009). Whilst the best described of these mechanisms is DNA methylation, other epigenetic mechanisms include physical and chemical changes to chromatin and regulation of gene expression by microRNAs (miRNAs).

DNA methylation plays an important role in DNA repair, recombination and replication, as well as regulating gene activity (see Fig. 1). DNA methylation involves the addition of a methyl group to the 5'-carbon of cytosine in CpG dinucleotide sequences, catalysed by a family DNA methyltransferases (DNMTs). CpG-rich regions, known as CpG islands are commonly found associated with the 5'-region of vertebrate genes (Gardiner-Garden and Frommer 1987) and are generally protected from methylation (Bird 2002). For many years, CpG islands have been implicated in gene regulation with their methylation strongly correlated with gene silencing (Illingworth and Bird 2009). DNA methylation can regulate gene activity via two mechanisms. Firstly, methylation of CpG dinucleotides within transcription factor binding sites can inhibit transcription factor binding and, therefore, directly influence gene activity (Hark et al. 2000). Secondly, methylated CpG dinucleotides act as binding sites for methyl CpG binding proteins, which are associated with other factors such as histone deacetylases, involved in establishing repressive chromatin structures (Jones et al. 1998; Nan et al. 1998).

Changes in DNA methylation patterns have been linked with cancer for many years now (Jones and Baylin 2007). However, the methylation changes are complex, with both hypomethylation and hypermethylation occurring in cancer cells. Aberrant DNA hypermethylation occurs when there is a gain of DNA methylation at regions that are normally unmethylated and when this occurs at gene promoters can lead to gene inactivation. Localised hypermethylation of gene promoters has been reported in virtually all types of cancers, including prostate cancer. On the other hand, DNA hypomethylation is the demethylation of normally methylated DNA and can lead to chromosomal instability and activation of proto-oncogenes (Dunn 2003; Eden et al. 2003; Sharma et al. 2010). Both global and gene-specific hypomethylation events have also been implicated in prostate cancer.

Whilst alterations in DNA methylation have long been linked to cancer, there is also mounting evidence that other epigenetic changes, such as changes to chromatin composition or structure, contribute to cancer. Within the eukaryotic nucleus, DNA is assembled into chromatin, the basic unit of which is the nucleosome. Nucleosomes are composed of approximately 147 base pairs of DNA wrapped around an octamer of core histone proteins, containing two each of histones H2A, H2B, H3 and H4 (Kornberg and Lorch 1999). The N termini of the core histones protrude from the nucleosome and are subjected to a range of covalent modifications, catalysed by various histone-modifying enzymes. At least ten different histone modifications have now been reported, including acetylation, methylation, phosphorylation and ubiquitination (Kouzarides 2007; Gardner et al. 2011). Each of these modifications affects the chromatin structure and function in a different way by either disrupting chromatin contacts or affecting the recruitment of other proteins to the chromatin (Kouzarides 2007). Acetylation of lysine residues in histone H3 and H4 (H3Ac and H4Ac) is in general associated with transcriptional activity, whereas histone methylation is associated with transcriptional activation or repression depending on the site of modification and the number of methyl groups added (see Fig. 1). For example, histone H3 lysine 4 (H3K4) methylation is generally associated with transcriptional activation, whereas histone H3 lysine 9 (H3K9) and histone H3 lysine 27 (H3K27) di- and tri-methylation is generally associated with transcriptional repression (Kouzarides 2007). As with DNA methylation, there is an increasing body of evidence that changes in histone modifications due to aberrant activity or mis-targeting of chromatin-modifying enzymes is involved in carcinogenesis (Hake et al. 2004).

More recently, miRNAs have been identified as important regulators of gene expression. miRNAs are endogenous small non-coding, single-stranded RNAs, 18–24 bases




Fig. 1 A model depicting epigenetic modifications associated with different transcriptional states and some of the inhibitors that target epigenetic modifiers and, therefore, influence transcriptional activity

long, which regulate gene expression by affecting the stability or the translation efficiency of target mRNAs (Pang et al. 2010). The miRNA binds to complementary sequences within the 3' mRNA tails, although usually only with partial base pair complementarity. When miRNAs bind to mRNA with partial complementarity, they function as translational repressors whereas perfect complementarity induces degradation of the target mRNA (He and Hannon 2004; Porkka et al. 2007). Recent studies have found that aberrations in miRNA expression are associated with various human cancers including prostate cancer. About half of the known miRNA genes are located at cancer-related genomic regions (Calin et al. 2004), and in cancer, miRNAs can function as either oncogenes or tumour suppressor genes (Hayashita et al. 2005; He et al. 2005). One miRNA can target hundreds of mRNAs and, therefore, alterations in a single miRNA can have dramatic effects on cell biology. Our understanding of the function and regulatory mechanisms of miRNAs has expanded enormously over recent years; however, we still have limited capacity to identify targets of miRNAs and have much to learn about their involvement in cellular processes and disease.

Epigenetic alterations in prostate cancer progression

DNA hypomethylation

There is considerable evidence that changes in methylation patterns occur in prostate cancer with DNA hypomethylation in tumour samples first documented more than 20 years ago. In 1987, Bedford and van Helden analysed DNA 5'-methylcytosine content in human prostate samples and reported a correlation between global hypomethylation and development of benign prostatic hyperplasia (BPH) and metastatic tumours (Bedford and van Helden 1987). Comparison of DNA methylation levels in tumour and normal prostate tissue by immunohistochemistry similarly detected global hypomethylation in prostate cancer (Brothman et al. 2005). In keeping with this, Santourlidis et al. (1999) examined methylation of LINE-1 repetitive sequences in prostate adenocarcinomas and found that LINE-1 methylation tended to decrease with tumour stage. A further study analysing tumour samples similarly demonstrated an association between DNA hypomethylation, tumour state and metastasis, with extensive hypomethylation of LINE-1 observed in 64% of cases with lymph node or distant

metastases but only in 21% of cases without detectable metastases (Schulz et al. 2002). All cases of hormone-refractory locally recurrent tumours displayed extensive LINE-1 hypomethylation. The study by Schulz et al. (2002) also found a strong association between LINE-1 DNA hypomethylation and chromosomal alterations, in support of the notion that DNA hypomethylation increases genomic instability (Schulz et al. 2002). Most prostate cancer samples with prominent DNA hypomethylation also exhibited a large number of chromosomal alterations and vice versa. These findings were supported by a further study which found increased prevalence of LINE-1 hypomethylation in later-stage prostate cancers and lymph-node-positive prostate cancers (Flori et al. 2004), in contrast to gene-specific hypermethylation events which did not correlate with tumour stage, being present in both early and late-stage prostate cancer. These data, therefore, suggest that hypomethylation is a later event in prostate cancer progression compared with gene-specific hypermethylation. Consistent with this, a recent study examined LINE-1 methylation in primary prostate cancers compared with normal prostate tissues and found a significantly higher level of LINE-1 hypomethylation in metastatic prostate cancer tissues, suggesting that global hypomethylation occurs late in prostate cancer progression, particularly at the metastatic disease stage (Yegnasubramanian et al. 2008).

Whilst global DNA hypomethylation can contribute to cancer by promoting genomic instability, the hypomethylation of individual gene promoters can also contribute to cancer development and progression by directing aberrant gene expression. Hypomethylation of a number of genes has been linked to prostate cancer. For example, a study by Yegnasubramanian et al. (2008), found that CpG islands associated with a class of cancer testis antigen genes were hypomethylated in prostate cancer, correlating with their overexpression in primary prostate cancers, and more so in metastatic prostate cancer. Similarly, hypomethylation of the promoter of the heparanase gene has been reported in prostate cancer, compared with BPH (Ogishima et al. 2005). This correlates with increased mRNA expression of the heparanase gene, which is associated with tumour invasion and metastasis (Vlodavsky et al. 1988; Hulett et al. 1999). Further, upregulation of urokinase expression in invasive prostate cancer cell lines correlates with hypomethylation of the promoter (Pakneshan et al. 2003). Upregulation of cytochrome P450 1B1 (CYP1B1) mRNA and protein levels in prostate cancer compared with BPH tissue was also found to correlate with hypomethylation of the regulatory regions of this gene (Tokizane et al. 2005). Other genes such as wingless-related MMTV integration site 5A (WNT5A), S100 calcium-binding protein P and cysteine-rich protein 1 were also found to be hypomethylated in primary prostate cancer tissues. These genes are similarly

associated with tumourigenesis and metastasis (Dissanayake et al. 2007; Wang et al. 2006). Further investigation of the WNT5A gene revealed three CpG sites in the promoter region which were consistently methylated in a normal prostate cell line and normal prostate tissues but not in a prostate cancer cell line and primary prostate cancer tissues (Wang et al. 2007). Therefore, hypomethylation and consequent upregulation of genes involved in metastasis and cell invasion may be an important factor in prostate cancer progression. In support of this, treatment of the PC3 prostate cancer cell line with reagents that prevent DNA hypomethylation has an inhibitory effect on cell invasion *in vitro* and tumour growth *in vivo* (Shukeir et al. 2006).

DNA hypermethylation

DNA hypermethylation is the most commonly reported epigenetic alteration observed in prostate cancer. Many genes have been identified as aberrantly hypermethylated in prostate cancer and these genes include tumour suppressor genes, DNA damage repair genes, hormonal response genes and genes involved in cell cycle control, tumour cell invasion and metastasis (Phé et al. 2010). Hypermethylation of DNA can lead to inappropriate gene silencing, disrupting gene function and, thus, contributing to tumour initiation, progression and metastasis (Li et al. 2004). This review highlights some of the best described and most frequently hypermethylated genes in prostate cancer.

The most frequently reported hypermethylated gene in prostate cancer is the π -class glutathione-*S*-transferase (GSTP1) gene. GSTP1 is an enzyme involved in the metabolism, detoxification and elimination of reactive chemical compounds and in this way protects cells from DNA damage and cancer initiation (Lee 2007). Hypermethylation of the GSTP1 gene in prostate cancer was first reported by Lee et al. (1994) who found hypermethylation of the gene in all 20 human prostate cancer tissue samples examined, but not in normal tissues or BPH. Following this, GSTP1 gene expression was analysed in 60 high-grade PIN samples, with all samples showing a loss of GSTP1 gene expression (Brooks et al. 1998). Further investigation found that this loss of GSTP1 expression was due to hypermethylation of the gene promoter. A large number of studies have now reported GSTP1 hypermethylation in prostate cancer samples, with methylation detected in up to 90% of samples, suggesting GSTP1 hypermethylation is a common epigenetic alteration in prostate cancer (Goessl et al. 2001; Chu et al. 2002; Jeronimo et al. 2002; Harden et al. 2003; Nakayama et al. 2003; Bastian et al. 2004).

Promoter hypermethylation of genes involved in tumour invasion has also been observed in prostate cancer. The maintenance of the normal cell architecture is regulated by

the cadherin-catenin system. Loss of E-cadherin (CDH1) gene expression is associated with the transition from adenoma to carcinoma and the acquisition of metastatic potential (Perl et al. 1998). CDH1 hypermethylation was observed in prostate cancer cell lines and treatment with the demethylating agent, 5-aza-2'-deoxycytidine restored CDH1 mRNA and protein expression in the cell lines, suggesting that promoter hypermethylation was responsible for CDH1 silencing in these cells (Graff et al. 1995). Furthermore, Kallakury et al. (2001) reported that the CDH1 gene promoter was methylated in eight out of ten prostate cancer tissues examined. Interestingly, the degree of CDH1 promoter methylation correlated with the pathological stage of the prostate tumour tissues, with CDH1 promoter methylation occurring in 30% of low-grade prostate cancer tissues, but increasing to 70% in high-grade tumours (Li et al. 2001). These results suggest that methylation of the CDH1 promoter is associated with prostate tumour progression.

CD44, which is a cell surface glycoprotein involved in cell matrix adhesion and signal transduction is also silenced in prostate cancer by methylation of the gene promoter. Lou et al. (1999) examined methylation levels of the CD44 gene in 84 matched normal and prostate cancer samples and found hypermethylation of CD44 in 31 out of 40 of the primary prostate cancer samples. Further investigation by Verkaik et al. (1999) showed that in CD44-negative prostate cancer cell lines (LNCaP and PC346C) the CD44 promoter was hypermethylated compared with CD44-positive prostate cancer cell lines (Du145, PC3 and TSU). A further study of CD44 silencing was conducted on human tissue samples and demonstrated that nine out of 11 lymph node metastases of prostate cancer displayed CD44 gene promoter methylation. These data, therefore, suggest that hypermethylation of the CD44 promoter resulting in downregulation of CD44 gene expression may be involved in prostate cancer progression and metastasis (Verkaik et al. 2000).

DNA methylation is also involved in regulation of the androgen receptor (AR). AR is activated by androgen, which plays a critical role in the development, growth and maintenance of the prostate (Jenster 1999). In the initial stages, prostate cancer is androgen dependent, but eventually becomes androgen independent, due to the loss of AR expression (Jarrard et al. 1998; Tekur et al. 2001; Takahashi et al. 2002; Suzuki et al. 2003). Data from a number of studies suggest that this loss of AR expression is at least partly due to hypermethylation of the AR gene promoter. Jarrard et al. (1998) found AR promoter hypermethylation in AR-negative prostate cancer cell lines (DU145, DuPro, TSU-PR1 and PPC1) whereas the promoter was unmethylated in AR-positive cell lines (LNCaP and PC3). Expression of the AR gene was restored in the AR-negative cell lines by treatment with the demethylating agent 5-aza-2'-deoxycytidine,

suggesting that promoter methylation was responsible for AR gene silencing in the AR-negative cell lines. Further, Suzuki et al. (2003) reported that promoter hypermethylation of AR leading to loss of AR expression occurs in 30% of hormone-refractory prostate cancers, suggesting that DNA hypermethylation contributes to loss of AR expression in at least some prostate cancers.

The silencing of cell cycle regulation genes by DNA hypermethylation has also been observed in prostate cancer. The Ras-associated domain family 1A gene (RASSF1A) is highly methylated in several human cancers, including prostate cancer (Liu et al. 2002; Kang et al. 2004; Pfeifer and Dammann 2005; Dammann et al. 2005; Aitchison et al. 2007; Hesson et al. 2007). Liu et al. (2002) examined methylation of the RASSF1A promoter in primary prostate tumours and reported methylation of the RASSF1A promoter in over 70% of the tumours. Further investigation found a correlation between the frequency of methylation and the Gleason score of the tumour, with highly aggressive tumours displaying more frequent DNA methylation compared with less aggressive tumours. Similarly, Kang et al. (2004) reported methylation of a number of genes, including RASSF1A in prostate cancer and PIN samples, with more frequent methylation correlating with higher PSA levels and Gleason score.

Silencing of tumour suppressor genes by DNA methylation is also often observed in prostate cancer. DNA hypermethylation of the adenomatous polyposis coli (APC) gene in prostate cancer individuals was observed in a study by Rosenbaum et al. (2005), which examined promoter methylation of a number of genes. Hypermethylation of APC alone, and hypermethylation of APC and the cell cycle regulation gene cyclin D2 in combination were found to be significant predictors of prostate cancer progression. In keeping with this, Henrique et al. (2007) analysed a small panel of gene promoters in prostate biopsy samples and similarly found that hypermethylation of APC is an independent predictor of poor prognosis in prostate cancer. Subsequent studies have similarly found APC hypermethylation to be a predictor of prostate cancer progression (Liu et al. 2011b; Richiardi et al. 2009).

Histone modifications

Compared with DNA methylation the involvement of histone modifications in prostate cancer is relatively poorly understood, even though these two epigenetic mechanisms are closely related (Sharma et al. 2010). However, some data relating particular histone modifications to prostate cancer have emerged in recent years. Seligson et al. (2005) analysed a range of histone modifications by immunohistochemistry including acetylated histone H3 lysine 9, acetylated histone H3 lysine 18 (H3K18Ac), H4K12Ac,

di-methylated H4 arginine 3 (H4R3me2) and di-methylated H3 lysine 4 (H3K4me2) in 183 primary prostate cancer tissue samples. In the individuals with low-grade tumours, the study found two subgroups with different risks of tumour recurrence based on the presence of similar combinations of global histone modifications. Individuals with a lower risk of tumour recurrence were those who were above the 60th percentile staining for H3K4me2 or above the 35th percentile staining for H3K18Ac and H3K4me2. However, these histone modification patterns did not correlate with the Gleason score. In contrast, a more recent study of primary and metastatic prostate cancer samples found that high global levels of H3K18Ac and H3K4me2 correlated with a threefold increased risk of prostate cancer recurrence (Bianco-Miotto et al. 2010). A further study showed that global levels of H3K4me1, H3K9me2, H3K9me3, H3Ac and H4Ac were significantly reduced in prostate cancer compared with BPH and normal prostate tissue (Ellinger et al. 2010), with H3Ac and H3K9me2 in particular discriminating between the malignant and non-malignant samples. They also found that individuals with high H3K4me1 levels were more likely to experience recurrence of the prostate cancer and thus suggested that analysis of H3K4me1 may provide predictive information regarding likelihood of tumour recurrence.

In addition to the studies outlined above that have documented changes in histone modifications associated with prostate cancer, there is accumulating evidence that expression of histone-modifying enzymes is altered in prostate cancer. Bianco-Miotto et al. (2010) identified a candidate gene signature consisting of six genes encoding epigenetic modifiers, including both DNA methyltransferases and histone methyltransferases (HMTs) that was associated with prostate cancer progression. In addition, a number of studies have documented alterations in the HMT EZH2, which is responsible for the repressive H3K27me3 modification, in prostate cancer. EZH2 is upregulated in hormone-refractory metastatic prostate cancer (Varambally et al. 2002). Overexpression of EZH2 in prostate cancer cell lines increases the invasive characteristics of the cells, whilst knockdown of EZH2 decreases the proliferative capacity of the cells, and more so in hormone-independent cell lines (Varambally et al. 2002; Karanikolas et al. 2010). Furthermore, microarray analysis of metastatic prostate cancer tissue identified a group of EZH2 repressed genes that were associated with prostate cancer progression (Yu et al. 2007).

A number of studies have also implicated lysine-specific demethylase 1 (LSD1) in prostate cancer. LSD1 was originally identified as a H3K4 demethylating enzyme (HDM). This enzyme was thought to function as a transcriptional co-repressor by reversing the H3K4me modifications associated with transcriptional activation

(Shi et al. 2004). However, genome-wide studies of LSD1 homologues in yeast suggest that these enzymes can act as both co-activators and co-repressors by targeting H3K4 or H3K9, respectively (Opel et al. 2007). In keeping with this, LSD1 has been demonstrated to form ligand-dependent, chromatin-associated complexes with AR, stimulating AR-dependent transcription (Metzger et al. 2005). In this case, LSD1 acts by demethylating histone H3K9, thus activating AR target genes by removing repressive H3K9me modifications. This study found that LSD1 and AR were co-localised in both normal prostate and prostate cancer. A further study by Kahl et al. (2006) correlated the expression patterns of AR, LSD1 and the AR co-activator FHL2 (four and a half LIM-domain protein 2) with Gleason score, Gleason grade and p53 expression in 153 prostate tumour samples with relapse after radical prostatectomy. The study found that increased LSD1, nuclear expression of FHL2, high Gleason score and grade and high levels of p53 were strongly associated with relapse during follow-up. In addition, upregulation of both LSD1 mRNA and protein levels was associated with high risk of relapse. Whilst LSD1 demethylates H3K9me1 and H3K9me2, a further study found that the Jumonji C domain containing protein JMJD2C can demethylate H3K9me3 (Wissmann et al. 2007). JMJD2C associates with AR and LSD1 in prostate cells, and this complex acts to demethylate H3K9me3 and increase AR-dependent gene transcription. A second Jumonji protein, JHDM2A, which also demethylates H3K9me1/2 has similarly been found to stimulate transcription of AR dependent genes (Yamane et al. 2006). Furthermore, Gaughan et al. (2011) showed that AR interacts with and is methylated by the HMT enzyme SET9. SET9 was originally identified as an enzyme responsible for catalysing the activating H3K4me1 modification, and, therefore, was associated with transcriptional activation (Nishioka et al. 2002). However, Gaughan et al. (2011) found that by methylating AR SET9 increased the transcriptional activation of AR itself. In doing so, SET9 was found to have pro-proliferative and anti-apoptotic activity in the AR-dependent LNCaP prostate cancer cell line.

Whilst the studies outlined above have highlighted differences in histone modifications and histone modifiers correlating with prostate cancer stage or recurrence, it should be noted that many of these studies have described global changes and further studies are, therefore, needed to investigate how these global changes relate to gene-specific loci, particularly at genes that have already been implicated in prostate cancer.

miRNA regulation

To date, there have been more than 50 miRNAs found to be deregulated in prostate cancer using sensitive miRNA array studies on primary tumour samples, normal prostate

samples and prostate cancer cell lines (Volinia et al. 2006; Tang et al. 2011; Wach et al. 2011). A miRNA expression profiling study using six prostate cancer cell lines (PC3, DU145, LNCaP, 22Rv1, VCaP and LAPC4), nine prostate xenograft samples and 12 clinical prostate tissue samples revealed a miRNA signature of prostate cancer (Porkka et al. 2007). A total of 51 miRNAs were differentially expressed in benign tumours and carcinoma tumours, of which 37 miRNAs were downregulated whilst 14 miRNAs were upregulated. Based on miRNA expression, the authors were able to accurately separate the carcinomas from the BPH samples and also classify the tumours according to their androgen dependence. This suggests that androgens might regulate, or be regulated by, some miRNAs. A further study investigating both miRNA and mRNA profiles in prostate cancer, similarly found evidence for androgen regulation of miRNAs (Ambs et al. 2008). This study reported deregulation of both miRNAs and miRNA processing pathways in prostate cancer as well as demonstrating regulation of a set of cancer-associated genes by miRNAs. Similarly, analysis of miRNA profiles in androgen-dependent and androgen-independent cell lines by next generation sequencing revealed a cohort of differentially expressed miRNAs, many of which target cell communication and signal transduction pathways (Xu et al. 2010).

Cao et al. (2010) identified miRNA-101 (miR-101) as an androgen responsive miRNA in prostate cancer cell lines, with the androgen agonist R1881 stimulating miR-101 expression. They also demonstrated regulation of EZH2 by miR-101. As mentioned earlier, EZH2 is overexpressed in prostate cancer and, therefore, acts as an oncogene. miR-101 was found to inhibit endogenous EZH2 in PC3, DU145 and LNCaP prostate cancer cell lines and decrease invasion and migration of DU145 and LNCaP cells, respectively. These findings are in keeping with another study which analysed miR-101 expression in human prostate tumour samples and found that miR-101 expression decreases during cancer progression, correlating with the increase in EZH2 expression (Varambally et al. 2008). Therefore, these studies strongly support the notion that miR-101 functions as a tumour suppressor by targeting EZH2.

Whilst some miRNAs are clearly regulated by androgens, recently, a study identified a number of miRNAs which target the 3'UTR of the AR mRNA and thus, regulate AR levels (Ostling et al. 2011). The study screened 1,129 miRNAs in a panel of prostate cancer cell lines quantifying changes in AR levels using protein lysate microarrays. The assays identified 13 miRNAs that inhibit AR expression. Further analysis on 47 clinical prostate tumour samples confirmed a negative correlation between expression of two of these, miR-34a and miR-34c and AR expression levels. miR-34a was also found to positively correlate with expression of the tumour suppressor p53,

which inhibits expression of genes involved in cell cycle and apoptosis (Fujita et al. 2008). miR-34a expression was found to be higher in p53 positive cell lines (PrEC and LNCaP) than in p53-defective cell lines (PC3 and DU145). This study further identified the histone deacetylase SIRT1, which had been implicated in diverse cellular processes including apoptosis as a direct target of miR-34a. Downregulation of SIRT1 was observed in PC3 and DU145 cells compared with LNCaP cells, and SIRT1 mRNA levels also decreased in response to ectopic expression of miR-34a. Ectopic miR-34a expression also resulted in cell cycle arrest at G1 phase and inhibited growth of PC3 cells. In addition, ectopic miR-34a expression attenuated chemoresistance to the anticancer drug camptothecin (a topoisomerase I inhibitor) by inducing apoptosis. These data suggest that modulation of miR-34a activity may represent a novel approach for treating malignant prostate cancers with aberrant p53 function.

Interestingly, CD44, which has also been implicated in prostate cancer, as noted above, has also recently been identified as a target of miR-34a (Liu et al. 2011a). CD44⁺ prostate cancer cell populations have previously been shown to possess cancer stem cell (CSC) properties, which includes an enhanced tumour initiating capacity, self-renewal capabilities, and resistance to conventional anticancer therapeutics. Thus, CSCs have been implicated in tumour recurrence and metastasis (Visvader and Lindeman 2008; Patrawala et al. 2006, 2007). A comparison of 137 miRNAs in CD44⁺ versus CD44⁻ prostate cancer cells purified from xenograft and primary tumours, suggest that miR-34a is underexpressed in the CD44⁺ populations. Furthermore, overexpression of miR-34a in CD44⁺ prostate cells and CD44 knockdown in CD44⁻ prostate cells inhibited clonogenic expansion, tumour regeneration, and metastasis. In a prostate-tumour-bearing mouse model, systemically delivered miR-34a-inhibited prostate cancer metastasis and extended survival of tumour bearing mice. This emerging role of miR-34a in regulating CSCs further supports the idea that miR-34a represents a potential therapeutic target in prostate cancer.

Several miRNAs have also been identified as oncogenic miRNAs in prostate cancer. Analysis of miR221 and miR-222 expression in three prostate cancer cell lines (PC3, 22RV1 and LNCaP) showed that miR-221 and miR-222 are overexpressed in the more aggressive prostate tumour line, PC3, compared with the less aggressive cell lines, LNCaP and 22RV1 (Galardi et al. 2007). Galardi et al. (2007), identified p27 mRNA as a target of miR-221 and miR-222, with overexpression of miR-221/222 resulting in p27 downregulation in LNCaP cells and increased cell proliferation. Deregulation of cyclin-dependent kinase inhibitor p27 has been implicated in prostate cancer progression and is a well-established marker of poor prognosis with decreased p27 expression associated with high tumour grade and poor prognosis of prostate cancer (Macri and

Loda 1998; Tsihlias et al. 1998; Yang et al. 1998). Furthermore in the study, knockdown of miR-221 and miR-222 increased p27 expression in PC3 cells and reduced their clonogenicity *in vitro*. Following this, a study was conducted to determine the role of miR-221 and miR-222 *in vivo* (Mercatelli et al. 2008). LNCaP cells stably transfected with miR-221 were subcutaneously injected into SCID mice. A significant increase in growth was observed in tumours overexpressing miR-221 correlating with a reduction in p27 expression. Furthermore, SCID mice injected with PC3 cells in which miR-221/222 were knocked down, displayed significantly reduced tumour growth associated with increased p27 expression.

There is currently intense interest in the action of miRNAs and their important role in cancer progression is increasingly recognised. It is clear that there are global changes to miRNAs in prostate cancer, and a number of these miRNAs, including those highlighted above, are already being recognised as potential biomarkers or therapeutic targets. However, we still have much to learn about the mechanism of action and target selection of miRNAs, and further research is clearly needed in order to understand the role of these molecules in prostate and other cancers.

Epigenetic modifications as potential prostate cancer biomarkers

The current challenge for prostate cancer is its early detection, as this provides the best chance of successful treatment. Since the introduction of PSA testing, there has been a significant increase in the diagnosis of prostate cancer. Although PSA is the best tumour marker available today, PSA screening has limited accuracy. Whilst a PSA level of more than 4.0 ng/mL has predictive value for the diagnosis of prostate cancer (Thompson et al. 2004), the Prostate Cancer Prevention Trial conducted by Thompson et al. (2004) showed that 15% of men with a serum PSA value less than 4.0 ng/mL have prostate cancer, with 25% of these men having cancer of Gleason score 7 or above. Further, the majority of those with elevated PSA do not have prostate cancer (Villers et al. 1997). Therefore, there is an urgent need for more accurate biomarkers that can identify men at high prostate cancer risk and therefore improve current treatments.

Recently, increased understanding of the involvement of epigenetic alterations in the early stages of cancer has led to the investigation of epigenetic changes, particularly DNA hypermethylation as potential biomarkers. DNA hypermethylation is a relatively stable modification and has therefore been investigated as a potential biomarker, as reviewed in Phé et al. (2010). Furthermore, DNA methylation can be easily

detected using highly sensitive PCR-based methods and bisulphite sequencing. In addition, as outlined above, many studies suggest that DNA hypermethylation is an important mechanism leading to gene loss of function in prostate cancer (Li et al. 2004; Whang et al. 1998; Nakayama et al. 2003). Further, recent studies suggest that DNA hypermethylation is involved in an early event in carcinogenesis (Feinberg et al. 2006) and, therefore, DNA hypermethylation may prove to be a useful tumour biomarker.

As described earlier, GSTP1 hypermethylation is a common occurrence in prostate cancer and has therefore been investigated as a potential biomarker. Jeronimo et al. (2004) examined promoter hypermethylation at GSTP1 and several other gene promoters in 118 prostate cancer samples, 38 paired high-grade PIN and 30 BPH, in order to determine the sensitivity of using GSTP1 hypermethylation as a biomarker to detect prostate cancer. Interestingly, the results showed that the frequency of methylation was different between the three groups with no GSTP1 promoter hypermethylation observed in BPH. This suggests that GSTP1 promoter hypermethylation is a sensitive prostate cancer marker with the ability to distinguish between the different stages of prostate cancer. In addition, the study found that using a combination of GSTP1 and APC methylation levels increased the detection rate of prostate cancer to almost 94%. Similarly, another study of promoter hypermethylation of GSTP1, APC and prostaglandin-endoperoxide synthase 2 (or cyclo-oxygenase 2 (COX-2)) reported that GSTP1 methylation analysis in combination with either or both of the other genes provided good correlation with tumour stage and Gleason score but not PSA levels (Bastian et al. 2007). Therefore, analysis of a combination of hypermethylated genes may provide increased sensitivity and specificity of prostate cancer detection and therefore may have potential as a biomarker in prostate cancer diagnosis.

In addition, a number of studies have demonstrated that DNA methylation may be detectable in urine samples, raising the possibility of detecting epigenetic changes using non-invasive screening methods. Cairns et al. (2001) demonstrated that GSTP1 hypermethylation can be detected in voided urine using methylation-specific PCR. Whilst GSTP1 hypermethylation was only detected in one third of the urine samples of which it was found in matched prostate cancer samples, it was not detected in any of the samples in which prostate tissue was negative for GSTP1 hypermethylation. This demonstrates that molecular diagnosis of prostate cancer by analysis of DNA methylation in urine samples may be feasible. In support of this, Goessl et al. (2001) analysed urine sediments from patients with BPH, PIN and various stages of prostate cancer and reported detection of GSTP1 promoter methylation with 98% specificity and 73% sensitivity. A further study investigated

methylation of a panel of genes in post-prostate massage-voided urine (Roupret et al. 2007). The study involved 95 individuals with early prostate cancer and found that a four-gene combination of GSTP1, RASSF1a, retinoic acid receptor β 2 (RAR β 2) and APC was the best discriminator of malignant compared with non-malignant samples with a sensitivity of 86% and an accuracy of 89%. These studies, therefore, suggest that post-prostate massage-voided urine may be a realistic alternative as a non-invasive method of prostate cancer screening, with multiple gene panels providing a better discrimination of malignant from non-malignant cells.

Another potential cancer biomarker is the cell-free circulating DNA in serum/plasma. Recent studies have shown that individuals with prostate cancer have higher levels of cell-free DNA and that the DNA fragmentation pattern from healthy individuals is different to that from individuals with benign prostate cancer (Ellinger et al. 2008b; Jung et al. 2004; Papadopoulou et al. 2004; Altamari et al. 2008). The cell-free circulating DNA seems to be released from solid malignant tumours that undergo cellular necrosis and apoptosis (Jahr et al. 2001). Allen et al. (2004) showed that individuals with prostate cancer and PIN had significantly higher levels of cell-free circulating DNA compared with those with BPH. However, this increase in cell-free DNA appears to be a common feature of cancer, as it is also observed in breast (Huang et al. 2006), cervical (Trejo-Becerril et al. 2003), lung (Sozzi et al. 2001) and bladder (Ellinger et al. 2008a) cancer. Therefore, detecting methylation of a panel of prostate cancer-specific genes in the cell-free circulating DNA may be useful to allow a more specific diagnosis. Cell-free DNA hypermethylation at GSTP1 detected in individuals with prostate cancer, in a number of studies have shown a diagnostic specificity of nearly 100% (Jeronimo et al. 2002; Bastian et al. 2005; Papadopoulou et al. 2004; Reibenwein et al. 2007; Chuang et al. 2007; Altamari et al. 2008). However, detection of methylation at a panel of prostate-cancer-specific genes (GSTP1, TIGI, COX-2 and RPRM) increased the sensitivity of diagnosis and the ability to predict prognosis of prostate cancer even further (Ellinger et al. 2008c). Therefore, cell-free DNA may be a promising non-invasive biomarker for prostate cancer.

Finally, very recently miRNAs have been suggested as potential biomarkers in prostate cancer. An expression profile study consisting of six prostate cancer-related miRNAs (miR-141, miR-16, miR-101, miR-34c, miR-21 and miR-125b) was conducted on human prostate samples and prostate cell lines of both benign and malignant tumours (Hao et al. 2011). These miRNAs were selected on the basis of their association with prostate cancer-related proteins, PSA, Bcl-2, COX-2, and p53. However, expression levels of all six miRNAs were different across cell lines,

corresponding xenograft tumours and human specimens. Interestingly, all the miRNAs showed a significantly different expression pattern when compared between the benign cell line, BPH1, and the tumourigenic subline, BPH1^{CAFTD}, with miR-21 and miR-125b highly expressed in BPH1^{CAFTD}, and lower expression of miR-16, miR-34c, miR-101 and miR-141 detected in BPH1 cells. Therefore, profiles of these miRNAs clearly distinguished the benign BPH1 from the tumourigenic BPH1^{CAFTD} cells. Analysis of human tissue samples showed a twofold upregulation of miR-21 and miR-141 in tumour samples compared with BPH samples, suggesting that miR-21 and miR-141 might be a potential biomarker useful for distinguishing tumours from benign samples. In addition, miRNAs have been detected in human plasma in a stable form and, thus, may be an attractive non-invasive biomarker (Mitchell et al. 2008). A panel of six miRNAs (miR-100, miR-125b, miR-141, miR-143, miR-205 and miR-296) were also analysed in serum samples from 25 individuals with metastatic prostate cancer and 25 healthy individuals as controls. Of all the miRNAs examined, miR-141 showed a significantly higher expression in individuals diagnosed with prostate cancer compared with individuals without prostate cancer, correlating well with the previous study by Hao et al. (2011). Therefore, miR-141 appears to be a potential non-invasive biomarker that can distinguish prostate cancer from benign or healthy controls.

Pharmacological reversal of epigenetic modifications in prostate cancer

As highlighted above, changes in epigenetic modifications are increasingly being associated with prostate cancer. Unlike genetic mutations, epigenetic alterations are potentially reversible and a range of pharmacological reagents that reverse epigenetic modifications have been characterised in recent years (see Fig. 1). This review highlights several of these agents and their potential use in the treatment of prostate cancer.

Pharmacological reversal of DNA hypermethylation in prostate cancer

DNA methylation can be reversed by DNA methyltransferase inhibitors (DNMTi), as indicated in Fig. 1. The azanucleoside, 5-azacytidine was the first DNMTi developed and was shown to demethylate DNA in human cell lines (Jones and Taylor 1980). This modified nucleoside is incorporated into DNA during replication and is recognised by the DNMT enzymes. However, the reaction is not able to be completed and 5-azacytidine traps the DNMT enzyme on the DNA (Santi et al. 1984). Consequently, the DNA remains unmethylated and the DNMT enzyme is rapidly depleted

within the cell and thus, genomic DNA is demethylated following DNA replication. However, 5-azacytidine is a ribose nucleoside which must be converted to a deoxyribonucleoside triphosphate before incorporation into DNA, and therefore, some of the 5-azacytidine is incorporated into RNA, and thus, it has cellular effects unrelated to DNA demethylation (Cihak 1974). Consequently, the analogue 5-aza-2'-deoxycytidine (decitabine) was developed, which is directly incorporated into DNA.

There have been a plethora of studies demonstrating the ability of the nucleoside inhibitors to re-express genes silenced by DNA methylation in prostate cancer cell lines, and due to space limitations only a few of these studies will be highlighted here. In 1997, two independent studies examined the effect of DNMTi on cyclin-dependent kinase 2a (CDKN2) gene expression (Chi et al. 1997; Jarrard et al. 1997). CDKN2 is a tumour suppressor gene which is suggested to play an important role in prostate cancer progression (Cairns et al. 1995). Re-expression of the CDKN2 gene in non-expressing prostate cancer cell lines (PC3, PPC1 and TSU-PR1) was observed following treatment with 5-azacytidine and 5-aza-2'-deoxycytidine. In both studies, morphological changes were observed in the cell lines following treatment with the demethylating agents. Further, both studies reported a decrease in growth rate in the PC3 cell lines treated with the DNMTi. Subsequently, Whang et al. (1998) examined the candidate tumour suppressor gene PTEN/MMAC1. This gene is implicated in multiple tumours, including prostate cancer. In this study, a LuCaP-35 xenograft was explanted from an animal, then propagated in culture and treated with 5-aza-2'-deoxycytidine. The treatment successfully restored PTEN/MMAC1 gene expression in the LuCaP-35 prostate cancer xenograft cells. Additionally, given the prevalence of GSTP1 hypermethylation in prostate cancer, as discussed earlier, demethylation of the GSTP1 gene following DNMTi treatment has been studied extensively. For example, studies conducted by Lin et al. (2001b) and Singal et al. (2001) demonstrated rescue of GSTP1 expression in LNCaP cells using 5-azacytidine and 5-aza-2'-deoxycytidine, respectively. Similarly, Vidanes et al. (2002) successfully restored expression of GSTP1 in the MDA PCa 2a and MDA PCa 2b prostate cancer cell lines by treatment with 5-azacytidine.

The effects of DNMTi have also been studied using *in vivo* models of prostate cancer, including the transgenic adenocarcinoma of the mouse prostate (TRAMP) mouse model. TRAMP is a prostate cancer mouse model, which closely mirrors the pathogenesis of human prostate cancer. Zorn et al. (2007) conducted a study to determine the effect of 5-azacytidine on pre-existing TRAMP mouse prostate cancers and its ability to prevent androgen-independent prostate cancer. In this study, 5-azacytidine treatment prevented prostate enlargement and prolonged the time

taken to develop androgen-independent status. Further, a recent study by Gravina et al. (2010) suggests 5-azacytidine in combination with anti-androgen therapy may have therapeutic potential in some individuals. This study, in which cells were treated with 5-azacytidine along with the anti-androgen bicalutamide (BCLT), was conducted using both cell lines and an *in vivo* model in which PC3 and 22RV1 cells were subcutaneously injected into the flank of CD1 nude mice. 5-Azacytidine acted synergistically with BCLT in both cases, upregulating cell cycle proteins and increasing the balance of pro-apoptotic compared with anti-apoptotic proteins. In addition, treatment with 5-azacytidine and BCLT in combination significantly repressed tumour growth *in vivo*, suggesting that the combined treatment has an anti-tumour effect. Further investigation of the *in vivo* model also detected upregulation of caspases, which suggests this effect may be mediated by the induction of apoptosis.

5-Azacytidine and 5-aza-2'-deoxycytidine are the most clinically advanced DNMTi and have been examined extensively in clinical trials in a range of cancers, with 5-aza-2'-deoxycytidine receiving US Food and Drug Administration (FDA) approval for the treatment of myelodysplastic syndrome (Kaminskas et al. 2005). However, a small number of clinical trials conducted in individuals diagnosed with prostate cancer suggest these inhibitors may have more limited use in prostate cancer. Although the *in vitro* studies of nucleoside inhibitors have been successful in restoring silenced gene expression in prostate cancer cell lines, as described above, a phase II study of 5-aza-2'-deoxycytidine conducted in 14 men with androgen-independent metastatic prostate cancer had limited success with only two of the 12 individuals displaying stable disease with delayed time to progression (Thibault et al. 1998). This study concluded that whilst 5-aza-2'-deoxycytidine is well tolerated, it has only moderate effects on hormone-independent prostate cancer. Further, although these inhibitors may reverse aberrant DNA hypermethylation, they have previously been found to have side effects such as myelosuppression (Kantarjian et al. 2003; Schrupp et al. 2006; Appleton et al. 2007; Cashen et al. 2008; Batty et al. 2010; Chuang et al. 2010), tumorigenesis (Walker and Nettesheim 1986; Schnekenburger et al. 2011; Hamm et al. 2009) and mutagenesis (Jackson-Grusby et al. 1997; Perry et al. 1992; Sauntharajah et al. 2003; Lavelle et al. 2007), which may limit their usefulness as a therapeutic drug in prostate cancer.

Non-nucleoside DNMTi have also been characterised and their ability to reverse DNA hypermethylation examined in prostate cancer. The non-nucleoside DNMTi procainamide acts mainly by inhibiting DNMT1 (Lee et al. 2005). Procainamide treatment was found to successfully reverse GSTP1 CpG island hypermethylation and restore GSTP1 expression in LNCaP prostate cancer cells (Lin et al. 2001a).

Procainamide was also tested in an *in vivo* model in which LNCaP prostate cancer cells were inoculated into the subcutaneous region of the flanks of athymic mice. Following treatment with procainamide for 2 weeks, the xenograft tumours were excised and found to display increased GSTP1 expression, correlating with a reduction in GSTP1 CpG island methylation. Whilst this study suggests that non-nucleoside DNMTi may effectively reverse methylation-dependent gene silencing in prostate cancer cells, a study by Chuang et al. (2005) which examined the effectiveness of procainamide and two other non-nucleoside demethylating agents; (–)-epigallocatechin-3-gallate and hydralazine, compared with the nucleoside inhibitor 5-aza-2'-deoxycytidine in a range of prostate cancer cell lines, found that 5-aza-2'-deoxycytidine is considerably more effective in demethylating DNA and reactivating genes compared with the non-nucleoside inhibitors.

Whilst agents that inhibit DNA methylation are therefore being actively investigated as potential therapeutic agents in prostate cancer, one issue that also needs to be considered is their potential to hypomethylate and therefore upregulate genes involved in metastasis. As outlined above, hypomethylation of a number of genes has been linked to prostate cancer progression and metastasis (Shukeir et al. 2006) and the effect of DNMTi on these genes also needs to be considered.

Use of histone deacetylase inhibitors in prostate cancer

Two types of enzymes are involved in determining the acetylation and deacetylation of the histone tails: histone acetyltransferases and histone deacetylases (HDACs). Studies on prostate cancer samples, cell lines and prostate cancer mouse models have shown that HDAC activity is upregulated in prostate cancer (Halkidou et al. 2004; Waltregny et al. 2004). HDAC complexes can be inhibited by a variety of agents known as HDAC inhibitors (HDACi), as indicated in Fig. 1. These fall into a number of categories, namely, hydroxamic acids, aliphatic acids, cyclic peptides, and benzamides. A range of these HDACi have been examined for their potential use in prostate cancer, and the most extensively examined and those with the most promising therapeutic potential will be highlighted here.

Trichostatin A (TSA) was one of the first natural compounds found to be a potent HDACi. A study by Huang et al. (1999) treated the prostate cancer cell line PC3 with the HDACi sodium butyrate and TSA. They found that PC3 cells undergo differentiation and apoptosis following sodium butyrate treatment. Another study conducted by Rashid et al. (2001), also treated a range of prostate cancer cell lines with sodium butyrate and TSA. This study investigated whether HDACi treatment modulated responses of prostate cancer cells to 1 alpha, 25-dihydroxyvitamin D(3). 1 Alpha, 25-dihydroxyvitamin D

(3) is involved in regulating the growth and differentiation of the prostate (Schwartz and Hulka 1990; Konety et al. 1996), but prostate cancer cells are less sensitive to this hormone (Miller et al. 1995; Krill et al. 1999). They found that sodium butyrate and TSA had little effect on their own at low doses but synergized with 1 alpha, 25-dihydroxyvitamin D(3) to inhibit the growth of the prostate cancer cell lines. These studies suggest that TSA and sodium butyrate inhibit prostate cancer cell growth by inducing apoptosis. However, the anti-proliferative effects of these HDACi are likely to be complicated as Suenaga et al. (2002), have also shown that TSA and sodium butyrate downregulate telomerase activity in both PC3 and LNCaP cell lines.

Suberoylanilide hydroxamic acid (SAHA; Vorinostat) was the first HDACi approved by the US FDA for the treatment of cancer, in particular the treatment of cutaneous T-cell lymphoma. This compound has been found to inhibit proliferation of the prostate cancer cell lines LNCaP, DU-145 and PC3 (Lakshmikanthan et al. 2006; Chinnaiyan et al. 2005; Gediya et al. 2008; Kulp et al. 2006; Marrocco et al. 2007; Schmudde et al. 2008). SAHA has also been tested in animal models of prostate cancer and found to have potential as an anti-tumour drug. SAHA was used to treat mice transplanted with CWR22 human prostate tumours (Butler et al. 2000). At a dose without detectable toxicity, a 97% reduction in tumour growth was detected. These results suggest that SAHA is relatively non-toxic and may be useful in treating prostate cancer. As most metastatic prostate cancer cells tend to be oxidatively stressed (Halliwell 2006), a recent study by Basu et al. (2011), using LNCaP and PC3 prostate cancer cell lines, showed that the response to SAHA was enhanced by combination treatment with an anti-oxidant like vitamin E and SAHA. Qian et al. (2006) also investigated the effect of another HDACi, panobinostat, in a prostate cancer mouse model. A PC3 xenograft mouse model was examined following a daily treatment with panobinostat and this treatment was found to inhibit tumour growth and angiogenesis.

The aliphatic acid, valproic acid (VPA) is another HDACi that has been investigated in both *in vitro* and *in vivo* prostate cancer models. The prostate cancer cell lines DU-145 and PC3 were treated with VPA and their viability and cell cycle status monitored along with expression of cell cycle and proliferation markers (Shabbeer et al. 2007). The anti-proliferative effect of VPA was dose and time dependent and an effect on cell cycle arrest was observed even 24 h after its removal. VPA was then administered in the drinking water of nude mice bearing xenografts of human prostate cancer cell lines, DU-145 and PC3. In this model, VPA was able to induce growth arrest, cell death and senescence. An increase in acetyl H3K9 was detected in VPA-treated xenografts, suggesting that VPA was functioning as an HDACi in this model.

The cyclic peptide HDACi, FK228 (depsipeptide; romidepsin) has also been extensively examined in experimental studies of prostate cancer. Lai et al. (2008) conducted a study to determine the effects FK228 has on prostate cancer cells *in vivo*. 22RV1 cells were subcutaneously inoculated into male non-obese diabetic-severe combined immune-deficient mice and were either left untreated or treated with FK228, given orally three times a week. All the untreated mice had died at 98 days whereas, 61% of the treated mice remained alive. In addition, in untreated mice the human prostate cancer cells metastasized to the lung, whereas FK228-treated mice showed normal lung morphology. These data, therefore, suggest that FK228 is efficient at suppressing prostate cancer metastases. Recently, Zhang et al. (2007) also conducted a study to determine whether FK228 can augment the effect of other chemotherapeutic agents against androgen-independent prostate cancer. They tested the combination of FK228 with docetaxel (a microtubule stabilizer and conventional chemotherapeutic agent), 5-aza-2'-deoxycytidine (the nucleoside DNMTi described above) and genistein (a protein kinase inhibitor) on three androgen-independent prostate cancer cell lines, PC3, DU-145 and C4-2. They found that FK228/docetaxel had greater cytotoxic effects than the other combinations. Furthermore, *in vivo* treatment of nude mice with PC3 tumours with FK228/docetaxel had greater inhibitory effects than the other compounds. Thus, FK228 augments the effect of docetaxel against androgen-independent prostate cancer, providing a promising treatment option.

MS-275 is an orally active synthetic benzamide derivative with HDACi activity. Qian et al. (2005) conducted a study to evaluate the effects of combination therapy of MS-275 with 13-*cis* retinoic acid to reactivate RAR β 2 gene expression in the human prostate cancer cell lines PC3 and DU-145. Retinoids have been used in the treatment of epithelial tumours but their use has been hindered by the development of resistance and is associated with the silencing of RAR β 2 gene expression due to DNA methylation and histone deacetylation. The effects of the combination therapy to reactivate the RAR β 2 gene expression was also examined in nude mice, which were injected with PC3 cells stably transfected with a RAR β 2 promoter-luciferase gene. The combination therapy was able to reactivate endogenous RAR β 2 gene expression by acetylation of the RAR β 2 promoter both *in vitro* and *in vivo*. The combination therapy also resulted in synergistic activation of the luciferase reporter gene both *in vitro* and *in vivo*. In addition, combination therapy restored retinoid sensitivity in prostate cancer cell lines and had a greater inhibitory effect on tumour cell growth compared with treatment with the agents by themselves. Therefore, this study suggests that HDACi and retinoid combination therapy may be a

promising treatment option for retinoid-resistant prostate cancer. Subsequently, Qian et al. (2007) conducted another study to examine the biological effects of MS-275 both *in vitro* (PC3, LNCaP and DU-145) and *in vivo* (PC3, LNCaP and DU-145 injected mice and TRAMP mice). MS-275 significantly upregulated histone H3 acetylation and p21 (a potent cyclin-dependent kinase inhibitor) gene expression in the PC3, LNCaP and DU-145 cell lines. Upregulation of p21 corresponded with growth arrest in PC3 and LNCaP cells and MS-275 treatment induced cell death in DU-145 cells. In the *in vivo* models, MS-275 successfully inhibited the growth of PC3, LNCaP and DU-145 in subcutaneous xenografts. Tumour samples from MS-275 treated mice had increased histone acetylation levels and increased expression of p21. Long-term treatment of MS-275 in TRAMP mice slowed progression of prostate carcinomas with a significant reduction in cell proliferation. Therefore, these data strongly suggest that MS-275 is a promising HDACi for the treatment of prostate cancer.

Clinical trials

There have been a number of clinical trials on HDACi conducted in individuals diagnosed with prostate cancer either alone or in combination with other anti-cancer agents. A phase I clinical study with oral panobinostat alone or in combination with docetaxel in castration-resistant prostate cancer was conducted in 16 individuals (Rathkopf et al. 2010). In this trial, a total of eight individuals were treated with oral panobinostat alone and another eight individuals were treated with oral panobinostat in combination with intravenous docetaxel. All individuals treated with panobinostat alone developed progressive disease. However, five individuals treated with the combination therapy achieved a partial response and this included an individual that had previously progressed on docetaxel alone. A phase I trial of the HDACi SAHA in combination with doxorubicin has also been conducted (Munster et al. 2009). This study involved 32 individuals with a range of solid tumours, including two with prostate cancer. Doxorubicin is a topoisomerase II inhibitor and a commonly used chemotherapy agent for solid tumours. The individuals with prostate cancer showed partial responses to the treatment as determined by PSA assessment and were found to have higher levels of histone acetylation in their tumour tissue samples. These studies suggest that HDACi in combination with other conventional anti-cancer agents may have potential for the treatment of prostate cancer.

Recently, a phase II clinical trial of FK228 was undertaken involving individuals with progressing, metastatic, castration-resistant prostate cancer (Molife et al. 2010). FK228 was intravenously administered to a total of 35 individuals. Two individuals achieved stable disease

lasting more than 6 months and with a decline in PSA levels of more than 50%. However, 11 individuals discontinued the treatment due to toxicity and reported side effects such as nausea, fatigue, vomiting and anorexia.

Use of histone methylation modulators in prostate cancer

The development of inhibitors of HMTs and HDMs is less advanced than for DNMTs and HDACs; however, there are a number of inhibitors of these molecules that have now been investigated in cancer models, including prostate cancer. Recently, a carbocyclic adenosine analogue, 3-deazaneplanocin-A (DZNep), was found to inhibit EZH2 HMT activity and, thus, inhibit H3K27 methylation (Tan et al. 2007; Chase and Cross 2011). Although subsequent studies suggest that this agent is a global HMT inhibitor rather than a specific EZH2 inhibitor (Miranda et al. 2009), several studies had demonstrated anti-tumourigenic activities in prostate cancer models. Crea et al. (2011) investigated the effects of DZNep on prostate CSCs. Treatment of two prostate cancer cell lines; the less aggressive LNCaP and the tumourigenic DU145 with 1 μ M DZNep inhibited EZH2 expression and reduced H3K27 methylation by 33%. Treatment of both cell lines with DZNep also inhibited formation of spheres, which are associated with CSC formation, by 95%. In addition, DZNep decreased the CSC enriched cell population (CD44⁺/24⁻) in LNCaP cells containing 0.1% CSC fraction and caused apoptosis in DU145 cells which are almost 100% CD44⁺ and more stem cell like. The effect of DZNep on *in vivo* tumour growth was examined by pre-treatment of LNCaP and Du145 with DZNep and then injection of these cells into NOD/SCID mice. In LNCaP, DZNep significantly reduced tumour formation in NOD/SCID mice but this effect was not observed in DU145. This suggests that DZNep is more effective in the early stage of prostate cancer by preventing prostate cancer metastasis rather than in the advanced stage of prostate cancer.

As noted above the HDM, LSD1 is upregulated in high-risk tumours and, therefore, inhibition of LSD1 is likely to be an attractive therapeutic target (Kahl et al. 2006). Since LSD1 is an amine oxidase, its activity can be inhibited by monoamine oxidase inhibitors, such as pargyline (Metzger et al. 2005). Pargyline was found to block demethylation of H3K9me1 and H3K9me2 during androgen-induced transcription in LNCaP cells. In addition, Kahl et al. (2006) showed that pargyline treatment of the LNCaP prostate cancer cell line conferred dose-dependent growth inhibition.

Currently, there are no studies of pargyline, DZNep or other modulators of histone methylation in clinical trials. However, modulation of histone methylation represents a promising therapeutic target, and therefore further investi-

gation of these, or other inhibitors, as they become available is clearly warranted.

Conclusions

Early and accurate diagnosis of prostate cancer is essential, and available studies suggest that gene-specific DNA methylation analysis may prove to be useful as a prostate cancer biomarker. Hypermethylation of the GSTP1 promoter, particularly in combination with other genes implicated in tumourigenesis, is proving to be a highly specific and sensitive measure of prostate cancer. Hypermethylation of particular genes also appears to be an early event in prostate cancer development. Further studies are, therefore, required to identify panels of genes that are aberrantly methylated in prostate cancer and may be useful in early detection of the disease.

In prostate cancer, as in other cancers, epigenetic changes are proving to be as common and important in tumour development and progression as genetic changes. Given the potential to reverse aberrant epigenetic modifications, and the promising results that have been reported for some inhibitors of epigenetic modifiers in prostate cancer animal models and clinical trials, further investigation of the mechanisms of action of these agents on prostate cancer cells is warranted. As recent studies suggest, these agents are most likely to be of use in the clinic when used in combination with conventional therapies, and further studies are therefore required to investigate their action in combination with other chemotherapeutic agents.

Acknowledgements Work in the authors' laboratories is supported by The Cancer Council Tasmania, The Australian Cancer Research Foundation and Cancer Australia.

Conflicts of interest The authors declare that they have no conflict of interest.

References

- Aitchison A, Warren A, Neal D, Rabbitts P (2007) RASSF1A promoter methylation is frequently detected in both pre-malignant and non-malignant microdissected prostatic epithelial tissues. *Prostate* 67(6):638–644
- Allen D, Butt A, Cahill D, Wheeler M, Popert R, Swaminathan R (2004) Role of cell-free plasma DNA as a diagnostic marker for prostate cancer. *Ann N Y Acad Sci* 1022:76–80
- Altimari A, Grigioni AD, Benedettini E, Gabusi E, Schiavina R, Martinelli A, Morselli-Labate AM, Martorana G, Grigioni WF, Fiorentino M (2008) Diagnostic role of circulating free plasma DNA detection in patients with localized prostate cancer. *Am J Clin Pathol* 129(5):756–762
- Ambis S, Prueitt RL, Yi M, Hudson RS, Howe TM, Petrocca F, Wallace TA, Liu CG, Volinia S, Calin GA, Yfantis HG, Stephens RM, Croce CM (2008) Genomic profiling of microRNA and

- messenger RNA reveals deregulated microRNA expression in prostate cancer. *Cancer Res* 68(15):6162–6170
- Appleton K, Mackay HJ, Judson I, Plumb JA, McCormick C, Strathdee G, Lee C, Barrett S, Reade S, Jadayel D, Tang A, Bellenger K, Mackay L, Setanoians A, Schatzlein A, Twelves C, Kaye SB, Brown R (2007) Phase I and pharmacodynamic trial of the DNA methyltransferase inhibitor decitabine and carboplatin in solid tumors. *J Clin Oncol* 25(29):4603–4609
- Bastian PJ, Nakayama M, De Marzo AM, Nelson WG (2004) GSTP1 CpG island hypermethylation as a molecular marker of prostate cancer. *Urologe A* 43(5):573–579
- Bastian PJ, Palapattu GS, Lin X, Yegnasubramanian S, Mangold LA, Trock B, Eisenberger MA, Partin AW, Nelson WG (2005) Preoperative serum DNA GSTP1 CpG island hypermethylation and the risk of early prostate-specific antigen recurrence following radical prostatectomy. *Clin Cancer Res* 11(11):4037–4043
- Bastian PJ, Ellinger J, Heukamp LC, Kahl P, Muller SC, von Rucker A (2007) Prognostic value of CpG island hypermethylation at PTGS2, RAR-beta, EDNRB, and other gene loci in patients undergoing radical prostatectomy. *Eur Urol* 51(3):665–674
- Basu HS, Mahlum A, Mehraein-Ghomi F, Kegel SJ, Guo S, Peters NR, Wilding G (2011) Pretreatment with anti-oxidants sensitizes oxidatively stressed human cancer cells to growth inhibitory effect of suberoylanilide hydroxamic acid (SAHA). *Cancer Chemother Pharmacol* 67(3):705–715
- Batty GN, Kantarjian H, Issa JP, Jabbour E, Santos FP, McCue D, Garcia-Manero G, Pierce S, O'Brien S, Cortes JE, Ravandi F (2010) Feasibility of therapy with hypomethylating agents in patients with renal insufficiency. *Clin Lymphoma Myeloma Leuk* 10(3):205–210
- Bedford MT, van Helden PD (1987) Hypomethylation of DNA in pathological conditions of the human prostate. *Cancer Res* 47(20):5274–5276
- Berger S, Kouzarides T, Shiekhhattar R, Shilatifard A (2009) An operational definition of epigenetics. *Genes Dev* 23(7):781–783
- Bianco-Miotto T, Chiam K, Buchanan G, Jindal S, Day TK, Thomas M, Pickering MA, O'Loughlin MA, Ryan NK, Raymond WA, Horvath LG, Kench JG, Stricker PD, Marshall VR, Sutherland RL, Henshall SM, Gerald WL, Scher HI, Risbridger GP, Clements JA, Butler LM, Tilley WD, Horsfall DJ, Ricciardelli C (2010) Global levels of specific histone modifications and an epigenetic gene signature predict prostate cancer progression and development. *Cancer Epidemiol Biomarkers Prev* 19(10):2611–2622
- Bird A (2002) DNA methylation patterns and epigenetic memory. *Genes Dev* 16(1):6–21
- Brooks JD, Weinstein M, Lin X, Sun Y, Pin SS, Bova GS, Epstein JI, Isaacs WB, Nelson WG (1998) CG island methylation changes near the GSTP1 gene in prostatic intraepithelial neoplasia. *Cancer Epidemiol Biomarkers Prev* 7(6):531–536
- Brothman AR, Swanson G, Maxwell TM, Cui J, Murphy KJ, Herrick J, Speights VO, Isaac J, Rohr LR (2005) Global hypomethylation is common in prostate cancer cells: a quantitative predictor for clinical outcome? *Cancer Genet Cytogenet* 156(1):31–36
- Bubendorf L, Schopfer A, Wagner U, Sauter G, Moch H, Willi N, Gasser TC, Mihatsch MJ (2000) Metastatic patterns of prostate cancer: an autopsy study of 1,589 patients. *Hum Pathol* 31(5):578–583
- Butler LM, Agus DB, Scher HI, Higgins B, Rose A, Cordon-Cardo C, Thaler HT, Rifkind RA, Marks PA, Richon VM (2000) Suberoylanilide hydroxamic acid, an inhibitor of histone deacetylase, suppresses the growth of prostate cancer cells in vitro and in vivo. *Cancer Res* 60(18):5165–5170
- Cairns P, Polascik TJ, Eby Y, Tokino K, Califano J, Merlo A, Mao L, Herath J, Jenkins R, Westra W et al (1995) Frequency of homozygous deletion at p16/CDKN2 in primary human tumours. *Nat Genet* 11(2):210–212
- Cairns P, Esteller M, Herman JG, Schoenberg M, Jeronimo C, Sanchez-Cespedes M, Chow NH, Grasso M, Wu L, Westra WB, Sidransky D (2001) Molecular detection of prostate cancer in urine by GSTP1 hypermethylation. *Clin Cancer Res* 7(9):2727–2730
- Calin GA, Sevignani C, Dumitru CD, Hyslop T, Noch E, Yendamuri S, Shimizu M, Rattan S, Bullrich F, Negrini M, Croce CM (2004) Human microRNA genes are frequently located at fragile sites and genomic regions involved in cancers. *Proc Natl Acad Sci USA* 101(9):2999–3004
- Cao P, Deng Z, Wan M, Huang W, Cramer SD, Xu J, Lei M, Sui G (2010) MicroRNA-101 negatively regulates Ezh2 and its expression is modulated by androgen receptor and HIF-1alpha/HIF-1beta. *Mol Cancer* 9:108–120
- Cashen AF, Shah AK, Todt L, Fisher N, DiPersio J (2008) Pharmacokinetics of decitabine administered as a 3-h infusion to patients with acute myeloid leukemia (AML) or myelodysplastic syndrome (MDS). *Cancer Chemother Pharmacol* 61(5):759–766
- Chase A, Cross NC (2011) Aberrations of EZH2 in Cancer. *Clin Cancer Res* 17(9):2613–2618
- Chi SG, deVere White RW, Muenzer JT, Gumerlock PH (1997) Frequent alteration of CDKN2 (p16(INK4A)/MTS1) expression in human primary prostate carcinomas. *Clin Cancer Res* 3(10):1889–1897
- Chinnaiyan P, Vallabhaneni G, Armstrong E, Huang SM, Harari PM (2005) Modulation of radiation response by histone deacetylase inhibition. *Int J Radiat Oncol Biol Phys* 62(1):223–229
- Chu DC, Chuang CK, Fu JB, Huang HS, Tseng CP, Sun CF (2002) The use of real-time quantitative polymerase chain reaction to detect hypermethylation of the CpG islands in the promoter region flanking the GSTP1 gene to diagnose prostate carcinoma. *J Urol* 167(4):1854–1858
- Chuang JC, Yoo CB, Kwan JM, Li TW, Liang G, Yang AS, Jones PA (2005) Comparison of biological effects of non-nucleoside DNA methylation inhibitors versus 5-aza-2'-deoxycytidine. *Mol Cancer Ther* 4(10):1515–1520
- Chuang CK, Chu DC, Tzou RD, Liou SI, Chia JH, Sun CF (2007) Hypermethylation of the CpG islands in the promoter region flanking GSTP1 gene is a potential plasma DNA biomarker for detecting prostate carcinoma. *Cancer Detect Prev* 31(1):59–63
- Chuang JC, Warner SL, Vollmer D, Vankayalapati H, Redkar S, Bearss DJ, Qiu X, Yoo CB, Jones PA (2010) S110, a 5-Aza-2'-deoxycytidine-containing dinucleotide, is an effective DNA methylation inhibitor in vivo and can reduce tumor growth. *Mol Cancer Ther* 9(5):1443–1450
- Cihak A (1974) Biological effects of 5-azacytidine in eukaryotes. *Oncology* 30(5):405–422
- Crea F, Hurt EM, Mathews LA, Cabarcas SM, Sun L, Marquez VE, Danesi R, Farrar WL (2011) Pharmacologic disruption of polycomb repressive complex 2 inhibits tumorigenicity and tumor progression in prostate cancer. *Mol Cancer* 10(1):40. doi:10.1186/1476-4598-10-40
- Dammann R, Schagdarsurengin U, Seidel C, Strunnikova M, Rastetter M, Baier K, Pfeifer GP (2005) The tumor suppressor RASSF1A in human carcinogenesis: an update. *Histol Histopathol* 20(2):645–663
- De Marzo AM, DeWeese TL, Platz EA, Meeker AK, Nakayama M, Epstein JI, Isaacs WB, Nelson WG (2004) Pathological and molecular mechanisms of prostate carcinogenesis: implications for diagnosis, detection, prevention, and treatment. *J Cell Biochem* 91(3):459–477
- Dissanayake S, Wade M, Johnson C, O'Connell M, Leotlela P, French A, Shah K, Hewitt K, Rosenthal D, Indig F (2007) The Wnt5A/

- protein kinase C pathway mediates motility in melanoma cells via the inhibition of metastasis suppressors and initiation of an epithelial to mesenchymal transition. *J Biol Chem* 282(23):17259–17271
- Dunn BK (2003) Hypomethylation: one side of a larger picture. *Ann N Y Acad Sci* 983:28–42
- Eden A, Gaudet F, Waghmare A, Jaenisch R (2003) Chromosomal instability and tumors promoted by DNA hypomethylation. *Science* 300(5618):455–455
- Ellinger J, Bastian PJ, Ellinger N, Kahl P, Perabo FG, Buttner R, Muller SC, Ruecker A (2008a) Apoptotic DNA fragments in serum of patients with muscle invasive bladder cancer: a prognostic entity. *Cancer Lett* 264(2):274–280
- Ellinger J, Bastian PJ, Haan KI, Heukamp LC, Buettner R, Fimmers R, Mueller SC, von Ruecker A (2008b) Noncancerous PTGS2 DNA fragments of apoptotic origin in sera of prostate cancer patients qualify as diagnostic and prognostic indicators. *Int J Cancer* 122(1):138–143
- Ellinger J, Bastian PJ, Jurgan T, Biermann K, Kahl P, Heukamp LC, Wernert N, Muller SC, von Ruecker A (2008c) CpG island hypermethylation at multiple gene sites in diagnosis and prognosis of prostate cancer. *Urology* 71(1):161–167
- Ellinger J, Kahl P, von der Gathen J, Rogenhofer S, Heukamp LC, Gutgemann I, Walter B, Hofstadter F, Buttner R, Muller SC, Bastian PJ, von Ruecker A (2010) Global levels of histone modifications predict prostate cancer recurrence. *Prostate* 70(1):61–69
- Esteller M (2008) Epigenetics in cancer. *N Engl J Med* 358(11):1148
- Feinberg A, Ohlsson R, Henikoff S (2006) The epigenetic progenitor origin of human cancer. *Nature Rev Genet* 7(1):21–33
- Flori AR, Steinhoff C, Muller M, Seifert HH, Hader C, Engers R, Ackermann R, Schulz WA (2004) Coordinate hypermethylation at specific genes in prostate carcinoma precedes LINE-1 hypomethylation. *Br J Cancer* 91(5):985–994
- Fujita Y, Kojima K, Hamada N, Ohhashi R, Akao Y, Nozawa Y, Deguchi T, Ito M (2008) Effects of miR-34a on cell growth and chemoresistance in prostate cancer PC3 cells. *Biochem Biophys Res Commun* 377(1):114–119
- Galardi S, Mercatelli N, Giorda E, Massalini S, Frajese GV, Ciafre SA, Farace MG (2007) miR-221 and miR-222 expression affects the proliferation potential of human prostate carcinoma cell lines by targeting p27Kip1. *J Biol Chem* 282(32):23716–23724
- Gardiner-Garden M, Frommer M (1987) CpG islands in vertebrate genomes. *J Mol Biol* 196(2):261–282
- Gardner KE, Allis CD, Strahl BD (2011) Operating on chromatin, a colorful language where context matters. *J Mol Biol* 409(1):36–46
- Gaughan L, Stockley J, Wang N, McCracken SR, Treumann A, Armstrong K, Shaheen F, Watt K, McEwan IJ, Wang C, Pestell RG, Robson CN (2011) Regulation of the androgen receptor by SET9-mediated methylation. *Nucleic Acids Res* 39(4):1266–1279
- Gediya LK, Belosay A, Khandelwal A, Purushottamachar P, Njar VC (2008) Improved synthesis of histone deacetylase inhibitors (HDIs) (MS-275 and CI-994) and inhibitory effects of HDIs alone or in combination with RAMBAs or retinoids on growth of human LNCaP prostate cancer cells and tumor xenografts. *Bioorg Med Chem* 16(6):3352–3360
- Goessl C, Muller M, Heicappell R, Krause H, Straub B, Schrader M, Miller K (2001) DNA-based detection of prostate cancer in urine after prostatic massage. *Urology* 58(3):335–338
- Graff JR, Herman JG, Lapidus RG, Chopra H, Xu R, Jarrard DF, Isaacs WB, Pitha PM, Davidson NE, Baylin SB (1995) E-cadherin expression is silenced by DNA hypermethylation in human breast and prostate carcinomas. *Cancer Res* 55(22):5195–5199
- Gravina G, Marampon F, Di Staso M, Bonfili P, Vitturini A, Jannini E, Pestell R, Tombolini V, Festuccia C (2010) 5 Azacitidine restores and amplifies the bicalutamide response on preclinical models of androgen receptor expressing or deficient prostate tumors. *Prostate* 70(11):1166–1178
- Hake SB, Xiao A, Allis CD (2004) Linking the epigenetic ‘language’ of covalent histone modifications to cancer. *Br J Cancer* 90(4):761–769
- Halkidou K, Gaughan L, Cook S, Leung HY, Neal DE, Robson CN (2004) Upregulation and nuclear recruitment of HDAC1 in hormone refractory prostate cancer. *Prostate* 59(2):177–189
- Halliwel B (2006) Oxidative stress and neurodegeneration: where are we now? *J Neurochem* 97(6):1634–1658
- Hamm CA, Xie H, Costa FF, Vanin EF, Seftor EA, Sredni ST, Bischoff J, Wang D, Bonaldo MF, Hendrix MJ, Soares MB (2009) Global demethylation of rat chondrosarcoma cells after treatment with 5-aza-2'-deoxycytidine results in increased tumorigenicity. *PLoS One* 4(12):e8340
- Hao Y, Zhao Y, Zhao X, He C, Pang X, Wu TC, Califano JA, Gu X (2011) Improvement of prostate cancer detection by integrating the PSA test with miRNA expression profiling. *Cancer Invest* 29(4):318–324
- Harden SV, Guo Z, Epstein JI, Sidransky D (2003) Quantitative GSTP1 methylation clearly distinguishes benign prostatic tissue and limited prostate adenocarcinoma. *J Urol* 169(3):1138–1142
- Hark AT, Schoenherr CJ, Katz DJ, Ingram RS, Levorse JM, Tilghman SM (2000) CTCF mediates methylation-sensitive enhancer-blocking activity at the H19/Igf2 locus. *Nature* 405(6785):486–489
- Hayashita Y, Osada H, Tatematsu Y, Yamada H, Yanagisawa K, Tomida S, Yatabe Y, Kawahara K, Sekido Y, Takahashi T (2005) A polycistronic microRNA cluster, miR-17-92, is overexpressed in human lung cancers and enhances cell proliferation. *Cancer Res* 65(21):9628–9632
- He L, Hannon GJ (2004) MicroRNAs: small RNAs with a big role in gene regulation. *Nat Rev Genet* 5(7):522–531
- He L, Thomson JM, Hemann MT, Hernando-Monge E, Mu D, Goodson S, Powers S, Cordon-Cardo C, Lowe SW, Hannon GJ, Hammond SM (2005) A microRNA polycistron as a potential human oncogene. *Nature* 435(7043):828–833
- Henrique R, Ribeiro FR, Fonseca D, Hoque MO, Carvalho AL, Costa VL, Pinto M, Oliveira J, Teixeira MR, Sidransky D, Jeronimo C (2007) High promoter methylation levels of APC predict poor prognosis in sextant biopsies from prostate cancer patients. *Clin Cancer Res* 13(20):6122–6129
- Hesson LB, Cooper WN, Latif F (2007) The role of RASSF1A methylation in cancer. *Dis Markers* 23(1–2):73–87
- Huang H, Reed CP, Zhang JS, Shridhar V, Wang L, Smith DI (1999) Carboxypeptidase A3 (CPA3): a novel gene highly induced by histone deacetylase inhibitors during differentiation of prostate epithelial cancer cells. *Cancer Res* 59(12):2981–2988
- Huang ZH, Li LH, Hua D (2006) Quantitative analysis of plasma circulating DNA at diagnosis and during follow-up of breast cancer patients. *Cancer Lett* 243(1):64–70
- Hulett MD, Freeman C, Hamdorf BJ, Baker RT, Harris MJ, Parish CR (1999) Cloning of mammalian heparanase, an important enzyme in tumor invasion and metastasis. *Nat Med* 5(7):803–809
- Illingworth R, Bird A (2009) CpG islands—‘a rough guide’. *FEBS Lett* 583(11):1713–1720
- Jackson-Grusby L, Laird PW, Magge SN, Moeller BJ, Jaenisch R (1997) Mutagenicity of 5-aza-2'-deoxycytidine is mediated by the mammalian DNA methyltransferase. *Proc Natl Acad Sci USA* 94(9):4681–4685
- Jahr S, Hentze H, Englisch S, Hardt D, Fackelmayer FO, Hesch RD, Knippers R (2001) DNA fragments in the blood plasma of cancer patients: quantitations and evidence for their origin from apoptotic and necrotic cells. *Cancer Res* 61(4):1659–1665

- Jarrard DF, Bova GS, Ewing CM, Pin SS, Nguyen SH, Baylin SB, Cairns P, Sidransky D, Herman JG, Isaacs WB (1997) Deletional, mutational, and methylation analyses of CDKN2 (p16/MTS1) in primary and metastatic prostate cancer. *Genes Chromosom Cancer* 19(2):90–96
- Jarrard DF, Kinoshita H, Shi Y, Sandefur C, Hoff D, Meisner LF, Chang C, Herman JG, Isaacs WB, Nassif N (1998) Methylation of the androgen receptor promoter CpG island is associated with loss of androgen receptor expression in prostate cancer cells. *Cancer Res* 58(23):5310–5314
- Jenster G (1999) The role of the androgen receptor in the development and progression of prostate cancer. *Semin Oncol* 26(4):407–421
- Jeronimo C, Usadel H, Henrique R, Silva C, Oliveira J, Lopes C, Sidransky D (2002) Quantitative GSTP1 hypermethylation in bodily fluids of patients with prostate cancer. *Urology* 60(6):1131–1135
- Jeronimo C, Henrique R, Hoque MO, Mambo E, Ribeiro FR, Vazrim G, Oliveira J, Teixeira MR, Lopes C, Sidransky D (2004) A quantitative promoter methylation profile of prostate cancer. *Clin Cancer Res* 10(24):8472–8480
- Jones PA, Baylin SB (2007) The epigenomics of cancer. *Cell* 128(4):683–692
- Jones P, Taylor S (1980) Cellular differentiation, cytidine analogs and DNA methylation. *Cell* 20(1):85–93
- Jones P, Veenstra G, Wade P, Vermaak D, Kass S, Landsberger N, Strouboulis J, Wolffe A (1998) Methylated DNA and MeCP2 recruit histone deacetylase to repress transcription. *Nat Genet* 19(2):187–191
- Jung K, Stephan C, Lewandowski M, Klotzek S, Jung M, Kristiansen G, Lein M, Loening SA, Schnorr D (2004) Increased cell-free DNA in plasma of patients with metastatic spread in prostate cancer. *Cancer Lett* 205(2):173–180
- Kahl P, Gullotti L, Heukamp LC, Wolf S, Friedrichs N, Vörreuther R, Solleder G, Bastian PJ, Ellinger J, Metzger E, Schule R, Buettner R (2006) Androgen receptor coactivators lysine-specific histone demethylase 1 and four and a half LIM domain protein 2 predict risk of prostate cancer recurrence. *Cancer Res* 66(23):11341–11347
- Kallakury BV, Sheehan CE, Winn-Deen E, Oliver J, Fisher HA, Kaufman RP Jr, Ross JS (2001) Decreased expression of catenins (alpha and beta), p120 CTN, and E-cadherin cell adhesion proteins and E-cadherin gene promoter methylation in prostatic adenocarcinomas. *Cancer* 92(11):2786–2795
- Kaminskas E, Farrell A, Abraham S, Baird A, Hsieh L, Lee S, Leighton J, Patel H, Rahman A, Sridhara R (2005) Approval summary: azacitidine for treatment of myelodysplastic syndrome subtypes. *Clin Cancer Res* 11(10):3604–3608
- Kang GH, Lee S, Lee HJ, Hwang KS (2004) Aberrant CpG island hypermethylation of multiple genes in prostate cancer and prostatic intraepithelial neoplasia. *J Pathol* 202(2):233–240
- Kantarjian HM, O'Brien S, Cortes J, Giles FJ, Faderl S, Issa JP, Garcia-Manero G, Rios MB, Shan J, Andreeff M, Keating M, Talpaz M (2003) Results of decitabine (5-aza-2'-deoxycytidine) therapy in 130 patients with chronic myelogenous leukemia. *Cancer* 98(3):522–528
- Karanikolas B, Figueiredo M, Wu L (2010) Comprehensive evaluation of the role of EZH2 in the growth, invasion, and aggression of a panel of prostate cancer cell lines. *Prostate* 70(6):675–688
- Konety BR, Schwartz GG, Acierno JS Jr, Becich MJ, Getzenberg RH (1996) The role of vitamin D in normal prostate growth and differentiation. *Cell Growth Differ* 7(11):1563–1570
- Kornberg RD, Lorch Y (1999) Twenty-five years of the nucleosome, fundamental particle of the eukaryote chromosome. *Cell* 98(3):285–294
- Kouzarides T (2007) Chromatin modifications and their function. *Cell* 128(4):693–705
- Krill D, Stoner J, Konety BR, Becich MJ, Getzenberg RH (1999) Differential effects of vitamin D on normal human prostate epithelial and stromal cells in primary culture. *Urology* 54(1):171–177
- Kulp SK, Chen CS, Wang DS, Chen CY (2006) Antitumor effects of a novel phenylbutyrate-based histone deacetylase inhibitor, (S)-HDAC-42, in prostate cancer. *Clin Cancer Res* 12(17):5199–5206
- Lai MT, Yang CC, Lin TY, Tsai FJ, Chen WC (2008) Depsipeptide (FK228) inhibits growth of human prostate cancer cells. *Urol Oncol* 26(2):182–189
- Lakshmikanthan V, Kaddour-Djebbar I, Lewis RW, Kumar MV (2006) SAHA-sensitized prostate cancer cells to TNFalpha-related apoptosis-inducing ligand (TRAIL): mechanisms leading to synergistic apoptosis. *Int J Cancer* 119(1):221–228
- Lavelle D, Chin J, Vaitkus K, Redkar S, Phiasivongsa P, Tang C, Will R, Hankewych M, Roxas B, Singh M (2007) Oral decitabine reactivates expression of the methylated globin gene in *Papio anubis*. *Am J Hematol* 82(11):981–985
- Lee J (2007) GSTP1 promoter hypermethylation is an early event in breast carcinogenesis. *Virchows Arch* 450(6):637–642
- Lee WH, Morton RA, Epstein JI, Brooks JD, Campbell PA, Bova GS, Hsieh WS, Isaacs WB, Nelson WG (1994) Cytidine methylation of regulatory sequences near the pi-class glutathione S-transferase gene accompanies human prostatic carcinogenesis. *Proc Natl Acad Sci USA* 91(24):11733–11737
- Lee BH, Yegnasubramanian S, Lin X, Nelson WG (2005) Procainamide is a specific inhibitor of DNA methyltransferase 1. *J Biol Chem* 280(49):40749–40756
- Lee B, Moussa A, Li J, Fareed K, Jones J (2011) Percentage of Free Prostate-specific Antigen: Implications in Modern Extended Scheme Prostate Biopsy. *Urology* 77(4):899–903
- Li LC, Zhao H, Nakajima K, Oh BR, Ribeiro Filho LA, Carroll P, Dahiya R (2001) Methylation of the E-cadherin gene promoter correlates with progression of prostate cancer. *J Urol* 166(2):705–709
- Li L, Okino S, Dahiya R (2004) DNA methylation in prostate cancer. *Biochim Biophys Acta* 1704(2):87–102
- Lin X, Asgari K, Putzi MJ, Gage WR, Yu X, Cornblatt BS, Kumar A, Piantadosi S, DeWeese TL, De Marzo AM, Nelson WG (2001a) Reversal of GSTP1 CpG island hypermethylation and reactivation of pi-class glutathione S-transferase (GSTP1) expression in human prostate cancer cells by treatment with procainamide. *Cancer Res* 61(24):8611–8616
- Lin X, Tascilar M, Lee WH, Vles WJ, Lee BH, Veeraswamy R, Asgari K, Freije D, van Rees B, Gage WR, Bova GS, Isaacs WB, Brooks JD, DeWeese TL, De Marzo AM, Nelson WG (2001b) GSTP1 CpG island hypermethylation is responsible for the absence of GSTP1 expression in human prostate cancer cells. *Am J Pathol* 159(5):1815–1826
- Liu L, Yoon JH, Dammann R, Pfeifer GP (2002) Frequent hypermethylation of the RASSF1A gene in prostate cancer. *Oncogene* 21(44):6835–6840
- Liu C, Kelnar K, Liu B, Chen X, Calhoun-Davis T, Li H, Patrawala L, Yan H, Jeter C, Honorio S, Wiggins JF, Bader AG, Fagin R, Brown D, Tang DG (2011a) The microRNA miR-34a inhibits prostate cancer stem cells and metastasis by directly repressing CD44. *Nat Med* 17(2):211–215
- Liu L, Kron KJ, Pethe VV, Demetrashvili N, Nesbitt ME, Trachtenberg J, Ozelik H, Fleshner NE, Briollais L, van der Kwast TH, Bapat B (2011b) Association of tissue promoter methylation levels of APC, TGFbeta2, HOXD3, and RASSF1A with prostate cancer progression. *Int J Cancer*. doi:10.1002/ijc.25908
- Lou W, Krill D, Dhir R, Becich MJ, Dong JT, Frierson HF Jr, Isaacs WB, Isaacs JT, Gao AC (1999) Methylation of the CD44 metastasis suppressor gene in human prostate cancer. *Cancer Res* 59(10):2329–2331

- Macri E, Loda M (1998) Role of p27 in prostate carcinogenesis. *Cancer Metastasis Rev* 17(4):337–344
- Marrocco DL, Tilley WD, Bianco-Miotto T, Evdokiou A, Scher HI, Rifkind RA, Marks PA, Richon VM, Butler LM (2007) Suberoylanilide hydroxamic acid (vorinostat) represses androgen receptor expression and acts synergistically with an androgen receptor antagonist to inhibit prostate cancer cell proliferation. *Mol Cancer Ther* 6(1):51–60
- Mercatelli N, Coppola V, Bonci D, Miele F, Costantini A, Guadagnoli M, Bonanno E, Muto G, Frajese GV, De Maria R, Spagnoli LG, Farace MG, Ciafre SA (2008) The inhibition of the highly expressed miR-221 and miR-222 impairs the growth of prostate carcinoma xenografts in mice. *PLoS One* 3(12):4029–4039
- Metzger E, Wissmann M, Yin N, Muller JM, Schneider R, Peters AH, Gunther T, Buettner R, Schule R (2005) LSD1 demethylates repressive histone marks to promote androgen-receptor-dependent transcription. *Nature* 437(7057):436–439
- Miller GJ, Stapleton GE, Hedlund TE, Moffat KA (1995) Vitamin D receptor expression, 24-hydroxylase activity, and inhibition of growth by 1 α ,25-dihydroxyvitamin D₃ in seven human prostatic carcinoma cell lines. *Clin Cancer Res* 1(9):997–1003
- Miranda TB, Cortez CC, Yoo CB, Liang G, Abe M, Kelly TK, Marquez VE, Jones PA (2009) DNep is a global histone methylation inhibitor that reactivates developmental genes not silenced by DNA methylation. *Mol Cancer Ther* 8(6):1579–1588
- Mitchell PS, Parkin RK, Kroh EM, Fritz BR, Wyman SK, Pogosova-Agadjanyan EL, Peterson A, Noteboom J, O'Brian KC, Allen A, Lin DW, Urban N, Drescher CW, Knudsen BS, Stirewalt DL, Gentleman R, Vessella RL, Nelson PS, Martin DB, Tewari M (2008) Circulating microRNAs as stable blood-based markers for cancer detection. *Proc Natl Acad Sci USA* 105(30):10513–10518
- Molife LR, Attard G, Fong PC, Karavasili V, Reid AH, Patterson S, Riggs CE Jr, Higano C, Stadler WM, McCulloch W, Dearnaley D, Parker C, de Bono JS (2010) Phase II, two-stage, single-arm trial of the histone deacetylase inhibitor (HDACi) romidepsin in metastatic castration-resistant prostate cancer (CRPC). *Ann Oncol* 21(1):109–113
- Munich PN, Marchion D, Thomas S, Egorin M, Minton S, Springett G, Lee JH, Simon G, Chiappori A, Sullivan D, Daud A (2009) Phase I trial of vorinostat and doxorubicin in solid tumours: histone deacetylase 2 expression as a predictive marker. *Br J Cancer* 101(7):1044–1050
- Nakayama M, Bennett CJ, Hicks JL, Epstein JI, Platz EA, Nelson WG, De Marzo AM (2003) Hypermethylation of the human glutathione S-transferase-pi gene (GSTP1) CpG island is present in a subset of proliferative inflammatory atrophy lesions but not in normal or hyperplastic epithelium of the prostate: a detailed study using laser-capture microdissection. *Am J Pathol* 163(3):923–933
- Nan X, Ng H, Johnson C, Laherty C, Turner B, Eisenman R, Bird A (1998) Transcriptional repression by the methyl-CpG-binding protein MeCP2 involves a histone deacetylase complex. *Nature* 393(6683):386–389
- Nishioka K, Chuikov S, Sarma K, Erdjument-Bromage H, Allis CD, Tempst P, Reinberg D (2002) Set9, a novel histone H3 methyltransferase that facilitates transcription by precluding histone tail modifications required for heterochromatin formation. *Genes Dev* 16(4):479–489
- Ogishima T, Shiina H, Breault JE, Tabatabai L, Bassett WW, Enokida H, Li LC, Kawakami T, Urakami S, Ribeiro-Filho LA, Terashima M, Fujime M, Igawa M, Dahiya R (2005) Increased heparanase expression is caused by promoter hypomethylation and up-regulation of transcriptional factor early growth response-1 in human prostate cancer. *Clin Cancer Res* 11(3):1028–1036
- Opel M, Lando D, Bonilla C, Treweek SC, Boukaba A, Wälfridsson J, Cauwood J, Werler PJ, Carr AM, Kouzarides T, Murzina NV, Allshire RC, Ekwall K, Laue ED (2007) Genome-wide studies of histone demethylation catalysed by the fission yeast homologues of mammalian LSD1. *PLoS One* 2(4):e386
- Ostling P, Leivonen SK, Aakula A, Kohonen P, Makela R, Hagman Z, Edsjo A, Kangaspeka S, Edgren H, Nicorici D, Bjartell A, Ceder Y, Perala M, Kallioniemi O (2011) Systematic analysis of microRNAs targeting the androgen receptor in prostate cancer cells. *Cancer Res* 71(5):1956–1967
- Pakneshan P, Xing R, Rabbani S (2003) Methylation status of uPA promoter as a molecular mechanism regulating prostate cancer invasion and growth in vitro and in vivo. *FASEB J* 17(9):1081–1088
- Pang Y, Young CY, Yuan H (2010) MicroRNAs and prostate cancer. *Acta Biochim Biophys Sin (Shanghai)* 42(6):363–369
- Papadopoulou E, Davilas E, Sotiriou V, Koliopoulos A, Aggelakis F, Dardoufas K, Agnanti NJ, Karydas I, Nasioulas G (2004) Cell-free DNA and RNA in plasma as a new molecular marker for prostate cancer. *Oncol Res* 14(9):439–445
- Patrawala L, Calhoun T, Schneider-Broussard R, Li H, Bhatia B, Tang S, Reilly JG, Chandra D, Zhou J, Claypool K, Coghlan L, Tang DG (2006) Highly purified CD44+ prostate cancer cells from xenograft human tumors are enriched in tumorigenic and metastatic progenitor cells. *Oncogene* 25(12):1696–1708
- Patrawala L, Calhoun-Davis T, Schneider-Broussard R, Tang DG (2007) Hierarchical organization of prostate cancer cells in xenograft tumors: the CD44+ α 2beta1+ cell population is enriched in tumor-initiating cells. *Cancer Res* 67(14):6796–6805
- Perl A, Wilgenbus P, Dahl U, Semb H, Christofori G (1998) A causal role for E-cadherin in the transition from adenoma to carcinoma. *Nature* 392(6672):190–193
- Perry ME, Rolfe M, McIntyre P, Commane M, Stark GR (1992) Induction of gene amplification by 5-aza-2'-deoxycytidine. *Mutat Res* 276(3):189–197
- Pfeifer GP, Dammann R (2005) Methylation of the tumor suppressor gene RASSF1A in human tumors. *Biochemistry (Mosc)* 70(5):576–583
- Phé V, Cussenot O, Rouprêt M (2010) Methylated genes as potential biomarkers in prostate cancer. *BJUI* 105(10):1364–1370
- Porkka KP, Pfeiffer MJ, Waltering KK, Vessella RL, Tammela TL, Visakorpi T (2007) MicroRNA expression profiling in prostate cancer. *Cancer Res* 67(13):6130–6135
- Qian DZ, Ren M, Wei Y, Wang X, van de Geijn F, Rasmussen C, Nakanishi O, Sacchi N, Pili R (2005) In vivo imaging of retinoic acid receptor beta2 transcriptional activation by the histone deacetylase inhibitor MS-275 in retinoid-resistant prostate cancer cells. *Prostate* 64(1):20–28
- Qian DZ, Kato Y, Shabbeer S, Wei Y, Verheul HM, Salumbides B, Sanni T, Atadja P, Pili R (2006) Targeting tumor angiogenesis with histone deacetylase inhibitors: the hydroxamic acid derivative LBH589. *Clin Cancer Res* 12(2):634–642
- Qian DZ, Wei YF, Wang X, Kato Y, Cheng L, Pili R (2007) Antitumor activity of the histone deacetylase inhibitor MS-275 in prostate cancer models. *Prostate* 67(11):1182–1193
- Rashid SF, Moore JS, Walker E, Driver PM, Engel J, Edwards CE, Brown G, Uskokovic MR, Campbell MJ (2001) Synergistic growth inhibition of prostate cancer cells by 1 α ,25-dihydroxyvitamin D₃ and its 19-nor-hexafluoride analogs in combination with either sodium butyrate or trichostatin A. *Oncogene* 20(15):1860–1872
- Rathkopf D, Wong BY, Ross RW, Anand A, Tanaka E, Woo MM, Hu J, Dzik-Jurasz A, Yang W, Scher HI (2010) A phase I study of oral panobinostat alone and in combination with docetaxel in patients with castration-resistant prostate cancer. *Cancer Chemother Pharmacol* 66(1):181–189
- Reibenwein J, Pils D, Horak P, Tomicek B, Goldner G, Worel N, Elandt K, Krainer M (2007) Promoter hypermethylation of

- GSTP1, AR, and 14-3-3sigma in serum of prostate cancer patients and its clinical relevance. *Prostate* 67(4):427–432
- Richiardi L, Fiano V, Vizzini L, De Marco L, Delsedime L, Akre O, Tos AG, Merletti F (2009) Promoter methylation in APC, RUNX3, and GSTP1 and mortality in prostate cancer patients. *J Clin Oncol* 27(19):3161–3168
- Rosenbaum E, Hoque MO, Cohen Y, Zahurak M, Eisenberger MA, Epstein JI, Partin AW, Sidransky D (2005) Promoter hypermethylation as an independent prognostic factor for relapse in patients with prostate cancer following radical prostatectomy. *Clin Cancer Res* 11(23):8321–8325
- Roupret M, Hupertan V, Yates DR, Catto JW, Rehman I, Meuth M, Ricci S, Lacave R, Cancel-Tassin G, de la Taille A, Rozet F, Cathelineau X, Vallancien G, Hamdy FC, Cussenot O (2007) Molecular detection of localized prostate cancer using quantitative methylation-specific PCR on urinary cells obtained following prostate massage. *Clin Cancer Res* 13(6):1720–1725
- Santi DV, Norment A, Garrett CE (1984) Covalent bond formation between a DNA-cytosine methyltransferase and DNA containing 5-azacytosine. *Proc Natl Acad Sci USA* 81(22):6993–6997
- Santourlidis S, Florl A, Ackermann R, Wirtz HC, Schulz WA (1999) High frequency of alterations in DNA methylation in adenocarcinoma of the prostate. *Prostate* 39(3):166–174
- Sauntharajah Y, Hillery C, Lavelle D, Molokie R, Dorn L, Bressler L, Gavazova S, Chen Y, Hoffman R, DeSimone J (2003) Effects of 5-aza-2'-deoxycytidine on fetal hemoglobin levels, red cell adhesion, and hematopoietic differentiation in patients with sickle cell disease. *Blood* 102(12):3865–3870
- Schmudde M, Braun A, Pende D, Sonnemann J, Klier U, Beck JF, Moretta L, Broker BM (2008) Histone deacetylase inhibitors sensitize tumour cells for cytotoxic effects of natural killer cells. *Cancer Lett* 272(1):110–121
- Schnekenburger M, Grandjenette C, Ghelfi J, Karius T, Foliguet B, Dicato M, Diederich M (2011) Sustained exposure to the DNA demethylating agent, 2'-deoxy-5-azacytidine, leads to apoptotic cell death in chronic myeloid leukemia by promoting differentiation, senescence, and autophagy. *Biochem Pharmacol* 81(3):364–378
- Schrump DS, Fischette MR, Nguyen DM, Zhao M, Li X, Kunst TF, Hancox A, Hong JA, Chen GA, Pishchik V, Figg WD, Murgo AJ, Steinberg SM (2006) Phase I study of decitabine-mediated gene expression in patients with cancers involving the lungs, esophagus, or pleura. *Clin Cancer Res* 12(19):5777–5785
- Schulz WA, Elo JP, Florl AR, Pennanen S, Santourlidis S, Engers R, Buchardt M, Seifert HH, Visakorpi T (2002) Genomewide DNA hypomethylation is associated with alterations on chromosome 8 in prostate carcinoma. *Genes Chromosom Cancer* 35(1):58–65
- Schwartz G, Hulka B (1990) Is vitamin D deficiency a risk factor for prostate cancer? *Anticancer Res* 10(5A):1307–1311
- Seligson D, Horvath S, Shi T, Yu H, Tze S, Grunstein M, Kurdiani S (2005) Global histone modification patterns predict risk of prostate cancer recurrence. *Nature* 435(7046):1262–1266
- Shabbeer S, Kortenhorst MS, Kachhap S, Galloway N, Rodriguez R, Carducci MA (2007) Multiple molecular pathways explain the anti-proliferative effect of valproic acid on prostate cancer cells in vitro and in vivo. *Prostate* 67(10):1099–1110
- Sharma S, Kelly T, Jones P (2010) Epigenetics in cancer. *Carcinogenesis* 31(1):27–36
- Shi Y, Lan F, Matson C, Mulligan P, Whetstone JR, Cole PA, Casero RA (2004) Histone demethylation mediated by the nuclear amine oxidase homolog LSD1. *Cell* 119(7):941–953
- Shukeir N, Pakneshan P, Chen G, Szyf M, Rabbani S (2006) Alteration of the methylation status of tumor-promoting genes decreases prostate cancer cell invasiveness and tumorigenesis in vitro and in vivo. *Cancer Res* 66(18):9202–9210
- Singal R, van Wert J, Bashambu M (2001) Cytosine methylation represses glutathione S-transferase P1 (GSTP1) gene expression in human prostate cancer cells. *Cancer Res* 61(12):4820–4826
- Sozzi G, Conte D, Mariani L, Lo Vullo S, Roz L, Lombardo C, Pierotti MA, Tavecchio L (2001) Analysis of circulating tumor DNA in plasma at diagnosis and during follow-up of lung cancer patients. *Cancer Res* 61(12):4675–4678
- Suenaga M, Soda H, Oka M, Yamaguchi A, Nakatomi K, Shiozawa K, Kawabata S, Kasai T, Yamada Y, Kamihira S, Tei C, Kohno S (2002) Histone deacetylase inhibitors suppress telomerase reverse transcriptase mRNA expression in prostate cancer cells. *Int J Cancer* 97(5):621–625
- Suzuki H, Ueda T, Ichikawa T, Ito H (2003) Androgen receptor involvement in the progression of prostate cancer. *Endocr Relat Cancer* 10(2):209–216
- Takahashi S, Inaguma S, Sakakibara M, Cho YM, Suzuki S, Ikeda Y, Cui L, Shirai T (2002) DNA methylation in the androgen receptor gene promoter region in rat prostate cancers. *Prostate* 52(1):82–88
- Tan J, Yang X, Zhuang L, Jiang X, Chen W, Lee PL, Karuturi RK, Tan PB, Liu ET, Yu Q (2007) Pharmacologic disruption of polycomb-repressive complex 2-mediated gene repression selectively induces apoptosis in cancer cells. *Genes Dev* 21(9):1050–1063
- Tang X, Gal J, Kyprianou N, Zhu H, Tang G (2011) Detection of microRNAs in prostate cancer cells by microRNA array. *Methods Mol Biol* 732:69–88
- Tekur S, Lau KM, Long J, Burnstein K, Ho SM (2001) Expression of RFG/ELE1alpha/ARA70 in normal and malignant prostatic epithelial cell cultures and lines: regulation by methylation and sex steroids. *Mol Carcinogen* 30(1):1–13
- Thibault A, Figg WD, Bergan RC, Lush RM, Myers CE, Tompkins A, Reed E, Samid D (1998) A phase II study of 5-aza-2'-deoxycytidine (decitabine) in hormone independent metastatic (D2) prostate cancer. *Tumori* 84(1):87–89
- Thompson I, Pauler D, Goodman P, Tangen C, Lucia M, Parnes H, Minasian L, Ford L, Lippman S, Crawford E (2004) Prevalence of prostate cancer among men with a prostate-specific antigen level 4.0 ng per milliliter. *N Engl J Med* 350(22):2239–2246
- Tokizane T, Shiina H, Igawa M, Enokida H, Urakami S, Kawakami T, Ogishima T, Okino ST, Li LC, Tanaka Y, Nonomura N, Okuyama A, Dahiya R (2005) Cytochrome P450 1B1 is overexpressed and regulated by hypomethylation in prostate cancer. *Clin Cancer Res* 11(16):5793–5801
- Trejo-Becerril C, Perez-Cardenas E, Trevino-Cuevas H, Taja-Chayeb L, Garcia-Lopez P, Segura-Pacheco B, Chavez-Blanco A, Lizano-Soberon M, Gonzalez-Fierro A, Mariscal I, Wegman-Ostrosky T, Duenas-Gonzalez A (2003) Circulating nucleosomes and response to chemotherapy: an in vitro, in vivo and clinical study on cervical cancer patients. *Int J Cancer* 104(6):663–668
- Tsihlias J, Kapusta LR, DeBoer G, Morava-Protzner I, Zbieranowski I, Bhattacharya N, Catzavelos GC, Klotz LH, Slingerland JM (1998) Loss of cyclin-dependent kinase inhibitor p27Kip1 is a novel prognostic factor in localized human prostate adenocarcinoma. *Cancer Res* 58(3):542–548
- Varambally S, Dhanasekaran S, Zhou M, Barrette T, Kumar-Sinha C, Sanda M, Ghosh D, Pienta K, Sewalt R, Otte A (2002) The polycomb group protein EZH2 is involved in progression of prostate cancer. *Nature* 419(6907):624–629
- Varambally S, Cao Q, Mani RS, Shankar S, Wang X, Ateeq B, Laxman B, Cao X, Jing X, Ramnarayanan K, Brenner JC, Yu J, Kim JH, Han B, Tan P, Kumar-Sinha C, Lonigro RJ, Palanisamy N, Maher CA, Chinnaiyan AM (2008) Genomic loss of microRNA-101 leads to overexpression of histone methyltransferase EZH2 in cancer. *Science* 322(5908):1695–1699
- Verkaik NS, Trapman J, Romijn JC, Van der Kwast TH, Van Steenbrugge GJ (1999) Down-regulation of CD44 expression in human prostatic carcinoma cell lines is correlated with DNA hypermethylation. *Int J Cancer* 80(3):439–443

- Verkaik NS, van Steenbrugge GJ, van Weerden WM, Bussemakers MJ, van der Kwast TH (2000) Silencing of CD44 expression in prostate cancer by hypermethylation of the CD44 promoter region. *Lab Invest* 80(8):1291–1298
- Vidanes GM, Paton V, Wallen E, Peehl DM, Navone N, Brooks JD (2002) Silencing of pi-class glutathione S-transferase in MDA PCa 2a and MDA PCa 2b cells. *Prostate* 51(4):225–230
- Villers A, Soulie M, Haillot O, Boccon-Gibod L (1997) Prostate cancer screening (III): risk factors, natural history, course without treatment. Characteristics of detected cancers. *Prog Urol* 7(4):655–661
- Visvader JE, Lindeman GJ (2008) Cancer stem cells in solid tumours: accumulating evidence and unresolved questions. *Nat Rev Cancer* 8(10):755–768
- Vlodavsky I, Michaeli RI, Bar-Ner M, Fridman R, Horowitz AT, Fuks Z, Biran S (1988) Involvement of heparanase in tumor metastasis and angiogenesis. *Isr J Med Sci* 24(9–10):464–470
- Volinia S, Calin GA, Liu CG, Ambros S, Cimmino A, Petrocca F, Visone R, Iorio M, Roldo C, Ferracin M, Prueitt RL, Yanaihara N, Lanza G, Scarpa A, Vecchione A, Negrini M, Harris CC, Croce CM (2006) A microRNA expression signature of human solid tumors defines cancer gene targets. *Proc Natl Acad Sci USA* 103(7):2257–2261
- Wach S, Nolte E, Szczyrba J, Stohr R, Hartmann A, Orntoft T, Dyrskjot L, Eltze E, Wieland W, Keck B, Ekici AB, Grasser F, Wullich B (2011) MiRNA profiles of prostate carcinoma detected by multi-platform miRNA screening. *Int J Cancer*. doi:10.1002/ijc.26064
- Walker C, Nettesheim P (1986) In vitro transformation of primary rat tracheal epithelial cells by 5-azacytidine. *Cancer Res* 46(12 Pt 1):6433–6437
- Wältregny D, North B, Van Mellaert F, de Leval J, Verdin E, Castronovo V (2004) Screening of histone deacetylases (HDAC) expression in human prostate cancer reveals distinct class I HDAC profiles between epithelial and stromal cells. *Eur J Histochem* 48(3):273–290
- Wang G, Platt-Higgins A, Carroll J, de Silva RS, Winstanley J, Barraclough R, Rudland P (2006) Induction of metastasis by S100P in a rat mammary model and its association with poor survival of breast cancer patients. *Cancer Res* 66(2):1199–1207
- Wang Q, Williamson M, Bott S, Brookman-Amisshah N, Freeman A, Nariculam J, Hubank MJ, Ahmed A, Masters JR (2007) Hypomethylation of WNT5A, CRIP1 and S100P in prostate cancer. *Oncogene* 26(45):6560–6565
- Wang YE, Wu X, Suzuki H, Reiter RE, Tran C, Vessella RL, Said JW, Isaacs WB, Sawyers CL (1998) Inactivation of the tumor suppressor PTEN/MMAC1 in advanced human prostate cancer through loss of expression. *Proc Natl Acad Sci USA* 95(9):5246–5250
- Wissmann M, Yin N, Muller JM, Greschik H, Fodor BD, Jenuwein T, Vogler C, Schneider R, Gunther T, Buettner R, Metzger E, Schule R (2007) Cooperative demethylation by JMJD2C and LSD1 promotes androgen receptor-dependent gene expression. *Nat Cell Biol* 9(3):347–353
- Xu G, Wu J, Zhou L, Chen B, Sun Z, Zhao F, Tao Z (2010) Characterization of the small RNA transcriptomes of androgen dependent and independent prostate cancer cell line by deep sequencing. *PLoS One* 5(11):e15519
- Yamane K, Toumazou C, Tsukada Y, Erdjument-Bromage H, Tempst P, Wong J, Zhang Y (2006) JHDM2A, a JmJc-containing H3K9 demethylase, facilitates transcription activation by androgen receptor. *Cell* 125(3):483–495
- Yang RM, Naitoh J, Murphy M, Wang HJ, Phillipson J, deKernion JB, Loda M, Reiter RE (1998) Low p27 expression predicts poor disease-free survival in patients with prostate cancer. *J Urol* 159(3):941–945
- Yegnasubramanian S, Haffner MC, Zhang Y, Gurel B, Cornish TC, Wu Z, Irizarry RA, Morgan J, Hicks J, DeWeese TL, Isaacs WB, Bova GS, De Marzo AM, Nelson WG (2008) DNA hypomethylation arises later in prostate cancer progression than CpG island hypermethylation and contributes to metastatic tumor heterogeneity. *Cancer Res* 68(21):8954–8967
- Yu J, Rhodes D, Tomlins S, Cao X, Chen G, Mehra R, Wang X, Ghosh D, Shah R (2007) A polycomb repression signature in metastatic prostate cancer predicts cancer outcome. *Cancer Res* 67(22):10657–10663
- Zhang Z, Stanfield J, Frenkel E, Kabbani W, Hsieh JT (2007) Enhanced therapeutic effect on androgen-independent prostate cancer by depsipeptide (FK228), a histone deacetylase inhibitor, in combination with docetaxel. *Urology* 70(2):396–401
- Zorn CS, Wojno KJ, McCabe MT, Kuefer R, Gschwend JE, Day ML (2007) 5-Aza-2'-deoxycytidine delays androgen-independent disease and improves survival in the transgenic adenocarcinoma of the mouse prostate mouse model of prostate cancer. *Clin Cancer Res* 13(7):2136–2143