

Hindawi Publishing Corporation
Advances in Difference Equations
Volume 2009, Article ID 734090, 16 pages
doi:10.1155/2009/734090

Research Article

Nonlocal Controllability for the Semilinear Fuzzy Integro-differential Equations in n -Dimensional Fuzzy Vector Space

Young Chel Kwun,¹ Jeong Soon Kim,¹ Min Ji Park,¹
and Jin Han Park²

¹ Department of Mathematics, Dong-A University, Pusan 604-714, South Korea

² Division of Mathematics Sciences, Pukyong National University, Pusan 608-737, South Korea

Correspondence should be addressed to Jin Han Park, jihpark@pknu.ac.kr

Received 23 February 2009; Revised 20 June 2009; Accepted 3 August 2009

Recommended by Točka Diagana

We study the existence and uniqueness of solutions and controllability for the semilinear fuzzy integro-differential equations in n -dimensional fuzzy vector space $(E_N)^n$ by using Banach fixed point theorem, that is, an extension of the result of J. H. Park et al. to n -dimensional fuzzy vector space.

Copyright © 2009 Young Chel Kwun et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

Many authors have studied several concepts of fuzzy systems. Diamond and Kloeden [1] proved the fuzzy optimal control for the following system:

$$\dot{x}(t) = a(t)x(t) + u(t), \quad x(0) = x_0, \quad (1.1)$$

where $x(\cdot)$ and $u(\cdot)$ are nonempty compact interval-valued functions on E^1 . Kwun and Park [2] proved the existence of fuzzy optimal control for the nonlinear fuzzy differential system with nonlocal initial condition in E_N^1 by using Kuhn-Tucker theorems. Fuzzy integro-differential equations are a field of interest, due to their applicability to the analysis of phenomena with memory where imprecision is inherent. Balasubramaniam and Muralisankar [3] proved the existence and uniqueness of fuzzy solutions for the semilinear fuzzy integro-differential equation with nonlocal initial condition. They considered the semilinear one-dimensional heat equation on a connected domain $(0, 1)$ for material with

memory. In one-dimensional fuzzy vector space E_N^1 , Park et al. [4] proved the existence and uniqueness of fuzzy solutions and presented the sufficient condition of nonlocal controllability for the following semilinear fuzzy integrodifferential equation with nonlocal initial condition:

$$\begin{aligned} \frac{dx(t)}{dt} &= A \left[x(t) + \int_0^t G(t-s)x(s)ds \right] + f(t, x) + u(t), \quad t \in J = [0, T], \\ x(0) + g(t_1, t_2, \dots, t_p, x(t_m)) &= x_0 \in E_N, \quad m = 1, 2, \dots, p, \end{aligned} \quad (1.2)$$

where $T > 0$, $A : J \rightarrow E_N$ is a fuzzy coefficient, E_N is the set of all upper semicontinuous convex normal fuzzy numbers with bounded α -level intervals, $f : J \times E_N \rightarrow E_N$ is a nonlinear continuous function, $g : J^p \times E_N \rightarrow E_N$ is a nonlinear continuous function, $G(t)$ is an $n \times n$ continuous matrix such that $dG(t)x/dt$ is continuous for $x \in E_N$ and $t \in J$ with $\|G(t)\| \leq K$, $K > 0$, with all nonnegative elements, $u : J \rightarrow E_N$ is control function.

In [5], Kwun et al. proved the existence and uniqueness of fuzzy solutions for the semilinear fuzzy integrodifferential equations by using successive iteration. In [6], Kwun et al. investigated the continuously initial observability for the semilinear fuzzy integrodifferential equations. Bede and Gal [7] studied almost periodic fuzzy-number-valued functions. Gal and N'Guérékata [8] studied almost automorphic fuzzy-number-valued functions.

In this paper, we study the the existence and uniqueness of solutions and controllability for the following semilinear fuzzy integrodifferential equations:

$$\begin{aligned} \frac{dx_i(t)}{dt} &= A_i \left[x_i(t) + \int_0^t G(t-s)x_i(s)ds \right] + f_i(t, x_i(t)) + u_i(t) \text{ on } E_N^i, \\ x_i(0) + g_i(x_i) &= x_{0_i} \in E_N^i \quad (i = 1, 2, \dots, n), \end{aligned} \quad (1.3)$$

where $A_i : [0, T] \rightarrow E_N^i$ is fuzzy coefficient, E_N^i is the set of all upper semicontinuously convex fuzzy numbers on R with $E_N^i \neq E_N^j$ ($i \neq j$), $f_i : [0, T] \times E_N^i \rightarrow E_N^i$ is a nonlinear regular fuzzy function, $g_i : E_N^i \rightarrow E_N^i$ is a nonlinear continuous function, $G(t)$ is $n \times n$ continuous matrix such that $dG(t)x_i/dt$ is continuous for $x_i \in E_N^i$ and $t \in [0, T]$ with $\|G(t)\| \leq k$, $k > 0$, $u_i : [0, T] \rightarrow E_N^i$ is control function and $x_{0_i} \in E_N^i$ is initial value.

2. Preliminaries

A fuzzy set of R^n is a function $u : R^n \rightarrow [0, 1]$. For each fuzzy set u , we denote by $[u]^\alpha = \{x \in R^n : u(x) \geq \alpha\}$ for any $\alpha \in [0, 1]$, its α -level set.

Let u, v be fuzzy sets of R^n . It is well known that $[u]^\alpha = [v]^\alpha$ for each $\alpha \in [0, 1]$ implies $u = v$.

Let E^n denote the collection of all fuzzy sets of R^n that satisfies the following conditions:

- (1) u is normal, that is, there exists an $x_0 \in R^n$ such that $u(x_0) = 1$;
- (2) u is fuzzy convex, that is, $u(\lambda x + (1 - \lambda)y) \geq \min\{u(x), u(y)\}$ for any $x, y \in R^n$, $0 \leq \lambda \leq 1$;

(3) $u(x)$ is upper semicontinuous, that is, $u(x_0) \geq \overline{\lim}_{k \rightarrow \infty} u(x_k)$ for any $x_k \in R^n$ ($k = 0, 1, 2, \dots$), $x_k \rightarrow x_0$;

(4) $[u]^0$ is compact.

We call $u \in E^n$ an n -dimension fuzzy number.

Wang et al. [9] defined n -dimensional fuzzy vector space and investigated its properties.

For any $u_i \in E$, $i = 1, 2, \dots, n$, we call the ordered one-dimension fuzzy number class u_1, u_2, \dots, u_n (i.e., the Cartesian product of one-dimension fuzzy number u_1, u_2, \dots, u_n) an n -dimension fuzzy vector, denote it as (u_1, u_2, \dots, u_n) , and call the collection of all n -dimension fuzzy vectors (i.e., the Cartesian product $\overbrace{E \times E \times \dots \times E}^n$) n -dimensional fuzzy vector space, and denote it as $(E)^n$.

Definition 2.1 (see [9]). If $u \in E^n$, and $[u]^\alpha$ is a hyperrectangle, that is, $[u]^\alpha$ can be represented by $\prod_{i=1}^n [u_{il}^\alpha, u_{ir}^\alpha]$, that is, $[u_{1l}^\alpha, u_{1r}^\alpha] \times [u_{2l}^\alpha, u_{2r}^\alpha] \times \dots \times [u_{nl}^\alpha, u_{nr}^\alpha]$ for every $\alpha \in [0, 1]$, where $u_{il}^\alpha, u_{ir}^\alpha \in R$ with $u_{il}^\alpha \leq u_{ir}^\alpha$ when $\alpha \in (0, 1]$, $i = 1, 2, \dots, n$, then we call u a fuzzy n -cell number. We denote the collection of all fuzzy n -cell numbers by $L(E^n)$.

Theorem 2.2 (see [9]). For any $u \in L(E^n)$ with $[u]^\alpha = \prod_{i=1}^n [u_{il}^\alpha, u_{ir}^\alpha]$ ($\alpha \in [0, 1]$), there exists a unique $(u_1, u_2, \dots, u_n) \in (E)^n$ such that $[u_i]^\alpha = [u_{il}^\alpha, u_{ir}^\alpha]$ ($i = 1, 2, \dots, n$ and $\alpha \in [0, 1]$).

Conversely, for any $(u_1, u_2, \dots, u_n) \in (E)^n$ with $[u_i]^\alpha = [u_{il}^\alpha, u_{ir}^\alpha]$ ($i = 1, 2, \dots, n$ and $\alpha \in [0, 1]$), there exists a unique $u \in L(E^n)$ such that $[u]^\alpha = \prod_{i=1}^n [u_{il}^\alpha, u_{ir}^\alpha]$ ($\alpha \in [0, 1]$).

Note 1 (see [9]). Theorem 2.2 indicates that fuzzy n -cell numbers and n -dimension fuzzy vectors can represent each other, so $L(E^n)$ and $(E)^n$ may be regarded as identity. If $(u_1, u_2, \dots, u_n) \in (E)^n$ is the unique n -dimension fuzzy vector determined by $u \in L(E^n)$, then we denote $u = (u_1, u_2, \dots, u_n)$.

Let $(E_N^i)^n = E_N^1 \times E_N^2 \times \dots \times E_N^n$, E_N^i ($i = 1, 2, \dots, n$) be fuzzy subset of R . Then $(E_N^i)^n \subseteq (E)^n$.

Definition 2.3 (see [9]). The complete metric D_L on $(E_N^i)^n$ is defined by

$$\begin{aligned}
 D_L(u, v) &= \sup_{0 < \alpha \leq 1} d_L([u]^\alpha, [v]^\alpha) \\
 &= \sup_{0 < \alpha \leq 1} \max_{1 \leq i \leq n} \{ |u_{il}^\alpha - v_{il}^\alpha|, |u_{ir}^\alpha - v_{ir}^\alpha| \}
 \end{aligned}
 \tag{2.1}$$

for any $u, v \in (E_N^i)^n$, which satisfies $d_L(u + w, v + w) = d_L(u, v)$.

Definition 2.4. Let $u, v \in C([0, T] : (E_N^i)^n)$, then

$$H_1(u, v) = \sup_{0 \leq t \leq T} D_L(u(t), v(t)).
 \tag{2.2}$$

Definition 2.5 (see [9]). The derivative $x'(t)$ of a fuzzy process $x \in (E_N^i)^n$ is defined by

$$[x'(t)]^\alpha = \prod_{i=1}^n \left[(x_{il}^\alpha)'(t), (x_{ir}^\alpha)'(t) \right] \quad (2.3)$$

provided that the equation defines a fuzzy $x'(t) \in (E_N^i)^n$.

Definition 2.6 (see [9]). The fuzzy integral $\int_b^a x(t)dt$, $a, b \in [0, T]$ is defined by

$$\left[\int_b^a x(t)dt \right]^\alpha = \prod_{i=1}^n \left[\int_b^a x_{il}^\alpha(t)dt, \int_b^a x_{ir}^\alpha(t)dt \right] \quad (2.4)$$

provided that the Lebesgue integrals on the right-hand side exist.

3. Existence and Uniqueness

In this section we consider the existence and uniqueness of the fuzzy solution for (1.3) ($u \equiv 0$).

We define

$$\begin{aligned} A &= (A_1, A_2, \dots, A_n), \\ x &= (x_1, x_2, \dots, x_n), \\ f &= (f_1, f_2, \dots, f_n), \\ u &= (u_1, u_2, \dots, u_n), \\ g &= (g_1, g_2, \dots, g_n), \\ x_0 &= (x_{0_1}, x_{0_2}, \dots, x_{0_n}). \end{aligned} \quad (3.1)$$

Then

$$A, x, f, x_0, u, g \in (E_N^i)^n. \quad (3.2)$$

Instead of (1.3), we consider the following fuzzy integrodifferential equations in $(E_N^i)^n$:

$$\begin{aligned} \frac{dx(t)}{dt} &= A \left[x(t) + \int_0^t G(t-s)x(s)ds \right] + f(t, x(t)) + u(t) \text{ on } (E_N^i)^n \\ x(0) + g(x) &= x_0 \in (E_N^i)^n \end{aligned} \quad (3.3)$$

with fuzzy coefficient $A : [0, T] \rightarrow (E_N^i)^n$, initial value $x_0 \in (E_N^i)^n$, and $u : [0, T] \rightarrow (E_N^i)^n$ is a control function. Given nonlinear regular fuzzy function $f : [0, T] \times (E_N^i)^n \rightarrow (E_N^i)^n$ satisfies a global Lipschitz condition, that is, there exists a finite $k > 0$ such that

$$d_L([f(s, x(s))]^\alpha, [f(s, y(s))]^\alpha) \leq kd_L([x(s)]^\alpha, [y(s)]^\alpha) \quad (3.4)$$

for all $x(s), y(s) \in (E_N^i)^n$. The nonlinear function $g : (E_N^i)^n \rightarrow (E_N^i)^n$ is a continuous function and satisfies the Lipschitz condition

$$d_L([g(x(\cdot))]^\alpha, [g(y(\cdot))]^\alpha) \leq hd_L([x(\cdot)]^\alpha, [y(\cdot)]^\alpha) \tag{3.5}$$

for all $x(\cdot), y(\cdot) \in (E_N^i)^n$, h is a finite positive constant.

Definition 3.1. The fuzzy process $x : I = [0, T] \rightarrow (E_N^i)^n$ with α -level set $[x(t)]^\alpha = \Pi_{i=1}^n [x_i]^\alpha = \Pi_{i=1}^n [x_{il}^\alpha, x_{ir}^\alpha]$ is a fuzzy solution of (3.3) without nonhomogeneous term if and only if

$$\begin{aligned} (x_{il}^\alpha)'(t) &= \min \left\{ A_{ij}^\alpha(t) \left[x_{ik}^\alpha(t) + \int_0^t G(t-s)x_{ik}^\alpha(s)ds \right] : j, k = l, r \right\}, \\ (x_{ir}^\alpha)'(t) &= \max \left\{ A_{ij}^\alpha(t) \left[x_{ik}^\alpha(t) + \int_0^t G(t-s)x_{ik}^\alpha(s)ds \right] : j, k = l, r \right\}, \\ x_{il}^\alpha(0) + g_{il}^\alpha(x_{il}^\alpha) &= x_{0il}^\alpha, \quad x_{ir}^\alpha(0) + g_{ir}^\alpha(x_{ir}^\alpha) = x_{0ir}^\alpha, \quad i = 1, 2, \dots, n. \end{aligned} \tag{3.6}$$

For the sequel, we need the following assumptions.

(H1) $S(t)$ is a fuzzy number satisfying, for $y \in (E_N^i)^n$, $(d/dt) S(t)y \in C^1(I : (E_N^i)^n) \cap C(I : (E_N^i)^n)$, the equation

$$\begin{aligned} \frac{d}{dt} S(t)y &= A \left[S(t)y + \int_0^t G(t-s)S(s)y ds \right] \\ &= S(t)Ay + \int_0^t S(t-s)AG(s)y ds, \quad t \in I, \end{aligned} \tag{3.7}$$

where

$$[S(t)]^\alpha = \prod_{i=1}^n [S_i(t)]^\alpha = \prod_{i=1}^n [S_{il}^\alpha(t), S_{ir}^\alpha(t)], \tag{3.8}$$

and $S_{ij}^\alpha(t)$ ($j = l, r$) is continuous with $|S_{ij}^\alpha(t)| \leq c, c > 0$, for all $t \in I = [0, T]$.

(H2) $c\{h(1 + T + cT) + kT(1 + cT)\} < 1$.

In view of Definition 3.1 and (H1), (3.3) can be expressed as

$$\begin{aligned} x(t) &= S(t)(x_0 - g(x)) + \int_0^t S(t-s)(f(s, x(s)) + u(s))ds, \\ x(0) + g(x) &= x_0. \end{aligned} \tag{3.9}$$

Theorem 3.2. Let $T > 0$. If hypotheses (H1)-(H2) are hold, then for every $x_0 \in (E_N^i)^n$, (3.9) ($u \equiv 0$) have a unique fuzzy solution $x \in C([0, T] : (E_N^i)^n)$.

Proof. For each $x(t) \in (E_N^i)^n$ and $t \in [0, T]$, define $(G_0x)(t) \in (E_N^i)^n$ by

$$(G_0x)(t) = S(t)(x_0 - g(x)) + \int_0^t S(t-s)f(s, x(s))ds. \quad (3.10)$$

Thus, $G_0x : [0, T] \rightarrow (E_N^i)^n$ is continuous, so G_0 is a mapping from $C([0, T] : (E_N^i)^n)$ into itself. By Definitions 2.3 and 2.4, some properties of d_L , and inequalities (3.4) and (3.5), we have following inequalities. For $x, y \in C([0, T] : (E_N^i)^n)$,

$$\begin{aligned} & d_L([(G_0x)(t)]^\alpha, [(G_0y)(t)]^\alpha) \\ &= d_L\left(\left[S(t)(x_0 - g(x)) + \int_0^t S(t-s)f(s, x(s))ds\right]^\alpha, \right. \\ &\quad \left.[S(t)(x_0 - g(y)) + \int_0^t S(t-s)f(s, y(s))ds\right]^\alpha\right) \\ &= d_L\left(\left[-S(t)g(x) + \int_0^t S(t-s)f(s, x(s))ds\right]^\alpha, \right. \\ &\quad \left.[-S(t)g(y) + \int_0^t S(t-s)f(s, y(s))ds\right]^\alpha\right) \\ &\leq d_L([S(t)g(x)]^\alpha, [S(t)g(y)]^\alpha) + \int_0^t d_L([S(t-s)f(s, x(s))]^\alpha, [S(t-s)f(s, y(s))]^\alpha)ds \\ &= \max_{1 \leq i \leq n} \{ |S_{il}^\alpha(t)(g_{il}^\alpha(x) - g_{il}^\alpha(y))|, |S_{ir}^\alpha(t)(g_{ir}^\alpha(x) - g_{ir}^\alpha(y))| \} \\ &\quad + \int_0^t \max_{1 \leq i \leq n} \{ |S_{il}^\alpha(t-s)(f_{il}^\alpha(s, x(s)) - f_{il}^\alpha(s, y(s)))|, |S_{ir}^\alpha(t-s)(f_{ir}^\alpha(s, x(s)) - f_{ir}^\alpha(s, y(s)))| \} ds \\ &\leq c \max_{1 \leq i \leq n} \{ |(g_{il}^\alpha(x) - g_{il}^\alpha(y))|, |(g_{ir}^\alpha(x) - g_{ir}^\alpha(y))| \} \\ &\quad + c \int_0^t \max_{1 \leq i \leq n} \{ |f_{il}^\alpha(s, x(s)) - f_{il}^\alpha(s, y(s))|, |f_{ir}^\alpha(s, x(s)) - f_{ir}^\alpha(s, y(s))| \} ds \\ &= cd_L([g(x)]^\alpha, [g(y)]^\alpha) + c \int_0^t d_L([f(s, x(s))]^\alpha, [f(s, y(s))]^\alpha)ds \\ &\leq chd_L([x(\cdot)]^\alpha, [y(\cdot)]^\alpha) + ck \int_0^t d_L([x(s)]^\alpha, [y(s)]^\alpha)ds. \end{aligned} \quad (3.11)$$

Therefore

$$\begin{aligned}
 &D_L((G_0x)(t), (G_0y)(t)) \\
 &= \sup_{0 < \alpha \leq 1} d_L([(G_0x)(t)]^\alpha, [(G_0y)(t)]^\alpha) \\
 &\leq ch \sup_{0 < \alpha \leq 1} d_L([x(\cdot)]^\alpha, [y(\cdot)]^\alpha) + ck \sup_{0 < \alpha \leq 1} \int_0^t d_L([x(s)]^\alpha, [y(s)]^\alpha) ds \quad (3.12) \\
 &\leq ch D_L(x(\cdot), y(\cdot)) + ck \int_0^t D_L(x(s), y(s)) ds.
 \end{aligned}$$

Hence

$$\begin{aligned}
 H_1(G_0x, G_0y) &= \sup_{0 \leq t \leq T} D_L((G_0x)(t), (G_0y)(t)) \\
 &\leq ch \sup_{0 \leq t \leq T} D_L(x(\cdot), y(\cdot)) + ck \sup_{0 \leq t \leq T} \int_0^t D_L(x(s), y(s)) ds \quad (3.13) \\
 &\leq ch H_1(x, y) + ckT H_1(x, y) \\
 &= c(h + kT) H_1(x, y).
 \end{aligned}$$

By hypothesis (H2), G_0 is a contraction mapping.

Using the Banach fixed point theorem, (3.9) have a unique fixed point $x \in C([0, T] : (E_N^i)^n)$. □

4. Controllability

In this section, we show the nonlocal controllability for the control system (1.3).

Definition 4.1. Equation (1.3) is nonlocal controllable. Then there exists $u(t)$ such that the fuzzy solution $x(t)$ for (3.9) as $x(T) = x^1 - g(x)$ (i.e., $[x(T)]^\alpha = [x^1 - g(x)]^\alpha$) where $x^1 \in (E_N^i)^n$ is target set.

Define the fuzzy mapping $\tilde{\beta} : \tilde{P}(R^n) \rightarrow (E_N^i)^n$ by

$$\tilde{\beta}^\alpha(v) = \begin{cases} \int_0^T S^\alpha(T-s)v(s)ds, & v \in \bar{\Gamma}_u, \\ 0, & \text{otherwise,} \end{cases} \quad (4.1)$$

where $\bar{\Gamma}_u$ is closed support of u . Then there exists

$$\tilde{\beta}_i : \tilde{P}(R) \longrightarrow E_N^i \quad (i = 1, 2, \dots, n) \quad (4.2)$$

such that

$$\tilde{\beta}_i^\alpha(v_i) = \begin{cases} \int_0^T S_i^\alpha(T-s)v_i(s)ds, & v_i(s) \subset \bar{\Gamma}_{u_i}, \\ 0, & \text{otherwise.} \end{cases} \quad (4.3)$$

Then $\tilde{\beta}_{ij}^\alpha$ ($j = l, r$) exists such that

$$\begin{aligned} \tilde{\beta}_{il}^\alpha(v_{il}) &= \int_0^T S_{il}^\alpha(T-s)v_{il}(s)ds, \quad v_{il}(s) \in [u_{il}^\alpha(s), u_i^1], \\ \tilde{\beta}_{ir}^\alpha(v_{ir}) &= \int_0^T S_{ir}^\alpha(T-s)v_{ir}(s)ds, \quad v_{ir}(s) \in [u_i^1, u_{ir}^\alpha(s)]. \end{aligned} \quad (4.4)$$

We assume that $\tilde{\beta}_{il}^\alpha, \tilde{\beta}_{ir}^\alpha$ are bijective mappings.

We can introduce α -level set of $u(s)$ of (3.4)-(3.5)

$$\begin{aligned} [u(s)]^\alpha &= \prod_{i=1}^n [u_i(s)]^\alpha \\ &= \prod_{i=1}^n [u_{il}^\alpha(s), u_{ir}^\alpha(s)] \\ &= \prod_{i=1}^n \left[(\tilde{\beta}_{il}^\alpha)^{-1} \left(\left((x^1)_{il}^\alpha - g_{il}^\alpha(x_{il}^\alpha) \right) - S_{il}^\alpha(T) \left(x_{0il}^\alpha - g_{il}^\alpha(x_{il}^\alpha) \right) \right. \right. \\ &\quad \left. \left. - \int_0^T S_{il}^\alpha(T-s)f_{il}^\alpha(s, x_{il}^\alpha(s))ds \right), \right. \\ &\quad \left. (\tilde{\beta}_{ir}^\alpha)^{-1} \left(\left((x^1)_{ir}^\alpha - g_{ir}^\alpha(x_{ir}^\alpha) \right) - S_{ir}^\alpha(T) \left(x_{0ir}^\alpha - g_{ir}^\alpha(x_{ir}^\alpha) \right) \right. \right. \\ &\quad \left. \left. - \int_0^T S_{ir}^\alpha(T-s)f_{ir}^\alpha(s, x_{ir}^\alpha(s))ds \right) \right]. \end{aligned} \quad (4.5)$$

Then substituting this expression into (3.9) yields α -level of $x(T)$.

For each $i = 1, 2, \dots, n$,

$$\begin{aligned}
 [x_i(T)]^\alpha &= \left[S_{il}^\alpha(T) \left(x_{0il}^\alpha - g_{il}^\alpha(x_{il}^\alpha) \right) + \int_0^T S_{il}^\alpha(T-s) f_{il}^\alpha(s, x_{il}^\alpha(s)) ds \right. \\
 &\quad + \int_0^T S_{il}^\alpha(T-s) \left(\tilde{\beta}_{il}^\alpha \right)^{-1} \left(\left((x^1)_{il}^\alpha - g_{il}^\alpha(x_{il}^\alpha) \right) - S_{il}^\alpha(T) \left(x_{0il}^\alpha - g_{il}^\alpha(x_{il}^\alpha) \right) \right. \\
 &\quad \left. \left. - \int_0^T S_{il}^\alpha(T-s) f_{il}^\alpha(s, x_{il}^\alpha(s)) ds \right) ds, \right. \\
 &\quad S_{ir}^\alpha(T) \left(x_{0ir}^\alpha - g_{ir}^\alpha(x_{ir}^\alpha) \right) + \int_0^T S_{ir}^\alpha(T-s) f_{ir}^\alpha(s, x_{ir}^\alpha(s)) ds \\
 &\quad \left. + \int_0^T S_{ir}^\alpha(T-s) \left(\tilde{\beta}_{ir}^\alpha \right)^{-1} \left(\left((x^1)_{ir}^\alpha - g_{ir}^\alpha(x_{ir}^\alpha) \right) - S_{ir}^\alpha(T) \left(x_{0ir}^\alpha - g_{ir}^\alpha(x_{ir}^\alpha) \right) \right. \right. \\
 &\quad \left. \left. - \int_0^T S_{ir}^\alpha(T-s) f_{ir}^\alpha(s, x_{ir}^\alpha(s)) ds \right) ds \right] \\
 &= \left[(x^1 - g(x))_{il}^\alpha, (x^1 - g(x))_{ir}^\alpha \right] = \left[(x^1 - g(x))_i \right]^\alpha.
 \end{aligned} \tag{4.6}$$

Therefore

$$[x(T)]^\alpha = \prod_{i=1}^n [x_i(T)]^\alpha = \prod_{i=1}^n \left[(x^1 - g(x))_i \right]^\alpha = \left[x^1 - g(x) \right]^\alpha. \tag{4.7}$$

We now set

$$\begin{aligned}
 \Phi x(t) &= S(t)(x_0 - g(x)) + \int_0^t S(t-s) f(s, x(s)) ds \\
 &\quad + \int_0^t S(t-s) \tilde{\beta}^{-1} \left(x^1 - g(x) - S(T)(x_0 - g(x)) - \int_0^T S(T-s) f(s, x(s)) ds \right) ds,
 \end{aligned} \tag{4.8}$$

where the fuzzy mapping $\tilde{\beta}^{-1}$ satisfies above statements.

Notice that $\Phi x(T) = x^1 - g(x)$, which means that the control $u(t)$ steers (3.9) from the origin to $x^1 - g(x)$ in time T provided that we can obtain a fixed point of the operator Φ .

(H3) Assume that the linear system of (3.9) ($f \equiv 0$) is controllable.

Theorem 4.2. *Suppose that hypotheses (H1)–(H3) are satisfied. Then (3.9) are nonlocal controllable.*

Proof. We can easily check that Φ is continuous function from $C([0, T] : (E_N^i)^n)$ to itself. By Definitions 2.3 and 2.4, some properties of d_L , and inequalities (3.4) and (3.5), we have the following inequalities. For any $x, y \in C([0, T] : (E_N^i)^n)$,

$$\begin{aligned}
& d_L([\Phi x(t)]^\alpha, [\Phi y(t)]^\alpha) \\
&= d_L\left(\left[S(t)(x_0 - g(x)) + \int_0^t S(t-s)f(s, x(s))ds + \int_0^t S(t-s)\tilde{\beta}^{-1}\right.\right. \\
&\quad \left.\left.\times\left(x^1 - g(x) - S(T)(x_0 - g(x)) - \int_0^T S(T-s)f(s, x(s))ds\right)ds\right]^\alpha, \right. \\
&\quad \left.[S(t)(x_0 - g(y)) + \int_0^t S(t-s)f(s, y(s))ds + \int_0^t S(t-s)\tilde{\beta}^{-1}\right. \\
&\quad \left.\times\left(x^1 - g(y) - S(T)(x_0 - g(y)) - \int_0^T S(T-s)f(s, y(s))ds\right)ds\right]^\alpha\bigg) \\
&\leq d_L([S(t)g(x)]^\alpha, [S(t)g(y)]^\alpha) + \int_0^t d_L([S(t-s)f(s, x(s))]^\alpha, [S(t-s)f(s, y(s))]^\alpha)ds \\
&\quad + \int_0^t d_L([S(t-s)\tilde{\beta}^{-1}g(x)]^\alpha, [S(t-s)\tilde{\beta}^{-1}g(y)]^\alpha)ds \\
&\quad + \int_0^t d_L([S(t-s)\tilde{\beta}^{-1}S(T)g(x)]^\alpha, [S(t-s)\tilde{\beta}^{-1}S(T)g(y)]^\alpha)ds \\
&\quad + \int_0^t d_L\left(\left[S(t-s)\tilde{\beta}^{-1}\int_0^T S(T-s)f(s, x(s))ds\right]^\alpha, \left[S(t-s)\tilde{\beta}^{-1}\int_0^T S(T-s)f(s, y(s))ds\right]^\alpha\right)ds \\
&= \max_{1 \leq i \leq n} \{|S_{il}^\alpha(t)(g_{il}^\alpha(x) - g_{il}^\alpha(y))|, |S_{ir}^\alpha(t)(g_{ir}^\alpha(x) - g_{ir}^\alpha(y))|\} \\
&\quad + \int_0^t \max_{1 \leq i \leq n} \{|S_{il}^\alpha(t-s)(f_{il}^\alpha(s, x(s)) - f_{il}^\alpha(s, y(s)))|, |S_{ir}^\alpha(t-s)(f_{ir}^\alpha(s, x(s)) - f_{ir}^\alpha(s, y(s)))|\} ds \\
&\quad + \int_0^t \max_{1 \leq i \leq n} \left\{ \left| S_{il}^\alpha(t-s)(\tilde{\beta}_{il}^\alpha)^{-1}(g_{il}^\alpha(x) - g_{il}^\alpha(y)) \right|, \left| S_{ir}^\alpha(t-s)(\tilde{\beta}_{ir}^\alpha)^{-1}(g_{ir}^\alpha(x) - g_{ir}^\alpha(y)) \right| \right\} ds \\
&\quad + \int_0^t \max_{1 \leq i \leq n} \left\{ \left| S_{il}^\alpha(t-s)(\tilde{\beta}_{il}^\alpha)^{-1} S_{il}^\alpha(T)(g_{il}^\alpha(x) - g_{il}^\alpha(y)) \right|, \right. \\
&\quad \quad \left. \left| S_{ir}^\alpha(t-s)(\tilde{\beta}_{ir}^\alpha)^{-1} S_{ir}^\alpha(T)(g_{ir}^\alpha(x) - g_{ir}^\alpha(y)) \right| \right\} ds \\
&\quad + \int_0^t \max_{1 \leq i \leq n} \left\{ \left| S_{il}^\alpha(t-s)(\tilde{\beta}_{il}^\alpha)^{-1} \left(\int_0^T S_{il}^\alpha(T-s)f_{il}^\alpha(s, x(s))ds - \int_0^T S_{il}^\alpha(T-s)f_{il}^\alpha(s, y(s))ds \right) \right|, \right. \\
&\quad \quad \left. \left| S_{ir}^\alpha(t-s)(\tilde{\beta}_{ir}^\alpha)^{-1} \left(\int_0^T S_{ir}^\alpha(T-s)f_{ir}^\alpha(s, x(s))ds - \int_0^T S_{ir}^\alpha(T-s)f_{ir}^\alpha(s, y(s))ds \right) \right| \right\} ds
\end{aligned}$$

$$\begin{aligned}
 &\leq c \max_{1 \leq i \leq n} \{ |g_{il}^\alpha(x) - g_{il}^\alpha(y)|, |g_{ir}^\alpha(x) - g_{ir}^\alpha(y)| \} \\
 &\quad + c \int_0^t \max_{1 \leq i \leq n} \{ |f_{il}^\alpha(s, x(s)) - f_{il}^\alpha(s, y(s))|, |f_{ir}^\alpha(s, x(s)) - f_{ir}^\alpha(s, y(s))| \} ds \\
 &\quad + c \int_0^t \max_{1 \leq i \leq n} \{ |g_{il}^\alpha(x) - g_{il}^\alpha(y)|, |g_{ir}^\alpha(x) - g_{ir}^\alpha(y)| \} ds \\
 &\quad + c^2 \int_0^t \max_{1 \leq i \leq n} \{ |g_{il}^\alpha(x) - g_{il}^\alpha(y)|, |g_{ir}^\alpha(x) - g_{ir}^\alpha(y)| \} ds \\
 &\quad + c^2 \int_0^t \int_0^T \max_{1 \leq i \leq n} \{ |f_{il}^\alpha(s, x(s)) - f_{il}^\alpha(s, y(s))|, |f_{ir}^\alpha(s, x(s)) - f_{ir}^\alpha(s, y(s))| \} ds ds \\
 &= c d_L([g(x)]^\alpha, [g(y)]^\alpha) + c \int_0^t d_L([f(s, x(s))]^\alpha, [f(s, y(s))]^\alpha) ds \\
 &\quad + c \int_0^t d_L([g(x)]^\alpha, [g(y)]^\alpha) ds + c^2 \int_0^t d_L([g(x)]^\alpha, [g(y)]^\alpha) ds \\
 &\quad + c^2 \int_0^t \int_0^T d_L([f(s, x(s))]^\alpha, [f(s, y(s))]^\alpha) ds ds \\
 &\leq ch \left\{ d_L([x(\cdot)]^\alpha, [y(\cdot)]^\alpha) + (1+c) \int_0^t d_L([x(\cdot)]^\alpha, [y(\cdot)]^\alpha) ds \right\} \\
 &\quad + ck \left\{ \int_0^t d_L([x(s)]^\alpha, [y(s)]^\alpha) ds + c \int_0^t \int_0^T d_L([x(s)]^\alpha, [y(s)]^\alpha) ds ds \right\}.
 \end{aligned} \tag{4.9}$$

Therefore

$$\begin{aligned}
 &D_L(\Phi x(t), \Phi y(t)) \\
 &= \sup_{0 < \alpha \leq 1} d_L([\Phi x(t)]^\alpha, [\Phi y(t)]^\alpha) \\
 &\leq ch \left\{ \sup_{0 < \alpha \leq 1} d_L([x(\cdot)]^\alpha, [y(\cdot)]^\alpha) + (1+c) \int_0^t \sup_{0 < \alpha \leq 1} d_L([x(\cdot)]^\alpha, [y(\cdot)]^\alpha) ds \right\} \\
 &\quad + ck \left\{ \int_0^t \sup_{0 < \alpha \leq 1} d_L([x(s)]^\alpha, [y(s)]^\alpha) ds + c \int_0^t \int_0^T \sup_{0 < \alpha \leq 1} d_L([x(s)]^\alpha, [y(s)]^\alpha) ds ds \right\} \tag{4.10} \\
 &= ch \left\{ D_L(x(\cdot), y(\cdot)) + (1+c) \int_0^t D_L(x(\cdot), y(\cdot)) ds \right\} \\
 &\quad + ck \left\{ \int_0^t D_L(x(s), y(s)) ds + c \int_0^t \int_0^T D_L(x(s), y(s)) ds ds \right\}.
 \end{aligned}$$

Hence

$$\begin{aligned}
H_1(\Phi x, \Phi y) &= \sup_{0 \leq t \leq T} D_L(\Phi x(t), \Phi y(t)) \\
&\leq ch \left\{ \sup_{0 \leq t \leq T} D_L(x(\cdot), y(\cdot)) + (1+c) \sup_{0 \leq t \leq T} \int_0^t D_L(x(\cdot), y(\cdot)) ds \right\} \\
&\quad + ck \left\{ \sup_{0 \leq t \leq T} \int_0^t D_L(x(s), y(s)) ds + c \sup_{0 \leq t \leq T} \int_0^t \int_0^T D_L(x(s), y(s)) ds ds \right\} \quad (4.11) \\
&\leq ch \{ H_1(x, y) + (1+c)T H_1(x, y) \} + ck \{ T H_1(x, y) + cT^2 H_1(x, y) \} \\
&= c \{ h(1+T+cT) + kT(1+cT) \} H_1(x, y).
\end{aligned}$$

By hypothesis (H2), Φ is a contraction mapping. Using the Banach fixed point theorem, (4.8) has a unique fixed point $x \in C([0, T] : (E_N^i)^n)$. \square

5. Example

Consider the two semilinear one-dimensional heat equations on a connected domain $(0, 1)$ for material with memory on E_N^i , $i = 1, 2$, boundary condition $x_i(t, 0) = x_i(t, 1) = 0$, $i = 1, 2$ and with initial conditions $x_i(0, z_i) + \sum_{k=1}^p (c_k)_i x_i(t_k, z_i) = x_{0i}(z_i)$, where $x_{0i}(z_i) \in E_N^i$, $\sum_{k=1}^p (c_k)_i x_i(t_k, z_i) = g_i(x_i)$, $i = 1, 2$. Let $x_i(t, z_i)$, $i = 1, 2$, be the internal energy and let $f_i(t, x_i(t, z_i)) = \tilde{2}tx_i(t, z_i)^2$, $i = 1, 2$, be the external heat.

Let

$$\begin{aligned}
A &= (A_1, A_2) = \left(\tilde{2} \frac{\partial^2}{\partial z_1^2}, \tilde{2} \frac{\partial^2}{\partial z_2^2} \right), \\
f(t, x(t)) &= (f_1(t, x_1(t)), f_2(t, x_2(t))) = \left(\tilde{2}tx_1(t, z_1)^2, \tilde{2}tx_2(t, z_2)^2 \right), \\
g(x) &= (g_1(x_1), g_2(x_2)) = \left(\sum_{k=1}^p (c_k)_1 x_1(t_k, z_1), \sum_{k=1}^p (c_k)_2 x_2(t_k, z_2) \right), \quad (5.1) \\
x(0) + g(x) &= (x_1(0) + g_1(x), x_2(0) + g_2(x)), \quad x_0 = (x_{01}, x_{02}) = (\tilde{0}, \tilde{0}), \\
G(t-s) &= (e^{-(t-s)}, e^{-(t-s)}),
\end{aligned}$$

then the balance equations become

$$\begin{aligned}
\frac{dx(t)}{dt} &= A \left[x(t) + \int_0^t G(t-s)x(s) ds \right] + f(t, x(t)) \text{ on } (E_N^i)^2, \quad (5.2) \\
x(0) + g(x) &= x_0 \in (E_N^i)^2.
\end{aligned}$$

The α -level sets of fuzzy numbers are the following: $[\tilde{0}]^\alpha = [\alpha - 1, 1 - \alpha]$, $[\tilde{2}]^\alpha = [\alpha + 1, 3 - \alpha]$ for all $\alpha \in [0, 1]$. Then α -level set of $f(t, x(t))$ is

$$\begin{aligned}
 & [f(t, x(t))]^\alpha \\
 &= [\tilde{2}tx_1(t)^2]^\alpha \times [\tilde{2}tx_2(t)^2]^\alpha \\
 &= [\tilde{2}]^\alpha \cdot t[x_1(t)^2]^\alpha \times [\tilde{2}]^\alpha \cdot t[x_2(t)^2]^\alpha \\
 &= [\alpha + 1, 3 - \alpha] \cdot t[(x_{1l}^\alpha(t))^2, (x_{1r}^\alpha(t))^2] \times [\alpha + 1, 3 - \alpha] \cdot t[(x_{2l}^\alpha(t))^2, (x_{2r}^\alpha(t))^2] \\
 &= [(\alpha + 1)t(x_{1l}^\alpha(t))^2, (3 - \alpha)t(x_{1r}^\alpha(t))^2] \times [(\alpha + 1)t(x_{2l}^\alpha(t))^2, (3 - \alpha)t(x_{2r}^\alpha(t))^2].
 \end{aligned} \tag{5.3}$$

Further, we have

$$\begin{aligned}
 & d_L([f(t, x(t))]^\alpha, f(t, y(t))^\alpha) \\
 &= d_L\left([\alpha + 1)t(x_{il}^\alpha(t))^2, (3 - \alpha)t(x_{ir}^\alpha(t))^2], [(\alpha + 1)t(y_{il}^\alpha(t))^2, (3 - \alpha)t(y_{ir}^\alpha(t))^2]\right) \\
 &= t \max_{1 \leq i \leq 2} \left\{ (\alpha + 1) \left| (x_{il}^\alpha(t))^2 - (y_{il}^\alpha(t))^2 \right|, (3 - \alpha) \left| (x_{ir}^\alpha(t))^2 - (y_{ir}^\alpha(t))^2 \right| \right\} \\
 &\leq T(3 - \alpha) \max_{1 \leq i \leq 2} \left\{ |x_{il}^\alpha(t) - y_{il}^\alpha(t)| |x_{il}^\alpha(t) + y_{il}^\alpha(t)|, |x_{ir}^\alpha(t) - y_{ir}^\alpha(t)| |x_{ir}^\alpha(t) + y_{ir}^\alpha(t)| \right\} \\
 &\leq 3T |x_{ir}^\alpha(t) + y_{ir}^\alpha(t)| \times \max_{1 \leq i \leq 2} \left\{ |x_{il}^\alpha(t) - y_{il}^\alpha(t)|, |x_{ir}^\alpha(t) - y_{ir}^\alpha(t)| \right\} \\
 &= kd_L([x(t)]^\alpha, [y(t)]^\alpha), \\
 & d_L([g(x(\cdot))]^\alpha, [g(y(\cdot))]^\alpha) \\
 &= d_L\left(\left[\sum_{k=1}^p c_k(x(t_k))\right]^\alpha, \left[\sum_{k=1}^p c_k(y(t_k))\right]^\alpha\right) \\
 &= \max_{1 \leq i \leq 2} \left\{ \left| \sum_{k=1}^p (c_k)_i (x_{il}^\alpha(t_k)) - \sum_{k=1}^p (c_k)_i (y_{il}^\alpha(t_k)) \right|, \left| \sum_{k=1}^p (c_k)_i (x_{ir}^\alpha(t_k)) - \sum_{k=1}^p (c_k)_i (y_{ir}^\alpha(t_k)) \right| \right\} \\
 &\leq \sum_{k=1}^p c_k \max_{1 \leq i \leq 2} \left\{ |x_{il}^\alpha(t_k) - y_{il}^\alpha(t_k)|, |x_{ir}^\alpha(t_k) - y_{ir}^\alpha(t_k)| \right\} \\
 &= \sum_{k=1}^p c_k d_L([x(t_k)]^\alpha, [y(t_k)]^\alpha) \\
 &\leq \sum_{k=1}^p c_k \max_k d_L([x(t_k)]^\alpha, [y(t_k)]^\alpha) \\
 &= hd_L([x(\cdot)]^\alpha, [y(\cdot)]^\alpha),
 \end{aligned} \tag{5.4}$$

where k and h satisfy the inequality (3.4) and (3.5), respectively. Choose T such that $T < (1 - ch)/ck$. Then all conditions stated in Theorem 3.2 are satisfied, so problem (5.2) has a unique fuzzy solution.

Let target set be $x^1 = (x_1^1, x_2^1) = (\tilde{2}, \tilde{3})$. The α -level set of fuzzy numbers is $\tilde{3}[\tilde{3}]^\alpha = [\alpha + 2, 4 - \alpha]$.

From the definition of fuzzy solution,

$$\begin{aligned} x_{il}^\alpha(t) &= S_{il}^\alpha(t) \left((x_0)_{il}^\alpha - \sum_{k=1}^p (c_k)_i (x_{il}^\alpha(t_k)) \right) \\ &\quad + \int_0^t S_{il}^\alpha(t-s) (\alpha+1) s (x_{il}^\alpha(s))^2 ds + \int_0^t S_{il}^\alpha(t-s) u_{il}^\alpha(s) ds, \\ x_{ir}^\alpha(t) &= S_{ir}^\alpha(t) \left((x_0)_{ir}^\alpha - \sum_{k=1}^p (c_k)_i (x_{ir}^\alpha(t_k)) \right) \\ &\quad + \int_0^t S_{ir}^\alpha(t-s) (3-\alpha) s (x_{ir}^\alpha(s))^2 ds + \int_0^t S_{ir}^\alpha(t-s) u_{ir}^\alpha(s) ds, \end{aligned} \tag{5.5}$$

where $i = 1, 2$.

Thus the α -level of $u(s)$ is

$$\begin{aligned} u_{1l}^\alpha(s) &= \left(\tilde{\beta}_{1l}^\alpha \right)^{-1} \left((\alpha+1) - \sum_{k=1}^p (c_k)_1 (x_{il}^\alpha(t_k)) \right. \\ &\quad \left. - \left[S_{1l}^\alpha(T) \left((x_0)_{1l}^\alpha - \sum_{k=1}^p (c_k)_1 (x_{1l}^\alpha(t_k)) \right) + \int_0^T (\alpha+1) S_{1l}^\alpha(T-s) s (x_{1l}^\alpha(s))^2 ds \right] \right), \\ u_{1r}^\alpha(s) &= \left(\tilde{\beta}_{1r}^\alpha \right)^{-1} \left((3-\alpha) - \sum_{k=1}^p (c_k)_1 (x_{ir}^\alpha(t_k)) \right. \\ &\quad \left. - \left[S_{1r}^\alpha(T) \left((x_0)_{1r}^\alpha - \sum_{k=1}^p (c_k)_1 (x_{1r}^\alpha(t_k)) \right) + \int_0^T (3-\alpha) S_{1r}^\alpha(T-s) s (x_{1r}^\alpha(s))^2 ds \right] \right), \\ u_{2l}^\alpha(s) &= \left(\tilde{\beta}_{2l}^\alpha \right)^{-1} \left((\alpha+2) - \sum_{k=1}^p (c_k)_2 (x_{il}^\alpha(t_k)) \right. \\ &\quad \left. - \left[S_{2l}^\alpha(T) \left((x_0)_{2l}^\alpha - \sum_{k=1}^p (c_k)_2 (x_{2l}^\alpha(t_k)) \right) + \int_0^T (\alpha+1) S_{2l}^\alpha(T-s) s (x_{2l}^\alpha(s))^2 ds \right] \right), \\ u_{2r}^\alpha(s) &= \left(\tilde{\beta}_{2r}^\alpha \right)^{-1} \left((4-\alpha) - \sum_{k=1}^p (c_k)_2 (x_{ir}^\alpha(t_k)) \right. \\ &\quad \left. - \left[S_{2r}^\alpha(T) \left((x_0)_{2r}^\alpha - \sum_{k=1}^p (c_k)_2 (x_{2r}^\alpha(t_k)) \right) + \int_0^T (3-\alpha) S_{2r}^\alpha(T-s) s (x_{2r}^\alpha(s))^2 ds \right] \right). \end{aligned} \tag{5.6}$$

Then α -level of $x(T) = (x_1(T), x_2(T))$ is

$$\begin{aligned}
 & [x_1(T)]^\alpha \\
 &= [x_{1l}^\alpha(T), x_{1r}^\alpha(T)] \\
 &= \left[S_{1l}^\alpha(T) \left((x_0)_{1l}^\alpha - \sum_{k=1}^p (c_k)_1 (x_{1l}^\alpha(t_k)) \right) + \int_0^T (\alpha + 1) S_{1l}^\alpha(T-s) s (x_{1l}^\alpha(s))^2 ds \right. \\
 &\quad \left. + \tilde{\beta}_{1l}^\alpha (\tilde{\beta}_{1l}^\alpha)^{-1} \left((\alpha + 1) - \sum_{k=1}^p (c_k)_1 (x_{1l}^\alpha(t_k)) \right) \right. \\
 &\quad \left. - \left\{ S_{1l}^\alpha(T) \left((x_0)_{1l}^\alpha - \sum_{k=1}^p (c_k)_1 (x_{1l}^\alpha(t_k)) \right) \right. \right. \\
 &\quad \left. \left. + \int_0^T (\alpha + 1) S_{1l}^\alpha(T-s) s (x_{1l}^\alpha(s))^2 ds \right\} \right] ds, \tag{5.7} \\
 & S_{1r}^\alpha(T) \left((x_0)_{1r}^\alpha - \sum_{k=1}^p (c_k)_1 (x_{1r}^\alpha(t_k)) \right) + \int_0^T (3 - \alpha) S_{1r}^\alpha(T-s) s (x_{1r}^\alpha(s))^2 ds \\
 &+ \tilde{\beta}_{1r}^\alpha (\tilde{\beta}_{1r}^\alpha)^{-1} \left((3 - \alpha) - \sum_{k=1}^p (c_k)_1 (x_{1r}^\alpha(t_k)) \right) \\
 &\quad - \left\{ S_{1r}^\alpha(T) \left((x_0)_{1r}^\alpha - \sum_{k=1}^p (c_k)_1 (x_{1r}^\alpha(t_k)) \right) \right. \\
 &\quad \left. + \int_0^T (3 - \alpha) S_{1r}^\alpha(T-s) s (x_{1r}^\alpha(s))^2 ds \right\} ds \Big] \\
 &= \left[(\alpha + 1) - \sum_{k=1}^p (c_k)_1 (x_{1l}^\alpha(t_k)), (3 - \alpha) - \sum_{k=1}^p (c_k)_1 (x_{1r}^\alpha(t_k)) \right] = \left[\tilde{2} - \sum_{k=1}^p (c_k)_1 (x_1(t_k)) \right]^\alpha.
 \end{aligned}$$

Similarly

$$[x_2(T)]^\alpha = [x_{2l}^\alpha(T), x_{2r}^\alpha(T)] = \left[\tilde{3} - \sum_{k=1}^p (c_k)_2 (x_2(t_k)) \right]^\alpha. \tag{5.8}$$

Hence

$$\begin{aligned}
 x(T) &= (x_1(T), x_2(T)) \\
 &= \left(\tilde{2} - \sum_{k=1}^p (c_k)_1 (x_1(t_k)), \tilde{3} - \sum_{k=1}^p (c_k)_2 (x_2(t_k)) \right) = x^1 - g(x). \tag{5.9}
 \end{aligned}$$

Then all the conditions stated in Theorem 4.2 are satisfied, so system (5.2) is nonlocal controllable on $[0, T]$.

Acknowledgment

This study was supported by research funds from Dong-A University.

References

- [1] P. Diamond and P. Kloeden, *Metric Spaces of Fuzzy Sets: Theory and Applications*, World Scientific, River Edge, NJ, USA, 1994.
- [2] Y. C. Kwun and D. G. Park, "Optimal control problem for fuzzy differential equations," in *Proceedings of the Korea-Vietnam Joint Seminar*, pp. 103–114, 1998.
- [3] P. Balasubramaniam and S. Muralisankar, "Existence and uniqueness of fuzzy solution for semilinear fuzzy integrodifferential equations with nonlocal conditions," *Computers & Mathematics with Applications*, vol. 47, no. 6-7, pp. 1115–1122, 2004.
- [4] J. H. Park, J. S. Park, and Y. C. Kwun, "Controllability for the semilinear fuzzy integrodifferential equations with nonlocal conditions," in *Fuzzy Systems and Knowledge Discovery*, vol. 4223 of *Lecture Notes in Computer Science*, pp. 221–230, Springer, Berlin, Germany.
- [5] Y. C. Kwun, M. J. Kim, B. Y. Lee, and J. H. Park, "Existence of solutions for the semilinear fuzzy integrodifferential equations using by successive iteration," *Journal of Korean Institute of Intelligent Systems*, vol. 18, pp. 543–548, 2008.
- [6] Y. C. Kwun, M. J. Kim, J. S. Park, and J. H. Park, "Continuously initial observability for the semilinear fuzzy integrodifferential equations," in *Proceedings of the 5th International Conference on Fuzzy Systems and Knowledge Discovery*, vol. 1, pp. 225–229, Jinan, China, October 2008.
- [7] B. Bede and S. G. Gal, "Almost periodic fuzzy-number-valued functions," *Fuzzy Sets and Systems*, vol. 147, no. 3, pp. 385–403, 2004.
- [8] S. G. Gal and G. M. N'Guérékata, "Almost automorphic fuzzy-number-valued functions," *Journal of Fuzzy Mathematics*, vol. 13, no. 1, pp. 185–208, 2005.
- [9] G. Wang, Y. Li, and C. Wen, "On fuzzy n -cell numbers and n -dimension fuzzy vectors," *Fuzzy Sets and Systems*, vol. 158, no. 1, pp. 71–84, 2007.