

RESEARCH

Open Access

On minimax programming problems involving right upper-Dini-derivative functions

Anurag Jayswal¹, Izhar Ahmad², Krishna Kumhari¹ and Suliman Al-Homidan^{2*}

*Correspondence:

homidan@kfupm.edu.sa

²Department of Mathematics and Statistics, King Fahd University of Petroleum and Minerals, Dhahran, 31261, Saudi Arabia

Full list of author information is available at the end of the article

Abstract

In this paper, we derive necessary and sufficient optimality conditions for a general minimax programming problem involving some classes of generalized convexities with the tool-right upper-Dini-derivative. Moreover, using the concept of optimality conditions, Mond-Weir type duality theory has been developed for such a minimax programming problem.

MSC: 26A51; 49J35; 90C32**Keywords:** minimax programming; upper-Dini-derivative; optimality; duality

1 Introduction

The minimax approach to optimization theory certainly is not new. It takes its origins in von Neumann's game theory. The broad spectrum of existing results and applications of minimax theory in the field of optimization is captured in the book edited by Du *et al.* [1]. Starting with work of Schmittendorf [2], minimax programming problems have been studied by several authors; for example, see [3–9] and the references cited therein.

Convexity is a sufficient but not necessary condition for many important results of mathematical programming, since there are diverse extensions of the notion of convexity bearing the same properties. Moreover, it is well known that a function is convex iff its restriction to each line segment in its domain is convex. This property inspired Ortega and Rheinboldt [10] to introduce an important generalization of convex functions by replacing a line segment joining two points by a continuous arc and called them arcwise connected functions defined on arcwise connected sets.

Following the idea of arcwise convexity, Avriel and Zang [11] introduced Q -connected (QCN) functions and P -connected (PCN) functions and also they have discussed necessary and sufficient local-global minimum properties of these functions. Some elementary properties of these functions in terms of their directional derivatives have been studied by Bhatia and Mehra [12]. Bhatia and Mehra [12] also established optimality conditions for scalar-valued nonlinear programming problems involving these functions.

To relax the definition of arcwise convexity in terms of directional derivative recently Yuan and Liu [13] introduced the concept of (α, ρ) -right upper-Dini-derivative locally arcwise connected with respect to the arc H and established optimality and duality results for a nonlinear multiobjective programming problem. In this paper, we use generalized

convex functions, in terms of the right upper-Dini-derivative to derive necessary and sufficient optimality conditions for a general minimax programming problem and duality results for its Mond-Weir type dual model.

This paper is structured as follows: Some preliminary concepts and properties regarding generalized convex functions are given in Section 2. In Section 3, we establish necessary and sufficient optimality conditions for a general minimax programming problem involving generalized convex functions. In Section 4, we establish appropriate duality theorems for a Mond-Weir type dual problem. Finally, in Section 5 we summarize our main results and also point out some further research opportunities.

2 Preliminaries

Let R^n denote the n -dimensional Euclidean space, R_+^n its nonnegative orthant and $X \subset R^n$. For a nonempty set Q in a topological vector space E , \bar{Q} denote the closure of Q and

$$Q^* = \{v \in E^* \mid v(q) \geq 0, \forall q \in Q\}$$

denotes the dual cone of Q , where E^* is the dual space of E .

For some nonempty subset Y , let $R^Y = \Pi_Y R$ denote the product space in a product topology. Then the topological dual space of R^Y is the generalized finite sequence space consisting of all the functions $u : Y \rightarrow R$ with finite support [14]. The set $R_+^Y = \Pi_Y R_+$ denote the convex cone of all nonnegative functions on Y . Then the topological dual of R_+^Y is given by

$$\begin{aligned} (R_+^Y)^* &= \wedge \\ &= \{ \lambda = (\lambda_y)_{y \in Y} \mid \exists \text{ a finite set } Y_0 \subseteq Y \text{ such that } \lambda_y = 0, \forall y \in Y \setminus Y_0 \\ &\quad \text{and } \lambda_y \geq 0, \forall y \in Y_0 \}. \end{aligned}$$

Now, we recall some well-known results and concepts which will be used in the sequel.

Definition 2.1 [15] A set $X \subset R^n$ is said to be an arcwise connected set if, for every $x_1 \in X$, $x_2 \in X$, there exists a continuous vector-valued function $H_{x_1, x_2} : [0, 1] \rightarrow X$, called an arc, such that

$$H_{x_1, x_2}(0) = x_1, \quad H_{x_1, x_2}(1) = x_2.$$

Definition 2.2 [13] Let φ be a real-valued function defined on an arcwise connected set $X \subset R^n$. Let $x_1, x_2 \in X$, and H_{x_1, x_2} be the arc connecting x_1 and x_2 in X . The right upper-Dini-derivative of φ with respect to $H_{x_2, x_1}(t)$ at $t = 0$ is defined as follows:

$$(d\varphi)^+(H_{x_2, x_1}(0^+)) = \limsup_{t \rightarrow 0^+} \frac{\varphi(H_{x_2, x_1}(t)) - \varphi(x_2)}{t}. \quad (1)$$

Using this upper-Dini-derivative concept, Yuan and Liu [13] introduced a class of functions, which called (α, ρ) -right upper-Dini-derivative function. For convenience, we use the following notations.

Definition 2.3 [13] A set $X \subset \mathbb{R}^n$ is said to be locally arcwise connected at \bar{x} if for any $x \in X$ and $x \neq \bar{x}$ there exist a positive number $a(x, \bar{x})$, with $0 < a(x, \bar{x}) \leq 1$, and a continuous arc $H_{\bar{x},x}$ such that $H_{\bar{x},x}(t) \in X$ for any $t \in (0, a(x, \bar{x}))$.

The set X is locally arcwise connected on X if X is locally arcwise connected at any $x \in X$.

Definition 2.4 [13] Let $X \subset \mathbb{R}^n$ be a locally arcwise connected set and $\varphi : X \rightarrow \mathbb{R}$ be a real function defined on X . The function φ is said to be (α, ρ) -right upper-Dini-derivative locally arcwise connected with respect to H at \bar{x} , if there exist real functions $\alpha : X \times X \rightarrow \mathbb{R}$, $\rho : X \times X \rightarrow \mathbb{R}$ such that

$$\varphi(x) - \varphi(\bar{x}) \geq \alpha(x, \bar{x})(d\varphi)^+(H_{\bar{x},x}(0^+)) + \rho(x, \bar{x}), \quad \forall x \in X.$$

If φ is (α, ρ) -right upper-Dini-derivative locally arcwise connected with respect to H at \bar{x} for any $\bar{x} \in X$, then φ is called (α, ρ) -right upper-Dini-derivative locally arcwise connected with respect to H on X .

Remark 2.1 It revealed by an example given in [13] that there exists a function, which is (α, ρ) -right upper-Dini-derivative locally arcwise connected but neither d - ρ - (η, θ) -invex [16] nor d -invex [17] nor directional differentially B-arcwise connected [15].

Now we define the notions of ρ -generalized-pseudo-right upper-Dini-derivative locally arcwise connected, strictly ρ -generalized-pseudo-right upper-Dini-derivative locally arcwise connected and ρ -generalized-quasi-right upper-Dini-derivative locally arcwise connected functions.

Definition 2.5 The function $\varphi : X \rightarrow \mathbb{R}$ is said to be ρ -generalized-pseudo-right upper-Dini-derivative locally arcwise connected (with respect to H) at \bar{x} , if there exists a real function $\rho : X \times X \rightarrow \mathbb{R}$ such that

$$(d\varphi)^+(H_{\bar{x},x}(0^+)) \geq -\rho(x, \bar{x}) \quad \Rightarrow \quad \varphi(x) \geq \varphi(\bar{x}), \quad \forall x \in X,$$

equivalently

$$\varphi(x) < \varphi(\bar{x}) \quad \Rightarrow \quad (d\varphi)^+(H_{\bar{x},x}(0^+)) < -\rho(x, \bar{x}), \quad \forall x \in X.$$

The function $\varphi : X \rightarrow \mathbb{R}$ is said to be ρ -generalized-pseudo-right upper-Dini-derivative locally arcwise connected (with respect to H) on X if it is ρ -generalized-pseudo-right upper-Dini-derivative locally arcwise connected (with respect to H) at any $\bar{x} \in X$.

The following example shows that there exists a function which is ρ -generalized-pseudo-right upper-Dini-derivative locally arcwise connected but not (α, ρ) -right upper-Dini-derivative locally arcwise connected with respect to the arc H .

Example 2.1 Let $X = (-1, 1)$ and the function $\varphi : X \rightarrow \mathbb{R}$ be defined by

$$\varphi(x) = \begin{cases} |x| \sin^2 \frac{1}{x}, & \text{if } x \in (-1, 0) \cup (0, 1), \\ 0, & \text{if } x = 0. \end{cases}$$

For any, $x, y \in R$, defining the arc $H : [0, 1] \rightarrow R$ by

$$H_{y,x}(t) = tx + (1 - t)y, \quad t \in [0, 1].$$

Note that, by the definition of right upper-Dini-derivative by (1), for $x \in (-1, 0) \cup (0, 1)$ we have

$$(d\varphi)^+(H_{0,x}(0^+)) = |x|.$$

Let $\rho : X \times X \rightarrow R$ be defined by

$$\rho(x, y) = \begin{cases} |x|; & \text{if } x \in (-1, 0) \cup (0, 1), \\ 0; & \text{if } x = 0. \end{cases}$$

Now, for $\bar{x} = 0$, it follows that

$$(d\varphi)^+(H_{\bar{x},x}(0^+)) \geq -\rho(x, \bar{x}) \implies \varphi(x) \geq \varphi(\bar{x}), \quad \forall x \in X.$$

This means that φ is ρ -generalized-pseudo-right upper-Dini-derivative locally arcwise connected (with respect to H) at $\bar{x} = 0$. But φ is not a (α, ρ) -right upper-Dini-derivative locally arcwise connected with respect to same arc H and ρ at $\bar{x} = 0$ because for $x \in (-1, 0) \cup (0, 1)$ and $\alpha(x, \bar{x}) = 1$, we can see that

$$\varphi(x) - \varphi(\bar{x}) - \alpha(x, \bar{x})(d\varphi)^+(H_{\bar{x},x}(0^+)) - \rho(x, \bar{x}) < 0.$$

Definition 2.6 The function $\varphi : X \rightarrow R$ is said to be strictly ρ -generalized-pseudo-right upper-Dini-derivative locally arcwise connected (with respect to H) at \bar{x} , if there exists a real function $\rho : X \times X \rightarrow R$ such that

$$(d\varphi)^+(H_{\bar{x},x}(0^+)) \geq -\rho(x, \bar{x}) \implies \varphi(x) > \varphi(\bar{x}), \quad \forall x \in X, x \neq \bar{x},$$

equivalently

$$\varphi(x) \leq \varphi(\bar{x}) \implies (d\varphi)^+(H_{\bar{x},x}(0^+)) < -\rho(x, \bar{x}), \quad \forall x \in X, x \neq \bar{x}.$$

The function $\varphi : X \rightarrow R$ is said to be strictly ρ -generalized-pseudo-right upper-Dini-derivative locally arcwise connected (with respect to H) on X if it is strictly ρ -generalized-pseudo-right upper-Dini-derivative locally arcwise connected (with respect to H) at any $\bar{x} \in X$.

Definition 2.7 The function $\varphi : X \rightarrow R$ is said to be ρ -generalized-quasi-right upper-Dini-derivative locally arcwise connected with respect to H at \bar{x} , if there exists a real function $\rho : X \times X \rightarrow R$ such that

$$\varphi(x) \leq \varphi(\bar{x}) \implies (d\varphi)^+(H_{\bar{x},x}(0^+)) \leq -\bar{\rho}(x, \bar{x}), \quad \forall x \in X,$$

equivalently

$$(d\varphi)^+(H_{\bar{x},x}(0^+)) > -\bar{\rho}(x, \bar{x}) \Rightarrow \varphi(x) > \varphi(\bar{x}), \quad \forall x \in X.$$

The function $\varphi : X \rightarrow R$ is said to be ρ -generalized-quasi-right upper-Dini-derivative locally arcwise connected (with respect to H) on X if it is ρ -generalized-quasi-right upper-Dini-derivative locally arcwise connected (with respect to H) at any $\bar{x} \in X$.

The next example shows that there exists a function which is ρ -generalized-quasi-right upper-Dini-derivative locally arcwise connected but neither (α, ρ) -right upper-Dini-derivative locally arcwise connected nor ρ -generalized-pseudo-right upper-Dini-derivative locally arcwise connected with respect to the arc H .

Example 2.2 Let $X = (-\frac{\pi}{2}, \frac{\pi}{2})$ and the function $\varphi : X \rightarrow R$ be defined by

$$\varphi(x) = \begin{cases} \pi|x| \sin^2 \frac{\pi}{x} - 2\pi|x|; & \text{if } x \in (-\frac{\pi}{2}, 0) \cup (0, \frac{\pi}{2}), \\ 0; & \text{if } x = 0. \end{cases}$$

For any, $x, y \in R$, defining the arc $H : [0, 1] \rightarrow R$ by

$$H_{y,x}(t) = tx + (1-t)y, \quad t \in [0, 1].$$

Clearly, for $x \in (-\frac{\pi}{2}, 0) \cup (0, \frac{\pi}{2})$ we have

$$(d\varphi)^+(H_{0,x}(0^+)) = -\pi|x|.$$

Let $\rho : X \times X \rightarrow R$ be defined by

$$\rho(x, y) = \begin{cases} \pi|x|; & \text{if } x \in (-\frac{\pi}{2}, 0) \cup (0, \frac{\pi}{2}), \\ 0; & \text{if } x = 0. \end{cases}$$

Now, we can easily verify that φ is ρ -generalized-quasi-right upper-Dini-derivative locally arcwise connected (with respect to H) at $\bar{x} = 0$. However, for $x \in (-\frac{\pi}{2}, 0) \cup (0, \frac{\pi}{2})$, $\alpha(x, \bar{x}) = 1$ and $\bar{x} = 0$, we can deduce that

$$\varphi(x) - \varphi(\bar{x}) - \alpha(x, \bar{x})(d\varphi)^+(H_{\bar{x},x}(0^+)) - \rho(x, \bar{x}) < 0,$$

and

$$\varphi(x) - \varphi(\bar{x}) < 0 \quad \text{implies} \quad (d\varphi)^+(H_{\bar{x},x}(0^+)) + \bar{\rho}(x, \bar{x}) = 0.$$

Hence, φ is neither (α, ρ) -right upper-Dini-derivative locally arcwise connected nor ρ -generalized-pseudo-right upper-Dini-derivative locally arcwise connected with respect to same arc H and ρ at $\bar{x} = 0$.

Definition 2.8 [13] A function $f : X \rightarrow R$ is called preinvex (with respect to $\eta : X \times X \rightarrow R^n$) on X if there exists a vector-valued function η such that

$$f(u + t\eta(x, u)) \leq tf(x) + (1 - t)f(u)$$

holds for all $x, u \in X$ and any $t \in [0, 1]$.

Definition 2.9 [13] A function $f : X \rightarrow R$ is said to be convexlike if for any $x, y \in X$ and $0 \leq \theta \leq 1$, there is $z \in X$ such that

$$f(z) \leq \theta f(x) + (1 - \theta)f(y).$$

Remark 2.2 The convex and the preinvex functions are convexlike functions.

In the next section we will use the following version of Theorem 2.3 from [9].

Lemma 2.1 Let $G : X \times Y \rightarrow R$ and $\psi : X \rightarrow R$, where X and Y are arbitrary nonempty sets. Let the pair (G, ψ) be convexlike on X . Assume that for some neighborhood U of 0 in R^Y and a constant $v > 0$, the set $\Omega_0 \cap \bar{U} \times (-\infty, v]$ is a nonempty closed subset of $R^Y \times R$, where

$$\Omega_0 = \{(u, r) | u : Y \rightarrow R \text{ and } \exists x \in X \text{ such that } \psi(x) \leq r, G(x, y) \leq u(y), \forall y \in Y\}.$$

Then exactly one of the following systems is solvable:

- (I) $G(x, y) \leq 0, \forall y \in Y, \psi(x) < 0$,
- (II) \exists an integer $s > 0$, scalars $\lambda_i \geq 0, 1 \leq i \leq s, \mu \geq 0$ and vectors $y^i \in Y, 1 \leq i \leq s$, such that $(\lambda_1, \dots, \lambda_s, \mu) \neq 0$ and $\sum_{i=1}^s \lambda_i G(x, y^i) + \mu \psi(x) \geq 0$.

3 Optimality conditions

Consider the following general minimax programming problem:

$$\begin{aligned} &\text{Minimize } \max_{y \in Y} f(x, y) \\ &\text{subject to } g(x) \leq 0, x \in X, \end{aligned} \tag{P}$$

where $f : X \times Y \rightarrow R, g = (g_1, g_2, \dots, g_m) : X \rightarrow R^m, X$ is an open arcwise connected subset of R^n, Y is a compact subset of R^m and $f(x, \cdot)$ is continuous on Y for every $x \in X. X_0 = \{x \in X | g_j(x) \leq 0, \forall 1 \leq j \leq m\}$ denote the set of feasible solutions of (P).

For $x \in X$, we define

$$\begin{aligned} I(x) &= \{j | g_j(x) = 0\}, \\ J(x) &= \{1, 2, \dots, m\} \setminus I(x), \\ Y(x) &= \left\{ y \in Y \mid f(x, y) = \sup_{z \in Y} f(x, z) \right\}. \end{aligned}$$

In view of the continuity of $f(x, \cdot)$ on Y and compactness of Y , it is clear that $Y(x)$ is nonempty compact subset of $Y, \forall x \in X$. Throughout this paper we assume that the right

upper-Dini-derivatives of the functions $f(\cdot, y), g_j(\cdot), j \in \{1, 2, \dots, m\}$ with respect to an arc $H_{\bar{x},x}$ at $t = 0$ exist $\forall \bar{x}, x \in X, \forall y \in Y$, and $(df)^+(H_{\bar{x},x}(0^+), \cdot)$ is continuous on $Y, \forall \bar{x}, x \in X$. Also assume that $g_j(\cdot), 1 \leq j \leq m$ is continuous on X .

The following lemma can be proved without difficulty on the same lines as in Lemma 3.1 (Mehra and Bhatia [9]).

Lemma 3.1 *Let \bar{x} be an optimal solution of (P). Then the system*

$$\begin{cases} (df)^+(H_{\bar{x},x}(0^+), y) < 0, & \forall y \in Y(\bar{x}), \\ (dg_j)^+(H_{\bar{x},x}(0^+)) < 0, & \forall j \in I(\bar{x}) \end{cases} \tag{2}$$

has no solution $x \in X$.

We now prove the following theorem by using Lemmas 2.1 and 3.1, which gives the necessary optimality conditions for an optimal solution of problem (P).

Theorem 3.1 (Necessary optimality conditions) *Let \bar{x} be an optimal solution of (P). Further, let $(df)^+(H_{\bar{x},x}(0^+)), (dg_j)^+(H_{\bar{x},x}(0^+)), j \in I(\bar{x})$ be convexlike functions of x on X and let there exist a neighborhood U of 0 in $R^{Y(\bar{x})}$ and a constant $v = (v_j)_{j \in I(\bar{x})}$ such that $\Omega(\bar{x}) \cap \bar{U} \times \prod_{j \in I(\bar{x})}(-\infty, v_j]$ is a nonempty closed set, where*

$$\Omega(\bar{x}) = \left\{ (u, r) \mid r = (r_j)_{j \in I(\bar{x})}, u : Y \rightarrow R \text{ and } \exists x \in X \text{ such that } f(x, y) \leq u(y), \forall y \in Y(\bar{x}), \right. \\ \left. g_j(x) \leq r_j, j \in I(\bar{x}) \right\}.$$

Then there exist an integer $s > 0$, scalars $\lambda_i \geq 0, 1 \leq i \leq s, \mu_j \geq 0, 1 \leq j \leq m$, and vectors $y^i \in Y(\bar{x}), 1 \leq i \leq s$, such that

$$\sum_{i=1}^s \lambda_i (df)^+(H_{\bar{x},x}(0^+), y^i) + \sum_{j=1}^m \mu_j (dg_j)^+(H_{\bar{x},x}(0^+)) \geq 0, \quad \forall x \in X, \\ \mu_j g_j(\bar{x}) = 0, \quad \forall 1 \leq j \leq m, \\ \sum_{i=1}^s \lambda_i + \sum_{j=1}^m \mu_j \neq 0.$$

Proof If \bar{x} is an optimal solution of (P) then, by Lemma 3.1, the system (2) has no solution $x \in X$. But the assumption of Lemma 2.1 also holds and since the system (2) has no solution $x \in X$, we obtain the result that there exist an integer $s > 0$, scalars $\lambda_i \geq 0, 1 \leq i \leq s, \mu_j \geq 0, j \in I(\bar{x})$, and vectors $y^i \in Y(\bar{x}), 1 \leq i \leq s$, such that

$$(\lambda_1, \lambda_2, \dots, \lambda_s, \mu_j)_{j \in I(\bar{x})} \neq 0, \tag{3}$$

and

$$\sum_{i=1}^s \lambda_i (df)^+(H_{\bar{x},x}(0^+), y^i) + \sum_{j \in I(\bar{x})} \mu_j (dg_j)^+(H_{\bar{x},x}(0^+)) \geq 0, \quad \forall x \in X. \tag{4}$$

If we put $\mu_j = 0$, for $j \in J(\bar{x})$, by (3) and (4) we obtain the required result. □

Now, we prove the following sufficient optimality conditions for the considered minimax problem (P) under generalized convexity with upper-Dini-derivative concept.

Theorem 3.2 (Sufficient optimality conditions) *Let $\bar{x} \in X_0$ and there exist an integer $s > 0$, scalars $\lambda_i \geq 0, 1 \leq i \leq s, \sum_{i=1}^s \lambda_i \neq 0, \mu_j \geq 0, 1 \leq j \leq m$, and vectors $y^i \in Y(\bar{x}), 1 \leq i \leq s$, such that $\alpha(x, \bar{x}) > 0$,*

$$\sum_{i=1}^s \lambda_i (df)^+(H_{\bar{x},x}(0^+), y^i) + \sum_{j=1}^m \mu_j (dg_j)^+(H_{\bar{x},x}(0^+)) \geq 0, \quad \forall x \in X, \tag{5}$$

$$\mu_j g_j(\bar{x}) = 0, \quad \forall 1 \leq j \leq m. \tag{6}$$

Also, assume that

- (i) for $1 \leq i \leq s, f(\cdot, y^i)$ is $(\alpha, \bar{\rho}_i)$ -right upper-Dini-derivative locally arcwise connected (with respect to H) at \bar{x} ,
- (ii) for $1 \leq j \leq m, g_j(\cdot)$ is $(\alpha, \check{\rho}_j)$ -right upper-Dini-derivative locally arcwise connected (with respect to H) at \bar{x} ,
- (iii) $\sum_{i=1}^s \lambda_i \bar{\rho}_i(x, \bar{x}) + \sum_{j=1}^m \mu_j \check{\rho}_j(x, \bar{x}) \geq 0$.

Then \bar{x} is an optimal solution of (P).

Proof Suppose to the contrary that \bar{x} is not an optimal solution of (P). Then there exists an $\tilde{x} \in X_0$ such that

$$\sup_{z \in Y} f(\tilde{x}, z) < \sup_{z \in Y} f(\bar{x}, z).$$

Further, as $y^i \in Y(\bar{x})$, we have

$$\sup_{z \in Y} f(\bar{x}, z) = f(\bar{x}, y^i), \quad 1 \leq i \leq s.$$

Also, $y^i \in Y, 1 \leq i \leq s$, we have

$$f(\tilde{x}, y^i) \leq \sup_{z \in Y} f(\tilde{x}, z), \quad 1 \leq i \leq s.$$

Thus, from the above three inequalities, we get

$$f(\tilde{x}, y^i) < f(\bar{x}, y^i), \quad 1 \leq i \leq s.$$

Using $\lambda_i \geq 0, 1 \leq i \leq s$ and $\sum_{i=1}^s \lambda_i \neq 0$, we obtain

$$\sum_{i=1}^s \lambda_i f(\tilde{x}, y^i) < \sum_{i=1}^s \lambda_i f(\bar{x}, y^i). \tag{7}$$

For $\tilde{x} \in X_0, \mu_j \geq 0, 1 \leq j \leq m$, we have $\mu_j g_j(\tilde{x}) \leq 0$, which in view of (6) implies that

$$\sum_{j=1}^m \mu_j g_j(\tilde{x}) \leq \sum_{j=1}^m \mu_j g_j(\bar{x}). \tag{8}$$

Now, by (7) and (8) we obtain

$$\sum_{i=1}^s \lambda_i f(\tilde{x}, y^i) + \sum_{j=1}^m \mu_j g_j(\tilde{x}) < \sum_{i=1}^s \lambda_i f(\bar{x}, y^i) + \sum_{j=1}^m \mu_j g_j(\bar{x}). \tag{9}$$

On the other hand, from the assumptions that $f(\cdot, y^i)$, $1 \leq i \leq s$ and $g_j(\cdot)$, $1 \leq j \leq m$ are $(\alpha, \bar{\rho}_i)$ and $(\alpha, \check{\rho}_j)$ -right upper-Dini-derivative locally arcwise connected (with respect to H) at \bar{x} , we have

$$f(\tilde{x}, y^i) - f(\bar{x}, y^i) \geq \alpha(\tilde{x}, \bar{x})(df)^+(H_{\tilde{x}, \bar{x}}(0^+), y^i) + \bar{\rho}_i(\tilde{x}, \bar{x}), \quad 1 \leq i \leq s, \tag{10}$$

$$g_j(\tilde{x}) - g_j(\bar{x}) \geq \alpha(\tilde{x}, \bar{x})(dg_j)^+(H_{\tilde{x}, \bar{x}}(0^+)) + \check{\rho}_j(\tilde{x}, \bar{x}), \quad 1 \leq j \leq m. \tag{11}$$

From (10) and (11) together with $\lambda_i \geq 0$, $1 \leq i \leq s$, and $\mu_j \geq 0$, $1 \leq j \leq m$, we get

$$\begin{aligned} & \left\{ \sum_{i=1}^s \lambda_i f(\tilde{x}, y^i) + \sum_{j=1}^m \mu_j g_j(\tilde{x}) \right\} - \left\{ \sum_{i=1}^s \lambda_i f(\bar{x}, y^i) + \sum_{j=1}^m \mu_j g_j(\bar{x}) \right\} \\ & \geq \alpha(\tilde{x}, \bar{x}) \left\{ \sum_{i=1}^s \lambda_i (df)^+(H_{\tilde{x}, \bar{x}}(0^+), y^i) + \sum_{j=1}^m \mu_j (dg_j)^+(H_{\tilde{x}, \bar{x}}(0^+)) \right\} \\ & \quad + \sum_{i=1}^s \lambda_i \bar{\rho}_i(\tilde{x}, \bar{x}) + \sum_{j=1}^m \mu_j \check{\rho}_j(\tilde{x}, \bar{x}). \end{aligned}$$

By (5) and using $\alpha(\tilde{x}, \bar{x}) > 0$, $\sum_{i=1}^s \lambda_i \bar{\rho}_i(\tilde{x}, \bar{x}) + \sum_{j=1}^m \mu_j \check{\rho}_j(\tilde{x}, \bar{x}) \geq 0$, it follows that

$$\left\{ \sum_{i=1}^s \lambda_i f(\tilde{x}, y^i) + \sum_{j=1}^m \mu_j g_j(\tilde{x}) \right\} - \left\{ \sum_{i=1}^s \lambda_i f(\bar{x}, y^i) + \sum_{j=1}^m \mu_j g_j(\bar{x}) \right\} \geq 0,$$

which is a contradiction to (9). Hence \bar{x} is an optimum solution for (P) and the theorem is proved. \square

Theorem 3.3 (Sufficient optimality conditions) *Let $\bar{x} \in X_0$ and there exist an integer $s > 0$, scalars $\lambda_i \geq 0$, $1 \leq i \leq s$, $\sum_{i=1}^s \lambda_i \neq 0$, $\mu_j \geq 0$, $1 \leq j \leq m$ and vectors $y^i \in Y(\bar{x})$, $1 \leq i \leq s$, such that the conditions (5) and (6) of Theorem 3.2 hold. Also, assume that*

- (i) $\sum_{i=1}^s \lambda_i f(\cdot, y^i)$ is $\bar{\rho}$ -generalized-pseudo-right upper-Dini-derivative locally arcwise connected (with respect to H) at \bar{x} ,
- (ii) $\sum_{j=1}^m \mu_j g_j(\cdot)$ is $\check{\rho}$ -generalized-quasi-right upper-Dini-derivative locally arcwise connected (with respect to H) at \bar{x} ,
- (iii) $\bar{\rho}(x, \bar{x}) + \check{\rho}(x, \bar{x}) \geq 0$.

Then \bar{x} is an optimal solution of (P).

Proof Suppose to the contrary that \bar{x} is not an optimal solution of (P) and following the proof of Theorem 3.2, we have

$$\sum_{i=1}^s \lambda_i f(\tilde{x}, y^i) < \sum_{i=1}^s \lambda_i f(\bar{x}, y^i),$$

which by $\bar{\rho}$ -generalized-pseudo-right upper-Dini-derivative locally arcwise connected (with respect to H) of $\sum_{i=1}^s \lambda_i f(\cdot, y^i)$ at \bar{x} , we have

$$\sum_{i=1}^s \lambda_i (df)^+(H_{\bar{x}, \bar{x}}(0^+), y^i) < -\bar{\rho}(\bar{x}, \bar{x}). \tag{12}$$

For $\tilde{x} \in X_0$, $\mu_j \geq 0$, $1 \leq j \leq m$, we have $\mu_j g_j(\tilde{x}) \leq 0$, which in view of (6) implies that

$$\sum_{j=1}^m \mu_j g_j(\tilde{x}) \leq \sum_{j=1}^m \mu_j g_j(\bar{x}),$$

which by $\check{\rho}$ -generalized-quasi-right upper-Dini-derivative locally arcwise connected (with respect to H) of $\sum_{j=1}^m \mu_j g_j(\cdot)$ at \bar{x} , we have

$$\sum_{j=1}^m \mu_j (dg_j)^+(H_{\bar{x}, \bar{x}}(0^+)) \leq -\check{\rho}(\bar{x}, \bar{x}). \tag{13}$$

By (12) and (13), we get

$$\sum_{i=1}^s \lambda_i (df)^+(H_{\bar{x}, \bar{x}}(0^+), y^i) + \sum_{j=1}^m \mu_j (dg_j)^+(H_{\bar{x}, \bar{x}}(0^+)) < -\bar{\rho}(\bar{x}, \bar{x}) - \check{\rho}(\bar{x}, \bar{x}) \leq 0,$$

where the last inequality is according to $\bar{\rho}(\bar{x}, \bar{x}) + \check{\rho}(\bar{x}, \bar{x}) \geq 0$. Therefore,

$$\sum_{i=1}^s \lambda_i (df)^+(H_{\bar{x}, \bar{x}}(0^+), y^i) + \sum_{j=1}^m \mu_j (dg_j)^+(H_{\bar{x}, \bar{x}}(0^+)) < 0,$$

which is a contradiction to (5). Hence \bar{x} is an optimum solution for (P) and the theorem is proved. \square

Theorem 3.4 (Sufficient optimality conditions) *Let $\bar{x} \in X_0$ and there exist an integer $s > 0$, scalars $\lambda_i \geq 0$, $1 \leq i \leq s$, $\sum_{i=1}^s \lambda_i \neq 0$, $\mu_j \geq 0$, $1 \leq j \leq m$, and vectors $y^i \in Y(\bar{x})$, $1 \leq i \leq s$, such that the conditions (5) and (6) of Theorem 3.2 hold. Also, assume that*

- (i) $\sum_{i=1}^s \lambda_i f(\cdot, y^i)$ is strictly $\bar{\rho}$ -generalized-pseudo-right upper-Dini-derivative locally arcwise connected (with respect to H) at \bar{x} ,
- (ii) $\sum_{j=1}^m \mu_j g_j(\cdot)$ is $\check{\rho}$ -generalized-quasi-right upper-Dini-derivative locally arcwise connected (with respect to H) at \bar{x} ,
- (iii) $\bar{\rho}(\bar{x}, \bar{x}) + \check{\rho}(\bar{x}, \bar{x}) \geq 0$.

Then \bar{x} is an optimal solution of (P).

Proof The proof follows along similar lines as the proof of Theorem 3.3 and hence is omitted. \square

Theorem 3.5 (Sufficient optimality conditions) *Let $\bar{x} \in X_0$ and there exist an integer $s > 0$, scalars $\lambda_i \geq 0$, $1 \leq i \leq s$, $\sum_{i=1}^s \lambda_i \neq 0$, $\mu_j \geq 0$, $1 \leq j \leq m$, and vectors $y^i \in Y(\bar{x})$, $1 \leq i \leq s$, such that the conditions (5) and (6) of Theorem 3.2 hold. Also, assume that*

- (i) $\sum_{i=1}^s \lambda_i f(\cdot, y^i)$ is $\bar{\rho}$ -generalized-quasi-right upper-Dini-derivative locally arcwise connected with respect to H at \bar{x} ,
- (ii) for $j \in I(\bar{x})$, $j \neq l$, $g_j(\cdot)$ is $\check{\rho}_j$ -generalized-quasi-right upper-Dini-derivative locally arcwise connected and $g_l(\cdot)$ is strictly $\check{\rho}_l$ -generalized-pseudo-right upper-Dini-derivative locally arcwise connected with respect to H at \bar{x} , with $\mu_l > 0$,
- (iii) $\bar{\rho}(x, \bar{x}) + \sum_{j=1}^m \mu_j \check{\rho}_j(x, \bar{x}) \geq 0$.

Then \bar{x} is an optimal solution of (P).

Proof Suppose to the contrary that \bar{x} is not an optimal solution of (P) and following the proof of Theorem 3.2, we have

$$\sum_{i=1}^s \lambda_i f(\tilde{x}, y^i) < \sum_{i=1}^s \lambda_i f(\bar{x}, y^i),$$

which by $\bar{\rho}$ -generalized-quasi-right upper-Dini-derivative locally arcwise connected of $\sum_{i=1}^s \lambda_i f(\cdot, y^i)$ with respect to H at \bar{x} , we have

$$\sum_{i=1}^s \lambda_i (df)^+(H_{\bar{x}, \bar{x}}(0^+), y^i) \leq -\bar{\rho}(\tilde{x}, \bar{x}). \tag{14}$$

Since for $\tilde{x} \in X_0$ and for $j \in I(\bar{x})$, we have

$$g_j(\tilde{x}) \leq 0 = g_j(\bar{x}),$$

which by $\check{\rho}_j$ -generalized-quasi-right upper-Dini-derivative locally arcwise connected of $g_j(\cdot)$, $j \in I(\bar{x})$, $j \neq l$ and strictly $\check{\rho}_l$ -generalized-pseudo-right upper-Dini-derivative locally arcwise connected of $g_l(\cdot)$ with respect to H at \bar{x} , we have

$$(dg_j)^+(H_{\bar{x}, \bar{x}}(0^+)) \leq -\check{\rho}_j(\tilde{x}, \bar{x}), \quad \text{for } j \in I(\bar{x}), j \neq l, \tag{15}$$

$$(dg_l)^+(H_{\bar{x}, \bar{x}}(0^+)) < -\check{\rho}_l(\tilde{x}, \bar{x}). \tag{16}$$

Since $\mu_j \geq 0$, $\forall j \in I(\bar{x})$, $\mu_l > 0$, and $\mu_j = 0$ for $j \in J(\bar{x})$, from (15) and (16), we get

$$\sum_{j=1}^m \mu_j (dg_j)^+(H_{\bar{x}, \bar{x}}(0^+)) < -\sum_{j=1}^m \mu_j \check{\rho}_j(\tilde{x}, \bar{x}). \tag{17}$$

By (14) and (17), we get

$$\sum_{i=1}^s \lambda_i (df)^+(H_{\bar{x}, \bar{x}}(0^+), y^i) + \sum_{j=1}^m \mu_j (dg_j)^+(H_{\bar{x}, \bar{x}}(0^+)) < -\bar{\rho}(\tilde{x}, \bar{x}) - \sum_{j=1}^m \mu_j \check{\rho}_j(\tilde{x}, \bar{x}) \leq 0,$$

where the last inequality is according to $\bar{\rho}(\tilde{x}, \bar{x}) + \sum_{j=1}^m \mu_j \check{\rho}_j(\tilde{x}, \bar{x}) \geq 0$. Therefore,

$$\sum_{i=1}^s \lambda_i (df)^+(H_{\bar{x}, \bar{x}}(0^+), y^i) + \sum_{j=1}^m \mu_j (dg_j)^+(H_{\bar{x}, \bar{x}}(0^+)) < 0,$$

which is a contradiction to (5). Hence \bar{x} is an optimum solution for (P) and the theorem is proved. \square

4 Duality

This section deals with the duality theorems for the following Mond-Weir type dual (D) of minimax problem (P):

$$\max_{(s,\lambda,y) \in \mathbb{K}} \sup_{(z,\mu) \in \mathbb{H}(s,\lambda,y)} \sum_{i=1}^s \lambda_i f(z, y^i), \tag{D}$$

where $\mathbb{K} = \{(s, \lambda, y) | s \text{ is an integer, } \lambda \in R_+^s, \sum_{i=1}^s \lambda_i = 1, y = (y^1, y^2, \dots, y^s), y^i \in Y(x) \text{ for some } x \in X, 1 \leq i \leq s\}$, and $\mathbb{H}(s, \lambda, y)$ denotes the set of all $(z, \mu) \in X \times R_+^m$ satisfying

$$\sum_{i=1}^s \lambda_i (df_i)^+(H_{z,x}(0^+), y^i) + \sum_{j=1}^m \mu_j (dg_j)^+(H_{z,x}(0^+)) \geq 0, \quad \text{for all } x \in X, \tag{18}$$

$$\sum_{j=1}^m \mu_j g_j(z) \geq 0. \tag{19}$$

If for a triplet (s, λ, y) in \mathbb{K} the set $\mathbb{H}(s, \lambda, y)$ is empty then we define the supremum over it to be $-\infty$.

Theorem 4.1 (Weak duality) *Let x and (z, μ, s, λ, y) be feasible solutions of (P) and (D), respectively. Assume that*

- (i) $\sum_{i=1}^s \lambda_i f(\cdot, y^i)$ is $\bar{\rho}$ -generalized-pseudo-right upper-Dini-derivative locally arcwise connected (with respect to H) at z ,
- (ii) $\sum_{j=1}^m \mu_j g_j(\cdot)$ is $\check{\rho}$ -generalized-quasi-right upper-Dini-derivative locally arcwise connected (with respect to H) at z ,
- (iii) $\bar{\rho}(x, z) + \check{\rho}(x, z) \geq 0$.

Then

$$\sup_{y \in Y} f(x, y) \geq \sum_{i=1}^s \lambda_i f(z, y^i).$$

Proof Suppose to the contrary that

$$\sup_{y \in Y} f(x, y) < \sum_{i=1}^s \lambda_i f(z, y^i).$$

Thus, we have

$$f(x, y^i) < \sum_{i=1}^s \lambda_i f(z, y^i), \quad \text{for all } y^i \in Y(x), 1 \leq i \leq s.$$

It follows from $\lambda_i \geq 0, 1 \leq i \leq s$, and $\sum_{i=1}^s \lambda_i = 1$, that

$$\sum_{i=1}^s \lambda_i f(x, y^i) < \sum_{i=1}^s \lambda_i f(z, y^i),$$

which by $\bar{\rho}$ -generalized-pseudo-right upper-Dini-derivative locally arcwise connected (with respect to H) of $\sum_{i=1}^s \lambda_i f(\cdot, y^i)$ at z , we have

$$\sum_{i=1}^s \lambda_i (df)^+(H_{z,x}(0^+), y^i) < -\bar{\rho}(x, z). \tag{20}$$

For $x \in X_0$, $\mu_j \geq 0$, $1 \leq j \leq m$, we have $\mu_j g_j(x) \leq 0$, which in view of (19) implies that

$$\sum_{j=1}^m \mu_j g_j(x) \leq \sum_{j=1}^m \mu_j g_j(z),$$

which by $\check{\rho}$ -generalized-quasi-right upper-Dini-derivative locally arcwise connected (with respect to H) of $\sum_{j=1}^m \mu_j g_j(\cdot)$ at z , we have

$$\sum_{j=1}^m \mu_j (dg_j)^+(H_{z,x}(0^+)) \leq -\check{\rho}(x, z). \tag{21}$$

By (20) and (21), we get

$$\sum_{i=1}^s \lambda_i (df)^+(H_{z,x}(0^+), y^i) + \sum_{j=1}^m \mu_j (dg_j)^+(H_{z,x}(0^+)) < -\bar{\rho}(x, z) - \check{\rho}(x, z) \leq 0,$$

where the last inequality is according to $\bar{\rho}(x, z) + \check{\rho}(x, z) \geq 0$. Therefore,

$$\sum_{i=1}^s \lambda_i (df)^+(H_{z,x}(0^+), y^i) + \sum_{j=1}^m \mu_j (dg_j)^+(H_{z,x}(0^+)) < 0,$$

which is a contradiction to (18). Hence the theorem is proved. □

Theorem 4.2 (Strong duality) *Let x^* be an optimal solution of (P). Assume that the conditions of Theorem 3.1 are satisfied. Then there exist $(s^*, \lambda^*, y^*) \in \mathbb{K}$, and $(x^*, \mu^*) \in \mathbb{H}(s^*, \lambda^*, y^*)$ such that $(x^*, \mu^*, s^*, \lambda^*, y^*)$ is a feasible solution of (D) and the two objectives have same values. If, in addition, the assumption of weak duality Theorem 4.1 hold for all feasible solutions of (D), then $(x^*, \mu^*, s^*, \lambda^*, y^*)$ is an optimal solution of (D).*

Proof Since x^* is an optimal solution for (P) and all the conditions of Theorem 3.1 are satisfied, there exist $(s^*, \lambda^*, y^*) \in \mathbb{K}$, and $(x^*, \mu^*) \in \mathbb{H}(s^*, \lambda^*, y^*)$ such that $(x^*, \mu^*, s^*, \lambda^*, y^*)$ is a feasible solution of (D) and the two objective values are equal. The optimality of $(x^*, \mu^*, s^*, \lambda^*, y^*)$ for (D) thus follows from Theorem 4.1. □

Theorem 4.3 (Strict converse duality) *Let x^* and $(z^*, \mu^*, s^*, \lambda^*, y^*)$ be optimal solutions of (P) and (D), respectively. Assume that the hypothesis of Theorem 4.2 is fulfilled. Also, assume that*

- (i) $\sum_{i=1}^{s^*} \lambda_i^* f(\cdot, y^{*i})$ is strictly $\bar{\rho}$ -generalized-pseudo-right upper-Dini-derivative locally arcwise connected (with respect to H) at z^* ,
- (ii) $\sum_{j=1}^m \mu_j^* g_j(\cdot)$ is $\check{\rho}$ -generalized-quasi-right upper-Dini-derivative locally arcwise connected (with respect to H) at z^* ,

$$(iii) \bar{\rho}(x^*, z^*) + \check{\rho}(x^*, z^*) \geq 0.$$

Then $z^* = x^*$.

Proof Suppose to the contrary that $z^* \neq x^*$. According to Theorem 4.2, we know that there exist $(s^*, \lambda^*, y^*) \in \mathbb{K}$, and $(x^*, \mu^*) \in \mathbb{H}(s^*, \lambda^*, y^*)$ such that $(x^*, \mu^*, s^*, \lambda^*, y^*)$ is a feasible solution of (D) and

$$\sup_{y^* \in Y} f(x^*, y^*) = \sum_{i=1}^{s^*} \lambda_i^* f(z^*, y^{*i}).$$

Thus, we have

$$f(x^*, y^{*i}) \leq \sum_{i=1}^{s^*} \lambda_i^* f(z^*, y^{*i}), \quad \text{for all } y^{*i} \in Y(x^*), 1 \leq i \leq s^*.$$

It follows from $\lambda_i^* \geq 0, 1 \leq i \leq s^*$, and $\sum_{i=1}^{s^*} \lambda_i^* = 1$, that

$$\sum_{i=1}^{s^*} \lambda_i^* f(x^*, y^{*i}) \leq \sum_{i=1}^{s^*} \lambda_i^* f(z^*, y^{*i}). \tag{22}$$

Now proceeding on the same lines as in Theorem 4.1, we get

$$\sum_{i=1}^{s^*} \lambda_i^* (df)^+(H_{z^*, x^*}(0^+), y^{*i}) + \sum_{j=1}^m \mu_j^* (dg_j)^+(H_{z^*, x^*}(0^+)) < 0,$$

which is a contradiction to (18). Hence the theorem is proved. \square

5 Conclusion

In this study we have established necessary and sufficient optimality conditions under generalized convexity using the tool-right upper-Dini-derivative for a general minimax programming problem. Mond-Weir type duality theory is also obtained. These results can be extended to the following semiinfinite minimax programming problem (SIP) with the tool-right upper-Dini-derivative:

$$\begin{aligned} & \text{Minimize } \max_{y \in Y} f(x, y) \\ & \text{subject to} \\ & G_j(x, t) \leq 0, \quad \text{for all } t \in T_j, j \in \underline{q} = \{1, 2, \dots, q\}, \\ & H_k(x, s) = 0, \quad \text{for all } s \in S_k, k \in \underline{r} = \{1, 2, \dots, r\}, \\ & x \in X, \end{aligned} \tag{SIP}$$

where $X \subset R^n$ is a nonempty open arcwise connected set, Y is a compact metrizable topological space, $f(\cdot, y)$ is a real-valued function defined on X . T_j and S_k are compact subsets of complete metric spaces, for each $j \in \underline{q}$, $G_j(\cdot, t)$ is a real-valued function defined on X for all $t \in T_j$, for each $k \in \underline{r}$, $H_k(\cdot, s)$ is a real-valued function defined on X for all $s \in S_k$,

for each $j \in \underline{q}$ and $k \in \underline{r}$, $G_j(x, \cdot)$, and $H_k(x, \cdot)$ are continuous real-valued functions defined, respectively, on T_j and S_k for all $x \in X$. We shall investigate this semiinfinite programming problem in subsequent papers.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors carried out the proof. All authors conceived of the study and participated in its design and coordination. All authors read and approved the final manuscript.

Author details

¹Department of Applied Mathematics, Indian School of Mines, Dhanbad, Jharkhand 826 004, India. ²Department of Mathematics and Statistics, King Fahd University of Petroleum and Minerals, Dhahran, 31261, Saudi Arabia.

Acknowledgements

The research of the second and fourth author is financially supported by King Fahd University of Petroleum and Minerals, Saudi Arabia under the Internal Research Project No. IN131026.

Received: 26 May 2014 Accepted: 29 July 2014 Published: 26 Aug 2014

References

1. Du, D, Pardalos, PM, Wu, WZ: Minimax and Applications. Kluwer Academic, Dordrecht (1995)
2. Schmittendorf, WE: Necessary conditions and sufficient conditions for static minimax problems. *J. Math. Anal. Appl.* **57**, 683-693 (1977)
3. Ahmad, I: Higher order duality in nondifferentiable minimax fractional programming involving generalized convexity. *J. Inequal. Appl.* **2012**, 306 (2012)
4. Ahmad, I: Second order nondifferentiable minimax fractional programming with square root terms. *Filomat* **27**, 135-142 (2013)
5. Ahmad, I, Husain, Z: Optimality conditions and duality in nondifferentiable minimax fractional programming with generalized convexity. *J. Optim. Theory Appl.* **129**, 255-275 (2006)
6. Gupta, SK, Dangar, D: On second order duality for nondifferentiable minimax fractional programming. *J. Comput. Appl. Math.* **255**, 878-886 (2014)
7. Jayswal, A, Prasad, AK, Kumhari, K: On nondifferentiable minimax fractional programming involving higher order generalized convexity. *Filomat* **27**, 1497-1504 (2013)
8. Jayswal, A, Stancu-Minasian, I, Ahmad, I, Kumhari, K: Generalized minimax fractional programming problems with generalized nonsmooth $(F, \alpha, \rho, d, \theta)$ -univex functions. *J. Nonlinear Anal. Optim.* **4**, 227-239 (2013)
9. Mehra, A, Bhatia, D: Optimality and duality for minmax problems involving arcwise connected and generalized arcwise connected functions. *J. Math. Anal. Appl.* **231**, 425-445 (1999)
10. Ortega, JM, Rheinboldt, WC: Iterative Solutions of Nonlinear Equations in Several Variables. Academic Press, New York (1970)
11. Avriel, M, Zang, I: Generalized arcwise connected functions and characterization of local-global minimum properties. *J. Optim. Theory Appl.* **32**, 407-425 (1980)
12. Bhatia, D, Mehra, A: Optimality conditions and duality involving arcwise connected and generalized arcwise connected functions. *J. Optim. Theory Appl.* **100**, 181-194 (1999)
13. Yuan, D, Liu, X: Mathematical programming involving (α, ρ) -right upper-Dini-derivative functions. *Filomat* **27**, 899-908 (2013)
14. Kelley, JL, Namioka, I: Linear Topological Spaces. Van Nostrand, Princeton (1963)
15. Zhang, QX: Optimality conditions and duality for semi-infinite programming involving B-arcwise connected functions. *J. Glob. Optim.* **45**, 615-629 (2009)
16. Nahak, C, Mohapatra, RN: d - ρ - (η, θ) -invexity in multiobjective optimization. *Nonlinear Anal.* **70**, 2288-2296 (2009)
17. Ye, YL: d -Invexity and optimality conditions. *J. Math. Anal. Appl.* **162**, 242-249 (1991)

10.1186/1029-242X-2014-326

Cite this article as: Jayswal et al.: On minimax programming problems involving right upper-Dini-derivative functions. *Journal of Inequalities and Applications* 2014, **2014**:326