

POSTER PRESENTATION

Open Access


Computational modeling of the development of detailed facial representations along ventral pathway

Akihiro Eguchi*, Simon M Stringer

From The Twenty Third Annual Computational Neuroscience Meeting: CNS*2014
Québec City, Canada. 26-31 July 2014

Experimental studies have shown that neurons at an intermediate stage of the primate ventral visual pathway encode the conformation and spatial relations of facial features [1],

while neurons in the later stages are selective to the full face [2]. In this study, we investigate how these cell firing properties may develop through visually-guided learning.



* Correspondence: akihiro.eguchi@psy.ox.ac.uk
Oxford Centre for theoretical neuroscience and artificial intelligence,
University of Oxford, OX1 3UD, UK

A hierarchical neural network model of the primate's ventral visual pathway is trained by presenting many randomly generated faces to the hierarchical competitive neural network while a local learning rule modifies the strengths of the synaptic connections between successive layers [3] (Figure 1A). After training, the model is found to have developed the experimentally observed cell firing properties. In particular, we have demonstrated how the primate brain learns to represent facial expression independently of facial identity as reported in [4] (Figure 1B). We have also shown how the visual system forms separate representations of facial features such as the eyes, nose and mouth (Figure 1C) as well as representations of spatial relationships between these facial features, as have been reported in single unit recording studies [1]. Therefore, this research makes an important contribution to understanding visual processing in the primate brain.

Published: 21 July 2014

References

1. Freiwald WA, Tsao DY, Livingstone MS: **A face feature space in the macaque temporal lobe.** *Nat Neurosci* 2009, **12**:1187-1196.
2. Tsao DY, Freiwald WA, Tootell RBH, Livingstone MS: **A cortical region consisting entirely of face-selective cells.** *Science* 2006, **311**:670-674.
3. Tromans JM, Harris M, Stringer SM: **A Computational Model of the Development of Separate Representations of Facial Identity and Expression in the Primate Visual System.** *PLoS ONE* 2011, **6**:e25616.
4. Hasselmo ME, Rolls ET, Baylis GC: **The role of expression and identity in the face-selective responses of neurons in the temporal visual cortex of the monkey.** *Behavioural Brain Research* 1989, **32**:203-218.

doi:10.1186/1471-2202-15-S1-P38

Cite this article as: Eguchi and Stringer: Computational modeling of the development of detailed facial representations along ventral pathway. *BMC Neuroscience* 2014 **15**(Suppl 1):P38.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

