



PUBLISHED FOR SISSA BY SPRINGER

RECEIVED: August 26, 2011

ACCEPTED: September 16, 2011

PUBLISHED: October 6, 2011

## Bessel-weighted asymmetries in semi-inclusive deep inelastic scattering

D. Boer,<sup>a</sup> L. Gamberg,<sup>b</sup> B.U. Musch<sup>c</sup> and A. Prokudin<sup>c</sup>

<sup>a</sup>*Theory group, KVI, University of Groningen,  
Zernikelaan 25, NL-9747 AA Groningen, The Netherlands*

<sup>b</sup>*Division of Science, Penn State University-Berks,  
Reading, Pennsylvania 19083, U.S.A.*

<sup>c</sup>*Jefferson Lab,  
12000 Jefferson Avenue, Newport News, VA 23606, U.S.A.*

*E-mail:* [d.boer@rug.nl](mailto:d.boer@rug.nl), [lpg10@psu.edu](mailto:lpg10@psu.edu), [bmusch@jlab.org](mailto:bmusch@jlab.org),  
[prokudin@jlab.org](mailto:prokudin@jlab.org)

**ABSTRACT:** The concept of weighted asymmetries is revisited for semi-inclusive deep inelastic scattering. We consider the cross section in Fourier space, conjugate to the outgoing hadron's transverse momentum, where convolutions of transverse momentum dependent parton distribution functions and fragmentation functions become simple products. Individual asymmetric terms in the cross section can be projected out by means of a generalized set of weights involving Bessel functions. Advantages of employing these Bessel weights are that they suppress (divergent) contributions from high transverse momentum and that soft factors cancel in (Bessel-)weighted asymmetries. Also, the resulting compact expressions immediately connect to previous work on evolution equations for transverse momentum dependent parton distribution and fragmentation functions and to quantities accessible in lattice QCD. Bessel-weighted asymmetries are thus model independent observables that augment the description and our understanding of correlations of spin and momentum in nucleon structure.

**KEYWORDS:** Deep Inelastic Scattering, QCD, Parton Model

**ARXIV EPRINT:** [1107.5294](https://arxiv.org/abs/1107.5294)

---

**Contents**

<b>1</b>	<b>Introduction</b>	<b>1</b>
1.1	Overview on weighted asymmetries	2
<b>2</b>	<b>The SIDIS cross section in Fourier space at tree level</b>	<b>3</b>
2.1	Elements of the SIDIS cross section	3
2.2	Representation in Fourier space	5
<b>3</b>	<b>Beyond tree level</b>	<b>8</b>
<b>4</b>	<b>TMD PDFs at the level of matrix elements</b>	<b>10</b>
<b>5</b>	<b>Bessel-weighted asymmetries</b>	<b>13</b>
<b>6</b>	<b>Average transverse momentum shift and Bessel-weighted counterpart</b>	<b>16</b>
<b>7</b>	<b>Conclusions</b>	<b>18</b>
<b>A</b>	<b>Conventions and useful relations</b>	<b>19</b>
<b>B</b>	<b>Multipole expansion and Fourier transform</b>	<b>20</b>
<b>C</b>	<b>Parameterization of the correlator in <math>b</math>-space</b>	<b>21</b>
<b>D</b>	<b>Structure functions in terms of Fourier transformed TMD PDFs and FFs</b>	<b>22</b>
<b>E</b>	<b>Cancellation of the soft factor in the Sivers asymmetry</b>	<b>23</b>
<b>F</b>	<b>Bessel-weighted asymmetries</b>	<b>24</b>
<b>G</b>	<b>Suppression of high transverse momentum</b>	<b>25</b>
G.1	Suppression of the tail in Bessel-weighted integrals	25
G.2	Fourier-transformed TMD PDFs, TMD FFs and their derivatives	27
G.3	Systematic errors from the region at large $P_{h\perp}$	27

---

**1 Introduction**

In the factorized picture of semi-inclusive processes, where the transverse momentum of the detected hadron  $P_{h\perp}$  is small compared to the photon virtuality  $Q^2$ , transverse momentum dependent (TMD) parton distribution functions (PDFs) characterize the spin and momentum structure of the proton [1–7]. At leading twist there are 8 TMD PDFs. They

can be studied experimentally by analyzing angular modulations in the differential cross section, so called spin and azimuthal asymmetries. These modulations are a function of the azimuthal angles of the final state hadron momentum about the virtual photon direction, as well as that of the target polarization (see e.g., ref. [8] for a review). TMD PDFs enter the SIDIS cross section in momentum space convoluted with transverse momentum dependent fragmentation functions (TMD FFs). However, after a two-dimensional Fourier transform of the cross section with respect to the transverse hadron momentum  $\mathbf{P}_{h\perp}$ , these convolutions become simple products of functions in Fourier  $\mathbf{b}_T$ -space. The usefulness of Fourier-Bessel transforms in studying the factorization as well as the scale dependence of transverse momentum dependent cross section has been known for some time [9–15]. In this paper we exhibit the structure of the cross section in  $\mathbf{b}_T$ -space and demonstrate how this representation results in model independent observables which are generalizations of the conventional weighted asymmetries [6, 7]. Further we explore the impact that these observables have in studying the scale dependence of the SIDIS cross section at small to moderate transverse momentum where the TMD framework is designed to give a good description of the cross section. In particular we study how the so called soft factor cancels from these observables. The soft factor [14–19] is an essential element of the cross section that emerges in the proofs of TMD factorization [11, 13–15]. It accounts for the collective effect of soft momentum gluons not associated with either the distribution or fragmentation part of the process and it is shown to be universal in hard processes [17]. Depending on the factorization framework, it appears explicitly in the structure functions and thus in the factorized cross section (see refs. [14, 18]), or it is completely absorbed in the definition of TMD PDFs and TMD FFs (see refs. [15, 19]). At tree level (zeroth order in  $\alpha_S$ ) the soft factor is unity, which explains its absence in the factorization formalism considered for example in ref. [8]. However, for a correct description of the energy scale dependence of the cross sections and asymmetries involving TMD PDFs, it is essential to include the soft factor. Yet, it is possible to consider observables where the soft factor is indeed absent or cancels out, these are precisely the weighted asymmetries.

### 1.1 Overview on weighted asymmetries

The concept of transverse momentum weighted single spin asymmetries (SSA) was proposed some time ago in refs. [6, 7]. Using the technique of weighting enables one to disentangle in a model independent way the cross sections and asymmetries in terms of the transverse (momentum) moments of TMD PDFs. A comprehensive list of such weights was derived in ref. [7] for semi-inclusive deep inelastic scattering (SIDIS). A prominent example is the weighted Sivers asymmetry, obtained from the differential cross section  $d\sigma$  according to

$$A_{UT,T}^{w_1 \sin(\phi_h - \phi_S)} = \frac{2 \int d|\mathbf{P}_{h\perp}| |\mathbf{P}_{h\perp}| d\phi_h d\phi_S w_1(|\mathbf{P}_{h\perp}|) \sin(\phi_h - \phi_S) \{d\sigma(\phi_h, \phi_S) - d\sigma(\phi_h, \phi_S + \pi)\}}{\int d|\mathbf{P}_{h\perp}| d\phi_h |\mathbf{P}_{h\perp}| d\phi_S w_0(|\mathbf{P}_{h\perp}|) \{d\sigma(\phi_h, \phi_S) + d\sigma(\phi_h, \phi_S + \pi)\}}, \tag{1.1}$$

where the integrations are performed over the observed transverse hadron momentum  $|\mathbf{P}_{h\perp}|$ , the hadron azimuthal angle  $\phi_h$  and the spin direction  $\phi_S$  of the transversely polarized target, and the weights are  $w_1 = |\mathbf{P}_{h\perp}|/zM$ ,  $w_0 = 1$ . At tree level and leading twist

the weighted Siverts asymmetry [7] then becomes,

$$A_{UT}^{\frac{|\mathbf{P}_{h\perp}|}{zM}} \sin(\phi_h - \phi_s) = -2 \frac{\sum_a e_a^2 f_{1T}^{\perp(1)a}(x) D_1^{(0)a}(z)}{\sum_a e_a^2 f_1^{(0)a}(x) D_1^{(0)a}(z)}, \quad (1.2)$$

where  $f_{1T}^{\perp(1)a}$ ,  $f_1^{(0)a}$  and  $D_1^{(0)a}$  are transverse momentum moments of TMD PDFs and TMD FFs, and  $e_a$  is the electric charge for a quark of flavor  $a$ . As explained in greater detail in section 5, the moments in eq. (1.2) are undefined without a subtraction prescription for the infinite contributions at large transverse momentum. Here, we propose generalized weights,  $w_n \propto J_n(|\mathbf{P}_{h\perp}|\mathcal{B}_T)$  with  $J_n$  denoting Bessel functions of the first kind, and where  $\mathcal{B}_T$  (in units  $(\text{GeV}/c)^{-1}$ ) is a free parameter that represents the Fourier conjugate to  $|\mathbf{P}_{h\perp}|$ . For the Siverts asymmetry,  $w_1 = 2J_1(|\mathbf{P}_{h\perp}|\mathcal{B}_T)/zM\mathcal{B}_T$  and  $w_0 = J_0(|\mathbf{P}_{h\perp}|\mathcal{B}_T)$ . This gives rise to the Bessel-weighted Siverts asymmetry, which reads

$$A_{UT}^{\frac{2J_1(|\mathbf{P}_{h\perp}|\mathcal{B}_T)}{zM\mathcal{B}_T}} \sin(\phi_h - \phi_s) = -2 \frac{\sum_a e_a^2 \tilde{f}_{1T}^{\perp(1)a}(x, z^2\mathcal{B}_T^2) \tilde{D}_1^{(0)a}(z, \mathcal{B}_T^2)}{\sum_a e_a^2 \tilde{f}_1^{(0)a}(x, z^2\mathcal{B}_T^2) \tilde{D}_1^{(0)a}(z, \mathcal{B}_T^2)}, \quad (1.3)$$

where  $\tilde{f}_{1T}^{\perp(1)a}$ ,  $\tilde{f}_1^{(0)a}$  and  $\tilde{D}_1^{(0)a}$  are TMD PDFs and TMD FFs Fourier transformed with respect to transverse momentum as defined in the next section. In the asymptotic limit  $\mathcal{B}_T \rightarrow 0$ , we recover the conventional weighted asymmetry eq. (1.2), and the Fourier transformed TMD PDFs and FFs can be identified with the moments in that equation. An advantage of the generalized weight relates to the asymptotic behavior of TMD PDFs (and TMD FFs). We will see that this provides a regularization of the infinite contributions at large transverse momentum as long as we keep  $\mathcal{B}_T^2$  non-zero. Moreover, our analysis will show that soft factors appearing beyond tree level cancel out of the weighted asymmetry.


The rest of the manuscript is organized as follows: In section 2 we write down the general form of the SIDIS cross section in the TMD factorization framework and show that the convolutions in momentum space appear as products in Fourier space. For simplicity, this discussion is presented at tree level. Modifications needed to go beyond tree level are discussed in section 3. Even though our arguments are quite general, for definiteness we use the framework of Ji, Ma, Yuan [14, 18], here referred to as the ‘‘JMY’’ framework. TMD PDFs at the level of matrix elements will be considered in section 4. In section 5 we will consider Bessel-weighted asymmetries, focusing on the Siverts asymmetry as an explicit example. Further asymmetries at leading twist are listed in appendix F. We will also consider  $x$  moments of TMD PDFs and introduce a method to study Fourier transformed moments in lattice QCD and compare with experiment. Our conclusions are presented in section 7.

## 2 The SIDIS cross section in Fourier space at tree level

### 2.1 Elements of the SIDIS cross section

The lepton-hadron cross section of SIDIS  $\ell(l) + N(P, S) \rightarrow \ell(l) + h(P_h) + X$  can be expressed [4, 8, 20, 21] in the notation of ref. [8] as

$$\frac{d\sigma}{dx_B dy d\psi dz_h d\phi_h |\mathbf{P}_{h\perp}| d|\mathbf{P}_{h\perp}|} = \frac{\alpha^2}{x_B y Q^2} \frac{y^2}{(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x_B}\right) L_{\mu\nu} W^{\mu\nu}, \quad (2.1)$$



**Figure 1.** Kinematics of the SIDIS process, compare refs. [8, 22].

where we assume one photon exchange.  $L_{\mu\nu}$  and  $W^{\mu\nu}$  are the leptonic and hadronic tensors respectively, and the vector  $\mathbf{P}_{h\perp}$  is the transverse momentum of the produced hadron in a frame where the virtual photon and the target are collinear, e.g. in the target rest frame or  $\gamma^*P$  center of mass frame. It makes an azimuthal angle  $\phi_h$  with the lepton scattering plane defined by the momenta of the incoming and the final leptons  $l$  and  $l'$  (see figure 1). We define  $q \equiv l - l'$ , and  $q^2 = -Q^2$  is the virtuality of the photon.  $\psi$  is the azimuthal angle of  $l'$  around the lepton beam axis relative to  $S_\perp$ , in DIS kinematics  $d\psi \approx d\phi_S$  [21]. The subscript “ $\perp$ ” denotes transverse projection in the target rest frame while the subscript “ $T$ ” denotes transverse projection in the light-cone frame. We use definitions for the kinematic variables and the ratio of longitudinal and transverse photon flux  $\epsilon$  as in ref. [8],

$$x_B = \frac{Q^2}{2P \cdot q}, \quad y = \frac{P \cdot q}{P \cdot l}, \quad z_h = \frac{P \cdot P_h}{P \cdot q}, \quad \gamma = \frac{2Mx}{Q}, \quad \epsilon = \frac{1 - y - \frac{1}{4}\gamma^2 y^2}{1 - y + \frac{1}{2}y^2 + \frac{1}{4}\gamma^2 y^2}, \quad (2.2)$$

where  $M$  is the mass of the target nucleon. We employ the standard light-cone decomposition of four-vectors  $\omega^\mu = \omega^+ n_+^\mu + \omega^- n_-^\mu + \omega_T^\mu$ . In the  $\gamma^*P$  center of mass frame with the proton three-momentum pointing in positive  $z$ -direction, the nucleon carries no transverse momentum,  $P_T = 0$ , and  $x \equiv p^+/P^+$  denotes the momentum fraction carried by the quark (parton) of momentum  $p$ . Further definitions of kinematic variables and details on the leptonic and hadronic tensor are given in appendix A and ref. [8].

At tree-level of the hard photon-quark scattering process, and to leading order in the  $1/Q$  expansion, the hadronic tensor can be written in factorized form as [5, 8, 23]

$$2MW^{\mu\nu} = \sum_a e_a^2 \int d^2\mathbf{p}_T d^2\mathbf{K}_T \delta^{(2)}(z\mathbf{p}_T + \mathbf{K}_T - \mathbf{P}_{h\perp}) \text{Tr} \left\{ \Phi(x, \mathbf{p}_T) \gamma^\mu \Delta(z, \mathbf{K}_T) \gamma^\nu \right\}. \quad (2.3)$$

The quark-quark correlator [11, 24] in the above equation is defined as

$$\Phi_{ij}(p, P, S) \equiv \int \frac{d^4b}{(2\pi)^4} e^{ip \cdot b} \langle P, S | \bar{\psi}_j(0) \mathcal{U}[\mathcal{C}_b] \psi_i(b) | P, S \rangle. \quad (2.4)$$

In eq. (2.3) an integration has been performed over the small component  $p^-$  of parton momentum to obtain a correlator that depends on light-cone fraction  $x$  and parton transverse momentum  $\mathbf{p}_T$ , that is,

$$\begin{aligned} \Phi_{ij}(x, \mathbf{p}_T) &\equiv \int dp^- \Phi_{ij}(p, P, S) \\ &= \int \frac{db^- d^2\mathbf{b}_T}{(2\pi)^3} e^{ixP^+ b^- - i\mathbf{p}_T \cdot \mathbf{b}_T} \langle P, S | \bar{\psi}_j(0) \mathcal{U}[\mathcal{C}_b] \psi_i(b) | P, S \rangle \Big|_{b^+=0}. \end{aligned} \quad (2.5)$$

The state  $|P, S\rangle$  represents a nucleon with four-momentum  $P$  and spin polarization vector  $S$ , and quark fields are located at position “0” and “ $b$ ” in coordinate space. The gauge link  $\mathcal{U}[\mathcal{C}_b]$  ensures gauge invariance of the correlator [23, 25]. It corresponds to a path in  $b$  space which is determined by the color flow in the hard sub-process [26, 27]. We will discuss the details of the definition of the correlator and the role of the gauge link  $\mathcal{U}[\mathcal{C}_b]$  in section 4. Analogous expressions define the fragmentation correlator  $\Delta_{ij}(z, \mathbf{p}_T)$  (see e.g. [8]).

## 2.2 Representation in Fourier space

In this section, we rewrite the SIDIS cross section and its transverse momentum dependent components in coordinate  $\mathbf{b}_T$  space, similar as previously done in ref. [28]. Here however, we take advantage of the rotational invariance of TMD PDFs and FFs.

First we use the representation of the  $\delta$ -function

$$\delta^{(2)}(z\mathbf{p}_T + \mathbf{K}_T - \mathbf{P}_{h\perp}) = \int \frac{d^2\mathbf{b}_T}{(2\pi)^2} e^{i\mathbf{b}_T(z\mathbf{p}_T + \mathbf{K}_T - \mathbf{P}_{h\perp})}, \quad (2.6)$$

along with the following definitions,

$$W^{\mu\nu}(\mathbf{P}_{h\perp}) \equiv \int \frac{d^2\mathbf{b}_T}{(2\pi)^2} e^{-i\mathbf{b}_T \cdot \mathbf{P}_{h\perp}} \tilde{W}^{\mu\nu}(\mathbf{b}_T), \quad (2.7)$$

$$\begin{aligned} \tilde{\Phi}_{ij}(x, z\mathbf{b}_T) &\equiv \int d^2\mathbf{p}_T e^{iz\mathbf{b}_T \cdot \mathbf{p}_T} \Phi_{ij}(x, \mathbf{p}_T) \\ &= \int \frac{db^-}{(2\pi)} e^{ixP^+b^-} \langle P, S | \bar{\psi}_j(0) \mathcal{U}[\mathcal{C}_b] \psi_i(b) | P, S \rangle \Big|_{b^+=0}, \end{aligned} \quad (2.8)$$

$$\tilde{\Delta}_{ij}(z, \mathbf{b}_T) \equiv \int d^2\mathbf{K}_T e^{i\mathbf{b}_T \cdot \mathbf{K}_T} \Delta_{ij}(z, \mathbf{K}_T), \quad (2.9)$$

to re-write the leading term in the hadronic tensor, eq. (2.3), in Fourier space

$$2M\tilde{W}^{\mu\nu} = \sum_a e_a^2 \text{Tr} \left( \tilde{\Phi}(x, z\mathbf{b}_T) \gamma^\mu \tilde{\Delta}(z, \mathbf{b}_T) \gamma^\nu \right). \quad (2.10)$$

The advantage of the  $\mathbf{b}_T$  space representation is clear: the hadronic tensor is no longer a convolution of  $\mathbf{p}_T$  and  $\mathbf{K}_T$  dependent functions but a simple product of  $\mathbf{b}_T$ -dependent functions. This motivates us to re-write the entire cross section in terms of the Fourier transform

$$\frac{d\sigma}{dx_B dy d\psi dz_h d\phi_h |\mathbf{P}_{h\perp}| d|\mathbf{P}_{h\perp}|} = \int \frac{d^2\mathbf{b}_T}{(2\pi)^2} e^{-i\mathbf{b}_T \cdot \mathbf{P}_{h\perp}} \left\{ \frac{\alpha^2}{x_B y Q^2} \frac{y^2}{(1-\varepsilon)} \left( 1 + \frac{\gamma^2}{2x_B} \right) L_{\mu\nu} \tilde{W}^{\mu\nu} \right\}. \quad (2.11)$$

Next, we decompose the correlators  $\tilde{\Phi}$  and  $\tilde{\Delta}$  into TMD PDFs and FFs in Fourier space. Using the trace notation (see also eqs. (A.8) and (A.9) in the appendix)

$$\tilde{\Phi}^{[\Gamma]} \equiv \frac{1}{2} \text{Tr}(\tilde{\Phi}\Gamma), \quad (2.12)$$

and restricting ourselves to leading twist projections, we obtain the following structures for  $\tilde{\Phi}$

$$\begin{aligned}
 \tilde{\Phi}^{[\gamma^+]}(x, \mathbf{b}_T) &= \tilde{f}_1(x, \mathbf{b}_T^2) - i \epsilon_T^{\rho\sigma} b_{T\rho} S_{T\sigma} M \tilde{f}_{1T}^{\perp(1)}(x, \mathbf{b}_T^2), \\
 \tilde{\Phi}^{[\gamma^+\gamma^5]}(x, \mathbf{b}_T) &= S_L \tilde{g}_{1L}(x, \mathbf{b}_T^2) + i \mathbf{b}_T \cdot \mathbf{S}_T M \tilde{g}_{1T}^{(1)}(x, \mathbf{b}_T^2), \\
 \tilde{\Phi}^{[i\sigma^{\alpha+}\gamma^5]}(x, \mathbf{b}_T) &= S_T^\alpha \tilde{h}_1(x, \mathbf{b}_T^2) + i S_L b_T^\alpha M \tilde{h}_{1L}^{\perp(1)}(x, \mathbf{b}_T^2) \\
 &\quad + \frac{1}{2} \left( b_T^\alpha b_T^\rho + \frac{1}{2} \mathbf{b}_T^2 g_T^{\alpha\rho} \right) M^2 S_{T\rho} \tilde{h}_{1T}^{\perp(2)}(x, \mathbf{b}_T^2) \\
 &\quad - i \epsilon_T^{\alpha\rho} b_{T\rho} M \tilde{h}_1^{\perp(1)}(x, \mathbf{b}_T^2),
 \end{aligned} \tag{2.13}$$

where  $\alpha = 1, 2$  and  $\rho = 1, 2$ . Similarly, we obtain the following structures for  $\tilde{\Delta}$

$$\begin{aligned}
 \tilde{\Delta}^{[\gamma^-]}(z, \mathbf{b}_T) &= \tilde{D}_1(z, \mathbf{b}_T^2) - i \epsilon_T^{\rho\sigma} b_{T\rho} S_{hT\sigma} z M_h \tilde{D}_{1T}^{\perp(1)}(z, \mathbf{b}_T^2), \\
 \tilde{\Delta}^{[\gamma^-\gamma^5]}(z, \mathbf{b}_T) &= S_{hL} \tilde{G}_{1L}(z, \mathbf{b}_T^2) - i \mathbf{b}_T \cdot \mathbf{S}_{hT} z M_h \tilde{G}_{1T}^{(1)}(z, \mathbf{b}_T^2), \\
 \tilde{\Delta}^{[i\sigma^{\alpha-}\gamma^5]}(z, \mathbf{b}_T) &= S_{hT}^\alpha \tilde{H}_1(z, \mathbf{b}_T^2) - i S_{hL} b_T^\alpha z M_h \tilde{H}_{1L}^{\perp(1)}(z, \mathbf{b}_T^2) \\
 &\quad + \frac{1}{2} \left( b_T^\alpha b_T^\rho + \frac{1}{2} \mathbf{b}_T^2 g_T^{\alpha\rho} \right) z^2 M_h^2 S_{hT\rho} \tilde{H}_{1T}^{\perp(2)}(z, \mathbf{b}_T^2) \\
 &\quad - i \epsilon_T^{\alpha\rho} b_{T\rho} z M_h \tilde{H}_1^{\perp(1)}(z, \mathbf{b}_T^2).
 \end{aligned} \tag{2.14}$$

$$\tag{2.15}$$

For future applications, we have written down the latter decomposition for the more general case of a spin- $\frac{1}{2}$  hadron; the expression for a spinless hadron is obtained by setting  $S_h = 0$ . The above decompositions can be deduced from the existing expressions for  $\Phi$  and  $\Delta$  in momentum space [5, 29], or starting from the symmetry properties of the correlators  $\tilde{\Phi}$  and  $\tilde{\Delta}$  and a parameterization in terms of Lorentz-invariant amplitudes, see also section 4 and appendix C. The functions  $\tilde{f}_1(x, \mathbf{b}_T^2)$ ,  $\tilde{g}_{1L}(x, \mathbf{b}_T^2)$ , ... are the Fourier transforms of the usual TMD PDFs  $f_1(x, \mathbf{p}_T^2)$ ,  $g_{1L}(x, \mathbf{p}_T^2)$ , ... For a generic TMD PDF called  $f$  and a generic TMD FF called  $D$ , this Fourier transform is given by

$$\begin{aligned}
 \tilde{f}(x, \mathbf{b}_T^2) &\equiv \int d^2 \mathbf{p}_T e^{i \mathbf{b}_T \cdot \mathbf{p}_T} f(x, \mathbf{p}_T^2) \\
 &= 2\pi \int d|\mathbf{p}_T| |\mathbf{p}_T| J_0(|\mathbf{b}_T| |\mathbf{p}_T|) f(x, \mathbf{p}_T^2),
 \end{aligned} \tag{2.16}$$

$$\tilde{D}(z, \mathbf{b}_T^2) \equiv \int d^2 \mathbf{K}_T e^{i \mathbf{b}_T \cdot \mathbf{K}_T} D(z, \mathbf{K}_T^2) = 2\pi \int d|\mathbf{K}_T| |\mathbf{K}_T| J_0(|\mathbf{b}_T| |\mathbf{K}_T|) D(z, \mathbf{K}_T^2). \tag{2.17}$$

Additionally, in eqs. (2.13) and (2.15) not only Fourier transformed TMD PDFs and TMD FFs, but also their  $\mathbf{b}_T^2$ -derivatives appear, which we denote as

$$\begin{aligned}
 \tilde{f}^{(n)}(x, \mathbf{b}_T^2) &\equiv n! \left( -\frac{2}{M^2} \partial_{\mathbf{b}_T^2} \right)^n \tilde{f}(x, \mathbf{b}_T^2) \\
 &= \frac{2\pi n!}{(M^2)^n} \int d|\mathbf{p}_T| |\mathbf{p}_T| \left( \frac{|\mathbf{p}_T|}{|\mathbf{b}_T|} \right)^n J_n(|\mathbf{b}_T| |\mathbf{p}_T|) f(x, \mathbf{p}_T^2),
 \end{aligned} \tag{2.18}$$

$$\begin{aligned}
 \tilde{D}^{(n)}(z, \mathbf{b}_T^2) &\equiv n! \left( -\frac{2}{z^2 M_h^2} \partial_{\mathbf{b}_T^2} \right)^n \tilde{D}(z, \mathbf{b}_T^2) \\
 &= \frac{2\pi n!}{(z^2 M_h^2)^n} \int d|\mathbf{K}_T| |\mathbf{K}_T| \left( \frac{|\mathbf{K}_T|}{|\mathbf{b}_T|} \right)^n J_n(|\mathbf{b}_T| |\mathbf{K}_T|) D(z, \mathbf{K}_T^2).
 \end{aligned} \tag{2.19}$$

The functions  $\tilde{f}$ ,  $\tilde{D}$ ,  $\tilde{f}^{(n)}$  and  $\tilde{D}^{(n)}$  are real valued and  $\tilde{f}^{(0)} = \tilde{f}$ ,  $\tilde{D}^{(0)} = \tilde{D}$ . Taking the ‘‘asymptotic limit’’  $|\mathbf{b}_T| \rightarrow 0$  on the right hand side of eqs. (2.19), we formally obtain the conventional moments of the TMD PDFs and TMD FFs,  $f^{(n)}(x)$  and  $D^{(n)}(z)$  respectively,

$$\begin{aligned}\tilde{f}^{(n)}(x, 0) &= \int d^2\mathbf{p}_T \left( \frac{\mathbf{p}_T^2}{2M^2} \right)^n f(x, \mathbf{p}_T^2) \equiv f^{(n)}(x), \\ \tilde{D}^{(n)}(z, 0) &= \int d^2\mathbf{K}_T \left( \frac{\mathbf{K}_T^2}{2z^2 M_h^2} \right)^n D(x, \mathbf{K}_T^2) \equiv D^{(n)}(z).\end{aligned}\quad (2.20)$$

Thus we find that the derivatives in  $\mathbf{b}_T$ -space are directly related to moments of TMD PDFs and FFs. Finally we re-write the SIDIS cross section of ref. [8] in the  $\gamma^*P$  center of mass frame with the proton three-momentum pointing in the negative  $z$ -direction (so called Trento conventions [22]), as

$$\begin{aligned}\frac{d\sigma}{dx_B dy d\phi_S dz_h d\phi_h |\mathbf{P}_{h\perp}| d|\mathbf{P}_{h\perp}|} &= \\ &\frac{\alpha^2}{x_B y Q^2} \frac{y^2}{(1-\varepsilon)} \left( 1 + \frac{\gamma^2}{2x_B} \right) \int \frac{d|\mathbf{b}_T|}{(2\pi)} |\mathbf{b}_T| \left\{ J_0(|\mathbf{b}_T||\mathbf{P}_{h\perp}|) \mathcal{F}_{UU,T} + \varepsilon J_0(|\mathbf{b}_T||\mathbf{P}_{h\perp}|) \mathcal{F}_{UU,L} \right. \\ &+ \sqrt{2\varepsilon(1+\varepsilon)} \cos\phi_h J_1(|\mathbf{b}_T||\mathbf{P}_{h\perp}|) \mathcal{F}_{UU}^{\cos\phi_h} + \varepsilon \cos(2\phi_h) J_2(|\mathbf{b}_T||\mathbf{P}_{h\perp}|) \mathcal{F}_{UU}^{\cos(2\phi_h)} \\ &+ \lambda_e \sqrt{2\varepsilon(1-\varepsilon)} \sin\phi_h J_1(|\mathbf{b}_T||\mathbf{P}_{h\perp}|) \mathcal{F}_{LU}^{\sin\phi_h} \\ &+ S_{\parallel} \left[ \sqrt{2\varepsilon(1+\varepsilon)} \sin\phi_h J_1(|\mathbf{b}_T||\mathbf{P}_{h\perp}|) \mathcal{F}_{UL}^{\sin\phi_h} + \varepsilon \sin(2\phi_h) J_2(|\mathbf{b}_T||\mathbf{P}_{h\perp}|) \mathcal{F}_{UL}^{\sin 2\phi_h} \right] \\ &+ S_{\parallel} \lambda_e \left[ \sqrt{1-\varepsilon^2} J_0(|\mathbf{b}_T||\mathbf{P}_{h\perp}|) \mathcal{F}_{LL} + \sqrt{2\varepsilon(1-\varepsilon)} \cos\phi_h J_1(|\mathbf{b}_T||\mathbf{P}_{h\perp}|) \mathcal{F}_{LL}^{\cos\phi_h} \right] \\ &+ |\mathbf{S}_{\perp}| \left[ \sin(\phi_h - \phi_S) J_1(|\mathbf{b}_T||\mathbf{P}_{h\perp}|) \left( \mathcal{F}_{UT,T}^{\sin(\phi_h - \phi_S)} + \varepsilon \mathcal{F}_{UT,L}^{\sin(\phi_h - \phi_S)} \right) \right. \\ &+ \varepsilon \sin(\phi_h + \phi_S) J_1(|\mathbf{b}_T||\mathbf{P}_{h\perp}|) \mathcal{F}_{UT}^{\sin(\phi_h + \phi_S)} \\ &+ \varepsilon \sin(3\phi_h - \phi_S) J_3(|\mathbf{b}_T||\mathbf{P}_{h\perp}|) \mathcal{F}_{UT}^{\sin(3\phi_h - \phi_S)} \\ &+ \sqrt{2\varepsilon(1+\varepsilon)} \sin\phi_S J_0(|\mathbf{b}_T||\mathbf{P}_{h\perp}|) \mathcal{F}_{UT}^{\sin\phi_S} \\ &+ \left. \sqrt{2\varepsilon(1+\varepsilon)} \sin(2\phi_h - \phi_S) J_2(|\mathbf{b}_T||\mathbf{P}_{h\perp}|) \mathcal{F}_{UT}^{\sin(2\phi_h - \phi_S)} \right] \\ &+ |\mathbf{S}_{\perp}| \lambda_e \left[ \sqrt{1-\varepsilon^2} \cos(\phi_h - \phi_S) J_1(|\mathbf{b}_T||\mathbf{P}_{h\perp}|) \mathcal{F}_{LT}^{\cos(\phi_h - \phi_S)} \right. \\ &+ \sqrt{2\varepsilon(1-\varepsilon)} \cos\phi_S J_0(|\mathbf{b}_T||\mathbf{P}_{h\perp}|) \mathcal{F}_{LT}^{\cos\phi_S} \\ &+ \left. \sqrt{2\varepsilon(1-\varepsilon)} \cos(2\phi_h - \phi_S) J_2(|\mathbf{b}_T||\mathbf{P}_{h\perp}|) \mathcal{F}_{LT}^{\cos(2\phi_h - \phi_S)} \right] \left. \right\} \quad (2.21)\end{aligned}$$

The structure of the cross section is what one gets from a multipole expansion in  $\mathbf{b}_T$ -space followed by a Fourier transform, see appendix B. Each of the structure functions  $\mathcal{F}_{\check{X}\check{Y}\check{Z}}$  in  $\mathbf{b}_T$ -space corresponds to the Hankel (or Fourier-Bessel) transform of the corresponding structure function  $F_{\check{X}\check{Y}\check{Z}}$  in the usual momentum space representation of the cross section. The combinations  $\sin(n\phi_h + \dots)J_n(|\mathbf{b}_T||\mathbf{P}_{h\perp}|)$  and  $\cos(n\phi_h + \dots)J_n(|\mathbf{b}_T||\mathbf{P}_{h\perp}|)$



act as basis functions of the combined transform to  $(|\mathbf{P}_{h\perp}|, \phi_h)$ -space. Due to the fact that the multipole expansion of the physical cross section terminates, only a finite number of terms appear in the cross section, with  $J_3$  being the Bessel function of highest order. The structures  $\mathcal{F}_{XY,Z}$  are functions of  $|\mathbf{b}_T|$ ,  $x$  and  $z$ , but no longer depend on the angular variables. Introducing a short-hand notation for products

$$\mathcal{P}[\tilde{f}^{(n)} \tilde{D}^{(m)}] \equiv x_B \sum_a e_a^2 (zM|\mathbf{b}_T|)^n (zM_h|\mathbf{b}_T|)^m \tilde{f}^{a(n)}(x, z^2 \mathbf{b}_T^2) \tilde{D}^{a(m)}(z, \mathbf{b}_T^2), \quad (2.22)$$

the leading twist tree level analysis in eqs. (2.10), (2.13) and (2.15) reveals that the Fourier transformed structures in the cross section are simple products of TMD PDFs and TMD FFs

$$\mathcal{F}_{UU,T} = \mathcal{P}[\tilde{f}_1^{(0)} \tilde{D}_1^{(0)}], \quad (2.23)$$

$$\mathcal{F}_{UT,T}^{\sin(\phi_h - \phi_S)} = -\mathcal{P}[\tilde{f}_{1T}^{\perp(1)} \tilde{D}_1^{(0)}], \quad (2.24)$$

$$\mathcal{F}_{LL} = \mathcal{P}[\tilde{g}_{1L}^{(0)} \tilde{D}_1^{(0)}], \quad (2.25)$$

$$\mathcal{F}_{LT}^{\cos(\phi_h - \phi_S)} = \mathcal{P}[\tilde{g}_{1T}^{(1)} \tilde{D}_1^{(0)}], \quad (2.26)$$

$$\mathcal{F}_{UT}^{\sin(\phi_h + \phi_S)} = \mathcal{P}[\tilde{h}_1^{(0)} \tilde{H}_1^{\perp(1)}], \quad (2.27)$$

$$\mathcal{F}_{UU}^{\cos(2\phi_h)} = \mathcal{P}[\tilde{h}_1^{\perp(1)} \tilde{H}_1^{\perp(1)}], \quad (2.28)$$

$$\mathcal{F}_{UL}^{\sin(2\phi_h)} = \mathcal{P}[\tilde{h}_{1L}^{\perp(1)} \tilde{H}_1^{\perp(1)}], \quad (2.29)$$

$$\mathcal{F}_{UT}^{\sin(3\phi_h - \phi_S)} = \frac{1}{4} \mathcal{P}[\tilde{h}_{1T}^{\perp(2)} \tilde{H}_1^{\perp(1)}]. \quad (2.30)$$

For completeness, we also list the above results in terms of the momentum-space structure functions  $F_{XY,Z}$  of ref. [8] in appendix D. Note that TMD evolution equations are typically derived in  $\mathbf{b}_T$ -space and are thus obtained in terms of the same (derivatives of) Fourier transformed TMD PDFs and TMD FFs that appear in the equations above, see, e.g., ref. [28], where a similar representation of the structure functions in Fourier space has been employed.

### 3 Beyond tree level

The formalism becomes more involved once diagrams beyond leading order in  $\alpha_s$  are taken into account. Various strategies have been proposed to address extra divergences that appear at the one loop level and higher order [15–19, 30–34]. The development of these frameworks for transverse momentum dependent factorization and the establishing of the corresponding factorization theorems is an active field of research (see e.g., refs. [15, 35]). The proposed strategies require the introduction of new variables that act as regularization scales, and most importantly as it pertains to the content of this paper, the so called soft factors coming from soft-gluon radiation. As stated in the introduction, depending on the framework, the soft factors appear explicitly in the structure functions [14, 18], or are absorbed into the definition of TMD PDFs and TMD FFs (see e.g., refs. [15, 19]). We will present general arguments that soft factors cancel in weighted asymmetries, independent

of the specific factorization framework; however for definiteness we work with the JMY framework [14, 18], which is based on the ideas of Collins, Soper, and Sterman for the factorization of  $e^+e^-$  and Drell Yan scattering [13, 30]. Again we consider the structure function giving rise to the Siverts asymmetry,

$$\mathcal{F}_{UT,T}^{\sin(\phi_h-\phi_S)} = H_{UT,T}^{\sin(\phi_h-\phi_S)}(Q^2, \mu^2, \rho) \tilde{S}^{(+)}(\mathbf{b}_T^2, \mu^2, \rho) \mathcal{P}[\tilde{f}_{1T}^{(1)} \tilde{D}_1^{(0)}] + \tilde{Y}_{UT,T}^{\sin(\phi_h-\phi_S)}(Q^2, \mathbf{b}_T^2) . \quad (3.1)$$

The first term in the following referred to as the ‘‘TMD expression’’, dominates in the region where  $|\mathbf{P}_{h\perp}|$  is small,  $|\mathbf{P}_{h\perp}|/z \approx Q_T \ll Q$ . The second term is necessary to properly describe the structure function for large transverse momentum, where  $Q_T \sim Q$ , and where fixed order perturbation theory and collinear factorization apply. Here  $H_{UT,T}^{\sin(\phi_h-\phi_S)}$  is the hard part, and  $\tilde{S}^{(+)}$  is a soft factor appearing explicitly in the structure function within the JMY formalism. It is the same in all the structure functions  $\mathcal{F}_{XY,Z}$ , see ref. [28]. All other structure functions of eqs. (2.23)–(2.30) need to be modified analogous to eq. (3.1).

The term  $\tilde{Y}_{UT,T}^{\sin(\phi_h-\phi_S)}(Q^2, \mathbf{b}_T^2)$  represents contributions that are relevant only in the region of large transverse momentum  $|\mathbf{P}_{h\perp}|$  [19, 36]. Qualitatively, this corresponds to the very small  $\mathbf{b}_T$  region,  $z|\mathbf{b}_T| \lesssim 1/Q$ . Since our aim is to study TMD PDFs, we want to focus on the region  $|\mathbf{P}_{h\perp}|/z \ll Q$  where we expect them to give the dominant contribution if  $z|\mathbf{b}_T| \gg 1/Q$ . Nevertheless, since we are considering weighted integrals of structure functions, the integrals do include the region of very large  $|\mathbf{P}_{h\perp}|$ . As a result, the  $\tilde{Y}$  term in eq. (3.1) is non-zero even if  $z|\mathbf{b}_T| \gg 1/Q$ . We note that the  $\tilde{Y}$  term is expected to be particularly important in the case of a ‘‘mismatch’’ between the tail of the TMD term and the  $|\mathbf{P}_{h\perp}|$ -behavior obtained from the collinear formalism in the regime of intermediate  $|\mathbf{P}_{h\perp}|$ , i.e.,  $M \ll Q_T \ll Q$ . Matches and mismatches between the collinear and TMD formalism have been discussed in detail in ref. [37]. An important example for the case of a mismatch is the  $\cos(2\phi_h)$  asymmetry. One possibility to avoid the discussion of the  $\tilde{Y}$ -term is to explicitly cut off the  $|\mathbf{P}_{h\perp}|$  integrals at some upper value  $\Lambda_{\text{TMD}}$ . This cutoff introduces an error in our extracted TMD expression, for which we give an estimate in appendix G.3. Another option is to simply ignore the  $\tilde{Y}$  term. This amounts to keeping the TMD term in the large  $|\mathbf{P}_{h\perp}|$  region, i.e., to include the large- $|\mathbf{P}_{h\perp}|$ -tail generated by the TMD term, which would otherwise be corrected by the  $Y$  term. In appendix G.3, we show that in the  $z|\mathbf{b}_T| \gg 1/Q$  region of interest this produces an error that falls off at least as a fractional inverse power with increasing  $|\mathbf{b}_T|$ . It should be mentioned that this estimate of the behavior of the error applies to the Bessel weighting which we discuss below. By contrast, no such error estimate exists for conventional weighting with powers of  $|\mathbf{P}_{h\perp}|$  since such integrals are divergent. Better error estimates, or equivalently, a better determination of the TMD region in  $\mathcal{B}_T$ , can be obtained by an explicit treatment of the  $\tilde{Y}$  term, which we will leave for future analyses.

In summary, we find that weighted integrals based on the TMD expression alone are valid only in a limited range of  $\mathcal{B}_T$ . Finally, beyond tree level, the product notation  $\mathcal{P}[fD]$  defined in eq. (2.22) has to be updated to include further dependences on the renormalization and cutoff parameters  $\mu^2$ ,  $\rho$ ,  $\zeta$  and  $\hat{\zeta}$  appearing in the JMY formalism

discussed in more detail below:<sup>1</sup>

$$\mathcal{P}[\tilde{f}^{(n)}\tilde{D}^{(m)}] \equiv x_B \sum_a e_a^2 (zM|\mathbf{b}_T|)^n (zM_h|\mathbf{b}_T|)^m \tilde{f}^{a(n)}(x, z^2\mathbf{b}_T^2, \mu^2, \zeta, \rho) \tilde{D}^{a(m)}(z, \mathbf{b}_T^2, \mu^2, \hat{\zeta}, \rho). \quad (3.2)$$

#### 4 TMD PDFs at the level of matrix elements

Apart from introducing the parameters  $\zeta$ ,  $\hat{\zeta}$  and  $\rho$  the purpose of this section is to review the formalism of Lorentz-invariant amplitudes underlying the decomposition of  $\tilde{\Phi}$  eq. (2.13). In the framework of JMY, the TMD correlator  $\Phi$  itself involves a soft factor  $S^{(+)}$  as already encountered above, i.e., eqs. (2.5) and (2.8) need to be modified. In the following, we label the unmodified correlators with the subscript “unsub”:

$$\Phi_{\text{unsub}}^{[\Gamma]}(p, P, S; v, \mu) = \int \frac{d^4b}{(2\pi)^4} e^{ip \cdot b} \frac{1}{2} \underbrace{\langle P, S | \bar{\psi}(0) \overbrace{\mathcal{U}[0, \infty v] \mathcal{U}[\infty v, b]}^{\mathcal{U}[\mathcal{C}_b]} \Gamma \psi(b) | P, S \rangle}_{\tilde{\Phi}_{\text{unsub}}^{[\Gamma]}(b, P, S; v, \mu)}. \quad (4.1)$$

The gauge link  $\mathcal{U}[\mathcal{C}_b]$  is essentially given by two parallel straight Wilson lines running out to infinity in the direction given by the four-vector  $v$  and back again. The definition of a straight Wilson line between two points  $a$  and  $b$  is

$$\mathcal{U}[a, b] \equiv \mathcal{P} \exp \left( -ig \int_a^b d\xi^\mu A_\mu(\xi) \right), \quad (4.2)$$

where  $A_\mu(\xi) = T^c A_\mu^c(\xi)$ ,  $c = 1 \dots 8$  is the (matrix valued) gauge field. A transverse link connecting these parallel Wilson lines at infinity can be omitted in the covariant gauge used by JMY. In case of SIDIS, the direction  $v = [v^-, v^+, 0]$  is slightly off the light-cone direction  $n_-$ , while for the Drell-Yan process  $v$  is slightly off the light cone direction  $-n_-$ . The shift away from the light cone is time-like in the JMY framework and specified in a Lorentz-invariant way by the parameter  $\zeta$ , defined by  $\zeta^2 = (2P \cdot v)^2/v^2$ . The parameter  $\zeta$  represents a rapidity cutoff parameter [30]. The above correlator can be parameterized in terms of real-valued Lorentz-invariant amplitudes. Here we restrict ourselves to the case  $\Gamma = \gamma^\mu$ . Reference [29] lists the following structures

$$\begin{aligned} \frac{1}{2} \Phi_{\text{unsub}}^{[\gamma^\mu]} &= P^\mu A_2^{(+)} + p^\mu A_3^{(+)} + \frac{1}{M} \epsilon^{\mu\nu\alpha\beta} P_\nu p_\alpha S_\beta A_{12}^{(+)} + \frac{M^2}{(v \cdot P)} v^\mu B_1^{(+)} \\ &+ \frac{M}{v \cdot P} \epsilon^{\mu\nu\alpha\beta} P_\nu v_\alpha S_\beta B_7^{(+)} + \frac{M}{v \cdot P} \epsilon^{\mu\nu\alpha\beta} p_\nu v_\alpha S_\beta B_8^{(+)} \\ &+ \frac{1}{M(v \cdot P)} (p \cdot S) \epsilon^{\mu\nu\alpha\beta} P_\nu p_\alpha v_\beta B_9^{(+)} + \frac{M}{(v \cdot P)^2} (v \cdot S) \epsilon^{\mu\nu\alpha\beta} P_\nu p_\alpha v_\beta B_{10}^{(+)}. \end{aligned} \quad (4.3)$$

The amplitudes  $B_i^{(+)}$  only appear when the dependence of the correlator on the direction  $v$  is explicitly taken into account, and were not listed in earlier works [1, 5]. Since  $v$  represents

---

<sup>1</sup>The framework of, e.g., ref. [19], would require analogous modifications within this formalism.

only a direction, the structures above should remain invariant under re-scaling of  $v$ , i.e., under the substitution  $v \rightarrow \eta v$ , for any positive real number  $\eta$ . This has been ensured by dividing by powers of  $v \cdot P$  in the expression above. The amplitudes can depend on  $p^2$ ,  $p \cdot P$ ,  $v \cdot p / (v \cdot P)$ ,  $v^2 / (v \cdot P)^2 = \zeta^{-2}$  and the sign of  $v \cdot P$ , which we denote with the superscript (+) (SIDIS case). For the Drell-Yan process,  $v \cdot P$  has the opposite sign (-).

For our discussion below, we make use of a similar decomposition as in eq. (4.3). However, instead of parameterizing the  $p$ -dependent correlator  $\Phi^{[\Gamma]}$ , we directly parameterize the  $b$ -dependent matrix elements  $\tilde{\Phi}^{[\Gamma]}$  of eq. (2.8) in terms of complex-valued amplitudes  $\tilde{A}_i^{(+)}$  and  $\tilde{B}_i^{(+)}$  that depend on  $b^2$ ,  $b \cdot P$ ,  $v \cdot b / (v \cdot P)$  and  $\zeta^{-2}$ . This parameterization in Fourier-space has already been employed in [38, 39].<sup>2</sup> As shown in appendix C, we can deduce this parameterization from eq. (4.3) using the substitution rule  $p \rightarrow -iM^2 b$ :

$$\begin{aligned} \frac{1}{2} \tilde{\Phi}_{\text{unsub}}^{[\Gamma]} = & P^\mu \tilde{A}_2^{(+)} - iM^2 b^\mu \tilde{A}_3^{(+)} - iM \epsilon^{\mu\nu\alpha\beta} P_\nu b_\alpha S_\beta \tilde{A}_{12}^{(+)} + \frac{M^2}{(v \cdot P)} v^\mu \tilde{B}_1^{(+)} \\ & + \frac{M}{v \cdot P} \epsilon^{\mu\nu\alpha\beta} P_\nu v_\alpha S_\beta \tilde{B}_7^{(+)} - \frac{iM^3}{v \cdot P} \epsilon^{\mu\nu\alpha\beta} b_\nu v_\alpha S_\beta \tilde{B}_8^{(+)} \\ & - \frac{M^3}{v \cdot P} (b \cdot S) \epsilon^{\mu\nu\alpha\beta} P_\nu b_\alpha v_\beta \tilde{B}_9^{(+)} - \frac{iM^3}{(v \cdot P)^2} (v \cdot S) \epsilon^{\mu\nu\alpha\beta} P_\nu b_\alpha v_\beta \tilde{B}_{10}^{(+)} . \end{aligned} \quad (4.4)$$

In order to connect to the framework of TMD PDFs, we integrate the correlator  $\Phi$  over the (suppressed) momentum component  $p^-$ . The integration with respect to  $p^-$  reduces the Fourier transform with respect to  $b^+$  to the evaluation of  $\tilde{\Phi}$  at  $b^+ = 0$ . Moreover, in the formalism of JMY, the defining correlator of TMD PDFs needs to be modified with a soft factor. The modified,  $p^-$ -integrated correlator reads

$$\begin{aligned} \Phi^{(+)[\Gamma]}(x, \mathbf{p}_T, P, S, \mu^2, \zeta, \rho) = & \int \frac{db^-}{(2\pi)} e^{ixb^- P^+} \int \frac{d^2 \mathbf{b}_T}{(2\pi)^2} e^{-i\mathbf{p}_T \cdot \mathbf{b}_T} \\ & \times \frac{1}{2} \langle P, S | \underbrace{\bar{\psi}(0) \mathcal{U}[\mathcal{C}_b] \Gamma \psi(b)}_{\tilde{\Phi}_{\text{unsub}}^{[\Gamma]}(b, P, S; v, \mu^2)} | P, S \rangle \Big/ \tilde{S}^{(+)}(\mathbf{b}_T^2, \mu^2, \rho) \Big|_{b^+ = 0}, \end{aligned} \quad (4.5)$$

where  $xP^+ = p^+$ . The soft factor is given as

$$\tilde{S}^{(+)}(\mathbf{b}_T^2, \mu^2, \rho) = \frac{1}{N_c} \langle 0 | \text{Tr}_c \{ \mathcal{U}[-\infty \tilde{v} + b_\perp, b_\perp] \mathcal{U}[b_\perp, b_\perp + \infty v] \mathcal{U}[\infty v, 0] \mathcal{U}[0, -\infty \tilde{v}] \} | 0 \rangle \quad (4.6)$$

and involves another time-like direction  $\tilde{v} = (\tilde{v}^-, \tilde{v}^+, 0)$  slightly off the light-cone direction  $n_+$ , controlled by the parameter  $\rho \equiv \sqrt{v^- \tilde{v}^+ / v^+ \tilde{v}^-}$ . Note that  $\rho^2 + 2 + \rho^{-2} = 4(v \cdot \tilde{v})^2 / v^2 \tilde{v}^2$  is a Lorentz-invariant expression. Here, the superscript (+) specifies the sign of  $v \cdot \tilde{v}$ , which is different for the SIDIS and the Drell-Yan process.

---

<sup>2</sup>In refs. [38, 39], a different convention for the position of the quark fields in the Fourier transformed correlator  $\tilde{\Phi}$  has been used. These references introduce  $\tilde{\Phi}$  as  $\tilde{\Phi}(l, P, S, \mathcal{C}) = \frac{1}{2} \langle P, S | \bar{\psi}(l) \mathcal{U}^\dagger[\mathcal{C}] \Gamma \psi(0) | P, S \rangle$ . In eq. (4.1) we stick to the more common convention of an operator  $\bar{\psi}(0) \dots \psi(b)$ . From translation invariance follows that the variable  $b$  corresponds to  $-l$  in refs. [38, 39]. In particular, our amplitudes  $\tilde{A}_i(b^2, b \cdot P, \dots)$  correspond to  $\tilde{A}_i(l^2, -l \cdot P, \dots)$  of refs. [38, 39].

In the formalism of JMY, the definition of the soft factor  $\tilde{S}^{(+)}$  above both applies to the occurrence of  $\tilde{S}^{(+)}$  in the TMD PDF correlator eq. (4.5) and in the structure function eq. (3.1). In the following, we will consider the case  $\Gamma = \gamma^+$ . The correlator  $\Phi^{(+)[\gamma^+]}$  can be decomposed into contributions from two distinct TMD PDFs:

$$\Phi^{(+)[\gamma^+]}(x, \mathbf{p}_T, P, S, \mu^2, \zeta, \rho) = f_1(x, \mathbf{p}_T^2; \mu^2, \zeta, \rho) - \frac{\epsilon_T^{ij} \mathbf{p}_{T i} \mathbf{S}_{T j}}{M} f_{1T}^\perp(x, \mathbf{p}_T^2; \mu^2, \zeta, \rho). \quad (4.7)$$

Strictly speaking,  $f_{1T}^\perp$  should also carry the superscript (+) since it has a different sign for Drell-Yan measurements [40]. We now use

$$\mathbf{b}_T^2 = -b^2 \Big|_{b^+=0}, \quad b^- = \frac{b \cdot P}{P^+} \Big|_{b^+=0}, \quad R(\zeta^2) \equiv \frac{M^2 v^+}{v \cdot P P^+} = 1 - \sqrt{1 - \frac{4M^2}{\zeta^2}}, \quad (4.8)$$

and insert the parameterization eq. (4.4) into eq. (4.5). Comparing with eq. (4.7) allows us to write the TMD PDFs  $f_1$  and  $f_{1T}^\perp$  as

$$f_1(x, \mathbf{p}_T^2; \mu^2, \zeta, \rho) = 2 \int \frac{d(b \cdot P)}{(2\pi)} e^{ix(b \cdot P)} \int_0^\infty \frac{d(-b^2)}{4\pi} J_0(\sqrt{-b^2 \mathbf{p}_T^2}) \times \frac{\tilde{A}_{2B}^{(+)} \left( b^2, b \cdot P, \frac{(b \cdot P)R(\zeta^2)}{M^2}, \zeta^{-2}, \mu^2 \right)}{\tilde{S}^{(+)}(-b^2, \mu^2, \rho)} \quad (4.9)$$

$$\begin{aligned} f_{1T}^\perp(x, \mathbf{p}_T^2; \mu^2, \zeta, \rho) &= 4M^2 \frac{\partial}{\partial(\mathbf{p}_T^2)} \int \frac{d(b \cdot P)}{(2\pi)} e^{ix(b \cdot P)} \int_0^\infty \frac{d(-b^2)}{4\pi} J_0(\sqrt{-b^2 \mathbf{p}_T^2}) \\ &\times \frac{\tilde{A}_{12B}^{(+)} \left( b^2, b \cdot P, \frac{(b \cdot P)R(\zeta^2)}{M^2}, \zeta^{-2}, \mu^2 \right)}{\tilde{S}^{(+)}(-b^2, \mu^2, \rho)} \\ &= 2M^2 \int \frac{d(b \cdot P)}{(2\pi)} e^{ix(b \cdot P)} \int_0^\infty \frac{d(-b^2)}{4\pi} \frac{J_1(\sqrt{-b^2 \mathbf{p}_T^2})}{\sqrt{-b^2 \mathbf{p}_T^2}} \\ &\times b^2 \frac{\tilde{A}_{12B}^{(+)} \left( b^2, b \cdot P, \frac{(b \cdot P)R(\zeta^2)}{M^2}, \zeta^{-2}, \mu^2 \right)}{\tilde{S}^{(+)}(-b^2, \mu^2, \rho)}, \end{aligned} \quad (4.10)$$

where

$$\tilde{A}_{2B}^{(+)} \equiv \tilde{A}_2^{(+)} + R(\zeta^2) \tilde{B}_1^{(+)}, \quad (4.11)$$

$$\tilde{A}_{12B}^{(+)} \equiv \tilde{A}_{12}^{(+)} - R(\zeta^2) \tilde{B}_8^{(+)}. \quad (4.12)$$

We observe that the amplitudes  $\tilde{B}_i$  give rise to structures in eqs. (4.9) and (4.10) that are suppressed by their explicit  $\zeta$ -dependence as  $\zeta \rightarrow \infty$ , i.e., in the limit of light-like  $v$ . The structures also disappear in the limit of vanishing nucleon mass  $M^2 \rightarrow 0$ . Notice that the two independent Fourier transforms in each of eqs. (4.9) and (4.10) naturally connect the TMD PDFs to a manifestly Lorentz-invariant framework and reveal  $x \leftrightarrow b \cdot P$  and  $\mathbf{p}_T^2 \leftrightarrow b^2$  to be pairs of conjugate variables.

The relations of the amplitudes  $\tilde{A}_{2B}^{(+)}$  and  $\tilde{A}_{12B}^{(+)}$  to the  $\mathbf{b}_T$ -Fourier-transformed TMD PDFs defined in eqs. (2.17) and (2.19) are given by

$$\tilde{f}_1^{(0)}(x, \mathbf{b}_T^2; \mu^2, \zeta, \rho) = \frac{2}{\tilde{S}^{(+)}(\mathbf{b}_T^2, \mu^2, \rho)} \int \frac{d(b \cdot P)}{(2\pi)} e^{ix(b \cdot P)} \tilde{A}_{2B}^{(+)} \left( -\mathbf{b}_T^2, b \cdot P, \frac{(b \cdot P)R(\zeta^2)}{M^2}, \zeta^{-2}, \mu^2 \right), \quad (4.13)$$

$$\tilde{f}_{1T}^{\perp(1)}(x, \mathbf{b}_T^2; \mu^2, \zeta, \rho) = \frac{-2}{\tilde{S}^{(+)}(\mathbf{b}_T^2, \mu^2, \rho)} \int \frac{d(b \cdot P)}{(2\pi)} e^{ix(b \cdot P)} \tilde{A}_{12B}^{(+)} \left( -\mathbf{b}_T^2, b \cdot P, \frac{(b \cdot P)R(\zeta^2)}{M^2}, \zeta^{-2}, \mu^2 \right). \quad (4.14)$$

We note that the soft factor would need to remain in the integrand if it were also dependent on  $v \cdot b / \sqrt{v^2}$ , i.e., the ‘‘angle’’ between the Wilson lines and the vector  $b$  separating the quark fields in the operator. The above result can also be obtained by comparison to the correlator  $\tilde{\Phi}^{[\gamma^+]}(x, \mathbf{b}_T)$  in eq. (2.13), where

$$\tilde{\Phi}^{(+)[\Gamma]}(x, \mathbf{b}_T, \mu^2, \zeta, \rho) \equiv \int \frac{d(b \cdot P)}{(2\pi)P^+} e^{ix(b \cdot P)} \frac{\tilde{\Phi}_{\text{unsub}}^{[\Gamma]}(b, P, S; v, \mu^2)}{\tilde{S}^{(+)}(-b^2, \mu^2, \rho)} \Big|_{b^+ = 0}. \quad (4.15)$$

## 5 Bessel-weighted asymmetries

As stated earlier, transverse momentum weighted asymmetries [5–7] provide a means to disentangle the convolutions in the cross section in a model independent way. Generally, the conventional weighted asymmetries are given by

$$A_{XY}^{\mathcal{W}} = \begin{cases} 2 \frac{\int d|\mathbf{P}_{h\perp}| |\mathbf{P}_{h\perp}| d\phi_h d\phi_S \mathcal{W}(|\mathbf{P}_{h\perp}|, \phi_h) d\sigma_{XY}}{\int d|\mathbf{P}_{h\perp}| |\mathbf{P}_{h\perp}| d\phi_h d\phi_S d\sigma_{XY}} & : \text{ for } XY = UU \\ 2 \frac{\int d|\mathbf{P}_{h\perp}| |\mathbf{P}_{h\perp}| d\phi_h d\phi_S \mathcal{W}(|\mathbf{P}_{h\perp}|, \phi_h, \phi_S) (d\sigma_{XY}^{\uparrow} - d\sigma_{XY}^{\downarrow})}{\int d|\mathbf{P}_{h\perp}| |\mathbf{P}_{h\perp}| d\phi_h d\phi_S (d\sigma_{XY}^{\uparrow} + d\sigma_{XY}^{\downarrow})} & : \text{ else,} \end{cases} \quad (5.1)$$

where the labels  $X, Y$  represent the polarization, ‘‘un’’ ( $U$ ), longitudinally ( $L$ ) and transversely ( $T$ ) of the beam and target, respectively. The angles  $\phi_S$  and  $\phi_h$  specify the directions of the hadron spin polarization and the transverse hadron momentum respectively, relative to the lepton scattering plane. In case of single or double spin asymmetries  $d\sigma_{XY}^{\downarrow}$  denotes the cross section with one of the polarizations opposite than for  $d\sigma_{XY}^{\uparrow}$ , such that the relevant structure function is projected out from eq. (2.21). We have introduced the short-hand notation  $\mathcal{W}$  which is a function containing various powers and  $\mathbf{P}_{h\perp}$  as well as angular dependences of the form  $\sin(m\phi_h \pm n\phi_S)$  or  $\cos(m\phi_h \pm n\phi_S)$ . For the conventional weighted Sivers asymmetry,  $\mathcal{W} \equiv w_1 \sin(\phi_h - \phi_S)$ , where  $w_1 = |\mathbf{P}_{h\perp}|/zM$  as in eq. (1.1).

Based on the expansion of the SIDIS cross section in terms of Bessel functions  $J_n$  of transverse momentum and impact parameter in eq. (2.21), we exploit the orthogonality to

generalize the weighting procedure. Now the weighting is of the form

$$A_{XY}^{\mathcal{W}}(\mathcal{B}_T) = \begin{cases} 2 \frac{\int d|\mathbf{P}_{h\perp}| |\mathbf{P}_{h\perp}| d\phi_h d\phi_S \mathcal{W}(|\mathbf{P}_{h\perp}|, \phi_h; \mathcal{B}_T) d\sigma_{XY}}{\int d|\mathbf{P}_{h\perp}| |\mathbf{P}_{h\perp}| d\phi_h d\phi_S J_0(|\mathbf{P}_{h\perp}| \mathcal{B}_T) d\sigma_{XY}} & : \text{ for } XY = UU \\ 2 \frac{\int d|\mathbf{P}_{h\perp}| |\mathbf{P}_{h\perp}| d\phi_h d\phi_S \mathcal{W}(|\mathbf{P}_{h\perp}|, \phi_h, \phi_S; \mathcal{B}_T) (d\sigma_{XY}^\uparrow - d\sigma_{XY}^\downarrow)}{\int d|\mathbf{P}_{h\perp}| |\mathbf{P}_{h\perp}| d\phi_h d\phi_S J_0(|\mathbf{P}_{h\perp}| \mathcal{B}_T) (d\sigma_{XY}^\uparrow + d\sigma_{XY}^\downarrow)} & : \text{ else,} \end{cases} \quad (5.2)$$

where the weight function  $\mathcal{W}$  corresponds to that of conventional weighted asymmetries, except that we replace

$$|\mathbf{P}_{h\perp}|^n \rightarrow J_n(|\mathbf{P}_{h\perp}| \mathcal{B}_T) n! \left( \frac{2}{\mathcal{B}_T} \right)^n. \quad (5.3)$$

As mentioned earlier, taking the asymptotic form of the Bessel function the conventional weights [6, 7] which are  $\propto |\mathbf{P}_{h\perp}|^n$  appear as the leading term of the Taylor expansion of the right hand side of eq. (5.3). Furthermore we note that the parameter  $\mathcal{B}_T > 0$  regularizes UV divergences in moments of TMD PDFs and FFs. More importantly, we will show that the parameter  $\mathcal{B}_T > 0$  allows us to scan TMD PDFs and TMD FFs in Fourier space. In fact, the form of eq. (5.2) already indicates that the weighting implements a Fourier-decomposition of the cross section in transverse momentum space.

Now we summarize the cancellation of the soft factor. We will illustrate this for the Sivvers Bessel-weighted asymmetry (for details see appendix F). One can see from eq. (2.21) that the appropriate weight for the Sivvers asymmetry is

$$\mathcal{W} = \frac{2 J_1(|\mathbf{P}_{h\perp}| \mathcal{B}_T)}{zM \mathcal{B}_T} \sin(\phi_h - \phi_S), \quad \text{i.e.,} \quad w_1 = \frac{2 J_1(|\mathbf{P}_{h\perp}| \mathcal{B}_T)}{zM \mathcal{B}_T}, \quad (5.4)$$

corresponding to  $|\mathbf{P}_{h\perp}|/zM$  in the limit  $|\mathbf{P}_{h\perp}| \ll 1/\mathcal{B}_T$ . Then the Bessel-weighted Sivvers asymmetry is

$$A_{UT}^{\frac{2 J_1(|\mathbf{P}_{h\perp}| \mathcal{B}_T)}{zM \mathcal{B}_T} \sin(\phi_h - \phi_S)}(\mathcal{B}_T) = 2 \frac{\int d|\mathbf{P}_{h\perp}| |\mathbf{P}_{h\perp}| d\phi_h d\phi_S \frac{2 J_1(|\mathbf{P}_{h\perp}| \mathcal{B}_T)}{zM \mathcal{B}_T} \sin(\phi_h - \phi_S) (d\sigma^\uparrow - d\sigma^\downarrow)}{\int d|\mathbf{P}_{h\perp}| |\mathbf{P}_{h\perp}| d\phi_h d\phi_S J_0(|\mathbf{P}_{h\perp}| \mathcal{B}_T) (d\sigma^\uparrow + d\sigma^\downarrow)}, \quad (5.5)$$

where the axially symmetric denominator is given by

$$\frac{2\alpha^2}{x_B y Q^2} \frac{y^2}{(1-\varepsilon)} \left( 1 + \frac{\gamma^2}{2x_B} \right) \int d|\mathbf{P}_{h\perp}| |\mathbf{P}_{h\perp}| d\phi_h d\phi_S J_0(|\mathbf{P}_{h\perp}| \mathcal{B}_T) \times \int \frac{d|\mathbf{b}_T|}{(2\pi)} |\mathbf{b}_T| J_0(|\mathbf{b}_T| |\mathbf{P}_{h\perp}|) \mathcal{F}_{UU,T}, \quad (5.6)$$

and from eq. (2.21) the numerator is

$$\frac{2\alpha^2}{x_B y Q^2} \frac{y^2}{(1-\varepsilon)} \left( 1 + \frac{\gamma^2}{2x_B} \right) \int d|\mathbf{P}_{h\perp}| |\mathbf{P}_{h\perp}| d\phi_h d\phi_S \frac{2 J_1(|\mathbf{P}_{h\perp}| \mathcal{B}_T)}{zM \mathcal{B}_T} \sin^2(\phi_h - \phi_S) \times \int \frac{d|\mathbf{b}_T|}{(2\pi)} |\mathbf{b}_T| J_1(|\mathbf{b}_T| |\mathbf{P}_{h\perp}|) \mathcal{F}_{UT}^{\sin(\phi_h - \phi_S)}. \quad (5.7)$$

Finally, making use of the closure relation of the Bessel function (see appendix E) we obtain for fixed  $x, y, z$ , cancellation of the soft factor  $S^{+(0)}(\mu^2, \rho)$  in eq. (3.1) from the Bessel-weighted Siverts asymmetry,

$$A_{UT,T}^{\frac{2J_1(|\mathbf{P}_{h\perp}| \mathcal{B}_T)}{zM\mathcal{B}_T} \sin(\phi_h - \phi_s)}(\mathcal{B}_T) = -2 \frac{\sum_a e_a^2 H_{UT,T}^{\sin(\phi_h - \phi_s)}(Q^2, \mu^2, \rho) \tilde{f}_{1T}^{\perp(1)a}(x, z^2 \mathcal{B}_T^2; \mu^2, \zeta, \rho) \tilde{D}_1^{(0)a}(z, \mathcal{B}_T^2; \mu^2, \hat{\zeta}, \rho)}{\sum_a e_a^2 H_{UU,T}(Q^2, \mu^2, \rho) \tilde{f}_1^{(0)a}(x, z^2 \mathcal{B}_T^2; \mu^2, \zeta, \rho) \tilde{D}_1^{(0)a}(z, \mathcal{B}_T^2; \mu^2, \hat{\zeta}, \rho)}. \quad (5.8)$$

Some comments are in order. First, if  $|\mathcal{B}_T|$  is large enough, the estimate  $\tilde{Y}_{UT,T}^{\sin(\phi_h - \phi_s)} \sim \mathcal{B}_T^{-1/2}$  can be applied and may indicate that the  $\tilde{Y}$ -terms are sufficiently suppressed to be neglected for practical purposes (see appendix G.3), which is what we have done in the above equation. The above result for the Siverts asymmetry can be generalized to any other asymmetry in the SIDIS cross section, eq. (2.21). We summarize those results with the full kinematic dependences in appendix F. Weighting with Bessel functions at various values of  $\mathcal{B}_T$  thus allows us to map out, ratios of Fourier-transformed TMD PDFs as well as azimuthal and spin asymmetries.

Secondly, the hard scattering factor  $H_{UT,T}^{\sin(\phi_h - \phi_s)}$  is expected to be the same as the unpolarized one  $H_{UU,T}$ , because the Siverts effect concerns unpolarized quarks which leads to unpolarized scattering on the partonic level. This expectation is confirmed in a recent calculation by Kang, Xiao and Yuan [41] at the one loop level, but should hold to all orders. Since this feature of the Siverts asymmetry is not shared by the other asymmetries, we will stick to writing  $H_{UT,T}^{\sin(\phi_h - \phi_s)}$  to avoid potential mistakes.

Thirdly, it is important to note that in the limit  $\mathcal{B}_T \rightarrow 0$ , the cancellation of the soft factor becomes trivial, since the soft factor  $\tilde{S}^+(\mathbf{b}_T, \mu, \rho)$  is unity at  $\mathbf{b}_T = 0$ . This has been shown in ref. [30], but it can also be seen easily from its formal definition in terms of Wilson lines given in eq. (4.6). Using  $\tilde{S}^{+(0)}(\mathbf{0}, \mu^2, \rho) = 1$  shows that the  $\mathbf{P}_{h\perp}$ -integrated cross section does not depend on the soft factor, as expected because the collinear factorization result should in principle be retrieved (after a proper regularization, which is a highly nontrivial matter as discussed in [31]). Due to the asymptotic properties of Bessel functions for small arguments, we recover conventional weighted asymmetries in the limit  $\mathcal{B}_T \rightarrow 0$

$$A_{UT,T}^{\frac{|\mathbf{P}_{h\perp}|}{zM} \sin(\phi_h - \phi_s)}(x_B, z, y) = -2 \frac{\sum_a e_a^2 H_{UT,T}^{\sin(\phi_h - \phi_s)}(Q^2, \mu^2, \rho) f_{1T}^{\perp(1)a}(x; \mu^2, \zeta, \rho) D_1^{a(0)}(z; \mu^2, \hat{\zeta}, \rho)}{\sum_a e_a^2 H_{UU,T}(Q^2, \mu^2, \rho) f_1^{a(0)}(x; \mu^2, \zeta, \rho) D_1^{a(0)}(z; \mu^2, \hat{\zeta}, \rho)}, \quad (5.9)$$

where  $f_{1T}^{\perp(1)a}$ ,  $f_1^{a(0)}$ , and  $D_1^{a(0)}$  are moments of TMD PDFs and fragmentation functions as defined in eq. (2.20). We caution the reader that these moments are not well-defined, since the corresponding integrals are known to fall off too slowly at large transverse momentum [37]. Furthermore, the arguments made earlier that the  $\tilde{Y}$ -terms are small are no longer applicable.



Lastly, we briefly address what is known about the energy scale dependence of the conventional weighted asymmetries. The current knowledge on this is limited to the one-loop level. Choosing the factorization scale  $\mu = Q$  removes the  $Q$  dependence from the hard scattering function  $H$  that is a function of  $\ln Q^2/\mu^2$ . This will lead to a  $Q$  dependence in the transverse moments of the TMD PDFs only [15, 19]. The scale dependence of  $f_1^{(0)}(x; Q^2)$  is known, assuming that a proper definition of the TMD PDF can be used, such that the zeroth moment corresponds to the collinear function  $f_1(x; Q^2)$  after the regularization is removed. The same applies to  $D_1^{(0)}(z; Q^2)$ . For the first moment of the Sivers function one can exploit that it is directly related to the Qiu-Sterman function  $T_F(x, x)$  [42] as shown in ref. [23]. The evolution equation of the Qiu-Sterman function has recently been obtained [43–46] allowing for evolution of the weighted Sivers asymmetry. The evolution of  $T_F(x, x)$  is not autonomous, since it depends not just on  $T_F(x, x)$  itself. This is true even in the large- $N_c$  limit, but in the large- $x$  limit it does become autonomous [46, 47]. It indicates that  $f_{1T}^{\perp(1)}(x)$  evolves logarithmically with  $Q^2$  just like  $f_1(x)$ , only falling off faster at a given  $x$  value as  $Q^2$  increases. The evolution has also been calculated for moments of other TMD PDFs such as  $h_1^{\perp(1)}$  [44, 48, 49] and is similar to that of  $f_{1T}^{\perp(1)}$  but simpler since nonsinglet. In addition, the evolution of the first moment of the Collins function,  $H_1^{\perp(1)}$  is calculated in [50, 51].

## 6 Average transverse momentum shift and Bessel-weighted counterpart

In a similar manner to section 5 we now consider the soft factor cancellation in the average transverse momentum shift of unpolarized quarks in a transversely polarized nucleon for a given longitudinal momentum fraction  $x$ . This shift is considered in [52] and defined by a ratio of the  $\mathbf{p}_T$ -weighted correlator:

$$\langle p_y(x) \rangle_{TU} = \frac{\int d^2\mathbf{p}_T \mathbf{p}_y \Phi^{(+)[\gamma^+]}(x, \mathbf{p}_T, P, S, \mu^2, \zeta, \rho)}{\int d^2\mathbf{p}_T \Phi^{(+)[\gamma^+]}(x, \mathbf{p}_T, P, S, \mu^2, \zeta, \rho)} \Bigg|_{S^\pm=0, \mathbf{S}_T=(1,0)} = M \frac{f_{1T}^{\perp(1)}(x; \mu^2, \zeta, \rho)}{f_1^{(0)}(x; \mu^2, \zeta, \rho)}, \quad (6.1)$$

where  $f_{1T}^{\perp(1)}$  and  $f_1^{(0)}$  are the moments defined in eqs. (2.20). Obviously, the average momentum shift is very similar in structure to the weighted asymmetry eq. (5.9). While the weighted asymmetries are accessible directly from the  $\mathbf{P}_{h\perp}$ -weighted cross section, the average transverse momentum shifts are obtained from the  $\mathbf{p}_T$ -weighted correlator and could in principle be accessible from weighted jet asymmetries. As already mentioned, the integrals defining the moments of TMD PDFs on the right hand side of the above equation are divergent without suitable regularization. In the following, we therefore generalize the above quantity, weighting with Bessel functions of  $|\mathbf{p}_T|$  instead. In particular, we replace

$$\mathbf{p}_y = |\mathbf{p}_T| \sin(\phi_p) \quad \longrightarrow \quad \frac{2J_1(|\mathbf{p}_T| \mathcal{B}_T)}{\mathcal{B}_T} \sin(\phi_p - \phi_S), \quad (6.2)$$

where  $\phi_S = 0$  for the choice  $\mathbf{S}_T = (1, 0)$  in eq. (6.1). The correlator  $\Phi^{(+)[\gamma^+]}$  reads in terms of the amplitudes  $\tilde{A}_i^{(+)}$  and  $B_i^{(+)}$ ,

$$\begin{aligned} \Phi^{(+)[\gamma^+]}(x, \mathbf{p}_T, P, S, \mu^2, \zeta, \rho) = & \not{x} \int_0^\infty \frac{d|\mathbf{b}_T|}{2\pi} |\mathbf{b}_T| \left\{ J_0(|\mathbf{b}_T| |\mathbf{p}_T|) 2\tilde{A}_{2B}^{(+)} / \tilde{S} \right. \\ & \left. - M |\mathbf{b}_T| |\mathbf{S}_T| \sin(\phi_p - \phi_S) J_1(|\mathbf{b}_T| |\mathbf{p}_T|) 2\tilde{A}_{12B}^{(+)} / \tilde{S} \right\}, \end{aligned} \quad (6.3)$$

where we abbreviate

$$\not{x} \equiv \int \frac{d(b \cdot P)}{(2\pi)} e^{ix(b \cdot P)}. \quad (6.4)$$

The Bessel-weighted analog of eq. (6.1) is thus

$$\begin{aligned} \langle p_y(x) \rangle_{TU}^{\mathcal{B}_T} & \equiv \frac{\int d|\mathbf{p}_T| |\mathbf{p}_T| \int d\phi_p \frac{2J_1(|\mathbf{p}_T| \mathcal{B}_T)}{\mathcal{B}_T} \sin(\phi_p - \phi_S) \Phi^{(+)[\gamma^+]}(x, \mathbf{p}_T, P, S, \mu^2, \zeta, \rho)}{\int d|\mathbf{p}_T| |\mathbf{p}_T| \int d\phi_p J_0(|\mathbf{p}_T| \mathcal{B}_T) \Phi^{(+)[\gamma^+]}(x, \mathbf{p}_T, P, S, \mu^2, \zeta, \rho)} \Big|_{|\mathbf{S}_T|=1} \\ & = -M \frac{\not{x} \tilde{A}_{12B}^{(+)} \left( -\mathcal{B}_T^2, b \cdot P, \frac{(b \cdot P) R(\zeta^2)}{M^2}, \zeta^{-2}, \mu^2 \right)}{\not{x} \tilde{A}_{2B}^{(+)} \left( -\mathcal{B}_T^2, b \cdot P, \frac{(b \cdot P) R(\zeta^2)}{M^2}, \zeta^{-2}, \mu^2 \right)} \\ & = M \frac{\tilde{f}_{1T}^{\perp(1)}(x, \mathcal{B}_T^2; \mu^2, \zeta, \rho)}{\tilde{f}_1^{(0)}(x, \mathcal{B}_T^2; \mu^2, \zeta, \rho)}. \end{aligned} \quad (6.5)$$

Again, the soft factors cancel. At this point, the independence of the soft factor on  $v \cdot b / \sqrt{v^2}$  is crucial. In the limit  $\mathcal{B}_T \rightarrow 0$ , we recover equation (6.1),  $\langle p_y(x) \rangle_{TU}^0 = \langle p_y(x) \rangle_{TU}$ , which we have thus shown to be formally free of any soft factor contribution. However, we caution the reader again that the expressions at  $\mathcal{B}_T = 0$  can be ill-defined without an additional regularization step.

We can go one step further and form ratios that are also integrated in  $x$ , with weights  $\exp(-ix\mathcal{B}_L)$ . For  $\mathcal{B}_L = 0$ , this is the same as taking the lowest  $x$ -moment that appears in the Burkardt sum rule [52]. The reason it is interesting to look at such quantities is their renormalization properties. Another motivation to discuss such quantities here is lattice QCD. Taking  $x$ -moments is a standard ingredient in lattice computations of nucleon structure, see e.g., ref. [53] for a review. First exploratory studies of TMD PDFs on the lattice [38, 39] focus to a large degree on computations of the lowest  $x$ -moment of distributions, but access to finite values of  $\mathcal{B}_L$  is also possible. By “integration over  $x$ ” we mean an integration over the entire support of the correlator; this includes contributions from negative  $x$  which correspond to anti-quark contributions, see e.g., ref. [5, 39] for details.

In particular, the  $x$ -integrals of the two TMD PDFs  $f_1$  and  $f_{1T}^\perp$  can be decomposed as

$$\begin{aligned} & \int_{-1}^1 dx e^{-ix\mathcal{B}_L} f_1(x, \mathbf{p}_T^2; \mu^2, \zeta, \rho) = \\ & \int_0^1 dx \{ e^{-ix\mathcal{B}_L} f_1(x, \mathbf{p}_T^2; \mu^2, \zeta, \rho) - e^{ix\mathcal{B}_L} \bar{f}_1(x, \mathbf{p}_T^2; \mu^2, \zeta, \rho) \}, \\ & \int_{-1}^1 dx e^{-ix\mathcal{B}_L} f_{1T}^\perp(x, \mathbf{p}_T^2; \mu^2, \zeta, \rho) = \\ & \int_0^1 dx \{ e^{-ix\mathcal{B}_L} f_{1T}^\perp(x, \mathbf{p}_T^2; \mu^2, \zeta, \rho) + e^{ix\mathcal{B}_L} \bar{f}_{1T}^\perp(x, \mathbf{p}_T^2; \mu^2, \zeta, \rho) \}, \end{aligned} \quad (6.6)$$

where  $\bar{f}_1$  and  $\bar{f}_{1T}^\perp$  are anti-quark distributions. In analogy to eq. (6.5), one can consider

$$\begin{aligned} \langle p_y \rangle_{TU}^{\mathcal{B}_L, \mathcal{B}_T} & \equiv \frac{\int d|\mathbf{p}_T| |\mathbf{p}_T| \int d\phi_p \frac{2J_1(|\mathbf{p}_T| \mathcal{B}_T)}{\mathcal{B}_T} \sin(\phi_p - \phi_S) \int dx e^{-ix\mathcal{B}_L} \Phi^{(+)[\gamma^+]}(x, \mathbf{p}_T, P, S, \mu^2, \zeta, \rho)}{\int d|\mathbf{p}_T| |\mathbf{p}_T| \int d\phi_p J_0(|\mathbf{p}_T| \mathcal{B}_T) \int dx e^{-ix\mathcal{B}_L} \Phi^{(+)[\gamma^+]}(x, \mathbf{p}_T, P, S, \mu^2, \zeta, \rho)} \\ & = M \frac{\int dx e^{-ix\mathcal{B}_L} \tilde{f}_{1T}^{\perp(1)}(x, \mathcal{B}_T^2; \mu^2, \zeta, \rho)}{\int dx e^{-ix\mathcal{B}_L} \tilde{f}_1^{(0)}(x, \mathcal{B}_T^2; \mu^2, \zeta, \rho)} \\ & = -M \frac{\tilde{A}_{12B}^{(+)} \left( -\mathcal{B}_T^2, \mathcal{B}_L, \mathcal{B}_L \frac{R(\zeta^2)}{M^2}, \zeta^{-2}, \mu^2 \right)}{\tilde{A}_{2B}^{(+)} \left( -\mathcal{B}_T^2, \mathcal{B}_L, \mathcal{B}_L \frac{R(\zeta^2)}{M^2}, \zeta^{-2}, \mu^2 \right)}. \end{aligned} \quad (6.7)$$

In this case, the cancellation of the soft factor occurs even for a soft factor that has a dependence on  $v \cdot b / \sqrt{v^2}$ . In the last line of eq. (6.7), the amplitudes in the numerator and denominator parameterize the same matrix element  $\tilde{\Phi}^{[\gamma^+]}(b, P, S; v, \mu)$ , hence they involve the same bi-local quark-quark operator. The work on renormalization properties of non-local operators involving Wilson lines in refs. [54–59] suggests that the operator  $\bar{\psi}(0) \mathcal{U}[\mathcal{C}_b] \psi(b)$  might renormalize multiplicatively for not too small  $|\mathbf{b}_T|$ , compare also [34]. As a result, the quantity  $\langle p_y \rangle_{TU}^{\mathcal{B}_T, \mathcal{B}_L}$  would be renormalization scheme and scale independent (up to the evolution with the rapidity cutoff parameter  $\zeta^2$ ), since all multiplicative renormalization factors would cancel in the ratio. This observation was already made in ref. [39] and is consistent with TMD factorization, which also involves only multiplicative renormalization for  $|\mathbf{P}_{h\perp}| \ll Q$  or  $\mathcal{B}_T \gg 1/Q$ . For smaller  $\mathcal{B}_T$ , mixing with gluonic operators is expected, as it is known that the Qiu-Sterman function for quarks and gluons (related to  $\tilde{f}_{1T}^{\perp(1)q,g}(x, \mathbf{b}_T^2 = 0)$ ) mix under changes of the scale [43–46], thereby preventing the cancellation of multiplicative factors in the ratio considered here. The properties of quantities like  $\langle p_y \rangle_{TU}^{\mathcal{B}_T, \mathcal{B}_L}$  remain to be studied more thoroughly. They could be interesting objects to make contact between theory predictions from, e.g., lattice QCD, and experiment.

## 7 Conclusions

We have shown that rewriting the SIDIS cross-section in coordinate space displays the important feature that structure functions become simple products of Fourier transformed TMD PDFs and FFs, or derivatives thereof. The angular structure of the cross section

naturally suggests weighting with Bessel functions in order to project out these Fourier-Bessel transformed distributions, which serve as well-defined replacements of the transverse moments entering conventional weighted asymmetries. In addition, Bessel-weighted asymmetries provide a unique opportunity to study nucleon structure in a model independent way due to the absence of the soft factor  $S^{+(0)}$ , which as we have shown cancels from these observables. This cancellation is based on the fact that the soft factor is flavor blind in hard processes, and it depends only on  $\mathbf{b}_T^2, \mu^2, \rho$ . Moreover, evolution equations for the distributions are typically calculated in terms of the (derivatives of) Fourier transformed TMD PDFs and FFs. As a result the study of the scale dependence of Bessel-weighted asymmetries should prove more straightforward. For the above stated reasons we propose Bessel-weighted asymmetries as clean observables to study the scale dependence of TMD PDFs and FFs at existing (HERMES, COMPASS, JLab) and future facilities (Electron Ion Collider, JLab 12 GeV). Our results are also easily generalized to other processes where TMD factorization is valid, such as  $e^+e^-$  annihilation and Drell-Yan processes.

### Acknowledgments

We thank Elke Aschenauer, Harut Avakian, Mert Aybat, Vladimir Braun, Matthias Burkardt, Maarten Buffing, John Collins, Markus Diehl, Rolf Ent, Philipp Hägler, Aram Kotzinian, Andreas Metz, Piet Mulders, Ted Rogers, and Feng Yuan for fruitful discussions. We are grateful for partial support from the Institute For Nuclear Theory (INT), University of Washington and to the organizers of the INT Workshop, "Gluons and the Quark Sea at High Energies: Distributions, Polarization, Tomography" where part of this work was undertaken. LG acknowledges support from U.S. Department of Energy under contract DE-FG02-07ER41460, and thanks the JLab theory group for support. Authored by Jefferson Science Associates, LLC under U.S. DOE Contract No. DE-AC05-06OR23177. The U.S. Government retains a non-exclusive, paid-up, irrevocable, world-wide license to publish or reproduce this manuscript for U.S. Government purposes.

### A Conventions and useful relations

In section 2 we use definitions for the kinematic variables and the ratio of longitudinal and transverse photon flux  $\epsilon$  as in ref. [8],

$$x_B = \frac{Q^2}{2P \cdot q}, \quad y = \frac{P \cdot q}{P \cdot l}, \quad z_h = \frac{P \cdot P_h}{P \cdot q}, \quad \gamma = \frac{2Mx}{Q}, \quad \epsilon = \frac{1 - y - \frac{1}{4}\gamma^2 y^2}{1 - y + \frac{1}{2}y^2 + \frac{1}{4}\gamma^2 y^2}, \quad (\text{A.1})$$

where  $M$  is the mass of the target nucleon. The off-collinearity of the process is characterized by the variable  $Q_T$  introduced through

$$q_T \equiv q + (1 + q_T^2/Q^2)xP - P_h/z, \quad Q_T \equiv \sqrt{-q_T^2} \quad (\text{A.2})$$

and for  $Q_T \ll Q$ , one finds  $|\mathbf{P}_{h\perp}| \approx zQ_T$ , see, e.g. [37]. The leptonic tensor is

$$L_{\mu\nu} = 2(l_\mu l'_\nu + l_\nu l'_\mu - (l \cdot l')g_{\mu\nu} + i\lambda_e \epsilon_{\mu\nu\alpha\beta} l^\alpha q^\beta), \quad (\text{A.3})$$

where we neglected the lepton mass. The hadronic tensor is

$$2MW^{\mu\nu} = \sum_X \int \frac{d^3\mathbf{P}_X}{(2\pi)^3 2P_X^0} \delta^{(4)}(q+P-P_X-P_h) \langle P, S | J_\mu(0) | P_X, P_h \rangle \langle P_X, P_h | J_\nu(0) | P, S \rangle. \quad (\text{A.4})$$

For an arbitrary four-vector  $\omega$ , we introduce the usual light-cone decomposition as

$$\omega^\mu = \omega^+ n_+^\mu + \omega^- n_-^\mu + \omega_T^\mu, \quad (\text{A.5})$$

where  $\omega^\pm = (\omega^0 \pm \omega^3)/\sqrt{2}$  and where the basis vectors  $n_+^\mu$  and  $n_-^\mu$  are

$$n_+^\mu = \frac{1}{\sqrt{2}}(1, 0, 0, 1), \quad n_-^\mu = \frac{1}{\sqrt{2}}(1, 0, 0, -1), \quad (\text{A.6})$$

such that  $n_\pm \cdot n_\mp = 1$ ,  $n_\pm \cdot n_\pm = 0$ ,  $n_\pm \cdot \omega_T = 0$ . Note that  $\omega_T \cdot \omega_T = -\omega_T^2$ . In the  $\gamma^*P$  center of mass frame with the proton three-momentum pointing in positive  $z$ -direction, we can decompose the proton and parton momenta as

$$\begin{aligned} P^\mu &= P^+ n_+^\mu + \frac{M^2}{2P^+} n_-^\mu, \\ p^\mu &= xP^+ n_+^\mu + \frac{p^2 + \mathbf{p}_T^2}{2xP^+} n_-^\mu + p_T^\mu, \end{aligned} \quad (\text{A.7})$$

where  $x = p^+/P^+$  is the quark light-cone momentum fraction.

Finally, using eq. (2.12) in section 2.2, we write the quark-quark correlator reconstructed from the trace projections

$$\begin{aligned} \tilde{\Phi} &= \frac{1}{2}\gamma_+ \tilde{\Phi}^{[\gamma^+]} - \frac{1}{2}\gamma_+ \gamma^5 \tilde{\Phi}^{[\gamma^+ \gamma^5]} - \frac{1}{4}i\sigma_{\alpha+} \gamma^5 \tilde{\Phi}^{[i\sigma^{\alpha+} \gamma^5]} + \frac{1}{2}\gamma_\beta \tilde{\Phi}^{[\gamma^\beta]} \\ &\quad - \frac{1}{2}\gamma_\beta \gamma^5 \tilde{\Phi}^{[\gamma^\beta \gamma^5]} - \frac{1}{4}i\sigma_{\alpha\beta} \gamma^5 \tilde{\Phi}^{[i\sigma^{\alpha\beta} \gamma^5]} + \frac{1}{2}\mathbf{1} \tilde{\Phi}^{[\mathbf{1}]}, \end{aligned} \quad (\text{A.8})$$

where  $\alpha = 1, 2$  and  $\beta = -, 1, 2$ . Here we have also included twist-3 and twist-4 terms. For the fragmentation correlator we have the following expression,

$$\begin{aligned} \tilde{\Delta} &= \frac{1}{2}\gamma_- \tilde{\Delta}^{[\gamma^-]} - \frac{1}{2}\gamma_- \gamma^5 \tilde{\Delta}^{[\gamma^- \gamma^5]} - \frac{1}{4}i\sigma_{\alpha-} \gamma^5 \tilde{\Delta}^{[i\sigma^{\alpha-} \gamma^5]} + \frac{1}{2}\gamma_\beta \tilde{\Delta}^{[\gamma^\beta]} \\ &\quad - \frac{1}{2}\gamma_\beta \gamma^5 \tilde{\Delta}^{[\gamma^\beta \gamma^5]} - \frac{1}{4}i\sigma_{\alpha\beta} \gamma^5 \tilde{\Delta}^{[i\sigma^{\alpha\beta} \gamma^5]} + \frac{1}{2}\mathbf{1} \tilde{\Delta}^{[\mathbf{1}]}. \end{aligned} \quad (\text{A.9})$$

## B Multipole expansion and Fourier transform

This appendix shows the simple underlying mathematical structure of eq. (2.21). Let us treat all kinematic variables except for  $\mathbf{P}_{h\perp}$  and  $\phi_h$  as constants. Consider the cross section  $\tilde{\sigma}$  in Fourier space as some arbitrary function that depends on  $\mathbf{b}_T$ . This dependence can be formulated in coordinate space ( $|\mathbf{b}_T|, \phi_b$ ),

$$\tilde{\sigma}(|\mathbf{b}_T|, \phi_b) = \sum_{n=-\infty}^{\infty} e^{in\phi_b} \tilde{\sigma}_n(|\mathbf{b}_T|) \quad (\text{B.1})$$

which is nothing but a multipole expansion with  $|\mathbf{b}_T|$  dependent coefficients  $\tilde{\sigma}_n$ . Performing a Fourier-transform of  $\tilde{\sigma}$  back to momentum space ( $|\mathbf{P}_{h\perp}|, \phi_h$ ), we obtain

$$\begin{aligned}
 \sigma(|\mathbf{P}_{h\perp}|, \phi_h) &= \int \frac{d^2\mathbf{b}_T}{(2\pi)^2} e^{-i\mathbf{P}_{h\perp}\cdot\mathbf{b}_T} \tilde{\sigma}(\mathbf{b}_T) \\
 &= \int \frac{d|\mathbf{b}_T|}{2\pi} |\mathbf{b}_T| \int_0^{2\pi} \frac{d\phi_b}{2\pi} e^{-i|\mathbf{P}_{h\perp}||\mathbf{b}_T|\cos(\phi_h-\phi_b)} \sum_{n=-\infty}^{\infty} e^{in\phi_b} \tilde{\sigma}_n(|\mathbf{b}_T|) \\
 &= \sum_{n=-\infty}^{\infty} e^{in\phi_h} \int \frac{d|\mathbf{b}_T|}{2\pi} |\mathbf{b}_T| (-i)^n J_n(|\mathbf{P}_{h\perp}||\mathbf{b}_T|) \tilde{\sigma}_n(|\mathbf{b}_T|). \tag{B.2}
 \end{aligned}$$

Using  $e^{in\phi_h} = \cos(n\phi_h) + i\sin(n\phi_h)$  and  $J_n = (-1)^n J_{-n}$ , it is evident that the last line of the above equation has exactly the form of the cross section eq. (2.21), where a finite number of the  $\tilde{\sigma}_{\pm n}$  is given by simple linear combinations of the structure functions  $\mathcal{F}_{XY,Z}^{\dots}$ , and the rest is zero. In our case, the Bessel function  $J_n$  with the highest  $n$  is  $J_3$ , which appears in combination with the angular  $\sin(3\phi_h - \phi_S)$  modulation and turns out to be associated with the quadrupole deformation of parton densities  $h_{1T}^\perp$ .

### C Parameterization of the correlator in $\mathbf{b}$ -space

First, we briefly review the relevant properties of the correlator under symmetry transformations. Applying Lorentz transformations ( $L$ ), parity transformation ( $P$ ), time-reversal ( $T$ ) and hermitian conjugation ( $\dagger$ ) to the matrix elements, we find that the correlator fulfills

$$(L) : \quad \Phi_{\text{unsub}}^{[\Gamma]}(p, P, S; v, \mu) = \Phi_{\text{unsub}}^{[\Lambda^{-1/2}\Gamma\Lambda_{1/2}]}(\Lambda p, \Lambda P, \Lambda S; \Lambda v, \mu), \tag{C.1}$$

$$(P) : \quad \Phi_{\text{unsub}}^{[\Gamma]}(p, P, S; v, \mu) = \Phi_{\text{unsub}}^{[\gamma^0\Gamma\gamma^0]}(\bar{p}, \bar{P}, -\bar{S}; \bar{v}, \mu), \tag{C.2}$$

$$(T) : \quad \left[ \Phi_{\text{unsub}}^{[\Gamma]}(p, P, S; v, \mu) \right]^* = \Phi_{\text{unsub}}^{[\gamma^1\gamma^3\Gamma^*\gamma^3\gamma^1]}(\bar{p}, \bar{P}, \bar{S}; -\bar{v}, \mu), \tag{C.3}$$

$$(\dagger) : \quad \left[ \Phi_{\text{unsub}}^{[\Gamma]}(p, P, S; v, \mu) \right]^* = \Phi_{\text{unsub}}^{[\gamma^0\Gamma^\dagger\gamma^0]}(p, P, S; v, \mu). \tag{C.4}$$

where we denote the sign change of spatial components of a given vector  $c$ ; that is,  $\bar{c} \equiv (c^0, -c^1, -c^2, -c^3)$ . From hermiticity ( $\dagger$ ) follows that the  $A_i^{(+)}$  and  $B_i^{(+)}$  in eq. (4.3) are real valued. Time reversal ( $T$ ) does not constrain the number of allowed structures, because it changes the sign of  $v\cdot P$ . Instead, time reversal ( $T$ ) establishes relations between SIDIS amplitudes  $A_i^{(+)}$ ,  $B_i^{(+)}$  and Drell-Yan amplitudes  $A_i^{(-)}$ ,  $B_i^{(-)}$ .

For any of the transformations  $\mathcal{T} \in \{L, P, T, \dagger\}$ , the eqs. (C.1)–(C.4) are of the general form

$$\mathcal{T}_\Phi(\Phi(p, w)) = \Phi(\mathcal{T}_p(p), \mathcal{T}_w(w)) \tag{C.5}$$

where we have omitted the subscript “unsub” and the renormalization scale  $\mu$ , and where the symbol  $w$  summarizes all dependences on  $\Gamma$ ,  $P$ ,  $S$  and  $v$ . Here  $\mathcal{T}_\Phi$  is either the identity function or complex conjugation. The transformation rule  $\mathcal{T}_p(p)$  maps onto  $\Lambda p$ ,  $p$  or  $\bar{p}$  and thus fulfills  $a\cdot b = \mathcal{T}_p(a)\cdot\mathcal{T}_p(b)$  for any two vectors  $a$  and  $b$ . The Fourier-transformed correlator

$$\tilde{\Phi}(b, w) = \int d^4p e^{-ip\cdot b} \Phi(p, w) \tag{C.6}$$

transforms according to

$$\begin{aligned}
 \mathcal{T}_\Phi \left( \tilde{\Phi}(b, w) \right) &= \int d^4 p \, e^{\mathcal{T}_\Phi(-i)p \cdot b} \mathcal{T}_\Phi \left( \Phi(p, w) \right) \\
 &= \int d^4 q \, e^{\mathcal{T}_\Phi(-i)\mathcal{T}_p^{-1}(q) \cdot b} \Phi(q, \mathcal{T}_w(w)) \\
 &= \int d^4 q \, e^{\mathcal{T}_\Phi(-i)q \cdot \mathcal{T}_p(b)} \Phi(q, \mathcal{T}_w(w)) \\
 &= \tilde{\Phi} \left( \frac{\mathcal{T}_\Phi(i)}{i} \mathcal{T}_p(b), \mathcal{T}_w(w) \right). \tag{C.7}
 \end{aligned}$$

For example,  $\tilde{\Phi}$  transforms under hermitian conjugation as

$$(\dagger) : \quad \left[ \tilde{\Phi}_{\text{unsub}}^{[\Gamma]}(b, P, S; v) \right]^* = \tilde{\Phi}_{\text{unsub}}^{[\gamma^0 \Gamma^\dagger \gamma^0]}(-b, P, S; v). \tag{C.8}$$

Let  $f(p, w)$  be any of the structures preceding the invariant amplitudes in the parameterization of  $\Phi$ . The structure  $f(p, w)$  is a homogeneous function of some degree  $n$  in  $p$ , i.e.,  $f(\alpha p, w) = \alpha^n f(p, w)$  for any number  $\alpha$ . For example, the structure  $f(p, w) = \frac{1}{M(v \cdot P)}(p \cdot S) \epsilon^{\mu\nu\alpha\beta} P_\nu p_\alpha v_\beta$  preceding  $B_9^{(+)}$  in eq. (4.3) has degree  $n = 2$ . If we define  $\tilde{f}(b, w) \equiv f(-iM^2 b, w)$ , then

$$\mathcal{T}_\Phi \left( \tilde{f}(b, w) \right) = \mathcal{T}_\Phi(-iM^2)^n \mathcal{T}_\Phi(f(b, w)) = f(\mathcal{T}_\Phi(-iM^2)\mathcal{T}_p(b), \mathcal{T}_w(w)) = \tilde{f} \left( \frac{\mathcal{T}_\Phi(i)}{i} b, w \right). \tag{C.9}$$

This shows that  $\tilde{f}$  transforms like  $\tilde{\Phi}$  in eq. (C.7). We conclude that the parameterization of  $\tilde{\Phi}$  can be found by the substitution  $p \rightarrow -iM^2 b$  in the structures parameterizing  $\Phi$ , and we arrive at eq. (4.4). The amplitudes  $\tilde{A}_i^{(+)}$  and  $\tilde{B}_i^{(+)}$  introduced this way are no longer constrained to be real valued functions. Instead, hermitian conjugation eq. (C.8) yields the relation

$$\left[ \tilde{A}_i^{(+)}(b^2, b \cdot P, v \cdot b / (v \cdot P), \zeta^{-2}, \mu^2) \right]^* = \tilde{A}_i^{(+)}(b^2, -b \cdot P, -v \cdot b / (v \cdot P), \zeta^{-2}, \mu^2). \tag{C.10}$$

## D Structure functions in terms of Fourier transformed TMD PDFs and FFs

The structure functions of ref. [8] can be expressed in terms of Fourier-transformed TMD PDFs and FFs as

$$F_{UU,T} = x_B \sum_a e_a^2 \int \frac{d|\mathbf{b}_T|}{(2\pi)} |\mathbf{b}_T| J_0(|\mathbf{b}_T| |\mathbf{P}_{h\perp}|) \tilde{f}_1^a(x, z^2 \mathbf{b}_T^2) \tilde{D}_1^a(z, \mathbf{b}_T^2), \tag{D.1}$$

$$F_{UT,T}^{\sin(\phi_h - \phi_s)} = -x_B \sum_a e_a^2 \int \frac{d|\mathbf{b}_T|}{(2\pi)} |\mathbf{b}_T|^2 J_1(|\mathbf{b}_T| |\mathbf{P}_{h\perp}|) Mz \tilde{f}_{1T}^{\perp a(1)}(x, z^2 \mathbf{b}_T^2) \tilde{D}_1^a(z, \mathbf{b}_T^2), \tag{D.2}$$

$$F_{LL} = x_B \sum_a e_a^2 \int \frac{d|\mathbf{b}_T|}{(2\pi)} |\mathbf{b}_T| J_0(|\mathbf{b}_T| |\mathbf{P}_{h\perp}|) \tilde{g}_{1L}^a(x, z^2 \mathbf{b}_T^2) \tilde{D}_1^a(z, \mathbf{b}_T^2), \tag{D.3}$$

$$F_{LT}^{\cos(\phi_h - \phi_s)} = x_B \sum_a e_a^2 \int \frac{d|\mathbf{b}_T|}{(2\pi)} |\mathbf{b}_T|^2 J_1(|\mathbf{b}_T| |\mathbf{P}_{h\perp}|) Mz \tilde{g}_{1T}^{\perp a(1)}(x, z^2 \mathbf{b}_T^2) \tilde{D}_1^a(z, \mathbf{b}_T^2), \tag{D.4}$$

$$F_{UT}^{\sin(\phi_h + \phi_S)} = x_B \sum_a e_a^2 \int \frac{d|\mathbf{b}_T|}{(2\pi)} |\mathbf{b}_T|^2 J_1(|\mathbf{b}_T| |\mathbf{P}_{h\perp}|) M_h z \tilde{h}_1^a(x, z^2 \mathbf{b}_T^2) \tilde{H}_1^{\perp a(1)}(z, \mathbf{b}_T^2), \quad (\text{D.5})$$

$$F_{UU}^{\cos(2\phi_h)} = x_B \sum_a e_a^2 \int \frac{d|\mathbf{b}_T|}{(2\pi)} |\mathbf{b}_T|^3 J_2(|\mathbf{b}_T| |\mathbf{P}_{h\perp}|) M M_h z^2 \tilde{h}_1^{\perp a(1)}(x, z^2 \mathbf{b}_T^2) \tilde{H}_1^{\perp a(1)}(z, \mathbf{b}_T^2), \quad (\text{D.6})$$

$$F_{UL}^{\sin(2\phi_h)} = x_B \sum_a e_a^2 \int \frac{d|\mathbf{b}_T|}{(2\pi)} |\mathbf{b}_T|^3 J_2(|\mathbf{b}_T| |\mathbf{P}_{h\perp}|) M M_h z^2 \tilde{h}_{1L}^{\perp a(1)}(x, z^2 \mathbf{b}_T^2) \tilde{H}_1^{\perp a(1)}(z, \mathbf{b}_T^2), \quad (\text{D.7})$$

$$F_{UT}^{\sin(3\phi_h - \phi_S)} = x_B \sum_a e_a^2 \int \frac{d|\mathbf{b}_T|}{(2\pi)} |\mathbf{b}_T|^4 J_3(|\mathbf{b}_T| |\mathbf{P}_{h\perp}|) \frac{M^2 M_h z^3}{4} \tilde{h}_{1T}^{\perp a(2)}(x, z^2 \mathbf{b}_T^2) \tilde{H}_1^{\perp a(1)}(z, \mathbf{b}_T^2). \quad (\text{D.8})$$

## E Cancellation of the soft factor in the Siverson asymmetry

Making use of the closure relation of the Bessel function

$$\int_0^\infty d|\mathbf{P}_{h\perp}| |\mathbf{P}_{h\perp}| J_n(|\mathbf{P}_{h\perp}| |\mathbf{b}_T|) J_n(|\mathbf{P}_{h\perp}| \mathcal{B}_T) = \frac{1}{\mathcal{B}_T} \delta(|\mathbf{b}_T| - \mathcal{B}_T), \quad (\text{E.1})$$

we obtain for the expression in eq. (5.6)

$$\begin{aligned} & \int d|\mathbf{P}_{h\perp}| |\mathbf{P}_{h\perp}| d\phi_h d\phi_S J_0(|\mathbf{P}_{h\perp}| \mathcal{B}_T) \int \frac{d|\mathbf{b}_T|}{(2\pi)} |\mathbf{b}_T| J_0(|\mathbf{b}_T| |\mathbf{P}_{h\perp}|) \mathcal{F}_{UU,T} \\ &= x_B \sum_a e_a^2 H_{UU,T}(Q^2, \mu^2, \rho) \int d|\mathbf{P}_{h\perp}| |\mathbf{P}_{h\perp}| \int d\phi_h \int d\phi_S J_0(|\mathbf{P}_{h\perp}| \mathcal{B}_T) \\ & \quad \times \int \frac{d|\mathbf{b}_T|}{(2\pi)} |\mathbf{b}_T| J_0(|\mathbf{P}_{h\perp}| |\mathbf{b}_T|) \tilde{f}_1^{(0)a}(x, z^2 \mathbf{b}_T^2; \mu^2, \zeta, \rho) \tilde{S}^{(+)}(\mathbf{b}_T^2; \mu^2, \rho) \tilde{D}_1^{(0)a}(z, \mathbf{b}_T^2; \mu, \hat{\zeta}, \rho) \\ &= 2\pi x_B \sum_a e_a^2 H_{UU,T}(Q^2, \mu^2, \rho) \tilde{f}_1^{(0)a}(x, z^2 \mathcal{B}_T^2; \mu^2, \zeta, \rho) \tilde{S}^{(+)}(\mathcal{B}_T^2; \mu^2, \rho) \tilde{D}_1^{(0)a}(z, \mathcal{B}_T^2; \mu, \hat{\zeta}, \rho) \end{aligned} \quad (\text{E.2})$$

Next, we consider the following expression in the numerator of the asymmetry, eq. (5.7),

$$\begin{aligned} & \int d|\mathbf{P}_{h\perp}| |\mathbf{P}_{h\perp}| \int d\phi_h \int d\phi_S \frac{2J_1(|\mathbf{P}_{h\perp}| \mathcal{B}_T)}{z M \mathcal{B}_T} \sin^2(\phi_h - \phi_S) \\ & \quad \times \int \frac{d|\mathbf{b}_T|}{(2\pi)} |\mathbf{b}_T|^2 J_1(|\mathbf{b}_T| |\mathbf{P}_{h\perp}|) \mathcal{F}_{UT,T}^{\sin(\phi_h - \phi_S)} \\ &= \int d|\mathbf{P}_{h\perp}| |\mathbf{P}_{h\perp}| \int d\phi_h \int d\phi_S \frac{2J_1(|\mathbf{P}_{h\perp}| \mathcal{B}_T)}{z M \mathcal{B}_T} \sin^2(\phi_h - \phi_S) \\ & \quad \times x_B \sum_a e_a^2 H_{UT,T}^{\sin(\phi_h - \phi_S)}(Q^2, \mu^2, \rho) \int \frac{d|\mathbf{b}_T|}{(2\pi)} |\mathbf{b}_T|^2 J_1(|\mathbf{b}_T| |\mathbf{P}_{h\perp}|) \\ & \quad \times M z \tilde{f}_{1T}^{\perp(1)a}(x, z^2 \mathbf{b}_T^2, \mu^2, \zeta, \rho) \tilde{S}^{(+)}(\mathbf{b}_T^2, \mu^2, \rho) \tilde{D}_1^{(0)a}(z, \mathbf{b}_T^2, \mu^2, \hat{\zeta}, \rho) \\ &= 2\pi x_B \sum_a e_a^2 H_{UT,T}^{\sin(\phi_h - \phi_S)}(Q^2, \mu^2, \rho) \tilde{f}_{1T}^{\perp(1)a}(x, z^2 \mathcal{B}_T^2, \mu^2, \zeta, \rho) \\ & \quad \times \tilde{S}^{(+)}(\mathcal{B}_T^2, \mu^2, \rho) \tilde{D}_1^{(0)a}(z, \mathcal{B}_T^2, \mu^2, \hat{\zeta}/z, \rho), \end{aligned} \quad (\text{E.3})$$



where we have used the closure relation eq. (E.1), and

$$\int_0^{2\pi} \cos^2(m\phi_h + n\phi_S) d\phi_S = \int_0^{2\pi} \sin^2(m\phi_h + n\phi_S) d\phi_S = \pi, \quad (\text{E.4})$$

for integer  $n$  and  $m$ . Thus, we obtain

$$\begin{aligned} A_{UT,T}^{\frac{2J_1(|\mathbf{P}_{h\perp}| \mathcal{B}_T)}{zM\mathcal{B}_T} \sin(\phi_h - \phi_S)}(\mathcal{B}_T) &= -2 \frac{\frac{\alpha^2}{yQ^2} \frac{y^2}{(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x_B}\right)}{\frac{\alpha^2}{yQ^2} \frac{y^2}{(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x_B}\right)} \\ &\times \frac{\sum_a e_a^2 H_{UT,T}^{\sin(\phi_h - \phi_S)}(Q^2, \mu^2, \rho) \tilde{f}_{1T}^{\perp(1)a}(x, z^2 \mathcal{B}_T^2; \mu^2, \zeta, \rho) \tilde{S}^+(\mathcal{B}_T^2, \mu^2, \rho) \tilde{D}_1^{(0)a}(z, \mathcal{B}_T^2; \mu^2, \hat{\zeta}, \rho)}{\sum_a e_a^2 H_{UU,T}(Q^2, \mu^2, \rho) \tilde{f}_1^{(0)a}(x, z^2 \mathcal{B}_T^2; \mu^2, \zeta, \rho) \tilde{S}^+(\mathcal{B}_T^2, \mu^2, \rho) \tilde{D}_1^{(0)a}(z, \mathcal{B}_T^2; \mu^2, \hat{\zeta}, \rho)}, \end{aligned} \quad (\text{E.5})$$

## F Bessel-weighted asymmetries

Here we introduce the Bessel weights  $w_n$

$$w_n \equiv J_n(|\mathbf{P}_{h\perp}| \mathcal{B}_T) n! \left(\frac{2}{\mathcal{B}_T}\right)^n, \quad (\text{F.1})$$

and summarize the Bessel-weighted asymmetries at leading twist:

Double Spin

$$\begin{aligned} A_{LL}^{J_0(|\mathbf{P}_{h\perp}| \mathcal{B}_T)}(\mathcal{B}_T) &= 2 \frac{\frac{\alpha^2}{yQ^2} \frac{y^2}{(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x_B}\right) \sqrt{1-\varepsilon^2}}{\frac{\alpha^2}{yQ^2} \frac{y^2}{(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x_B}\right)} \\ &\times \frac{\sum_a e_a^2 H_{LL}(Q^2, \mu^2, \rho) \tilde{g}_{1L}^{(0)a}(x, z^2 \mathcal{B}_T^2; \mu^2, \zeta, \rho) \tilde{D}_1^{(0)a}(z, \mathcal{B}_T^2; \mu^2, \hat{\zeta}, \rho)}{\sum_a e_a^2 H_{UU,T}(Q^2, \mu^2, \rho) \tilde{f}_1^{(0)a}(x, z^2 \mathcal{B}_T^2; \mu^2, \zeta, \rho) \tilde{D}_1^{(0)a}(z, \mathcal{B}_T^2; \mu^2, \hat{\zeta}, \rho)}, \end{aligned} \quad (\text{F.2})$$

Worm Gear

$$\begin{aligned} A_{LT}^{\frac{2J_1(|\mathbf{P}_{h\perp}| \mathcal{B}_T)}{zM\mathcal{B}_T} \cos(\phi_h - \phi_S)}(\mathcal{B}_T) &= 2 \frac{\frac{\alpha^2}{yQ^2} \frac{y^2}{(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x_B}\right) \sqrt{1-\varepsilon^2}}{\frac{\alpha^2}{yQ^2} \frac{y^2}{(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x_B}\right)} \\ &\times \frac{\sum_a e_a^2 H_{LT}^{\cos(\phi_h - \phi_S)}(Q^2, \mu^2, \rho) \tilde{g}_{1T}^{(1)a}(x, z^2 \mathcal{B}_T^2; \mu^2, \zeta, \rho) \tilde{D}_1^{(0)a}(z, \mathcal{B}_T^2; \mu^2, \hat{\zeta}, \rho)}{\sum_a e_a^2 H_{UU,T}(Q^2, \mu^2, \rho) \tilde{f}_1^{(0)a}(x, z^2 \mathcal{B}_T^2; \mu^2, \zeta, \rho) \tilde{D}_1^{(0)a}(z, \mathcal{B}_T^2; \mu^2, \hat{\zeta}, \rho)}, \end{aligned} \quad (\text{F.3})$$

Collins

$$\begin{aligned} A_{UT}^{\frac{2J_1(|\mathbf{P}_{h\perp}| \mathcal{B}_T)}{zM\mathcal{B}_T} \sin(\phi_h + \phi_S)}(\mathcal{B}_T) &= 2 \frac{\frac{\alpha^2}{yQ^2} \frac{y^2}{(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x_B}\right) \varepsilon}{\frac{\alpha^2}{yQ^2} \frac{y^2}{(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x_B}\right)} \\ &\times \frac{\sum_a e_a^2 H_{UT}^{\sin(\phi_h + \phi_S)}(Q^2, \mu^2, \rho) \tilde{h}_1^{(0)a}(x, z^2 \mathcal{B}_T^2; \mu^2, \zeta, \rho) \tilde{H}_1^{\perp(1)a}(z, \mathcal{B}_T^2; \mu^2, \hat{\zeta}, \rho)}{\sum_a e_a^2 H_{UU,T}(Q^2, \mu^2, \rho) \tilde{f}_1^{(0)a}(x, z^2 \mathcal{B}_T^2; \mu^2, \zeta, \rho) \tilde{D}_1^{(0)a}(z, \mathcal{B}_T^2; \mu^2, \hat{\zeta}, \rho)}, \end{aligned} \quad (\text{F.4})$$

Boer – Mulders

$$\begin{aligned}
\frac{2 J_2(|\mathbf{P}_{h\perp}| \mathcal{B}_T)}{z^2 M M_h \mathcal{B}_T^2} \cos(2\phi_h) \\
A_{UU}(\mathcal{B}_T) &= 2 \frac{\frac{\alpha^2}{y Q^2} \frac{y^2}{(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x_B}\right) \varepsilon}{\frac{\alpha^2}{y Q^2} \frac{y^2}{(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x_B}\right)} \\
&\times \frac{\sum_a e_a^2 H_{UU}^{\cos(2\phi_h)}(Q^2, \mu^2, \rho) \tilde{h}_1^{\perp(1)a}(x, z^2 \mathcal{B}_T^2; \mu^2, \zeta, \rho) \tilde{H}_1^{\perp(1)a}(z, \mathcal{B}_T^2; \mu^2, \hat{\zeta}, \rho)}{\sum_a e_a^2 H_{UU,T}(Q^2, \mu^2, \rho) \tilde{f}_1^{(0)a}(x, z^2 \mathcal{B}_T^2; \mu^2, \zeta, \rho) \tilde{D}_1^{(0)a}(z, \mathcal{B}_T^2; \mu^2, \hat{\zeta}, \rho)}, \quad (\text{F.5})
\end{aligned}$$

Kotzinian – Mulders

$$\begin{aligned}
\frac{2 J_2(|\mathbf{P}_{h\perp}| \mathcal{B}_T)}{z^2 M M_h \mathcal{B}_T^2} \sin(2\phi_h) \\
A_{UL}(\mathcal{B}_T) &= 2 \frac{\frac{\alpha^2}{y Q^2} \frac{y^2}{(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x_B}\right) \varepsilon}{\frac{\alpha^2}{y Q^2} \frac{y^2}{(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x_B}\right)} \\
&\times \frac{\sum_a e_a^2 H_{UL}^{\sin(2\phi_h)}(Q^2, \mu^2, \rho) \tilde{h}_{1L}^{\perp(1)a}(x, z^2 \mathcal{B}_T^2; \mu^2, \zeta, \rho) \tilde{H}_1^{\perp(1)a}(z, \mathcal{B}_T^2; \mu^2, \hat{\zeta}, \rho)}{\sum_a e_a^2 H_{UU,T}(Q^2, \mu^2, \rho) \tilde{f}_1^{(0)a}(x, z^2 \mathcal{B}_T^2; \mu^2, \zeta, \rho) \tilde{D}_1^{(0)a}(z, \mathcal{B}_T^2; \mu^2, \hat{\zeta}, \rho)}, \quad (\text{F.6})
\end{aligned}$$

Pretzelosity

$$\begin{aligned}
\frac{8 J_3(|\mathbf{P}_{h\perp}| \mathcal{B}_T)}{z^3 M^2 M_h \mathcal{B}_T^3} \sin(3\phi_h - \phi_s) \\
A_{UT}(\mathcal{B}_T) &= 2 \frac{\frac{\alpha^2}{y Q^2} \frac{y^2}{(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x_B}\right) \varepsilon}{\frac{\alpha^2}{y Q^2} \frac{y^2}{(1-\varepsilon)} \left(1 + \frac{\gamma^2}{2x_B}\right)} \\
&\times \frac{\sum_a e_a^2 H_{UT}^{\sin(3\phi_h - \phi_s)}(Q^2, \mu^2, \rho) \tilde{h}_{1T}^{\perp(2)a}(x, z^2 \mathcal{B}_T^2; \mu^2, \zeta, \rho) \tilde{H}_1^{\perp(1)a}(z, \mathcal{B}_T^2; \mu^2, \hat{\zeta}, \rho)}{\sum_a e_a^2 H_{UU,T}(Q^2, \mu^2, \rho) \tilde{f}_1^{(0)a}(x, z^2 \mathcal{B}_T^2; \mu^2, \zeta, \rho) \tilde{D}_1^{(0)a}(z, \mathcal{B}_T^2; \mu^2, \hat{\zeta}, \rho)}. \quad (\text{F.7})
\end{aligned}$$

## G Suppression of high transverse momentum

### G.1 Suppression of the tail in Bessel-weighted integrals

In this section, we are interested in the contribution from the high-momentum region to Bessel-weighted integrals. We begin by deriving convergence criteria and upper bounds for an integral of the form

$$I_{n,A}(\xi, \Lambda_\omega) \equiv \int_{\Lambda_\omega}^{\infty} d\omega \omega J_n(\omega \xi) A(\omega) = \xi^{-2} \int_{\Lambda_\omega \xi}^{\infty} d\nu \nu J_n(\nu) A(\nu/\xi). \quad (\text{G.1})$$

In the subsections to follow,  $\omega$  will assume the role of a momentum,  $|\mathbf{p}_T|$ ,  $|\mathbf{K}_T|$  or  $|\mathbf{P}_{h\perp}|$ , while  $\xi$  will represent  $|\mathbf{b}_T|$  or  $\mathcal{B}_T$ . In the equation above,  $A$  is a placeholder for a given function of  $\omega$ . We restrict our discussion of the integral  $I_{n,A}(\xi, \Lambda_\omega)$  to the region  $\Lambda_\omega \xi \gg 1$ , where the Bessel function in the integrand can be approximated by

$$J_n(\nu) \approx \sqrt{\frac{2}{\pi \nu}} \sin\left(\nu + \frac{\pi}{4} - \frac{n\pi}{2}\right) \quad \text{for } \nu \gg 1 \quad (\text{G.2})$$

First, let us consider a function  $A$  that fulfills the condition

$$|A(\omega)| \leq c \omega^{-\alpha} \quad \text{for any } \omega \geq \Lambda_\omega \quad (\text{G.3})$$

where  $c > 0$  and  $\alpha$  are two real valued constants. Using the envelope of the Bessel function  $|J_n(\nu)| \lesssim \sqrt{2/\pi\nu}$ , we obtain

$$\begin{aligned} |I_{n,A}(\xi, \Lambda_\omega)| &\leq \xi^{-2} \int_{\Lambda_\omega \xi}^{\infty} d\nu \left| \nu J_n(\nu) A(\nu/\xi) \right| \\ &\lesssim \xi^{-2} \int_{\Lambda_\omega \xi}^{\infty} d\nu \sqrt{\frac{2\nu}{\pi}} c \frac{\xi^\alpha}{\nu^\alpha} = \frac{1}{\alpha - \frac{3}{2}} \sqrt{\frac{2\Lambda_\omega^3}{\pi\xi}} c \Lambda_\omega^{-\alpha} \sim \xi^{-1/2}, \end{aligned} \quad (\text{G.4})$$

where the last equal sign only holds if the integral is convergent, i.e., if  $\alpha > 3/2$ .

Next, consider a function  $B$  for which  $\sqrt{\omega}B(\omega)$  is monotonously falling in the region  $\omega \geq \Lambda_\omega$  and converging to zero for  $\omega \rightarrow \infty$ . In this case, we can make use of the oscillatory behavior of the Bessel function to show convergence of the integral  $I_{n,B}(\xi, \Lambda_\omega)$ . Let  $N$  denote the smallest possible integer such that  $\Lambda_\omega \xi + \pi/4 - n\pi/2 \leq N\pi$ . We decompose the integration according to

$$\begin{aligned} I_{n,B}(\xi, \Lambda_\omega) &= \underbrace{\xi^{-2} \int_{\Lambda_\omega \xi}^{N\pi - \pi/4 + n\pi/2} d\nu \nu J_n(\nu) B(\nu/\xi)}_{T_1} \\ &+ \underbrace{\xi^{-2} \sum_{j=0}^{\infty} \int_{(N+j)\pi - \pi/4 + n\pi/2}^{(N+j+1)\pi - \pi/4 + n\pi/2} d\nu \nu J_n(\nu) B(\nu/\xi)}_{T_2}. \end{aligned} \quad (\text{G.5})$$

Applying the mean value theorem, we can find  $\bar{\nu} \in [\Lambda_\omega \xi, N\pi - \pi/4 + 2\pi/2]$  such that

$$\begin{aligned} T_1 &= \xi^{-2} \sqrt{\bar{\nu}} B(\bar{\nu}/\xi) \int_{\Lambda_\omega \xi}^{N\pi - \pi/4 + n\pi/2} d\nu \sqrt{\bar{\nu}} J_n(\nu) \\ &\approx \xi^{-2} \sqrt{\bar{\nu}} B(\bar{\nu}/\xi) \sqrt{\frac{2}{\pi}} \left( \underbrace{\cos(\Lambda_\omega \xi + \pi/4 - n\pi/4)}_{\in [-1, 1]} - \underbrace{\cos(N\pi)}_{(-1)^N} \right) \end{aligned} \quad (\text{G.6})$$

from which we derive bounds for the first term:

$$0 \leq (-1)^{N+1} T_1 \leq 2\sqrt{\frac{2}{\pi}} \xi^{-2} \sqrt{\bar{\nu}} B(\bar{\nu}/\xi) \leq 2\sqrt{\frac{2}{\pi}} \xi^{-2} \sqrt{\Lambda_\omega \xi + 2\pi} B(\Lambda_\omega). \quad (\text{G.7})$$

Using the mean value theorem again to determine the points  $\bar{\nu}_j$ , the second term becomes

$$\begin{aligned} T_2 &\approx \xi^{-2} \sum_{j=0}^{\infty} \sqrt{\bar{\nu}_j} B(\bar{\nu}_j/\xi) \int_{(N+j)\pi - \pi/4 + n\pi/2}^{(N+j+1)\pi - \pi/4 + n\pi/2} d\nu \sqrt{\frac{2}{\pi}} \sin(\nu + \pi/4 - n\pi/2) \\ &= 2\sqrt{\frac{2}{\pi}} \xi^{-2} \sum_{j=0}^{\infty} \sqrt{\bar{\nu}_j} B(\bar{\nu}_j/\xi) (-1)^{N+j}. \end{aligned} \quad (\text{G.8})$$

The expression on the right is an alternating series that fulfills the Leibnitz-test for convergence and is bounded by the size of its first term,

$$0 \leq (-1)^N T_2 \leq 2\sqrt{\frac{2}{\pi}} \xi^{-2} \sqrt{\bar{\nu}_0} B(\bar{\nu}_0/\xi) \leq 2\sqrt{\frac{2}{\pi}} \xi^{-2} \sqrt{\Lambda_\omega \xi + 2\pi} B(\Lambda_\omega). \quad (\text{G.9})$$

Since  $T_1$  and  $T_2$  have opposite signs, we arrive at a combined upper bound

$$\left| I_{n,B}(\xi, \Lambda_\omega) \right| \leq 2\sqrt{\frac{2}{\pi}}\xi^{-2}\sqrt{\Lambda_\omega\xi + 2\pi} B(\Lambda_\omega) \approx 2\sqrt{\frac{2\Lambda_\omega}{\pi\xi^3}} B(\Lambda_\omega) \sim \xi^{-3/2}. \quad (\text{G.10})$$

In summary, we find that the integral eq. (G.1) converges for any function  $A(\omega)$  that decays faster than  $\omega^{-3/2}$  which in turn determines an upper bound of the integral of order  $\xi^{-1/2}$ , eq. (G.4). The requirement for convergence can be relaxed to functions decaying faster than  $\omega^{-1/2}$  if monotony of  $\sqrt{\omega}$  times the function is ensured. In this case, eq. (G.10) gives an estimate of an upper bound that decays with  $\xi^{-3/2}$ . We remind the reader that these bounds are only valid for  $\xi \gg \Lambda_\omega^{-1}$ , but for all  $n$ .

### G.2 Fourier-transformed TMD PDFs, TMD FFs and their derivatives

Using the mathematical results from the previous sub-section, we investigate which of the (derivatives of) Fourier-transformed TMD PDFs  $\tilde{f}^{(n)}(x, \mathbf{b}_T^2)$  and TMD FFs  $\tilde{D}^{(n)}(x, \mathbf{b}_T^2)$  are well-defined by the right hand sides of eq. (2.19). Their behavior in the high transverse momentum region has been studied in detail in ref. [37]. They find power-suppressed tails of the form

$$f(x, |\mathbf{p}_T^2|) \sim \frac{1}{|\mathbf{p}_T|^m} \times \text{“logarithmic modifications”}, \quad (\text{G.11})$$

for integer powers  $m$ . Analogous expressions hold for the TMD FFs. Comparing the right hand side of eq. (2.19) with the criterion for functions of type  $B$ , we find that convergence is maintained if  $n < m - 1/2$ . The logarithmic modifications do not play a significant role since logarithms grow more slowly than any polynomial.

The analysis of ref. [37] reveals that (up to logarithmic modifications)  $f_1, g_{1L}, h_1, f^\perp, g_L^\perp, h_T, h_T^\perp, f_T, g_T, h_L, h, e_L, e, f_L, g^\perp, e_T, e_T^\perp, D_1, D^\perp, G^\perp, H, E \sim 1/\mathbf{p}_T^2$ . For these functions, the corresponding zero-derivative and single-derivative Fourier-transforms  $\tilde{f}^{(0)}(x, \mathbf{b}_T^2)$ ,  $\tilde{f}^{(1)}(x, \mathbf{b}_T^2)$ ,  $\tilde{D}^{(0)}(z, \mathbf{b}_T^2)$  and  $\tilde{D}^{(1)}(z, \mathbf{b}_T^2)$  exist. A second group of distributions exhibits the high-momentum behavior  $f_{1T}^\perp, g_{1T}, h_{1L}^\perp, h_1^\perp, f_T^\perp, g_T^\perp, h_{1T}^\perp, H_1^\perp \sim 1/\mathbf{p}_T^4$ . For these latter functions, the existence of  $n$ -derivative Fourier-transforms  $\tilde{f}^{(n)}(x, \mathbf{b}_T^2)$  and  $\tilde{D}^{(n)}(z, \mathbf{b}_T^2)$  is ensured up to  $n = 3$ . Again, we point out that these results are only valid for  $|\mathbf{b}_T| > 0$ , while the limiting case  $|\mathbf{b}_T| = 0$  leads to divergent integrals [37].

### G.3 Systematic errors from the region at large $\mathbf{P}_{h\perp}$

TMD frameworks have been designed to give a good description of the cross section at low transverse momentum, i.e., for  $|\mathbf{P}_{h\perp}|/z \ll Q$ . However, in weighted asymmetries we integrate over the whole range of  $|\mathbf{P}_{h\perp}|$ . The contributions from high  $|\mathbf{P}_{h\perp}|$  thus lead to theoretical errors in the results if one does not have a description of the cross section that is valid there, even when one restricts to the region  $z|\mathbf{b}_T| \gg 1/Q$ . The  $Y$  term can in principle be included to eliminate those errors, but its Fourier transform is expected to be power suppressed in the region  $z|\mathbf{b}_T| \gg 1/Q$ , because it was shown to be power suppressed at small  $|\mathbf{P}_{h\perp}|$  [13, 36]. Dropping the  $Y$  term means that we approximate the full result by the large  $|\mathbf{P}_{h\perp}|$ -tail of the TMD expression. This in general may be a bad approximation, but the question is whether it will affect the result much for  $z|\mathbf{b}_T| \gg 1/Q$ . In addition,

extending the integrals to arbitrarily large transverse momenta ignores the fact that the physical cross section should vanish above a certain maximum transverse momentum value  $|\mathbf{P}_{h\perp}|_{\max}$  (see also refs. [12, 36]). In this appendix we are going to estimate the effect of these various simplifications.

The  $Y$  term will be significant only in a finite region of  $|\mathbf{P}_{h\perp}|$ : between a scale  $\Lambda_{\text{TMD}}$  and  $|\mathbf{P}_{h\perp}|_{\max}$ . Note that both these scales will depend on  $Q$ . We can bound the error from neglecting the  $Y$  term in terms of its maximal value. As long as  $|\mathbf{b}_T| \gg \Lambda_{\text{TMD}}^{-1} > |\mathbf{P}_{h\perp}|_{\max}^{-1}$ , we can approximate the Bessel-function as in eq. (G.2) to obtain,

$$\begin{aligned} \tilde{Y}_{XY,Z}^{\sin/\cos(N\phi_h+\dots)}(Q^2, \mathbf{b}_T^2) &\equiv \int d|\mathbf{P}_{h\perp}| |\mathbf{P}_{h\perp}| 2\pi J_N(|\mathbf{b}_T||\mathbf{P}_{h\perp}|) Y_{XY,Z}^{\sin/\cos(N\phi_h+\dots)}(Q^2, \mathbf{P}_{h\perp}^2) \\ &\approx \int_{\Lambda_{\text{TMD}}}^{|\mathbf{P}_{h\perp}|_{\max}} d|\mathbf{P}_{h\perp}| |\mathbf{P}_{h\perp}| 2\pi J_N(|\mathbf{b}_T||\mathbf{P}_{h\perp}|) Y_{XY,Z}^{\sin/\cos(N\phi_h+\dots)}(Q^2, \mathbf{P}_{h\perp}^2) \\ &\lesssim (|\mathbf{P}_{h\perp}|_{\max} - \Lambda_{\text{TMD}}) 2\sqrt{\frac{2\pi}{|\mathbf{b}_T|\Lambda_{\text{TMD}}}} \left| Y_{XY,Z}^{\sin/\cos(N\phi_h+\dots)} \right|_{\max}. \end{aligned} \quad (\text{G.12})$$


Here  $|Y_{XY,Z}^{\sin/\cos(N\phi_h+\dots)}|_{\max}$  is the maximum absolute value of  $Y$  in the range between  $\Lambda_{\text{TMD}}$  and  $|\mathbf{P}_{h\perp}|_{\max}$ . It can be estimated from the (perturbatively calculable)  $Y$ -term. Thus, eq. (G.12) shows that the theoretical error from neglecting the  $Y$  term is (at least) suppressed as  $|\mathbf{b}_T|^{-1/2}$ . An explicit treatment of the  $Y$ -term in eq. (3.1) could eliminate this theoretical error to a given order in  $\alpha_s$  in the Fourier transformed TMD PDFs and TMD FFs extracted using Bessel-weighting. We will not do this here.

The second error coming from extending the TMD expression beyond  $|\mathbf{P}_{h\perp}|_{\max}$  is more suppressed and therefore less of a concern. Following a similar procedure as before we can estimate it to be suppressed as  $|\mathbf{b}_T|^{-3/2}$ . Let  $[F_{XY,Z}^{\sin/\cos(N\phi_h+\dots)}]_{\text{TMD}}$  denote the structure functions as determined purely within the TMD framework, i.e., from convolutions of TMD PDFs, TMD FFs and a potential soft factor. The contribution to its Fourier transform coming from the large  $|\mathbf{P}_{h\perp}|$  region can be bounded using that the TMD expression (times  $|\mathbf{P}_{h\perp}|^{1/2}$ ) is a monotonically decreasing function of  $|\mathbf{P}_{h\perp}|$ . Thus, applying eq. (G.10),

$$\begin{aligned} &\int_{|\mathbf{P}_{h\perp}|_{\max}}^{\infty} d|\mathbf{P}_{h\perp}| |\mathbf{P}_{h\perp}| 2\pi J_N(|\mathbf{b}_T||\mathbf{P}_{h\perp}|) [F_{XY,Z}^{\sin/\cos(N\phi_h+\dots)}]_{\text{TMD}}(Q^2, \mathbf{P}_{h\perp}^2) \\ &\lesssim 4\sqrt{\frac{2\pi|\mathbf{P}_{h\perp}|_{\max}}{|\mathbf{b}_T|^3}} \left| [F_{XY,Z}^{\sin/\cos(N\phi_h+\dots)}]_{\text{TMD}}(Q^2, |\mathbf{P}_{h\perp}|_{\max}^2) \right|, \end{aligned} \quad (\text{G.13})$$

where the upper bound applies as long as  $|\mathbf{b}_T| \gg |\mathbf{P}_{h\perp}|_{\max}^{-1}$ . This second error is therefore far less important than neglecting the  $Y$  term. The reason this same behavior could not be obtained for the  $Y$  term is that it is not expected to be a monotonically falling function of  $|\mathbf{P}_{h\perp}|$ .

Finally, let us consider what error would be introduced if all  $|\mathbf{P}_{h\perp}|$  integrations of the experimental data were to be cut off at  $\Lambda_{\text{TMD}}$ . In this case, we would be able to use eq. (G.13) as an error estimate, except that  $|\mathbf{P}_{h\perp}|_{\max}^{-1}$  would need to be replaced by  $\Lambda_{\text{TMD}}$ . Again the error estimate would be valid provided  $|\mathbf{b}_T| \gg \Lambda_{\text{TMD}}^{-1}$  and provided the structure function times  $|\mathbf{P}_{h\perp}|^{1/2}$  is monotonically falling, i.e., in its tail region, beyond  $\Lambda_{\text{TMD}}$ . This



**Figure 2.** Schematic illustration of important scales for Bessel-weighted asymmetries before and after the Fourier-transform.

simple cutoff method is expected to be useful when  $Q^2$  is very large, such that  $\Lambda_{\text{TMD}}$  can be chosen large with confidence.

In figure 2 we illustrate how the contributions from the TMD and the  $Y$  dominated regions contribute to the Fourier transform. The contributions from the region  $|P_{h\perp}| > \Lambda_{\text{TMD}}$  are only suppressed in the region of large  $B_T > 1/\Lambda_{\text{TMD}}$ . Therefore, an analysis without  $Y$  term at too low values of  $B_T$  has to be considered with caution. However, also the region of large  $B_T$  has to be treated with care in case of Bessel weighting, as one starts to probe the oscillations of the Bessel function. This is relevant whenever  $1/B_T$  becomes smaller than the experimental resolution in transverse momentum. A finite transverse momentum resolution  $|P_{h\perp}|_{\text{res}}$  can, for example, be a result of binning of the experimental data, as indicated in the figure.

**Open Access.** This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

## References

- [1] J.P. Ralston and D.E. Soper, *Production of Dimuons from High-Energy Polarized Proton Proton Collisions*, *Nucl. Phys. B* **152** (1979) 109 [SPIRES].
- [2] D.W. Sivers, *Single Spin Production Asymmetries from the Hard Scattering of Point-Like Constituents*, *Phys. Rev. D* **41** (1990) 83 [SPIRES].
- [3] J.C. Collins, *Fragmentation of transversely polarized quarks probed in transverse momentum distributions*, *Nucl. Phys. B* **396** (1993) 161 [hep-ph/9208213] [SPIRES].
- [4] A. Kotzinian, *New quark distributions and semiinclusive electroproduction on the polarized nucleons*, *Nucl. Phys. B* **441** (1995) 234 [hep-ph/9412283] [SPIRES].

- [5] P.J. Mulders and R.D. Tangerman, *The complete tree-level result up to order  $1/Q$  for polarized deep-inelastic leptonproduction*, *Nucl. Phys. B* **461** (1996) 197 [[hep-ph/9510301](#)] [[SPIRES](#)].
- [6] A.M. Kotzinian and P.J. Mulders, *Probing transverse quark polarization via azimuthal asymmetries in leptonproduction*, *Phys. Lett. B* **406** (1997) 373 [[hep-ph/9701330](#)] [[SPIRES](#)].
- [7] D. Boer and P.J. Mulders, *Time-reversal odd distribution functions in leptonproduction*, *Phys. Rev. D* **57** (1998) 5780 [[hep-ph/9711485](#)] [[SPIRES](#)].
- [8] A. Bacchetta et al., *Semi-inclusive deep inelastic scattering at small transverse momentum*, *JHEP* **02** (2007) 093 [[hep-ph/0611265](#)] [[SPIRES](#)].
- [9] G. Parisi and R. Petronzio, *Small Transverse Momentum Distributions in Hard Processes*, *Nucl. Phys. B* **154** (1979) 427 [[SPIRES](#)].
- [10] H.F. Jones and J. Wyndham, *On the equivalence between the impact parameter and transverse momentum space formalism for the Drell-Yan process*, *J. Phys. A* **14** (1981) 1457 [[SPIRES](#)].
- [11] J.C. Collins and D.E. Soper, *Parton Distribution and Decay Functions*, *Nucl. Phys. B* **194** (1982) 445 [[SPIRES](#)].
- [12] S.D. Ellis, N. Fleishon and W.J. Stirling, *Logarithmic approximations, quark form-factors and Quantum Chromodynamics*, *Phys. Rev. D* **24** (1981) 1386 [[SPIRES](#)].
- [13] J.C. Collins, D.E. Soper and G.F. Sterman, *Transverse Momentum Distribution in Drell-Yan Pair and W and Z Boson Production*, *Nucl. Phys. B* **250** (1985) 199 [[SPIRES](#)].
- [14] X.-d. Ji, J.-P. Ma and F. Yuan, *QCD factorization for spin-dependent cross sections in DIS and Drell-Yan processes at low transverse momentum*, *Phys. Lett. B* **597** (2004) 299 [[hep-ph/0405085](#)] [[SPIRES](#)].
- [15] J.C. Collins, *Foundations of Perturbative QCD*, Cambridge University Press, Cambridge U.K. (2011).
- [16] J.C. Collins and F. Hautmann, *Infrared divergences and non-lightlike eikonal lines in Sudakov processes*, *Phys. Lett. B* **472** (2000) 129 [[hep-ph/9908467](#)] [[SPIRES](#)].
- [17] J.C. Collins and A. Metz, *Universality of soft and collinear factors in hard-scattering factorization*, *Phys. Rev. Lett.* **93** (2004) 252001 [[hep-ph/0408249](#)] [[SPIRES](#)].
- [18] X.-d. Ji, J.-p. Ma and F. Yuan, *QCD factorization for semi-inclusive deep-inelastic scattering at low transverse momentum*, *Phys. Rev. D* **71** (2005) 034005 [[hep-ph/0404183](#)] [[SPIRES](#)].
- [19] S.M. Aybat and T.C. Rogers, *TMD Parton Distribution and Fragmentation Functions with QCD Evolution*, *Phys. Rev. D* **83** (2011) 114042 [[arXiv:1101.5057](#)] [[SPIRES](#)].
- [20] M. Gourdin, *Semiinclusive reactions induced by leptons*, *Nucl. Phys. B* **49** (1972) 501 [[SPIRES](#)].
- [21] M. Diehl and S. Sapeta, *On the analysis of lepton scattering on longitudinally or transversely polarized protons*, *Eur. Phys. J. C* **41** (2005) 515 [[hep-ph/0503023](#)] [[SPIRES](#)].
- [22] A. Bacchetta, U. D'Alesio, M. Diehl and C.A. Miller, *Single-spin asymmetries: The Trento conventions*, *Phys. Rev. D* **70** (2004) 117504 [[hep-ph/0410050](#)] [[SPIRES](#)].
- [23] D. Boer, P.J. Mulders and F. Pijlman, *Universality of T-odd effects in single spin and azimuthal asymmetries*, *Nucl. Phys. B* **667** (2003) 201 [[hep-ph/0303034](#)] [[SPIRES](#)].

- [24] D.E. Soper, *Partons and Their Transverse Momenta in Quantum Chromodynamics*, *Phys. Rev. Lett.* **43** (1979) 1847 [[SPIRES](#)].
- [25] A.V. Belitsky, X. Ji and F. Yuan, *Final state interactions and gauge invariant parton distributions*, *Nucl. Phys.* **B 656** (2003) 165 [[hep-ph/0208038](#)] [[SPIRES](#)].
- [26] C.J. Bomhof, P.J. Mulders and F. Pijlman, *Gauge link structure in quark quark correlators in hard processes*, *Phys. Lett.* **B 596** (2004) 277 [[hep-ph/0406099](#)] [[SPIRES](#)].
- [27] C.J. Bomhof, P.J. Mulders and F. Pijlman, *The construction of gauge-links in arbitrary hard processes*, *Eur. Phys. J.* **C 47** (2006) 147 [[hep-ph/0601171](#)] [[SPIRES](#)].
- [28] A. Idilbi, X.-d. Ji, J.-P. Ma and F. Yuan, *Collins-Soper equation for the energy evolution of transverse-momentum and spin dependent parton distributions*, *Phys. Rev.* **D 70** (2004) 074021 [[hep-ph/0406302](#)] [[SPIRES](#)].
- [29] K. Goeke, A. Metz and M. Schlegel, *Parameterization of the quark-quark correlator of a spin-1/2 hadron*, *Phys. Lett.* **B 618** (2005) 90 [[hep-ph/0504130](#)] [[SPIRES](#)].
- [30] J.C. Collins and D.E. Soper, *Back-To-Back Jets in QCD*, *Nucl. Phys.* **B 193** (1981) 381 [[SPIRES](#)].
- [31] J.C. Collins, *What exactly is a parton density?*, *Acta Phys. Polon.* **B 34** (2003) 3103 [[hep-ph/0304122](#)] [[SPIRES](#)].
- [32] F. Hautmann, *Endpoint singularities in unintegrated parton distributions*, *Phys. Lett.* **B 655** (2007) 26 [[hep-ph/0702196](#)] [[SPIRES](#)].
- [33] J. Chay, *Transverse-momentum-dependent parton distribution function in soft-collinear effective theory*, [arXiv:0711.4295](#) [[SPIRES](#)].
- [34] I.O. Cherednikov and N.G. Stefanis, *Wilson lines and transverse-momentum dependent parton distribution functions: A renormalization-group analysis*, *Nucl. Phys.* **B 802** (2008) 146 [[arXiv:0802.2821](#)] [[SPIRES](#)].
- [35] I.O. Cherednikov, A.I. Karanikas and N.G. Stefanis, *Wilson lines in transverse-momentum dependent parton distribution functions with spin degrees of freedom*, *Nucl. Phys.* **B 840** (2010) 379 [[arXiv:1004.3697](#)] [[SPIRES](#)].
- [36] J.C. Collins and D.E. Soper, *Back-To-Back Jets: Fourier Transform from B to K-Transverse*, *Nucl. Phys.* **B 197** (1982) 446 [[SPIRES](#)].
- [37] A. Bacchetta, D. Boer, M. Diehl and P.J. Mulders, *Matches and mismatches in the descriptions of semi-inclusive processes at low and high transverse momentum*, *JHEP* **08** (2008) 023 [[arXiv:0803.0227](#)] [[SPIRES](#)].
- [38] P. Hägler, B.U. Musch, J.W. Negele and A. Schäfer, *Intrinsic quark transverse momentum in the nucleon from lattice QCD*, *Europhys. Lett.* **88** (2009) 61001 [[arXiv:0908.1283](#)] [[SPIRES](#)].
- [39] B.U. Musch, P. Hägler, J.W. Negele and A. Schäfer, *Exploring quark transverse momentum distributions with lattice QCD*, *Phys. Rev.* **D 83** (2011) 094507 [[arXiv:1011.1213](#)] [[SPIRES](#)].
- [40] J.C. Collins, *Leading-twist Single-transverse-spin asymmetries: Drell- Yan and Deep-Inelastic Scattering*, *Phys. Lett.* **B 536** (2002) 43 [[hep-ph/0204004](#)] [[SPIRES](#)].
- [41] Z.-B. Kang, B.-W. Xiao and F. Yuan, *QCD Resummation for Single Spin Asymmetries*, [arXiv:1106.0266](#) [[SPIRES](#)].



- [42] J.-w. Qiu and G.F. Sterman, *Single transverse spin asymmetries*, *Phys. Rev. Lett.* **67** (1991) 2264 [SPIRES].
- [43] Z.-B. Kang and J.-W. Qiu, *Evolution of twist-3 multi-parton correlation functions relevant to single transverse-spin asymmetry*, *Phys. Rev. D* **79** (2009) 016003 [arXiv:0811.3101] [SPIRES].
- [44] J. Zhou, F. Yuan and Z.-T. Liang, *QCD Evolution of the Transverse Momentum Dependent Correlations*, *Phys. Rev. D* **79** (2009) 114022 [arXiv:0812.4484] [SPIRES].
- [45] W. Vogelsang and F. Yuan, *Next-to-leading Order Calculation of the Single Transverse Spin Asymmetry in the Drell-Yan Process*, *Phys. Rev. D* **79** (2009) 094010 [arXiv:0904.0410] [SPIRES].
- [46] V.M. Braun, A.N. Manashov and B. Pirnay, *On the scale dependence of twist-three contributions to single spin asymmetries*, *Phys. Rev. D* **80** (2009) 114002 [arXiv:0909.3410] [SPIRES].
- [47] P.G. Ratcliffe and O. Teryaev, *Non-Universality and Evolution of the Sivers Function*, *Mod. Phys. Lett. A* **24** (2009) 2984 [arXiv:0910.5348] [SPIRES].
- [48] J. Zhou, F. Yuan and Z.-T. Liang, *Hyperon Polarization in Unpolarized Scattering Processes*, *Phys. Rev. D* **78** (2008) 114008 [arXiv:0808.3629] [SPIRES].
- [49] J. Zhou, F. Yuan and Z.-T. Liang, *Transverse momentum dependent quark distributions and polarized Drell-Yan processes*, *Phys. Rev. D* **81** (2010) 054008 [arXiv:0909.2238] [SPIRES].
- [50] F. Yuan and J. Zhou, *Collins Fragmentation and the Single Transverse Spin Asymmetry*, *Phys. Rev. Lett.* **103** (2009) 052001 [arXiv:0903.4680] [SPIRES].
- [51] Z.-B. Kang, *QCD evolution of naive-time-reversal-odd fragmentation functions*, *Phys. Rev. D* **83** (2011) 036006 [arXiv:1012.3419] [SPIRES].
- [52] M. Burkardt, *Chromodynamic lensing and transverse single spin asymmetries*, *Nucl. Phys. A* **735** (2004) 185 [hep-ph/0302144] [SPIRES].
- [53] P. Hägler, *Hadron structure from lattice quantum chromodynamics*, *Phys. Rept.* **490** (2010) 49 [arXiv:0912.5483] [SPIRES].
- [54] V.S. Dotsenko and S.N. Vergeles, *Renormalizability of Phase Factors in the Nonabelian Gauge Theory*, *Nucl. Phys. B* **169** (1980) 527 [SPIRES].
- [55] N.S. Craigie and H. Dorn, *On the renormalization and short distance properties of hadronic operators in QCD*, *Nucl. Phys. B* **185** (1981) 204 [SPIRES].
- [56] I.Y. Arefeva, *Quantum contour field equations*, *Phys. Lett. B* **93** (1980) 347 [SPIRES].
- [57] S. Aoyama, *The renormalization of the string operator in QCD*, *Nucl. Phys. B* **194** (1982) 513 [SPIRES].
- [58] N.G. Stefanis, *Gauge invariant quark two-point green's function through connector insertion to  $\mathcal{O}(\alpha_s)$* , *Nuovo Cim. A* **83** (1984) 205 [SPIRES].
- [59] H. Dorn, *Renormalization of path ordered phase factors and related hadron operators in gauge field theories*, *Fortsch. Phys.* **34** (1986) 11 [SPIRES].