



PUBLISHED FOR SISSA BY SPRINGER

RECEIVED: May 25, 2015

ACCEPTED: June 10, 2015

PUBLISHED: June 26, 2015

Diphoton production at hadron colliders: transverse-momentum resummation at next-to-next-to-leading logarithmic accuracy

Leandro Cieri,^a Francesco Coradeschi^b and Daniel de Florian^c

^a*Dipartimento di Fisica, Università di Roma “La Sapienza” and
INFN — Sezione di Roma, I-00185 Rome, Italy*

^b*INFN — Sezione di Firenze and Dipartimento di Fisica,
Università di Firenze, I-50019 Sesto Fiorentino, Florence, Italy*

^c*Departamento de Física, FCEYN, Universidad de Buenos Aires,
(1428) Pabellón 1 Ciudad Universitaria, Capital Federal, Argentina*

E-mail: leandrocieri@gmail.com, fr.coradeschi@gmail.com,
deflo@df.uba.ar

ABSTRACT: We consider the transverse-momentum (q_T) distribution of a diphoton pair produced in hadron collisions. At small values of q_T , we resum the logarithmically-enhanced perturbative QCD contributions up to next-to-next-to-leading logarithmic accuracy. At intermediate and large values of q_T , we consistently combine resummation with the known next-to-leading order perturbative result. All perturbative terms up to order α_s^2 are included in our computation which, after integration over q_T , reproduces the known next-to-next-to-leading order result for the diphoton pair production total cross section. We present a comparison with LHC data and an estimate of the perturbative accuracy of the theoretical calculation by performing the corresponding variation of scales. In general we observe that the effect of the resummation is not only to recover the predictivity of the calculation at small transverse momentum, but also to improve substantially the agreement with the experimental data.

KEYWORDS: QCD Phenomenology, Hadronic Colliders

ARXIV EPRINT: [1505.03162](https://arxiv.org/abs/1505.03162)

Contents

1	Introduction	1
2	Transverse-momentum resummation	3
3	Numerical results for diphoton production at the LHC	6
4	Summary	16

1 Introduction

Diphoton production at hadronic colliders is a very relevant process, both from the point of view of testing the Standard Model (SM) predictions [1–6] as for new physics searches.

While a general understanding of QCD processes at hadronic collisions poses a serious challenge due the complicated environment, *direct* or *prompt* photons provide an ideal probe since they constitute a theoretically and experimentally *clean* final state. From the theory side, because they do not have QCD interactions with other final state particles and, from the experimental side, because their energies and momenta can be measured with high precision in modern electromagnetic calorimeters.

Besides purely QCD-related considerations, diphoton final states have played a crucial role in the recent discovery of a new boson at the LHC [7, 8], whose properties are compatible with those of the SM Higgs. They are also important in many new physics scenarios [9, 10], in particular in the search for extra-dimensions [11] or supersymmetry [12].

In this paper we are interested in the process $pp \rightarrow \gamma\gamma X$, and in particular in the transverse-momentum (q_T) spectrum of the diphoton pair. The lowest-order process ($\mathcal{O}(\alpha_S^0)$) occurs *via* the quark annihilation subprocess $q\bar{q} \rightarrow \gamma\gamma$. The QCD corrections at the first order in the strong coupling α_S involve quark annihilation and a new partonic channel, *via* the subprocess $qg \rightarrow \gamma\gamma q$. First order corrections have been computed and implemented in several fully-differential Monte Carlo codes [13–16]. At the second order in the strong coupling α_S the gg channel starts to contribute, and the large gluon-gluon luminosity makes this channel potentially sizeable.

The amplitudes needed to evaluate the corrections at the second order in the strong coupling α_S , for diphoton production, have been presented in [17–22], and first put together in a complete and consistent $\mathcal{O}(\alpha_S^2)$ calculation in the $2\gamma\text{NNLO}$ code [23]. The next-order gluonic corrections to the *box contribution* (which are part of the N^3LO QCD corrections to diphoton production) were also computed in ref. [14] and found to have a moderate quantitative effect.

The calculation of the q_T spectrum poses an additional challenge with respect to more inclusive calculations, such as the total cross section. In the large- q_T region ($q_T \sim M_{\gamma\gamma}$), where the transverse momentum is of the order of the diphoton invariant mass $M_{\gamma\gamma}$, calculations based on the truncation of the perturbative series at a fixed order in α_S are

theoretically justified. In this region, the QCD radiative corrections are known up to the next-to-leading order (NLO), including the corresponding partonic scattering amplitudes with $X = 2$ partons (at the tree level [18, 19]) and the partonic scattering amplitudes with $X = 1$ parton (up to the one-loop level [20, 21]). We remind the reader that at least one additional parton is needed in order to have $q_T \neq 0$ for the diphoton pair. The q_T spectrum of the diphoton pair has been calculated in fully-differential Monte Carlo codes at LO [13–16] and at NLO [23–25]. Recently, first calculations for diphoton production in association with two [26–28] and three [28] jets at NLO became available.

The bulk of the diphoton events is produced in the small- q_T region ($q_T \ll M_{\gamma\gamma}$), where the convergence of the fixed-order expansion is spoiled by the presence of large logarithmic terms, $\alpha_S^n \ln^m(M_{\gamma\gamma}^2/q_T^2)$. In order to obtain reliable predictions these logarithmically-enhanced terms have to be systematically resummed to all perturbative orders [29, 30]–[43]. The resummed calculation, valid at small values of q_T , and the fixed-order one at large q_T have then to be consistently matched to obtain a pQCD prediction for the entire range of transverse momenta.

We use the transverse-momentum resummation formalism proposed in refs. [42, 44, 45] (see also [46] for processes initiated by gg annihilation). The formalism is valid for a generic process in which a high-mass system of non strongly-interacting particles is produced in hadron-hadron collisions. The method has so far been applied to the production of the Standard Model (SM) Higgs boson [44, 45, 47–49], Higgs boson production in bottom quark annihilation [50], Higgs boson production via gluon fusion in the MSSM [51], single vector bosons at NLL+LO [52] and at NNLL+NLO [53], WW [54, 55] and ZZ [56] pairs, slepton pairs [57], and DY lepton pairs in polarized collisions [58–61].

Finally, note that besides the direct photon production from the hard subprocess, photons can also arise from the fragmentation of QCD partons. The computation of fragmentation subprocesses requires (the poorly known) non-perturbative information, in the form of parton fragmentation functions of the photon (the complete single- and double-fragmentation contributions are implemented in DIPHOX [13] for diphoton production at the first order in α_S). However, the effect of the fragmentation contributions is sizeably reduced by the *photon isolation* criteria that are necessarily applied in hadron collider experiments to suppress the very large irreducible background (e.g., photons that are faked by jets or produced by hadron decays). Two such criteria are the so-called “standard” cone isolation and the “smooth” cone isolation proposed by Frixione [62]. The standard cone isolation is easily implemented in experiments, but it only suppresses a fraction of the fragmentation contribution. By contrast, the smooth cone isolation (formally) eliminates the entire fragmentation contribution. For all of the results presented in this paper we rely on the smooth isolation prescription, which, for the parameters used in the experimental analysis reproduces the standard result within a 1% accuracy [63].

The paper is organized as follows. In section 2 we briefly review the resummation formalism of refs. [42, 44, 45]. In section 3 we present numerical results and we comment on their comparison with the LHC data [6]. We also study the scale dependence of our results with the purpose of estimating the corresponding perturbative uncertainty. In section 4 we summarize our results.

2 Transverse-momentum resummation

We briefly recall the main points of the transverse-momentum resummation formalism of refs. [42, 44, 45], referring to the original papers for the full details. The formalism is general, as long as the measured final state is composed of non strongly-interacting particles (transverse-momentum resummation for strongly-interacting final states, such as heavy-quark production, has been developed in refs. [64, 65]). Here we specialize to the case of diphoton production only for ease of reading. The inclusive hard-scattering process considered is

$$h_1(p_1) + h_2(p_2) \rightarrow \gamma\gamma(M_{\gamma\gamma}, q_T, y) + X, \quad (2.1)$$

where h_1 and h_2 are the colliding hadrons with momenta p_1 and p_2 , $\gamma\gamma$ is the diphoton pair with invariant mass $M_{\gamma\gamma}$, transverse momentum q_T and rapidity y , and X is an arbitrary and undetected final state.

The corresponding fully differential cross section, in q_T , $M_{\gamma\gamma}$ and y , which we denote for simplicity (since our focus is on the q_T distribution) by $d\sigma_{\gamma\gamma}/dq_T^2$, can be written using the factorization formula as

$$\frac{d\sigma_{\gamma\gamma}}{dq_T^2} = \sum_{a,b} \int_0^1 dx_1 \int_0^1 dx_2 f_{a/h_1}(x_1, \mu_F^2) f_{b/h_2}(x_2, \mu_F^2) \frac{d\hat{\sigma}_{ab}^{\gamma\gamma}}{dq_T^2}(q_T, M_{\gamma\gamma}, y, \hat{s}; \alpha_S, \mu_R^2, \mu_F^2) \quad (2.2)$$

(up to power-suppressed corrections), where the $f_{a/h}(x, \mu_F^2)$ ($a = q, \bar{q}, g$) are the parton densities of the hadron h at the factorization scale μ_F , $\alpha_S \equiv \alpha_S(\mu_R^2)$, $d\hat{\sigma}_{ab}^{\gamma\gamma}/dq_T^2$ is the pQCD *partonic cross section*, s ($\hat{s} = x_1 x_2 s$) is the square of the hadronic (partonic) centre-of-mass energy, and μ_R is the renormalization scale.

In the region where $q_T \sim M_{\gamma\gamma}$ the QCD perturbative series is controlled by a small expansion parameter, $\alpha_S(M_{\gamma\gamma})$, and a fixed-order calculation of the partonic cross section is theoretically justified. In this region, the QCD radiative corrections are known up to next-to-leading order (NLO) [17–22].

In the small- q_T region ($q_T \ll M_{\gamma\gamma}$), the convergence of the fixed-order perturbative expansion is spoiled by the presence of powers of large logarithmic terms, $\alpha_S^n \ln^m(M_{\gamma\gamma}^2/q_T^2)$. To obtain reliable predictions these terms have to be resummed to all orders.

To perform the resummation, we start by decomposing the partonic cross section as

$$\frac{d\hat{\sigma}_{ab}^{\gamma\gamma}}{dq_T^2} = \frac{d\hat{\sigma}_{\gamma\gamma ab}^{(\text{res.})}}{dq_T^2} + \frac{d\hat{\sigma}_{\gamma\gamma ab}^{(\text{fin.})}}{dq_T^2}. \quad (2.3)$$

The first term on the right-hand side contains all the logarithmically-enhanced contributions, which have to be resummed to all orders in α_S , while the second term is free of such contributions and can thus be evaluated at fixed order in perturbation theory. Using the Fourier transformation between the conjugate variables q_T and b (b is the impact parameter), the resummed component $d\hat{\sigma}_{\gamma\gamma ab}^{(\text{res.})}$ can be expressed as

$$\frac{d\hat{\sigma}_{\gamma\gamma ab}^{(\text{res.})}}{dq_T^2}(q_T, M_{\gamma\gamma}, y, \hat{s}; \alpha_S, \mu_R^2, \mu_F^2) = \frac{M_{\gamma\gamma}^2}{\hat{s}} \int_0^\infty db \frac{b}{2} J_0(bq_T) \mathcal{W}_{ab}^{\gamma\gamma}(b, M_{\gamma\gamma}, y, \hat{s}; \alpha_S, \mu_R^2, \mu_F^2), \quad (2.4)$$

where $J_0(x)$ is the 0th-order Bessel function. The form factor $\mathcal{W}^{\gamma\gamma}$ is best expressed in terms of its *double* Mellin moments $\mathcal{W}_{N_1 N_2}^{\gamma\gamma}$, taken with respect to the variables z_1, z_2 at fixed $M_{\gamma\gamma}$, with

$$z_1 z_2 \equiv z = \frac{M_{\gamma\gamma}^2}{\hat{s}}, \quad \frac{z_1}{z_2} = e^{2y}; \quad (2.5)$$

the resummation structure of $\mathcal{W}_{N_1 N_2}^{\gamma\gamma}$ can be organized in an exponential form¹

$$\mathcal{W}_{N_1 N_2}^{\gamma\gamma}(b, M_{\gamma\gamma}, y; \alpha_S, \mu_R^2, \mu_F^2) = \mathcal{H}_{N_1 N_2}^{\gamma\gamma}(M_{\gamma\gamma}, \alpha_S; M_{\gamma\gamma}^2/\mu_R^2, M_{\gamma\gamma}^2/\mu_F^2, M_{\gamma\gamma}^2/\mu_{res}^2) \times \exp\{\mathcal{G}_{N_1 N_2}(\alpha_S, L; M_{\gamma\gamma}^2/\mu_R^2, M_{\gamma\gamma}^2/\mu_{res}^2)\} \quad , \quad (2.6)$$

where we have defined the logarithmic expansion parameter $L \equiv \ln(\mu_{res}^2 b^2/b_0^2)$, and $b_0 = 2e^{-\gamma_E}$ ($\gamma_E = 0.5772\dots$ is the Euler number). The scale μ_{res} ($\mu_{res} \sim M_{\gamma\gamma}$), which appears on the right-hand side of eq. (2.6), is the resummation scale [44]. Although $\mathcal{W}_{N_1 N_2}^{\gamma\gamma}$ (i.e., the product $\mathcal{H}_{N_1 N_2}^{\gamma\gamma} \times \exp\{\mathcal{G}_{N_1 N_2}\}$) does not depend on μ_{res} when evaluated to all perturbative orders, its explicit dependence on μ_{res} appears when $\mathcal{W}_{N_1 N_2}^{\gamma\gamma}$ is computed by truncation of the resummed expression at some level of logarithmic accuracy (see eq. (2.7) below). Variations of μ_{res} around $M_{\gamma\gamma}$ can thus be used to estimate the size of yet uncalculated higher-order logarithmic contributions.

The form factor $\exp\{\mathcal{G}_{N_1 N_2}\}$ is universal² and contains all the terms that order-by-order in α_S are logarithmically divergent as $b \rightarrow \infty$ (or, equivalently, $q_T \rightarrow 0$). The resummed logarithmic expansion of the exponent $\mathcal{G}_{N_1 N_2}$ is defined as follows:

$$\mathcal{G}_{N_1 N_2}(\alpha_S, L; M_{\gamma\gamma}^2/\mu_R^2, M_{\gamma\gamma}^2/\mu_{res}^2) = L g^{(1)}(\alpha_S L) + g_{N_1 N_2}^{(2)}(\alpha_S L; M_{\gamma\gamma}^2/\mu_R^2, M_{\gamma\gamma}^2/\mu_{res}^2) + \frac{\alpha_S}{\pi} g_{N_1 N_2}^{(3)}(\alpha_S L, M_{\gamma\gamma}^2/\mu_R^2, M_{\gamma\gamma}^2/\mu_{res}^2) + \dots \quad (2.7)$$

where the term $L g^{(1)}$ collects the leading logarithmic (LL) $\mathcal{O}(\alpha_s^{p+n} L^{n+1})$ contributions, the function $g_{N_1 N_2}^{(2)}$ includes the next-to-leading leading logarithmic (NLL) $\mathcal{O}(\alpha_s^{p+n} L^n)$ contributions [35–37], $g_{N_1 N_2}^{(3)}$ controls the NNLL $\mathcal{O}(\alpha_s^{p+n} L^{n-1})$ terms [38, 39, 66–68] and so forth; p is the number of powers of α_s in the LO (Born) process. In eq. (2.7), $\alpha_S L$ is formally of order 1, so there is an explicit $\mathcal{O}(\alpha_S)$ suppression between different logarithmic orders. The explicit form of the functions $g^{(1)}$, $g_{N_1 N_2}^{(2)}$ and $g_{N_1 N_2}^{(3)}$ can be found in ref. [44]. The process dependent function $\mathcal{H}_{N_1 N_2}^{\gamma\gamma}$ does not depend on the impact parameter b and it includes all the perturbative terms that behave as constants as $b \rightarrow \infty$. It can thus be expanded in powers of α_S :

$$\begin{aligned} & \mathcal{H}_{N_1 N_2}^{\gamma\gamma}(M_{\gamma\gamma}, \alpha_S; M_{\gamma\gamma}^2/\mu_R^2, M_{\gamma\gamma}^2/\mu_F^2, M_{\gamma\gamma}^2/\mu_{res}^2) \\ &= \sigma_{\gamma\gamma}^{(0)}(\alpha_s, M_{\gamma\gamma}) \left[1 + \frac{\alpha_S}{\pi} \mathcal{H}_{N_1 N_2}^{\gamma\gamma(1)}(M_{\gamma\gamma}^2/\mu_F^2, M_{\gamma\gamma}^2/\mu_{res}^2) \right. \\ & \quad \left. + \left(\frac{\alpha_S}{\pi} \right)^2 \mathcal{H}_{N_1 N_2}^{\gamma\gamma(2)}(M_{\gamma\gamma}^2/\mu_R^2, M_{\gamma\gamma}^2/\mu_F^2, M_{\gamma\gamma}^2/\mu_{res}^2) + \dots \right], \quad (2.8) \end{aligned}$$

¹For the sake of simplicity we consider here only the case of the diagonal terms in the flavour space of the partonic indices a, b . For a detailed discussion, we refer to refs. [44, 45].

²The form factor does not depend on the final state; all the hard-scattering processes initiated by $q\bar{q}$ ($g\bar{g}$) annihilation have the same form factor.

where $\sigma_{\gamma\gamma}^{(0)}$ is the partonic cross section at the Born level. Since the formalism applies to non strongly-interacting final states, in general the Born cross-section can only correspond to a $q\bar{q}$ or gg initial state. In the specific case of the diphoton production, both channels contribute, but at different orders in α_s : the $q\bar{q}$ subprocess initiates as a pure QED process ($\mathcal{O}(\alpha_s^0)$), while the gg one requires a fermion loop, starting at $\mathcal{O}(\alpha_s)^2$.

In the present work, we keep contributions up to an uniform order in α_s (and all orders in $\alpha_s L$), namely up to $\alpha_s^n L^{n-1}$. For the $q\bar{q}$ channel, this requires the inclusion of the \mathcal{H} coefficients of eq. (2.8) up to order 2: the first-order coefficients $\mathcal{H}_{N_1 N_2}^{\gamma\gamma(1)}$ are known since a long time [66, 67], while the second-order coefficients $\mathcal{H}_{N_1 N_2}^{\gamma\gamma(2)}$ were computed only relatively recently [23, 43]. For the gg channel, it is sufficient to include the leading \mathcal{H} contribution (that is, the Born cross-section) and the appropriate \mathcal{G} in the exponential of eq. (2.7). Since it does not require any additional numerical effort, we decided, in all the plots presented in the paper, to include all the terms up to $g_{N_1 N_2}^{(3)}$ in the exponential \mathcal{G} factor also for this channel. In this way, we technically include some terms which are of higher order in α_s with respect to those in the $q\bar{q}$ channel; however we checked that those terms result in a negligible numerical effect (at 1% accuracy), that is, the difference produced by including the higher order terms is within the error bands obtained by the scale variations, which verifies the stability of the calculation.

Within a straightforward (‘naive’) implementation of eq. (2.6), the resummation of the large logarithmic contributions would affect not only the small- q_T region, but also the region of large values of q_T . This can easily be understood by observing that the logarithmic expansion parameter L diverges also when $b \rightarrow 0$. To reduce the impact of unjustified higher-order contributions in the large- q_T region, the logarithmic variable L in eq. (2.6) is actually replaced by $\tilde{L} \equiv \ln(\mu_{res}^2 b^2 / b_0^2 + 1)$ [44, 47]. This (unitarity related) replacement has an additional and relevant consequence: after inclusion of the finite component (see eq. (2.9)), we exactly recover the fixed-order perturbative value of the total cross section upon integration of the q_T distribution over q_T (i.e., the resummed terms give a vanishing contribution upon integration over q_T).

We now turn to consider the finite component of the transverse-momentum cross section (see eq. (2.3)). Since $d\sigma_{\gamma\gamma}^{(\text{fin.})}$ does not contain large logarithmic terms in the small- q_T region, it can be evaluated by truncation of the perturbative series at a given fixed order. In practice, the finite component is computed starting from the usual fixed-order perturbative truncation of the partonic cross section and subtracting the expansion of the resummed part at the same perturbative order. Introducing the subscript f.o. to denote the perturbative truncation of the various terms, we have:

$$\left[\frac{d\hat{\sigma}_{\gamma\gamma ab}^{(\text{fin.})}}{dq_T^2} \right]_{\text{f.o.}} = \left[\frac{d\hat{\sigma}_{\gamma\gamma ab}}{dq_T^2} \right]_{\text{f.o.}} - \left[\frac{d\hat{\sigma}_{\gamma\gamma ab}^{(\text{res.})}}{dq_T^2} \right]_{\text{f.o.}}. \tag{2.9}$$

This matching procedure between resummed and finite contributions guarantees to achieve uniform theoretical accuracy over the region from small to intermediate values of transverse momenta. At large values of q_T , the resummation (and matching) procedure is eventually superseded by the customary fixed-order calculations (their theoretical accuracy in the

large- q_T region cannot be improved by resummation of the logarithmic terms that dominate in the small- q_T region).

In summary, the inclusion of the functions $g^{(1)}$, $g_{N_1 N_2}^{(2)}$, $\mathcal{H}_{N_1 N_2}^{\gamma\gamma(1)}$ in the resummed component, together with the evaluation of the finite component at LO (i.e. at $\mathcal{O}(\alpha_S)$), allows us to perform the resummation at NLL+LO accuracy. This is the theoretical accuracy used in previous studies [16, 69–71] of the diphoton q_T distribution. Including also the functions $g_{N_1 N_2}^{(3)}$ and $\mathcal{H}_{N_1 N_2}^{\gamma\gamma(2)}$, together with the finite component at NLO (i.e. at $\mathcal{O}(\alpha_S^2)$) leads to full NNLL+NLO accuracy.

Using the $\mathcal{H}_{N_1 N_2}^{\gamma\gamma(2)}$ coefficient [23, 43], we are thus able to present the complete result for the diphoton q_T -distribution up to NNLL+NLO accuracy. We point out that the NNLL+NLO (NLL+LO) result includes the *full* NNLO (NLO) perturbative contribution in the small- q_T region. In particular, the NNLO (NLO) result for the total cross section is exactly recovered upon integration over q_T of the differential cross section $d\sigma_{\gamma\gamma}/dq_T$ at NNLL+NLO (NLL+LO) accuracy.

We conclude this section with some comments on the numerical implementation of our calculation. Within our formalism, the resummation factor $\mathcal{W}_{N_1 N_2}^{\gamma\gamma}(b, M_{\gamma\gamma})$ is directly defined, at fixed $M_{\gamma\gamma}$, in the space of the conjugate variables b and N_1, N_2 . To obtain the hadronic cross section, we have to perform inverse integral transformations: the Bessel transformation in eq. (2.4) and an inverse Mellin transformation, implemented following the prescription introduced in ref. [72]. These integrals are carried out numerically. The Mellin inversion requires the numerical evaluation of some basic N -moment functions that appear in the expression of the the second-order coefficients $\mathcal{H}_{N_1 N_2}^{\gamma\gamma(2)}$ [23, 43]: this evaluation has to be performed for complex values of N_1 and N_2 ; to evaluate some of the needed special function at complex value, we use the numerical routines of refs. [73–76]. We recall [44] that the resummed form factor $\exp\{\mathcal{G}_{N_1 N_2}(\alpha_S(\mu_R^2), \tilde{L})\}$ is singular at the values of b where $\alpha_S(\mu_R^2)\tilde{L} \geq \pi/\beta_0$ (β_0 is the first-order coefficient of the QCD β function). We avoid this singularity by introducing a smooth effective cut-off at small b , which is shown to have a negligible effect in the final result.

It is known that at small values of q_T , the perturbative QCD approach has to be supplemented with non-perturbative contributions, since they become relevant as q_T decreases. A discussion on non-perturbative effects on the q_T distribution is presented in ref. [44], and related quantitative results are shown in section 3.

3 Numerical results for diphoton production at the LHC

In this section we consider diphoton production in pp collisions at LHC energies ($\sqrt{s} = 7$ TeV). We present our resummed results at NNLL+NLO accuracy, and compare them with NLL+LO predictions and with available LHC data [6]. Since the present formulation of the q_T resummation formalism, is restricted to the production of colourless systems F , it does not treat parton fragmentation subprocesses (here F includes one or two coloured partons that fragment); therefore, we concentrate on the direct production of diphotons, and we rely on the smooth cone isolation criterion proposed by Frixione [62] (see also

refs. [77, 78]) which is defined by requesting

$$\sum E_T^{had} \leq E_{T\ max} \chi(r) ,$$

$$\text{inside any } r^2 = (y - y_\gamma)^2 + (\phi - \phi_\gamma)^2 \leq R^2 , \tag{3.1}$$

with a suitable choice for the function $\chi(r)$. This function has to vanish smoothly when its argument goes to zero ($\chi(r) \rightarrow 0$, if $r \rightarrow 0$), and it has to verify $0 < \chi(r) < 1$, if $0 < r < R$. One possible choice is

$$\chi(r) = \left(\frac{1 - \cos(r)}{1 - \cos R} \right)^n , \tag{3.2}$$

where n is typically chosen as $n = 1$. This condition implies that, closer to the photon, less hadronic activity is allowed inside the cone. At $r = 0$, when the parton and the photon are exactly collinear, the energy deposited inside the cone is required to be exactly equal to zero, and the fragmentation component (which is a purely collinear phenomenon in perturbative QCD) vanishes completely. Since no region of the phase space is forbidden, the cancellation of soft gluon effects takes place as in ordinary infrared-safe cross sections. That is the main advantage of this criterion: it eliminates all the fragmentation component in an infrared-safe way. By contrast, it can not be implemented within the usual experimental conditions; the standard way of implementing isolation in experiments is to use the prescription of eq. (3.2) with a constant $\chi(r) = 1$. In any case, from a purely pragmatic point of view, it has been recently shown [63] that if the isolation parameters are tight enough (e.g., $E_{T\ max} < 6$ GeV, $R = 0.4$), the standard and the smooth cone isolation prescription coincide at the 1% level, which is well within the theoretical uncertainty of our predictions.

The acceptance criteria used in this analysis ($\sqrt{s} = 7$ TeV) are those implemented by the ATLAS collaboration analysis [6]; in all the numerical results presented in this paper, we require $p_T^{\text{harder}} \geq 25$ GeV, $p_T^{\text{softer}} \geq 22$ GeV, and we restrict the rapidity of both photons to satisfy $|y_\gamma| < 1.37$ and $1.52 < |y_\gamma| \leq 2.37$. The isolation parameters are set to the values $E_{T\ max} = 4$ GeV, $n = 1$ and $R = 0.4$, and the minimum angular separation between the two photons is $R_{\gamma\gamma} = 0.4$. We use the Martin-Stirling-Thorne-Watt (MSTW) 2008 [79] sets of parton distributions, with densities and α_S evaluated at each corresponding order (i.e., we use $(n + 1)$ -loop α_S at N^n LO, with $n = 0, 1, 2$), and we consider $N_f = 5$ massless quarks/antiquarks and gluons in the initial state. The default renormalization (μ_R) and factorization (μ_F) scales are set to the value of the invariant mass of the diphoton system, $\mu_R = \mu_F = M_{\gamma\gamma}$, while the default resummation scale (μ_{res}) is set to $\mu_{res} = M_{\gamma\gamma}/2$. The QED coupling constant α is fixed to $\alpha = 1/137$.

This choice of the order of the parton densities and α_S is fully justified both in the small- q_T region (where the calculation of the partonic cross section includes the complete NNLO (NLO) result and is controlled by NNLL (NLL) resummation) and in the intermediate- q_T region (where the calculation is constrained by the value of the NNLO (NLO) total cross section).

Non-perturbative (NP) effects are expected to be important at very small q_T . In this paper we follow the strategy of ref. [44], implementing them by multiplying the b -space




Figure 1. The q_T spectrum of the photon pair (solid red lines) at the LHC (7 TeV): results at NLL+LO (left panel) and NNLL+NLO (right panel) accuracy. Each result is compared to the corresponding fixed-order result (dashed lines) and to the finite component (dotted lines) in eq. (2.9). The resummed spectrum includes a non-perturbative (NP) contribution parametrized as in eq. (3.3) and is obtained within the 3-body approach.

form factor $\mathcal{W}^{\gamma\gamma}$ of eq. (2.4) by a ‘NP factor’ which consists of a gaussian smearing of the form

$$S_{NP}^a = \exp(-C_a g_{NP} b^2), \quad (3.3)$$

where a denotes the initial state channel, $a = F$ for $q\bar{q}$ and $a = A$ for gg (as usual, $C_F = (N_c^2 - 1)/(2N_c)$ and $C_A = N_c$). In order to assess the importance of the NP contributions, we vary g_{NP} in the interval from $g_{NP} = 0 \text{ GeV}^2$ (no NP contributions) to $g_{NP} = 2 \text{ GeV}^2$, corresponding to *moderate* NP effects [44].

An additional and potentially important source of theoretical uncertainty arises from an ambiguity in the definition of the photon momenta in the resummation formalism. In fact, in the main resummation formula (2.4), which is used to define both the resummed and (via the subtraction eq. (2.9)) the finite contributions to the partonic cross-section, the diphoton pair total transverse momentum q_T is not associated with the recoil of any extra physical particles in the final state. After q_T resummation the angular distributions of the photons are still provided by the Born level functions ($\sigma_{\gamma\gamma}^{(0)}$, $\mathcal{H}_{N_1 N_2}^{\gamma\gamma}$), which appear as multiplicative factors in front of the Sudakov form factor of eq. (2.7). At this point there are two strategies to follow, which differ by corrections that are of $\mathcal{O}(q_T/M_{\gamma\gamma})$ order-by-order in the perturbative expansion [81] (after having matched the resummed calculation with a complete $N^k LO$ calculation, these corrections start to contribute at the $N^{k+1} LO$).

One of them is to just use the Born phase space for all the angular distributions in front of the Sudakov form factor of eq. (2.7). In this case the transverse momentum q_T is neglected in the calculation of the photon momenta, while the momenta q_i , ($i = 1, 2$) of the colliding partons are given by:

$$q_1^\mu = x_1 P_1^\mu \quad q_2^\mu = x_2 P_2^\mu, \quad (3.4)$$

where x_i ($i = 1, 2$) are the parton momentum fractions and P_i ($i = 1, 2$) are the momenta of the colliding hadrons. The momenta q_i respects the Born level kinematics

$$q_1 + q_2 = q_{\gamma 1} + q_{\gamma 2}, \quad (3.5)$$




Figure 2. Uncertainties in the q_T spectrum of diphoton production at the LHC ($\sqrt{s} = 7$ TeV). In the left panel we show the different predictions associated with the 2-body and 3-body parametrization of the photon momenta at the central scale $\mu_F = \mu_R = M_{\gamma\gamma}$, $\mu_{res} = M_{\gamma\gamma}/2$. In the right panel, the bands (solid and dotted lines) are obtained by varying μ_F and μ_R as described in the text; the prediction at the central scale with the NP contribution turned off ($g_{NP} = 0$ GeV² in eq. (3.3)) is also shown.

where $q_{\gamma i}$, ($i = 1, 2$) are the momenta of the two photons in the final state. In this case the two photons are always in the *back-to-back* configuration, and therefore the kinematic effects of the transverse-momentum recoil, are not included in the momentum of each single photon. We call this approach the 2-body phase space; all the differential distributions which use this method have a (2-body) label.

A more elaborate, and arguably more physical, approach is to consider the effects of the transverse-momentum recoil in the two-photon final state. Therefore these kinematic effects have to be ‘absorbed’ by the incoming parton momenta q_1 and q_2 , in order to respect the Born level kinematics

$$q_1(q_T) + q_2(q_T) = q_{\gamma 1}(q_T) + q_{\gamma 2}(q_T) , \tag{3.6}$$

and the LO kinematics of eq. (3.4). There are different consistent implementations of this approach, all of them giving equivalent (up to higher order corrections) results.³ The implementation that we use in the following is essentially equivalent to define the separate photon momenta according to the Born level angular distribution computed in the Collins-Soper frame [82] of the diphoton system (in practice, this is the same procedure used in ref. [16]). We call this method the 3-body phase space, and distributions obtained using this method are labelled (3-body). The modifications introduced by this phase space parametrization in the momentum fractions x_1 , x_2 of the incoming partons are neglected because they produce negligible effects ($\mathcal{O}(q_T/M_{\gamma\gamma})$) in the cross-section. Notice that it is not the formal 3-body phase space (the 3-body effect is produced by the Lorentz transformation with finite q_T of the Collins-Soper frame). We have checked that if the formal 3-body phase space is used,⁴ the changes in the cross section (comparing the 2-

³The details of the implementation of these kinematic effects are discussed in a forthcoming paper [81].

⁴Which violates the Born level kinematics of eq. (3.5).




Figure 3. The q_T spectrum of diphoton production at the LHC ($\sqrt{s} = 7$ TeV). Here, the scales $\mu_F = \mu_R = M_{\gamma\gamma}$ are kept fixed while we vary the resummation scale μ_{res} to establish its impact on the cross section. In the left panel we show the range $0 \text{ GeV} < q_{T\gamma\gamma} < 40 \text{ GeV}$, and in the right panel the full spectra in logarithmic scale.

body and 3-body cases) are of order $\mathcal{O}(q_T/M_{\gamma\gamma})$. The ambiguity in the treatment of the photon momenta is considered as an additional source of theoretical uncertainty.

In figure 1, left panel, we present the NLL+LO q_T spectrum at the LHC ($\sqrt{s} = 7$ TeV). The NLL+LO result (solid line) at the default scales ($\mu_F = \mu_R = M_{\gamma\gamma}$; $\mu_{res} = M_{\gamma\gamma}/2$) is compared with the corresponding LO result (dashed line). We use the 3-body phase space and a NP parameter $g_{NP} = 2 \text{ GeV}^2$. The LO finite component of the spectrum (see eq. (2.3)), is also shown for comparison (dotted line). We observe that the LO result diverges to $+\infty$ as $q_T \rightarrow 0$, as expected. The finite component is regular over the full q_T range, it smoothly vanishes as $q_T \rightarrow 0$ and gives an important contribution to the NLL+LO result in the low- q_T region. That is mostly originated by the qg channel, which starts at NLO and provides a *subleading* correction in terms of logs (single logarithmic terms) but contributes considerably to the cross-section due to the huge partonic luminosity compared to the formally leading $q\bar{q}$ channel. The resummation of the small- q_T logarithms leads to a well-behaved distribution: it vanishes as $q_T \rightarrow 0$ and approaches the corresponding LO result at large values of q_T .

The results in the right panel of figure 1 are systematically at one order higher: the q_T spectrum at NNLL+NLO accuracy (solid line) is compared with the NLO result (dashed line) and with the NLO finite component of the spectrum (dotted line). The NLO result diverges to $-\infty$ as $q_T \rightarrow 0$ and, at small values of q_T , it has an unphysical peak that is produced by the compensation of negative leading and positive subleading logarithmic contributions. The contribution of the NLO finite component to the NNLL+NLO result is of the order of the 50% at the peak and becomes more important as q_T increases. A similar quantitative behaviour is observed by considering the contribution of the NLO finite component to the NLO result. At large values of q_T the contribution of the NLO finite component tends to the NLO result. This behaviour indicates that the logarithmic terms are no longer dominant and that the resummed calculation cannot improve upon the predictivity of the fixed-order expansion.




Figure 4. The q_T spectrum of diphoton pairs at the LHC. The NNLL+NLO result is compared with the NLL+LO result, for the window $0 \text{ GeV} < q_T < 40 \text{ GeV}$ (left panel) and the full spectra (right panel). We use the 3-body parametrization in the resummed cross-section, and set $g_{NP} = 2 \text{ GeV}^2$. The bands are obtained by varying μ_R and μ_F as explained in the text.




Figure 5. Comparison of the theoretical prediction for the q_T spectrum of diphoton pairs at the LHC to the experimental data. The NNLL+NLO result is compared with the ATLAS data of ref. [6], for the window $0 \text{ GeV} < q_T < 40 \text{ GeV}$ (left panel) and the full spectra (right panel). In the right panel the NLL+LO distribution at central scale (dotted line) is also shown in order to compare it with the data and the NNLL+NLO result. In the theoretical curves we use the 2-body parametrization for the resummed cross-section and set $g_{NP} = 2 \text{ GeV}^2$; the bands (solid and dashed lines) are obtained by varying μ_R and μ_F as explained in the text.

We also observe that the position of the peak in the NNLL+NLO q_T distribution is slightly harder than the corresponding NLL+LO q_T distribution. This effect is (in part) due to the large transverse-momentum dependence of the fixed order corrections.

As discussed in section 2, the resummed calculation depends on the factorization and renormalization scales and on the resummation scale μ_{res} . Our convention to compute factorization and renormalization scale uncertainties is to consider independent variations of μ_F and μ_R by a factor of two around the central values $\mu_F = \mu_R = M_{\gamma\gamma}$ in independent way in order to maximise them: $(\mu_F = 2 M_{\gamma\gamma}, \mu_R = M_{\gamma\gamma}/2, \mu_{res} = M_{\gamma\gamma}/2)$ and $(\mu_R = 2 M_{\gamma\gamma}, \mu_F = M_{\gamma\gamma}/2, \mu_{res} = M_{\gamma\gamma}/2)$. The uncertainty due to the resummation scale variation is assessed separately by varying it between $\mu_{res} = M_{\gamma\gamma}/4$ and $\mu_{res} = M_{\gamma\gamma}$ at fixed μ_F and μ_R .




Figure 6. Comparison of the theoretical prediction for the q_T spectrum of diphoton pairs at the LHC with the experimental data. The NNLL+NLO result is compared with the ATLAS data of ref. [6], for the window $0 \text{ GeV} < q_T < 40 \text{ GeV}$ (left panel) and the full spectra (right panel). In the right panel the NNLL+LO distribution at central scale (dotted line) is also shown in order to compare it with the data and the NNLL+NLO result. In the theoretical curves we use the 3-body parametrization for the resummed cross-section and set $g_{NP} = 2 \text{ GeV}^2$. The bands (solid and dashed lines) are obtained by varying μ_R and μ_F as explained in the text.




Figure 7. Individual photon spectra in diphoton production at the LHC. Left panel: the $q_T^H_\gamma$ spectrum of the hard photon at the LHC [6] ($\sqrt{s} = 7 \text{ TeV}$). Right panel: the $q_T^S_\gamma$ spectrum of the softer photon at the LHC [6] ($\sqrt{s} = 7 \text{ TeV}$). In both panels, we compare the resummed prediction using the 3-body parametrization in the resummed cross-section ($g_{NP} = 2 \text{ GeV}^2$) with the fixed order prediction.

In figure 2 we compare the impact of the various sources of theoretical error for the NNLL+NLO predictions: the 2-body and 3-body parametrizations of the photon phase space, the effect of the variation of the non-perturbative contribution, and the variation of the factorization and renormalization scales. In the left panel of figure 2 we use only the central scale ($\mu_F = \mu_R = M_{\gamma\gamma}$, $\mu_{res} = M_{\gamma\gamma}/2$), and we note that the 3-body method results in a slightly larger cross section (about 10%) around the peak ($q_T \sim 5 \text{ GeV}$) than the 2-body one. For $q_T > 20 \text{ GeV}$ all the contributions in the left panel coincide, which is consistent with the fact that at these values the resummed component starts to vanish and the fixed order result dominates the cross section, as we can anticipate from figure 1. Also we note that if we use a NP parameter, the peak of the distribution is located at

larger values of q_T ($q_T \sim 5$ GeV) and the shape of the distribution is slightly different (for values $0 \text{ GeV} < q_T < 22 \text{ GeV}$) from the case in which the NP parameter is not implemented. These differences, which are stronger for $q_T < 10$ GeV, have their origin in the resummed component, which is the only contribution that depends on g_{NP} . In the right panel of figure 2 we show the comparison between the variation of the scales⁵ in the 3-body approach with the central scale results of the 2-body and 3-body frameworks (with and without non perturbative parameter, respectively). Evidently, the uncertainty due to the ambiguity of the parametrization of the photon momenta turns out to be subdominant with respect to the one arising from the variation of the scales which provides, by itself, a reasonable estimate of theoretical uncertainties.

In figure 3 we show the NNLL+NLO transverse momentum distribution for three different implementations of the μ_{res} parameter ($\mu_{res} = M_{\gamma\gamma}/4; M_{\gamma\gamma}/2; M_{\gamma\gamma}$) at fixed $\mu_F = \mu_R = M_{\gamma\gamma}$. The 2-body phase space and a non-perturbative parameter $g_{NP} = 2 \text{ GeV}^2$ were used. We notice the small impact of the variation of the μ_{res} scale in the cross section (at per-cent level). In the left panel of figure 3 we present the transverse momentum distribution for values of q_T within the interval $0 \text{ GeV} < q_T < 40 \text{ GeV}$, and in the right panel of figure 3 the full spectra. We also notice that the strongest effect of the variation of the μ_{res} scale appears in the last bin of right panel of figure 3. This is expected since the resummation scale effectively *sets* the value of transverse momentum at which the logarithms are dominant. A choice of a very large resummation scale affects the distribution at larger transverse momentum and might in general result in a mismatch with the fixed order prediction due to the artificial introduction of unphysically large logarithmic contributions in that region. Similar results are obtained if the 3-body phase space is used instead of the 2-body one.

In figure 4 we compare the variation of the scales of the NNLL+NLO and NLL+LO predictions (3-body phase space), for the interval $0 \text{ GeV} < q_T < 40 \text{ GeV}$ (left panel) and the full spectra (right panel). We notice that the dependence on the scales is not reduced when going from NLL+LO to NNLL+NLO. This is mostly because at NNLL+NLO a new channel (gg) opens, in which the box contribution (effectively “LO” but formally $\mathcal{O}(\alpha_S^2)$) ruins the reduction of the scale dependence usually expected when adding second order corrections for the $q\bar{q}$ channel and first order corrections for the qg channel. This effect has the same origin that the reported behaviour of the diphoton production at NNLO of ref. [23], when the variation of the scales is implemented. Since NNLL+NLO is the first order at which all partonic channels contribute, it is possible to argue that this is the first order at which estimates of theoretical uncertainties through scale variations can be considered as reliable. The same results are obtained if the 2-body phase space is used instead of the 3-body one.

The peak observed in the right panel of figure 4 is the so called *Guillet shoulder* [80], which is a real radiation effect and has its origin in the fixed order contribution. It appears stronger in the NNLL+NLO q_T distribution than in the NLL+LO, due to the larger size of the real contributions at NLO.

⁵We vary the scales from ($\mu_F = 2M_{\gamma\gamma}; \mu_R = M_{\gamma\gamma}/2$) to ($\mu_R = 2M_{\gamma\gamma}; \mu_F = M_{\gamma\gamma}/2$) at fixed $\mu_{res} = M_{\gamma\gamma}/2$.

In the small q_T region ($q_T < 4 \text{ GeV}$) the real radiation effects are no longer dominant in the q_T distribution. The finite and resummed component (which vanish as $q_T \rightarrow 0$) are of the same order for $q_T < 4 \text{ GeV}$. In absence of the strong real radiation effects the contributions are almost completely Born like. This is the main reason why the NNLL+NLO bands overlaps with those at the previous order.

In figures 5 and 6 we compare the LHC data ($\sqrt{s} = 7 \text{ TeV}$) from ATLAS [6] with our resummed theoretical predictions (at NNLL+NLO and NLL+LO) using the 2-body and 3-body approaches, respectively. In both cases we use a NP parameter different from zero ($g_{NP} = 2 \text{ GeV}^2$) and we estimate the theoretical uncertainty by the variation of the μ_R and μ_F scales. In the left panels we show the q_T distribution in the window ($0 \text{ GeV} < q_T < 40 \text{ GeV}$), while in the right ones we show the full spectra in logarithmic scale.

We observe in general an excellent agreement between the resummed NNLL+NLO prediction and the experimental data, that is accurately described within the theoretical uncertainty bands in the whole kinematic range. Also we observe that the NLL+LO result is not enough to describe the phenomenology of the transverse-momentum distribution of the LHC data (figures 5 and 6, right panel). By direct comparison to the fixed order prediction, we notice that the effect of resummation is not only to recover the predictivity of the calculation at small transverse momentum, but also to improve substantially the agreement with LHC data [6].

While the resummation performed in this work reaches NNLL accuracy formally only for the diphoton transverse momentum distribution, its predictions can be extended to other observables as well, since at least the leading logarithmic contributions have a common origin from soft and collinear emission. Note that the 3-body approach is more suitable for a consistent implementation of these leading logarithmic effects on the observables. In figure 7 we show results on more exclusive observables: the q_T distributions of the harder (left-hand plot) and softer (right-hand plot) photon at the central scale. We compare the results obtained with the 2-body and 3-body phase space approaches, using a non-perturbative parameter $g_{NP} = 2 \text{ GeV}^2$ with the fixed order result at NNLO for diphoton production [23].

The 2-body phase space transverse-momentum distribution at NNLL+NLO provides the same result than the NNLO fixed order cross section for diphoton production [23]. This is consistent with two following related facts: *i*) the single photon momentum does not carry any information about the recoil due to the transverse momentum q_T in the 2-body approach; *ii*) the NNLO (NLO) result for the total cross section is exactly recovered upon integration over q_T of the differential cross section $d\sigma_{\gamma\gamma}/dq_T$ at NNLL+NLO (NLL+LO) accuracy.

The effects of resummation are only present in these more exclusive observables if the recoil due to q_T are absorbed by the photons in the final state,⁶ which is equivalent to the implementation of a 3-body like phase space. In this way, because the single photon momentum depends on q_T , the integral over q_T of the NNLL+NLO (NLL+LO) distribution

⁶And also the recoil due to q_T has to be absorbed in the initial state to restore the momentum conservation (see eq. (3.6)).




Figure 8. The $\Delta\Phi_{\gamma\gamma}$ distribution of diphoton pairs at the LHC. In the left panel, we show the fixed order prediction, the complete resummed prediction and just the resummed contribution at the central scale. In the right, the full NNLL+NLO result (using the extreme values for μ_R and μ_F to estimate the theoretical uncertainty) is compared with the ATLAS data of ref. [6].

does not recover the NNLO (NLO) result as we can observe in figure 7. Here the last integral over the single photon momentum of the NNLL+NLO (NLL+LO) distribution, is required in order to recover the NNLO (NLO) result for the total cross section.

We comment on the q_T distribution of the softer photon in the region around the back-to-back threshold $q_{T\gamma}^S \sim 25$ GeV (see figure 7 right panel). The NLO fixed order result has a step-like behaviour, and this necessarily produces [83] integrable logarithmic singularities at each subsequent perturbative order. The peak of the NLO fixed order distribution at $q_{T\gamma}^S \sim 25$ GeV is an artifact of these perturbative instabilities. The instability is cured by all-order perturbative resummation, which leads to a smooth q_T distribution with a shoulder-like behaviour [83] in the vicinity of the back-to-back threshold.

In figure 8 we present the results of the cross section as a function of the azimuthal angle $\Delta\Phi_{\gamma\gamma}$. In the left panel we compare the fixed order (NLO), finite (NLO) and full (NNLL+NLO) $\Delta\Phi_{\gamma\gamma}$ distributions. The fixed order component dominates the cross section over the whole $\Delta\Phi_{\gamma\gamma}$ range. However, as could be expected, the effect of resummation is stronger for kinematic configurations near the $\Delta\Phi_{\gamma\gamma} \sim \pi$ which correspond to $q_T \sim 0$ GeV. As in the case of the fixed order q_T distribution, the $\Delta\Phi_{\gamma\gamma}$ fixed order differential cross section is not well-behaved near the back-to-back configuration: it actually diverges as $\Delta\Phi_{\gamma\gamma} \rightarrow \pi$ ($q_T \rightarrow 0$). The finite contribution (eq. (2.9)) is well-behaved near the back-to-back configuration, and the full result (NNLL+NLO) improves the description in the region near $\Delta\Phi_{\gamma\gamma} \sim 0$.

In the right panel of figure 8 we compare our theoretical prediction at NNLL+NLO level of accuracy with the LHC data [6] using the variation of the μ_R and μ_F scales to estimate the theoretical uncertainty. We observe that the transverse momentum resummation provides a better description of the data with respect to the fixed order result. In both panels of figure 8 we used the 3-body approach to describe the diphoton phase-space. In fact, in this kind of observables (also see figure 7), the 2-body parametrization again reproduces the result of the fixed order cross section.

4 Summary

In this paper we performed the transverse momentum resummation for diphoton production at NNLL accuracy in hadron collisions. At small values of q_T , the calculation includes the resummation of all logarithmically-enhanced perturbative QCD contributions, up to next-to-next-to-leading logarithmic accuracy; at intermediate and large values of q_T , it combines the resummation with the fixed next-to-leading order perturbative result. The combination is performed in such a way as to exactly reproduce the known next-to-next-to-leading order result for the total cross section; in the end, the calculation consistently includes all perturbative terms up to formal order α_S^2 . The theoretical uncertainty was estimated by varying the various scales (renormalization, factorization and resummation) introduced by the formalism as well as the parametrization of the diphoton phase-space. The result was compared to experimental data, showing good agreement between theory and experiment over the whole q_T range. With respect to the fixed-order calculation, the present implementation provides a better description of the data and recovers the correct physical behaviour in the small q_T region, with the spectrum smoothly going to zero. The same set-up also allows the calculation of more exclusive observable distributions; the q_T spectrum of the individual photons and the $\Delta\Phi_{\gamma\gamma}$ distribution are given as examples.

Acknowledgments

We are very grateful to Stefano Catani and Giancarlo Ferrera, for taking a careful look to this paper and making extremely valuable comments about it. LC would like to thank the INFN of Florence for kind hospitality, where parts of the project were carried out. FC would like to thank the Departamento de Física of the Universidad de Buenos Aires and the FCEYN, which supported him during the initial stages of this work.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] CMS collaboration, *Measurement of the production cross section for pairs of isolated photons in pp collisions at $\sqrt{s} = 7$ TeV*, *JHEP* **01** (2012) 133 [[arXiv:1110.6461](https://arxiv.org/abs/1110.6461)] [[INSPIRE](#)].
- [2] CDF collaboration, T. Aaltonen et al., *Measurement of the cross section for prompt isolated diphoton production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV*, *Phys. Rev. D* **84** (2011) 052006 [[arXiv:1106.5131](https://arxiv.org/abs/1106.5131)] [[INSPIRE](#)].
- [3] CMS collaboration, *Measurement of differential cross sections for the production of a pair of isolated photons in pp collisions at $\sqrt{s} = 7$ TeV*, *Eur. Phys. J. C* **74** (2014) 3129 [[arXiv:1405.7225](https://arxiv.org/abs/1405.7225)] [[INSPIRE](#)].
- [4] D0 collaboration, V.M. Abazov et al., *Measurement of the differential cross sections for isolated direct photon pair production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV*, *Phys. Lett. B* **725** (2013) 6 [[arXiv:1301.4536](https://arxiv.org/abs/1301.4536)] [[INSPIRE](#)].

- [5] ATLAS collaboration, *Measurement of the isolated di-photon cross-section in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector*, *Phys. Rev. D* **85** (2012) 012003 [[arXiv:1107.0581](#)] [[INSPIRE](#)].
- [6] ATLAS collaboration, *Measurement of isolated-photon pair production in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector*, *JHEP* **01** (2013) 086 [[arXiv:1211.1913](#)] [[INSPIRE](#)].
- [7] CMS collaboration, *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, *Phys. Lett. B* **716** (2012) 30 [[arXiv:1207.7235](#)] [[INSPIRE](#)].
- [8] ATLAS collaboration, *Observation of a new particle in the search for the standard model Higgs boson with the ATLAS detector at the LHC*, *Phys. Lett. B* **716** (2012) 1 [[arXiv:1207.7214](#)] [[INSPIRE](#)].
- [9] ATLAS collaboration, *Search for diphoton events with large missing transverse momentum in 7 TeV proton-proton collision data with the ATLAS detector*, *Phys. Lett. B* **718** (2012) 411 [[arXiv:1209.0753](#)] [[INSPIRE](#)].
- [10] CMS collaboration, *Search for new physics in events with photons, jets and missing transverse energy in pp collisions at $\sqrt{s} = 7$ TeV*, *JHEP* **03** (2013) 111 [[arXiv:1211.4784](#)] [[INSPIRE](#)].
- [11] ATLAS collaboration, *Search for extra dimensions in diphoton events using proton-proton collisions recorded at $\sqrt{s} = 7$ TeV with the ATLAS detector at the LHC*, *New J. Phys.* **15** (2013) 043007 [[arXiv:1210.8389](#)] [[INSPIRE](#)].
- [12] CMS collaboration, *Search for supersymmetry in events with photons and low missing transverse energy in pp collisions at $\sqrt{s} = 7$ TeV*, *Phys. Lett. B* **719** (2013) 42 [[arXiv:1210.2052](#)] [[INSPIRE](#)].
- [13] T. Binoth, J.P. Guillet, E. Pilon and M. Werlen, *A full next-to-leading order study of direct photon pair production in hadronic collisions*, *Eur. Phys. J. C* **16** (2000) 311 [[hep-ph/9911340](#)] [[INSPIRE](#)].
- [14] Z. Bern, L.J. Dixon and C. Schmidt, *Isolating a light Higgs boson from the diphoton background at the CERN LHC*, *Phys. Rev. D* **66** (2002) 074018 [[hep-ph/0206194](#)] [[INSPIRE](#)].
- [15] J.M. Campbell, R.K. Ellis and C. Williams, *Vector boson pair production at the LHC*, *JHEP* **07** (2011) 018 [[arXiv:1105.0020](#)] [[INSPIRE](#)].
- [16] C. Balázs, E.L. Berger, P.M. Nadolsky and C.-P. Yuan, *Calculation of prompt diphoton production cross-sections at Tevatron and LHC energies*, *Phys. Rev. D* **76** (2007) 013009 [[arXiv:0704.0001](#)] [[INSPIRE](#)].
- [17] D.A. Dicus and S.S.D. Willenbrock, *Photon pair production and the intermediate mass Higgs boson*, *Phys. Rev. D* **37** (1988) 1801 [[INSPIRE](#)].
- [18] V.D. Barger, T. Han, J. Ohnemus and D. Zeppenfeld, *Pair production of W^\pm , γ and Z in association with jets*, *Phys. Rev. D* **41** (1990) 2782 [[INSPIRE](#)].
- [19] V. Del Duca, W.B. Kilgore and F. Maltoni, *Multiphoton amplitudes for next-to-leading order QCD*, *Nucl. Phys. B* **566** (2000) 252 [[hep-ph/9910253](#)] [[INSPIRE](#)].
- [20] Z. Bern, L.J. Dixon and D.A. Kosower, *One loop corrections to two quark three gluon amplitudes*, *Nucl. Phys. B* **437** (1995) 259 [[hep-ph/9409393](#)] [[INSPIRE](#)].
- [21] A. Signer, *One loop corrections to five parton amplitudes with external photons*, *Phys. Lett. B* **357** (1995) 204 [[hep-ph/9507442](#)] [[INSPIRE](#)].

- [22] C. Anastasiou, E.W.N. Glover and M.E. Tejeda-Yeomans, *Two loop QED and QCD corrections to massless fermion boson scattering*, *Nucl. Phys. B* **629** (2002) 255 [[hep-ph/0201274](#)] [[INSPIRE](#)].
- [23] S. Catani, L. Cieri, D. de Florian, G. Ferrera and M. Grazzini, *Diphoton production at hadron colliders: a fully-differential QCD calculation at NNLO*, *Phys. Rev. Lett.* **108** (2012) 072001 [[arXiv:1110.2375](#)] [[INSPIRE](#)].
- [24] V. Del Duca, F. Maltoni, Z. Nagy and Z. Trócsányi, *QCD radiative corrections to prompt diphoton production in association with a jet at hadron colliders*, *JHEP* **04** (2003) 059 [[hep-ph/0303012](#)] [[INSPIRE](#)].
- [25] T. Gehrmann, N. Greiner and G. Heinrich, *Photon isolation effects at NLO in $\gamma\gamma$ + jet final states in hadronic collisions*, *JHEP* **06** (2013) 058 [Erratum *ibid.* **06** (2014) 076] [[arXiv:1303.0824](#)] [[INSPIRE](#)].
- [26] Z. Bern et al., *Next-to-leading order $\gamma\gamma$ + 2-jet production at the LHC*, *Phys. Rev. D* **90** (2014) 054004 [[arXiv:1402.4127](#)] [[INSPIRE](#)].
- [27] T. Gehrmann, N. Greiner and G. Heinrich, *Precise QCD predictions for the production of a photon pair in association with two jets*, *Phys. Rev. Lett.* **111** (2013) 222002 [[arXiv:1308.3660](#)] [[INSPIRE](#)].
- [28] S. Badger, A. Guffanti and V. Yundin, *Next-to-leading order QCD corrections to di-photon production in association with up to three jets at the Large Hadron Collider*, *JHEP* **03** (2014) 122 [[arXiv:1312.5927](#)] [[INSPIRE](#)].
- [29] Y.L. Dokshitzer, D. Diakonov and S.I. Troian, *On the transverse momentum distribution of massive lepton pairs*, *Phys. Lett. B* **79** (1978) 269 [[INSPIRE](#)].
- [30] Y.L. Dokshitzer, D. Diakonov and S.I. Troian, *Hard processes in quantum chromodynamics*, *Phys. Rept.* **58** (1980) 269 [[INSPIRE](#)].
- [31] G. Parisi and R. Petronzio, *Small transverse momentum distributions in hard processes*, *Nucl. Phys. B* **154** (1979) 427 [[INSPIRE](#)].
- [32] G. Curci, M. Greco and Y. Srivastava, *QCD jets from coherent states*, *Nucl. Phys. B* **159** (1979) 451 [[INSPIRE](#)].
- [33] J.C. Collins and D.E. Soper, *Back-To-Back Jets in QCD*, *Nucl. Phys. B* **193** (1981) 381 [Erratum *ibid.* **B 213** (1983) 545] [[INSPIRE](#)].
- [34] J.C. Collins and D.E. Soper, *Back-to-back jets: Fourier transform from B to K-transverse*, *Nucl. Phys. B* **197** (1982) 446 [[INSPIRE](#)].
- [35] J. Kodaira and L. Trentadue, *Summing soft emission in QCD*, *Phys. Lett. B* **112** (1982) 66 [[INSPIRE](#)].
- [36] J. Kodaira and L. Trentadue, *Soft gluon effects in perturbative quantum chromodynamics*, SLAC-PUB-2934 (1982).
- [37] J. Kodaira and L. Trentadue, *Single logarithm effects in electron-positron annihilation*, *Phys. Lett. B* **123** (1983) 335 [[INSPIRE](#)].
- [38] C.T.H. Davies and W.J. Stirling, *Nonleading corrections to the Drell-Yan cross-section at small transverse momentum*, *Nucl. Phys. B* **244** (1984) 337 [[INSPIRE](#)].
- [39] C.T.H. Davies, B.R. Webber and W.J. Stirling, *Drell-Yan cross-sections at small transverse momentum*, *Nucl. Phys. B* **256** (1985) 413 [[INSPIRE](#)].

- [40] G. Altarelli, R.K. Ellis, M. Greco and G. Martinelli, *Vector boson production at colliders: a theoretical reappraisal*, *Nucl. Phys. B* **246** (1984) 12 [INSPIRE].
- [41] J.C. Collins, D.E. Soper and G.F. Sterman, *Transverse momentum distribution in Drell-Yan pair and W and Z boson production*, *Nucl. Phys. B* **250** (1985) 199 [INSPIRE].
- [42] S. Catani, D. de Florian and M. Grazzini, *Universality of nonleading logarithmic contributions in transverse momentum distributions*, *Nucl. Phys. B* **596** (2001) 299 [[hep-ph/0008184](#)] [INSPIRE].
- [43] S. Catani, L. Cieri, D. de Florian, G. Ferrera and M. Grazzini, *Universality of transverse-momentum resummation and hard factors at the NNLO*, *Nucl. Phys. B* **881** (2014) 414 [[arXiv:1311.1654](#)] [INSPIRE].
- [44] G. Bozzi, S. Catani, D. de Florian and M. Grazzini, *Transverse-momentum resummation and the spectrum of the Higgs boson at the LHC*, *Nucl. Phys. B* **737** (2006) 73 [[hep-ph/0508068](#)] [INSPIRE].
- [45] G. Bozzi, S. Catani, D. de Florian and M. Grazzini, *Higgs boson production at the LHC: transverse-momentum resummation and rapidity dependence*, *Nucl. Phys. B* **791** (2008) 1 [[arXiv:0705.3887](#)] [INSPIRE].
- [46] S. Catani and M. Grazzini, *QCD transverse-momentum resummation in gluon fusion processes*, *Nucl. Phys. B* **845** (2011) 297 [[arXiv:1011.3918](#)] [INSPIRE].
- [47] G. Bozzi, S. Catani, D. de Florian and M. Grazzini, *The $q(T)$ spectrum of the Higgs boson at the LHC in QCD perturbation theory*, *Phys. Lett. B* **564** (2003) 65 [[hep-ph/0302104](#)] [INSPIRE].
- [48] D. de Florian, G. Ferrera, M. Grazzini and D. Tommasini, *Transverse-momentum resummation: Higgs boson production at the Tevatron and the LHC*, *JHEP* **11** (2011) 064 [[arXiv:1109.2109](#)] [INSPIRE].
- [49] D. de Florian, G. Ferrera, M. Grazzini and D. Tommasini, *Higgs boson production at the LHC: transverse momentum resummation effects in the $H \rightarrow 2\gamma$, $H \rightarrow WW \rightarrow l\nu l\nu$ and $H \rightarrow ZZ \rightarrow 4l$ decay modes*, *JHEP* **06** (2012) 132 [[arXiv:1203.6321](#)] [INSPIRE].
- [50] R.V. Harlander, A. Tripathi and M. Wiesemann, *Higgs production in bottom quark annihilation: Transverse momentum distribution at NNLO+NNLL*, *Phys. Rev. D* **90** (2014) 015017 [[arXiv:1403.7196](#)] [INSPIRE].
- [51] R.V. Harlander, H. Mantler and M. Wiesemann, *Transverse momentum resummation for Higgs production via gluon fusion in the MSSM*, *JHEP* **11** (2014) 116 [[arXiv:1409.0531](#)] [INSPIRE].
- [52] G. Bozzi, S. Catani, G. Ferrera, D. de Florian and M. Grazzini, *Transverse-momentum resummation: A Perturbative study of Z production at the Tevatron*, *Nucl. Phys. B* **815** (2009) 174 [[arXiv:0812.2862](#)] [INSPIRE].
- [53] G. Bozzi, S. Catani, G. Ferrera, D. de Florian and M. Grazzini, *Production of Drell-Yan lepton pairs in hadron collisions: Transverse-momentum resummation at next-to-next-to-leading logarithmic accuracy*, *Phys. Lett. B* **696** (2011) 207 [[arXiv:1007.2351](#)] [INSPIRE].
- [54] M. Grazzini, *Soft-gluon effects in WW production at hadron colliders*, *JHEP* **01** (2006) 095 [[hep-ph/0510337](#)] [INSPIRE].

- [55] P. Meade, H. Ramani and M. Zeng, *Transverse momentum resummation effects in W^+W^- measurements*, *Phys. Rev. D* **90** (2014) 114006 [[arXiv:1407.4481](#)] [[INSPIRE](#)].
- [56] R. Frederix and M. Grazzini, *Higher-order QCD effects in the $h \rightarrow ZZ$ search channel at the LHC*, *Phys. Lett. B* **662** (2008) 353 [[arXiv:0801.2229](#)] [[INSPIRE](#)].
- [57] G. Bozzi, B. Fuks and M. Klasen, *Transverse-momentum resummation for slepton-pair production at the CERN LHC*, *Phys. Rev. D* **74** (2006) 015001 [[hep-ph/0603074](#)] [[INSPIRE](#)].
- [58] H. Kawamura, J. Kodaira, H. Shimizu and K. Tanaka, *The dilepton $Q(T)$ spectrum in transversely polarized Drell-Yan process in QCD*, *Prog. Theor. Phys.* **115** (2006) 667 [[hep-ph/0512137](#)] [[INSPIRE](#)].
- [59] H. Kawamura, J. Kodaira and K. Tanaka, *Soft gluon corrections to double transverse-spin asymmetries for small- $Q(T)$ dilepton production at RHIC and J-PARC*, *Nucl. Phys. B* **777** (2007) 203 [[hep-ph/0703079](#)] [[INSPIRE](#)].
- [60] H. Kawamura, J. Kodaira and K. Tanaka, *Transversely polarized Drell-Yan process and soft gluon resummation in QCD*, *Prog. Theor. Phys.* **118** (2007) 581 [[arXiv:0709.1752](#)] [[INSPIRE](#)].
- [61] H. Kawamura, J. Kodaira and K. Tanaka, *Double-spin asymmetries for small- $Q(T)$ Drell-Yan pair production in transversely polarized $p\bar{p}$ collisions*, *Phys. Lett. B* **662** (2008) 139 [[arXiv:0801.0026](#)] [[INSPIRE](#)].
- [62] S. Frixione, *Isolated photons in perturbative QCD*, *Phys. Lett. B* **429** (1998) 369 [[hep-ph/9801442](#)] [[INSPIRE](#)].
- [63] J. Butterworth et al., *Les Houches 2013: physics at TeV colliders: standard model working group report*, [arXiv:1405.1067](#) [[INSPIRE](#)].
- [64] H.T. Li, C.S. Li, D.Y. Shao, L.L. Yang and H.X. Zhu, *Top quark pair production at small transverse momentum in hadronic collisions*, *Phys. Rev. D* **88** (2013) 074004 [[arXiv:1307.2464](#)] [[INSPIRE](#)].
- [65] S. Catani, M. Grazzini and A. Torre, *Transverse-momentum resummation for heavy-quark hadroproduction*, *Nucl. Phys. B* **890** (2014) 518 [[arXiv:1408.4564](#)] [[INSPIRE](#)].
- [66] D. de Florian and M. Grazzini, *Next-to-next-to-leading logarithmic corrections at small transverse momentum in hadronic collisions*, *Phys. Rev. Lett.* **85** (2000) 4678 [[hep-ph/0008152](#)] [[INSPIRE](#)].
- [67] D. de Florian and M. Grazzini, *The structure of large logarithmic corrections at small transverse momentum in hadronic collisions*, *Nucl. Phys. B* **616** (2001) 247 [[hep-ph/0108273](#)] [[INSPIRE](#)].
- [68] T. Becher and M. Neubert, *Drell-Yan production at small q_T , transverse parton distributions and the collinear anomaly*, *Eur. Phys. J. C* **71** (2011) 1665 [[arXiv:1007.4005](#)] [[INSPIRE](#)].
- [69] C. Balázs, E.L. Berger, P.M. Nadolsky and C.-P. Yuan, *All-orders resummation for diphoton production at hadron colliders*, *Phys. Lett. B* **637** (2006) 235 [[hep-ph/0603037](#)] [[INSPIRE](#)].
- [70] P.M. Nadolsky and C.R. Schmidt, *Diphoton production in gluon fusion at small transverse momentum*, *Phys. Lett. B* **558** (2003) 63 [[hep-ph/0211398](#)] [[INSPIRE](#)].
- [71] C. Balázs, P.M. Nadolsky, C. Schmidt and C.P. Yuan, *Diphoton background to Higgs boson production at the LHC with soft gluon effects*, *Phys. Lett. B* **489** (2000) 157 [[hep-ph/9905551](#)] [[INSPIRE](#)].

- [72] A. Kulesza, G.F. Sterman and W. Vogelsang, *Joint resummation in electroweak boson production*, *Phys. Rev. D* **66** (2002) 014011 [[hep-ph/0202251](#)] [[INSPIRE](#)].
- [73] J. Blumlein, *Analytic continuation of Mellin transforms up to two loop order*, *Comput. Phys. Commun.* **133** (2000) 76 [[hep-ph/0003100](#)] [[INSPIRE](#)].
- [74] J. Blumlein and S. Kurth, *Harmonic sums and Mellin transforms up to two loop order*, *Phys. Rev. D* **60** (1999) 014018 [[hep-ph/9810241](#)] [[INSPIRE](#)].
- [75] J. Blumlein and V. Ravindran, *Mellin moments of the next-to-next-to leading order coefficient functions for the Drell-Yan process and hadronic Higgs-boson production*, *Nucl. Phys. B* **716** (2005) 128 [[hep-ph/0501178](#)] [[INSPIRE](#)].
- [76] J. Blumlein and S.-O. Moch, *Analytic continuation of the harmonic sums for the 3-loop anomalous dimensions*, *Phys. Lett. B* **614** (2005) 53 [[hep-ph/0503188](#)] [[INSPIRE](#)].
- [77] S. Frixione and W. Vogelsang, *Isolated photon production in polarized pp collisions*, *Nucl. Phys. B* **568** (2000) 60 [[hep-ph/9908387](#)] [[INSPIRE](#)].
- [78] S. Catani et al., *QCD*, in *Geneva 1999, Standard model physics (and more) at the LHC*, May 25–29, Geneva, Switzerland (1999), [[hep-ph/0005025](#)] [[INSPIRE](#)].
- [79] A.D. Martin, W.J. Stirling, R.S. Thorne and G. Watt, *Parton distributions for the LHC*, *Eur. Phys. J. C* **63** (2009) 189 [[arXiv:0901.0002](#)] [[INSPIRE](#)].
- [80] T. Binoth, J.P. Guillet, E. Pilon and M. Werlen, *Beyond leading order effects in photon pair production at the Tevatron*, *Phys. Rev. D* **63** (2001) 114016 [[hep-ph/0012191](#)] [[INSPIRE](#)].
- [81] S. Catani, G. Ferrera, D. de Florian and M. Grazzini, in preparation.
- [82] J.C. Collins and D.E. Soper, *Angular distribution of dileptons in high-energy hadron collisions*, *Phys. Rev. D* **16** (1977) 2219 [[INSPIRE](#)].
- [83] S. Catani and B.R. Webber, *Infrared safe but infinite: Soft gluon divergences inside the physical region*, *JHEP* **10** (1997) 005 [[hep-ph/9710333](#)] [[INSPIRE](#)].