

## Understanding comorbidity: from epidemiological designs and model-fitting approaches to systems biology as a new tool

J. K. Buitelaar

Published online: 23 December 2011

© The Author(s) 2011. This article is published with open access at [Springerlink.com](http://Springerlink.com)

It is a common observation that persons with one particular psychiatric disorder often present with other disorders. This so-called co-occurrence, co-existence or comorbidity of disorders occurs at all ages, and cannot be explained away simply by sampling bias. Co-occurrence of disorders may lead to a higher rate of clinical referrals and use of services, and this could suggest inflated rates of co-occurrence. However, population-based studies confirm that many psychiatric disorders cluster in the same individual more than to be expected by chance [1–3]. Traditional explanatory models for this co-occurrence specify the following possibilities: artificial factors, such as overlapping diagnostic criteria or population stratification (the risk factors that influence the two disorders maybe more common in a part of the population); one disorder being a risk factor for the other; one disorder being a cause of the other; risk factors that are being shared between the two disorders; and both disorders being alternative expressions or different phases of the same overarching disorder [4]. Comorbidity has found to be associated with greater severity of illness, increased use of services and health care costs, lower response to treatment and an overall worse prognosis. Therefore, it is important to study the causes of comorbidity. This can have then a major impact on future research examining the two disorders, including research on classification, treatment, and etiology.

This issue contains a number of papers that focus on the co-occurrence of disorders. Polier et al. report on the

significant comorbidity between conduct disorder problems and anxiety and depression in both a clinical and an epidemiological sample. Children with combined conduct and internalizing problems were more impaired and had more social problems than children with only conduct problems. In fact, the social problems of these children were more closely linked to increased levels of anxiety and depression than to conduct problems. The theoretical implication is that internalizing problems do not protect from developing high rates of conduct problems. In clinical practice, the disruptive effects of aggressive and antisocial behaviors rather than the less manifest internalizing problems easily get the attention of parents, teachers and health care professionals. Therefore, the practical consequence should be that more active screening is needed for the presence of internalizing problems among aggressive youngsters and that interventions for these internalizing problems should be offered proactively.

Not only more than one psychiatric disorder but also psychiatric and somatic disorders tend to cluster in the same individuals. This may be due to common risk factors, such as unhealthy lifestyle, adverse psychosocial circumstances, and personality characteristics as neuroticism [5]. These factors, however, cannot probably wholly explain that for example patients with schizophrenia have a baseline decreased insulin tolerance and increased risk for developing diabetes mellitus type 2, even before any exposure to treatment with antipsychotics [6]. Certainly, patients with such severe psychiatric disorders tend to be physically less active, may have less access to foods lower in calories and higher in nutrients and are more inclined to smoke. Smoking in turn can increase insulin resistance. In addition to these lifestyle factors, it is likely that common genetic or metabolic abnormalities underlie the vulnerability to develop both schizophrenia and diabetes mellitus and metabolic syndrome.

---

J. K. Buitelaar (✉)

Department of Cognitive Neuroscience,  
Radboud University Nijmegen Medical Center,  
Karakter Child and Adolescent Psychiatry University Center,  
P.O. Box 9101 (204), HB 6500 Nijmegen, The Netherlands  
e-mail: [J.Buitelaar@psy.umcn.nl](mailto:J.Buitelaar@psy.umcn.nl)

These psychiatric–somatic comorbidities have been less well studied in children and adolescents. Kastner et al. discuss in this issue an example of somatic complications of psychiatric disease. About one-third of patients with anorexia nervosa had pericardial effusion on echocardiography that had no hemodynamic consequences and remitted in the majority of cases after normalisation of weight. Predictors of the pericardial effusion were a low body mass index and a low level of thyroid hormone T3. The authors do not recommend routine echocardiography in patients with anorexia but their findings should make clinicians even more cautious about handling the somatic status of these patients.

Erhart et al. were able to confirm that the association between ADHD and obesity reported in clinical samples is also found in population-based designs. Children with overweight in the general population were twice as likely to have significantly increased levels of ADHD symptoms compared to normal weight children. Vice versa, children with ADHD were twice as likely to have overweight. Their results are in agreement with at least two other studies in pediatric community samples, and with studies in adults with ADHD [7]. Erhart et al. discuss several explanations for their findings, such as increased impulsivity or delay aversion which may lead to uncontrolled eating behavior or even binge eating. Another possibility is that ADHD is related to increased risk for food addiction, next to addiction risk to various substances, through shared abnormalities in the dopaminergic and other biochemical systems of reward in the ventral striatum [8, 9]. We know for a longtime that acute intake of sweets and sugar is not a significant cause of symptoms of ADHD or cognitive dysfunctions in either low risk or high risk children [10, 11]. However, it could well be that increased long-term intake of high caloric foods sensitizes the dopaminergic systems in the ventral striatum and leads to food addiction. Obesity is a major factor in the development of numerous medical problems. Therefore, we should develop a more active stance toward identifying overweight in children with ADHD and integrate managing psychiatric symptoms and overweight and other aspects of somatic health in our care as usual.

Ekinici et al. did examine the phenomenon of autistic regression in a sample of children with autism spectrum disorders. Children with regression had much more often sleeping problems than those without regression, and those with regression type-2 (i.e., with regression of language and social behavior after already initially abnormal development) had higher rates of gastrointestinal complaints. They wisely refrain from speculating about the potential mechanisms underlying these links between psychiatric and gastrointestinal problems. The history about the relationship between autism spectrum disorders and gastrointestinal disease has been complicated by the

Wakefield paper in Lancet [12]. This paper mixed up the issue with unsupported claims about a connection to mumps, measles and rubella (MMR) vaccination and had to be withdrawn after several years. Recent studies have confirmed the association between autism and gastrointestinal complaints [13], and a recent review offers several interesting hypotheses about this link, such as abnormal composition of gut microbiotica, abnormal cellular immunity, and increased food allergy [14].

It is now time to complement the traditional methods to study comorbidity as cross-sectional and longitudinal epidemiologic designs [15] and model-fitting approaches in genetically sensitive designs [16] with more rigorous biological system approaches [17]. These require to deconstruct existing psychiatric and somatic disease categories and start with bottom-up measures of key neural and body systems. These may include tests of cognitive control, reward sensitivity, and empathy, but also measure the status of the autonomous and endocrine stress systems, cellular and humoral immunology, cardiovascular reactivity, mitochondrial energy metabolism, and components of the gut–brain axis. These systems' functions should be studied at several different levels of analysis, i.e., genes, proteins, physiological activity, brain imaging, and reports of symptoms, to define the key constructs. Boosted by the results of cross-disorder genome-wide association studies, this may lead to organize psychiatric and somatic symptoms into new classes with more homogeneous pathophysiology and etiology. This would provide a more solid basis to develop new diagnostic and therapeutic approaches of these comorbid and complex problems.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

## References

1. Angold A, Costello EJ, Erkanli A (1999) Comorbidity. *J Child Psychol Psychiatry* 40:57–88
2. Neale MC, Kendler KS (1995) Models of comorbidity for multifactorial disorders. *Am J Hum Genet* 57(4):935–953
3. Klein DN, Riso LP (1993) Psychiatric disorders: problems of boundaries and comorbidity. In: Costello CG (ed) *Basic issues in psychopathology*. New York, Guilford Press, pp 19–66
4. Caron Ch, Rutter M (1991) Comorbidity in child psychopathology: concepts, issues and research strategies. *J Child Psychol Psychiatry* 32:1063–1080
5. Neeleman J, Sytama S, Wadsworth M (2002) Propensity to psychiatric and somatic ill-health: evidence from a birth cohort. *Psychol Med* 32(5):793–803
6. Megna JL, Schwartz TL, Siddiqui UA, Herrera RM (2011) Obesity in adults with serious and persistent mental illness: a review of postulated mechanisms and current interventions. *Ann Clin Psychiatry* 23(2):131–140

7. Pagoto SL, Curtin C, Lemon SC, Bandini LG, Schneider KL, Bodenlos JS, Ma Y (2009) Association between adult attention deficit/hyperactivity disorder and obesity in the US population. *Obesity (Silver Spring)* 17(3):539–544
8. Corwin RL, Avena NM, Boggiano MM (2011) Feeding and reward: perspectives from three rat models of binge eating. *Physiol Behav* 104(1):87–97
9. Davis JF, Tracy AL, Schurdak JD, Tschop MH, Lipton JW, Clegg DJ, Benoit SC (2008) Exposure to elevated levels of dietary fat attenuates psychostimulant reward and mesolimbic dopamine turnover in the rat. *Behav Neurosci* 122(6):1257–1263
10. Wolraich ML, Wilson DB, White JW (1995) The effect of sugar on behavior or cognition in children. A meta-analysis. *JAMA* 274(20):1617–1621
11. Wolraich ML, Lindgren SD, Stumbo PJ, Stegink LD, Appelbaum MI, Kiritsy MC (1994) Effects of diets high in sucrose or aspartame on the behavior and cognitive performance of children. *N Engl J Med* 330(5):301–307
12. Wakefield AJ, Murch SH, Anthony A, Linnell J, Casson DM, Malik M, Berelowitz M, Dhillon AP, Thomson MA, Harvey P, Valentine A, Davies SE, Walker Smith JA (1998) Ileal-lymphoid-nodular hyperplasia, non-specific colitis, and pervasive developmental disorder in children [see comments]. *Lancet* 351(9103):637–641
13. Adams JB, Audhya T, Donough-Means S, Rubin RA, Quig D, Geis E, Gehn E, Loresto M, Mitchell J, Atwood S, Barnhouse S, Lee W (2011) Nutritional and metabolic status of children with autism vs. neurotypical children, and the association with autism severity. *Nutr Metab (Lond)* 8(1):34
14. de Theije CG, Wu J, da Silva SL, Kamphuis PJ, Garssen J, Korte SM, Kraneveld AD (2011) Pathways underlying the gut-to-brain connection in autism spectrum disorders as future targets for disease management. *Eur J Pharmacol* 668(Suppl 1):S70–S80
15. Fergusson DM, Boden JM, Horwood LJ (2011) Structural models of the comorbidity of internalizing disorders and substance use disorders in a longitudinal birth cohort. *Soc Psychiatry Psychiatr Epidemiol* 46(10):933–942
16. Dellava JE, Kendler KS, Neale MC (2011) Generalized anxiety disorder and anorexia nervosa: evidence of shared genetic variation. *Depress Anxiety* 28(8):728–733
17. Mardinoglu A, Nielsen J (2011) Systems medicine and metabolic modelling. *J Intern Med* 10:2796