

# Constraining proton lifetime in SO(10) with stabilized doublet-triplet splitting

K.S. Babu,<sup>a</sup> Jogesh C. Pati<sup>b</sup> and Zurab Tavartkiladze<sup>a,c</sup>

<sup>a</sup>*Department of Physics, Oklahoma State University,  
Stillwater, OK 74078, U.S.A.*

<sup>b</sup>*SLAC National Accelerator Laboratory,  
2575 Sand Hill Road, Menlo Park, CA 94025, U.S.A.*

<sup>c</sup>*E. Andronikashvili Institute of Physics,  
Tamarashvili 6, Tbilisi 01777, Georgia*

*E-mail:* [babu@okstate.edu](mailto:babu@okstate.edu), [pati@slac.stanford.edu](mailto:pati@slac.stanford.edu),  
[zurab.tavartkiladze@gmail.com](mailto:zurab.tavartkiladze@gmail.com)

**ABSTRACT:** We present a class of realistic unified models based on supersymmetric SO(10) wherein issues related to natural doublet-triplet (DT) splitting are fully resolved. Using a minimal set of low dimensional Higgs fields which includes a single adjoint, we show that the Dimopoulos-Wilzcek mechanism for DT splitting can be made stable in the presence of all higher order operators without having pseudo-Goldstone bosons and flat directions. The  $\mu$  term of order TeV is found to be naturally induced. A  $Z_2$ -assisted anomalous  $\mathcal{U}(1)_A$  gauge symmetry plays a crucial role in achieving these results. The threshold corrections to  $\alpha_3(M_Z)$ , somewhat surprisingly, are found to be controlled by only a few effective parameters. This leads to a very predictive scenario for proton decay. As a novel feature, we find an interesting correlation between the  $d = 6$  ( $p \rightarrow e^+\pi^0$ ) and  $d = 5$  ( $p \rightarrow \bar{\nu}K^+$ ) decay amplitudes which allows us to derive a constrained upper limit on the inverse rate of the  $e^+\pi^0$  mode. Our results show that both modes should be observed with an improvement in the current sensitivity by about a factor of five to ten.

**KEYWORDS:** Supersymmetric gauge theory, GUT

**ARXIV EPRINT:** [1003.2625](https://arxiv.org/abs/1003.2625)

---

## Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>Stabilizing doublet-triplet splitting</b>	<b>2</b>
<b>3</b>	<b>Novel correlation between <math>d = 5</math> and <math>d = 6</math> proton decays</b>	<b>7</b>
<b>4</b>	<b>Nucleon decay</b>	<b>10</b>
4.1	Fermion masses and $p \rightarrow \bar{\nu}K^+$ decay rate	11

---

## 1 Introduction

Although yet to be seen, proton decay is an indispensable tool to probe nature at truly high energies ( $\sim 10^{16}$  GeV). It still remains as the missing piece of grand unification [1–4]. In the light of a new set of planned detectors including those at the forthcoming deep underground laboratory DUSEL and the HyperKamiokande, we propose to address in this paper certain well known but partially unresolved theoretical issues of supersymmetric (SUSY) grand unification (GUT) which are especially relevant to proton decay.

Strong empirical support for grand unification arises not only from the observed quantum numbers of quarks and leptons and the quantization of electric charge, but in particular from the meeting of the three gauge couplings at a scale  $M_{\text{GUT}} \approx 2 \cdot 10^{16}$  GeV that occurs in the context of low energy SUSY [5–9], and the tiny neutrino masses, as observed in neutrino oscillation experiments. The latter fit extremely well with GUT symmetries that include the symmetry  $SU(2)_L \times SU(2)_R \times SU(4)_C$  [1, 2], the minimal such symmetry being  $SO(10)$  [10, 11]. We will therefore discuss proton decay in the context of supersymmetric  $SO(10)$ . The purpose of the present paper is to pay special attention to the problem of the so-called doublet-triplet (DT) splitting and to study the implications of its resolution for proton decay.

The DT splitting problem is common to all grand unified theories based on simple gauge groups. In SUSY  $SO(10)$  models the two Higgs doublets of MSSM, a color triplet and an anti-triplet lie (typically) in a 10-dimensional representation  $H(10)$ . The color triplets need to be superheavy so as to avoid rapid proton decay and also to preserve gauge coupling unification. Keeping the doublets light and the triplets superheavy self consistently is the doublet-triplet splitting problem.

A natural solution to this problem, avoiding severe fine-tuning is realized in SUSY  $SO(10)$  by the so called Dimopoulos-Wilczek (or the missing VEV) mechanism [12, 13]. It involves a coupling of two 10-plets of the form  $H(10)A(45)H'(10)$  with the adjoint  $A(45)$  having a GUT scale VEV in the  $(B - L)$ -preserving direction:

$$\langle A \rangle = i\sigma_2 \otimes \text{Diag}(a, a, a, 0, 0) . \tag{1.1}$$

This structure contributes to the triplet and not to the doublet masses, and thereby can lead to natural DT splitting without fine-tuning.

Given the very large hierarchy between the doublet and triplet masses, however, one must ensure: (i) that the missing VEV pattern for  $A(45)$  in eq. (1.1) is stable to a high enough accuracy in the presence of all allowed higher dimensional operators; (ii) that there are no undesirable pseudo-Goldstone bosons; and (iii) that there are no flat directions which would lead to VEVs of fields undetermined. Furthermore, (iv) one must also examine, by including all GUT-scale threshold corrections to the gauge couplings, the implication of the doublet-triplet splitting on coupling unification and on proton decay. To our knowledge, while some of these issues have been partially addressed in the literature (e.g. see [14–19]), and major progress was made in ref. [20] with regard to the issues (i) and (ii), *simultaneous resolution* of all four issues has so far remained a challenge.

In this paper we present a predictive class of  $SO(10)$  models, based on a minimal Higgs system, in which all the issues of DT splitting mentioned above are resolved, and where the threshold corrections to the gauge couplings and their implications for proton decay are properly studied as well. The Higgs sector we consider has a single adjoint, along with vectors and spinors. Such a low dimensional Higgs system would lead to smaller threshold effects [8,10], unlike in models [14, 15, 17–19] which employ multiple adjoints and/or 54 dimensional Higgs.<sup>1</sup> A postulated  $Z_2$ -assisted anomalous  $U(1)_A$  symmetry (which may have a string origin [31–33]) plays a crucial role in obtaining our results. We find somewhat surprisingly that the GUT scale threshold corrections to  $\alpha_3(M_Z)$  are determined in terms of a very few parameters. This makes the model rather predictive for proton decay. As a novel feature, we find an intriguing *correlation* between the  $d = 6$  and  $d = 5$  proton decays, which respectively lead to  $p \rightarrow e^+\pi^0$  and  $p \rightarrow \bar{\nu}K^+$  as the dominant decay modes. The correlation is such that the empirical lower limit on  $\Gamma^{-1}(p \rightarrow \bar{\nu}K^+)$  provides a *constrained upper limit* on  $\Gamma^{-1}(p \rightarrow e^+\pi^0)$ . Our results show that both decay modes should in fact be discovered with an improvement in the current limits on lifetimes by about a factor of five to ten.

## 2 Stabilizing doublet-triplet splitting

In order to break SUSY  $SO(10)$  to the supersymmetric standard model with a stabilized DT sector, and for the subsequent breaking of the electro-weak symmetry, we shall use a minimal low dimensional Higgs system. It consists of a single adjoint  $A(45)$ , two pairs of spinor-antispinor superfields  $\{C(16) + \bar{C}(\overline{16})\}$  and  $\{C'(16) + \bar{C}'(\overline{16})\}$ , two 10-plets  $H(10)$  and  $H'(10)$ , as well as two  $SO(10)$  singlets  $S$  and  $Z$ . The second spinorial pair  $C' + \bar{C}'$  is introduced, following ref. [20], to avoid pseudo-Goldstone degrees of freedom while maintaining the Dimopoulos-Wilczek VEV structure for  $A$  (cf: eq. (1.1)). The  $S$  and

---

<sup>1</sup>An alternative class of  $SO(10)$  models utilizing larger dimensional (e.g. 126) Higgs fields has been studied in ref. [21–29]. These models have the interesting feature that  $R$ -parity is automatic, being part of the gauge symmetry. However, threshold corrections are rather large in these models, making quantitative predictions for  $\alpha_3(m_Z)$  and proton decay difficult (see attempts in this regard by Aulakh and Garg [30]).

	$A(45)$	$H(10)$	$H'(10)$	$C(16)$	$\bar{C}(\overline{16})$	$Z$	$S$	$C'(16)$	$\bar{C}'(\overline{16})$	$16_{1,2}$	$16_3$
$Q$	0	1	-1	$\frac{k+4}{2k}$	$-\frac{1}{2}$	$\frac{2}{k}$	$\frac{2}{k}$	$\frac{k-4}{2k}$	$-\frac{k+8}{2k}$	$q_{1,2}$	$-\frac{1}{2}$
$\omega$	1	0	1	0	0	1	0	0	0	$P_{1,2}$	0

**Table 1.**  $\mathcal{U}(1)_A$  and  $Z_2$  charges  $Q_i$  and  $\omega_i$  of the superfield  $\phi_i$ . The transformations under  $\mathcal{U}(1)_A$  and  $Z_2$  are respectively  $\phi_i \rightarrow e^{iQ_i}\phi_i$  and  $\phi_i \rightarrow e^{i\frac{2\pi}{2}\omega_i}\phi_i$ .

$Z$  superfields are needed to fix various VEVs in the required directions through their superpotential couplings.

We supplement the gauge symmetry by a  $Z_2$ -assisted anomalous  $\mathcal{U}(1)_A$  symmetry in order to stabilize the VEV pattern of eq. (1.1) [17–19, 34, 35]. The charges of the Higgs fields and those of the three matter families  $16_i$  under  $\mathcal{U}(1)_A \times Z_2$  are listed in table 1. Here  $k$  is a positive integer which is unspecified for the moment. The superpotential of the symmetry breaking sector, consistent with these symmetries, is  $W = W_1 + W_2 + W_3$ , where

$$W_1 = M_A \text{tr} A^2 + \frac{\lambda_A}{M_*} (\text{tr} A^2)^2 + \frac{\lambda'_A}{M_*} \text{tr} A^4, \quad (2.1)$$

$$W_2 = C \left( \frac{a_1}{M_*} ZA + \frac{b_1}{M_*} C\bar{C} + c_1 S \right) \bar{C}' + C' \left( \frac{a_2}{M_*} ZA + \frac{b_2}{M_*} C\bar{C} + c_2 S \right) \bar{C}, \quad (2.2)$$

$$W_3 = \lambda_1 HAH' + \left( \lambda_{H'} SZ^{k-1} + \lambda'_{H'} Z^k \right) \frac{(H')^2}{M_*^{k-1}} + \lambda_2 H\bar{C}\bar{C} + \frac{\lambda_3}{M_*} AH'CC'. \quad (2.3)$$

For simplicity we assume that the  $\text{SO}(10)$  contractions in the  $C\bar{C}$  terms with coefficients  $b_{1,2}$  in eq. (2.2) are in the singlet channel. In the second term of eq. (2.3) the operator  $Z^k$  can appear only when  $k$  is even. While our consideration of DT splitting will hold for all  $k$ , if  $k$  is odd, matter parity is automatic, being part of  $\mathcal{U}(1)_A$ . The choice of  $k = 5$ , which we will use, is phenomenologically preferred, in particular for suppressing adequately all  $d = 5$  proton decay operators including those induced by Planck scale physics. Higher order operators such as  $A^6/M_*^3$  etc. are not exhibited in eqs. (2.1)–(2.3) because they are inconsequential for our purposes. The charges  $q_{1,2}$  and the parities  $P_{1,2}$  of the first two families are left unspecified for the present. They will however be relevant for the generation of quark and lepton masses.

Typically, we expect that the non-renormalizable operators such as  $\lambda_A$  and  $\lambda'_A$ -terms would be induced by quantum gravity effects involving exchange of heavy states in the string tower. Thus, we expect the cut-off scale  $M_* \sim M_{\text{Pl}}$  or  $M_{\text{String}} \sim 10^{18}$  GeV. We shall take all dimensionless couplings to be of order unity, i.e., in the range  $(1/4 - 2)$ .

Using the SUSY preserving condition  $F_Z = F_S = 0$ , together with the choice  $\langle C \rangle = \langle \bar{C} \rangle = c, \langle A \rangle \neq 0$  (which is one allowed option among the discrete set of degenerate vacuum solutions), we get  $\langle C\bar{C}' \rangle = \langle \bar{C}C' \rangle = 0$  and  $\langle C' \rangle = \langle \bar{C}' \rangle = 0$ . The VEV of  $A$  is then determined entirely by  $W_1$  of eq. (2.1). Setting  $F_A = 0$ , we find a solution in the  $B - L$  direction as in eq. (1.1), with

$$a^2 = \frac{M_A M_*}{2(6\lambda_A + \lambda'_A)}. \quad (2.4)$$

With  $\lambda_A, \lambda'_A \sim 1$  and  $M_* \sim 10^{18}$  GeV, we need to choose  $M_A \sim 10^{15}$  GeV to obtain  $a \sim M_{\text{GUT}} \approx 2 \cdot 10^{16}$  GeV. Demanding  $F$ -flatness conditions  $F_{C'} = F_{\bar{C}'} = 0$  and using the notations  $z = \langle Z \rangle$  and  $s = \langle S \rangle$  we get  $s = \frac{c^2}{M_*} \rho_1$  and  $z = \frac{c^2}{3a} \rho_2$ , where  $\rho_1 = \frac{b_1 a_2 - b_2 a_1}{a_1 c_2 - a_2 c_1}$ ,  $\rho_2 = \frac{b_1 c_2 - b_2 c_1}{a_1 c_2 - a_2 c_1}$ . We note that for all dimensionless couplings in the Lagrangian being in the range  $(1/4 - 2)$ , the effective couplings  $\rho_{1,2}$  can naturally take values as small as about  $1/50$ .

The sum of the VEVs get further constrained as follows. The anomalous  $\mathcal{U}(1)_A$  symmetry, presumed to have a string origin, generates the Fayet-Iliopoulos term  $\xi$  through quantum gravity, which is given by [31–33]  $\xi = \frac{g_{\text{st}}^2 M_{\text{Pl}}^2}{192\pi^2} \text{Tr} Q_A$ , where  $g_{\text{st}}$  denotes the string coupling and  $M_{\text{Pl}} \simeq 2.4 \cdot 10^{18}$  GeV is the reduced Planck mass. In our model, the particle spectrum of table 1 would lead to  $\text{Tr}(Q_A) = -8 - 60/k + 16(q_1 + q_2) = -84/5$ , (for  $k = 5$ ,  $q_{1,2} = -1/2 + 3/k$ , see later). This value will however be modified if there are additional singlets in the full theory. (Semi-realistic string solutions [36–38], possessing an anomalous  $\mathcal{U}(1)_A$ , typically lead to  $|\text{Tr} Q_A| \approx 30 - 100$ .) With the charges in table 1, the vanishing of  $D_A = \xi + \sum_i Q_i |\langle \phi_i \rangle|^2 = 0$  (required for preserving SUSY), yields  $c^2 + |z|^2 + |s|^2 = -\frac{k}{2} \xi$ . Thus, the VEVs of all the fields get determined. We see that quite naturally, the VEVs  $c, z \sim (\text{few} - 10) \times M_{\text{GUT}}$ , and  $s \sim (10^{-2} - 10^{-1}) \times M_{\text{GUT}}$  can arise, with the precise values depending on the order one couplings. Let us note that this setup also allows for additional singlet fields  $\{P_i\}$  which can play a role in the  $D_A = 0$  condition (for  $P_i$  with positive  $\mathcal{U}(1)_A$  charges) and can modify these estimates somewhat, without upsetting the stability of DT splitting.

Substituting the VEVs of the heavy fields in eqs. (2.2) and (2.3), we derive the mass matrices  $M_D$  and  $M_T$  for the  $\text{SU}(2)_L$  doublets and  $\text{SU}(3)_c$ -color triplets (written in the  $\text{SU}(5)$  notation):

$$M_{D,T} = \begin{matrix} & \mathbf{5}_H & \mathbf{5}_{H'} & \mathbf{5}_{\bar{C}} & \mathbf{5}_{\bar{C}'} \\ \begin{matrix} \bar{\mathbf{5}}_H \\ \bar{\mathbf{5}}_{H'} \\ \bar{\mathbf{5}}_{\bar{C}} \\ \bar{\mathbf{5}}_{\bar{C}'} \end{matrix} & \begin{pmatrix} 0 & \eta_{D,T} \lambda_1 a & \lambda_2 c & 0 \\ -\eta_{D,T} \lambda_1 a & M_{H'} & 0 & 0 \\ 0 & 0 & 0 & \kappa_{D,T} Y_1 \\ 0 & Y_{D,T} & \kappa_{D,T} Y_2 & M_{C'} \end{pmatrix} \end{matrix}, \quad (2.5)$$

with  $(\eta_D, \eta_T) = (0, 1)$ ,  $(\kappa_D, \kappa_T) = (3, 2)$ . Here  $M_{H'} = (\lambda_{H'} S Z^{k-1} + \lambda'_{H'} Z^k) / M_*^{k-1}$  (see eq. (2.3)),  $Y_{1,2} = 2a_{1,2} z a / (M_*)$  and  $Y_{D,T} \sim \lambda_3 \langle A \rangle c / (M_*)$ . For  $k = 5$  the dominant contribution to  $M_{H'}$  comes from the operator  $\lambda_{H'} s z^4 / M_*^4$ , which is in the range  $(10^{11} - 10^{12})$  GeV. The suppressed mass of  $H'$  will be crucial for an adequate suppression of  $d = 5$  proton decay. The entry  $M_{C'}$  in eq. (2.5) (allowed by the stability of Higgs doublet mass) would arise from the operator  $S Z^2 C' \bar{C}' / M^2$  and yields  $M_{C'} \sim (10^{-2} \text{ to } 10^{-1}) \times M_{\text{GUT}}$  if  $M \sim z$ , which happens if the superfields that are integrated out have GUT scale masses.

The zeros in the first column of eq. (2.5) are ensured, in the presence of *all higher dimensional operators*, for the doublet mass matrix by the  $\mathcal{U}(1)_A \times Z_2$  symmetry. The main reason for this all-order stability of the Higgs doublet masses is that all the effective Higgs fields (i.e. any positive power of  $Z, S$  and  $\bar{C}C$ ) which have super-large VEVs are positively charged under  $\mathcal{U}(1)_A$ , and can not couple to  $H^2$ , which is also positively charged. Thus, with  $\eta_D = 0$  one pair of the Higgs doublets will be massless, while the remaining

three pairs of doublets become superheavy. The role of the  $Z_2$  symmetry is that it allows the coupling of  $H$  to  $H'$  only through  $A$  (or odd powers of  $A$ ). Such couplings, however, do not generate a doublet mass due to the VEV structure in eq. (1.1) of  $\langle A \rangle$ . The VEV pattern of  $\langle A \rangle$  along the  $B - L$  direction is also guaranteed to be stable because of the  $U(1)_A$  symmetry. Indeed, note that the symmetry  $U(1)_A$  does not allow any superpotential coupling involving  $A, C$  and  $\bar{C}$  of the form  $A^n(C\bar{C})^m$ . It is only these couplings which, if allowed, would have upset the missing VEV pattern of eq. (1.1). Their absence to all orders thus guarantees that the pattern of eq. (1.1) is absolutely stable (barring of course SUSY breaking at the TeV scale which is safe). As far as the color-triplets are concerned, since  $\eta_T \neq 0$  in eq. (2.5) for the triplets, all four pairs become super-heavy, just as desired.

The two massless Higgs doublets which emerge from eq. (2.5) represent the MSSM doublets  $h_u$  and  $h_d$  which acquire light masses after SUSY breaking. Let us denote the down type doublets in  $H, H', C$  and  $C'$  by  $H_d, H_d', C_d$  and  $C_d'$  respectively, and likewise the up-type doublets. It is easy to see from eq. (2.5) that  $h_u$  is composed entirely of  $H$  - i.e.  $h_u = H_u$ , while  $h_d$  is a mixture of four components  $H_d, H_d', C_d$  and  $C_d'$ . In particular, the weights of  $h_d$  in  $H, H', C$  and  $C'$  are given by  $H \supset \cos \gamma \cdot h_d$ ,  $H' \supset \frac{\lambda_2 c Y_D}{3 Y_2 M_{H'}} \cos \gamma \cdot h_d$ ,  $C \supset \frac{\lambda_2 c M_{C'}}{9 Y_1 Y_2} \cos \gamma \cdot h_d$  and  $C' \supset \frac{\lambda_2 c}{3 Y_2} \cos \gamma \cdot h_d$ . The angle  $\gamma$  is determined in terms of the parameters of the superpotential. It is related to the MSSM parameter  $\tan \beta$  as  $\tan \beta = \frac{m_t}{m_b} \cos \gamma$ . Note that, unlike in many SO(10) models, the MSSM parameter  $\tan \beta$  is not required to be large here. It would turn out that conservative upper limits on proton lifetime would correspond to smaller values of  $\tan \beta$ .

The  $\mu$ -term, the coefficient of  $h_u h_d$  term of MSSM superpotential, is generated within our model in a simple way. In the unbroken SUSY limit,  $\mu$ -term is zero since terms such as  $H^2$  are forbidden. After SUSY breaking, the adjoint  $A(45)$  develops a VEV  $\sim m_{susy}$  along its  $I_{3R}$  direction, correcting the zeros of eq. (1.1), which generates the  $\mu$ -term. This occurs since the inclusion of the soft SUSY breaking terms induces VEVs  $\sim m_{susy}$  for the fields  $C'$  and  $\bar{C}'$  along their  $\nu^c$ -like scalar components. These will trigger the VEV ( $\sim m_{susy}$ ) of  $A(45)$  in the  $I_{3R}$  direction. From eq. (2.3) we obtain  $\lambda_1 H A H' \rightarrow m_{susy} h_u h_d$ , and thus  $\mu \sim m_{susy} \sim \text{TeV}$ , independent of the integer  $k$ . Thus the present setup provides a simple and elegant solution to the  $\mu$  problem without any new ingredients.

Using eq. (2.5), for the four heavy triplets  $T_i$  and three heavy doublets  $D_i$  (coming from four pairs of  $(5 + \bar{5})$ 's of SU(5) in  $H, H', C, C', \bar{C}, \bar{C}'$ ) we derive the following mass relations:

$$\frac{M_{D_1} M_{D_2} M_{D_3}}{M_{T_1} M_{T_2} M_{T_3} M_{T_4}} = \frac{9}{4 M_{\text{eff}} \cos \gamma}, \quad \frac{1}{M_{\text{eff}}} = (M_T^{-1})_{11} = \frac{M_{H'}}{\lambda_1^2 a^2}. \quad (2.6)$$

In our model, for  $k = 5$  we have  $M_{\text{eff}} = \frac{\lambda_1^2 a^2}{\lambda_{H' s z^4}} M_*^4$ . Putting  $a^2 = M_X^2 / g^2$  with  $g^2 \approx g_{\text{GUT}}^2 \approx 0.63(1 \pm 0.10)$ ,  $M_X \approx (0.6 - 1) \times 10^{16} \text{ GeV}$  (see discussion after eq. (4.1)), and taking an explicit solution<sup>2</sup> for the VEVs,  $z \sim 0.17 M_*$  and  $s \sim M_*/70$ , together with natural values

---

<sup>2</sup>Details of estimating the VEVs based on explicit solutions to the  $D_A = 0$  condition will be presented in a forthcoming longer paper [39].

of the couplings  $\lambda_1 \approx (1/4 - \sqrt{2})$  and  $\lambda_{H'} \approx (1/4 - 2)$ , we estimate

$$M_{\text{eff}} \sim (5 \times 10^{16} - 6 \times 10^{19}) \text{ GeV}. \quad (2.7)$$

The mass scale  $M_{\text{eff}}$  will control the  $d = 5$  proton decay amplitude (see e.g. ref. [16]). It would also enter the threshold corrections. Note that  $M_{\text{eff}}$ , which does not represent the physical mass of any particle, can naturally exceed even  $10^{19}$  GeV.

Now, the multiplets  $A, C, C', \bar{C}$  and  $\bar{C}'$  contain three pairs of  $(10 + \bar{10})$ 's of SU(5), which get masses through eqs. (2.1) and (2.2). Their mass matrix is given by:


$$M(\Psi^{10}) = \begin{matrix} \bar{\Psi}_A^{10} & \bar{\Psi}_C^{10} & \bar{\Psi}_{C'}^{10} \\ \Psi_A^{10} & \Psi_C^{10} & \Psi_{C'}^{10} \end{matrix} \begin{pmatrix} M_\Psi & 0 & X_1 \\ 0 & 0 & \kappa_\Psi Y_1 \\ X_2 & \kappa_\Psi Y_2 & M_{C'} \end{pmatrix}, \quad (2.8)$$

with  $\Psi = (u^c, q, e^c)$ ,  $\kappa_\Psi = (2, 1, 0)$ ,  $M_\Psi = (0, 0, M_\Sigma/2)$ , where  $X_{1,2} = 4a_{1,2}zc/M_*$  and  $Y_{1,2}$  are defined after eq. (2.5).  $M_\Sigma = 2\lambda'_A M_A / (6\lambda_A + \lambda'_A)$  yields the mass of the color octet and SU(2)<sub>L</sub> triplet in A(45):  $M_\Sigma \equiv M_8 = 2M_3$ . We see from eq. (2.8) that two pairs of  $(u^c, q, e^c)$ -like states are massive. The third massless pair  $(10 + \bar{10}$  of SU(5)) is eaten by the corresponding massive gauge superfields of SO(10)/SU(5). Denoting the masses of the components of  $(10 + \bar{10})$ 's by curly symbols (e.g.  $\mathcal{U}_1^c \equiv M(u_1^c)$  etc.), we derive from eq. (2.8):  $\mathcal{U}_1^c \mathcal{U}_2^c = Y_1 Y_2 (4 + p^2)$ ,  $\mathcal{Q}_1 \mathcal{Q}_2 = Y_1 Y_2 (1 + p^2)$ ,  $\mathcal{E}_1^c \mathcal{E}_2^c = Y_1 Y_2 \hat{p}^2$  where  $p^2 = \frac{4c^2}{a^2} = \frac{|X_1|^2}{|Y_1|^2} = \frac{|X_2|^2}{|Y_2|^2}$ ,  $\hat{p}^2 = p^2 \left| 1 - \frac{M_\Sigma M_{C'}}{2X_1 X_2} \right|$ . These expressions will be useful for the computation of threshold corrections in the model.

The masses of the heavy gauge boson superfields corresponding to the broken generators of SO(10) are given by (see e.g., the second paper in ref. [14, 15] and [16]):  $M^2(X, Y) = g^2 a^2 \equiv M_X^2$ ,  $M^2(X', Y') = M_X^2 (1 + p^2)$ ,  $M^2(V_{u^c, \bar{u}^c}) = M_X^2 (4 + p^2)$ ,  $M^2(V_{e^c, \bar{e}^c}) = M_X^2 p^2$  where  $g$  is the unified gauge coupling at the GUT scale. Given the  $p$  and  $\hat{p}$  dependence of the masses given above, we see that, except for the  $e^c$ -like states, threshold corrections from all other states in the  $(10 + \bar{10})$  matter sector cancel precisely against those from the corresponding states in the gauge sector. An accidental  $N = 4$  supersymmetry present in the model (the gauge bosons and three pairs of matter fields in the  $(10 + \bar{10})$  sector form an  $N = 4$  SUSY gauge multiplet) is responsible for this cancelation. This results in an enormous reduction of the parameters, rendering the model very predictive for proton decay.

We have presented the whole spectrum of the theory, except for the singlet sector, which is not relevant for the calculation of threshold corrections. We have however, analyzed the singlet sector and verified that there are no unwanted pseudo-Goldstone states in the model. While it might appear that there is a U(1) symmetry associated with the ‘‘integer’’  $k$  in table 1, it turns out that this is a linear combination of  $\mathcal{U}(1)_A$  and  $B - L$ , and its breaking does not lead to a pseudo-Goldstone boson.

The evolution of the three gauge couplings in the model with momentum is shown in figure 1, which takes into account all the threshold effects. It is clear from figure 1 that the three couplings merge into one at a unification scale  $M_{\text{GUT}} \sim 10^{16}$  GeV. Furthermore,



**Figure 1.** Evolution of the three standard model gauge couplings in the present SO(10) model including threshold corrections. We have used  $\alpha_3(M_Z) = 0.1176$  and assumed an mSUGRA spectrum with  $\{\tan \beta, m_0, m_{1/2}, \mu\} = \{3, 1448.2 \text{ GeV}, 155.93 \text{ GeV}, 1 \text{ TeV}\}$  (corresponding to  $m_{\tilde{q}} = 1.5 \text{ TeV}$ ,  $m_{\tilde{W}} = 130 \text{ GeV}$ ), and have taken  $p = 4$ ,  $r = 1/250$ ,  $M_{\text{eff}} = 4 \cdot 10^{19} \text{ GeV}$ ,  $Y_{1,2} = 2M_X/45$  for generating this plot.

we see that the unified SO(10) gauge coupling remains perturbative to scales well above  $M_{\text{GUT}}$ . This is a desirable feature which not all SO(10) models have.

### 3 Novel correlation between $d = 5$ and $d = 6$ proton decays

Writing down the three RG equations for  $\alpha_{1,2,3}^{-1}$  and eliminating the unified gauge coupling  $\alpha_G$  we obtain

$$\ln \frac{M_{\text{eff}} \cos \gamma}{M_Z} = \frac{5\pi}{6} \left( 3 \left( \alpha_2^{-1} + \Delta_{2,w}^{(2)} - \frac{1}{6\pi} \right) - 2 \left( \alpha_3^{-1} + \Delta_{3,w}^{(2)} - \frac{1}{4\pi} \right) - \left( \alpha_1^{-1} + \Delta_{1,w}^{(2)} \right) \right) - \ln \frac{4\kappa^{5/2}}{9} + \ln \frac{p}{\hat{p}}, \quad (3.1)$$

$$\ln \frac{(M_X^2 M_\Sigma)^{1/3}}{M_Z} = \frac{\pi}{18} \left( 5 \left( \alpha_1^{-1} + \Delta_{1,w}^{(2)} \right) - 3 \left( \alpha_2^{-1} + \Delta_{2,w}^{(2)} - \frac{1}{6\pi} \right) - 2 \left( \alpha_3^{-1} + \Delta_{3,w}^{(2)} - \frac{1}{4\pi} \right) \right) + \frac{1}{6} \ln \kappa - \frac{1}{3} \ln \frac{p}{\hat{p}}. \quad (3.2)$$

We have taken GUT scale threshold corrections in one loop approximation. The quantities  $\Delta_{i,w}^{(2)}$  include weak scale threshold corrections and 2-loop running effects for the



gauge couplings, including Yukawa interactions. Their values depend on the SUSY particle spectrum. We carry out our analysis within the minimal  $N = 1$  SUGRA scenario [40, 41] with family universal parameters. While we vary these parameters to draw our conclusions, for concreteness, we consider the set of values:  $\{\tan\beta, m_0, m_{1/2}, \mu\} = \{3, 1448.2 \text{ GeV}, 155.93 \text{ GeV}, 1 \text{ TeV}\}$ , which corresponds to  $m_{\tilde{q}} \simeq 1.5 \text{ TeV}$  and  $m_{\tilde{W}} \simeq 130 \text{ GeV}$ . For these values we obtain  $\Delta_{i,w}^{(2)} = (0.6093, 0.4079, 1.167)$ . Using the third RGE equation for the  $\alpha_i^{-1}$  we obtain for the present model  $\alpha_G(M_X) \simeq \alpha_3(M_X) \simeq 1/20$ . The parameter  $\kappa \equiv M_8/M_3 = 2$  in our model (as opposed to  $\kappa = 1$  in SUSY SU(5)).

It is important that the ratio  $r \equiv M_\Sigma/M_X$  entering into eq. (3.2) is constrained by symmetries of the model. Using expressions for  $M_\Sigma$ ,  $M_A$  and  $M_X$  presented earlier, we obtain:


$$r = \frac{M_\Sigma}{M_X} = \frac{4\lambda'_A}{g^2} \frac{M_X}{M_*} \approx \left( \frac{1}{15} - \frac{1}{300} \right). \quad (3.3)$$

The range for  $r$  is obtained by noting that  $\lambda'_A$  is allowed by symmetries of the model and thus naturally expected to be of order one. We have thus taken,  $1/5 \lesssim \lambda'_A \lesssim 2$  (say), and have set  $g^2 \simeq g_{\text{GUT}}^2 \approx 0.63(1 \pm 0.10)$ ,  $M_X \approx (0.6 - 1) \times 10^{16} \text{ GeV}$  (see discussion after eq. (4.1)), while  $M_* \approx M_{\text{Pl}} \simeq 2 \times 10^{18} \text{ GeV}$ . This restriction on  $r$  will be an important ingredient in the derivation of an upper limit on  $\Gamma^{-1}(p \rightarrow e^+\pi^0)$ .

Eliminating  $p/\hat{p}$  from eqs. (3.1) and (3.2) we obtain a correlation between  $M_{\text{eff}}$  and  $M_X$  for a given  $r$ :

$$M_{\text{eff}} \simeq 10^{19} \text{ GeV} \cdot \left( \frac{10^{16} \text{ GeV}}{M_X} \right)^3 \left( \frac{1/100}{r} \right) \left( \frac{3}{\tan\beta} \right) \left( \frac{0.6}{\eta_\gamma} \right) \left\{ \frac{\exp[2\pi(\Delta_{2,w}^{(2)} - \Delta_{3,w}^{(2)} - \delta\alpha_3^{-1})]}{2.54 \cdot 10^{-2}} \right\}, \quad (3.4)$$

where  $\eta_\gamma \simeq 0.6$  accounts for the running of  $\cos\gamma$ , and  $\delta\alpha_3^{-1}$  denotes the deviation of  $\alpha_3^{-1}$  from its central value of  $1/0.1176$ . Note that the curly bracket on the right side of eq. (3.4) is fully determined for any given choice of the SUSY parameters and  $\alpha_3(M_Z)$ . It turns out to be only mildly dependent on variations of  $m_0$  and  $m_{1/2}$ . Since  $M_{\text{eff}}$  and  $M_X$  respectively control  $d = 5$  ( $p \rightarrow \bar{\nu}K^+$ ) and  $d = 6$  ( $p \rightarrow e^+\pi^0$ ) decay amplitudes, eq. (3.4) in turn provides a correlation between the rates of these two otherwise unrelated decay modes. Such a correlation exists in minimal SUSY SU(5) as well [42, 43], but that leads to predictions for  $\alpha_3(M_Z)$  and  $d = 5$  proton decay rate which are inconsistent with experiments [44]. Exceptions to this conclusion has been suggested in ref. [45] which uses higher dimensional operators. This however leads to large threshold corrections, making the apparent unification of gauge couplings with low energy SUSY somewhat coincidental. For review of proton decay in SU(5) and in alternative scenarios see ref. [46].



**Figure 2.** Correlations between  $M_{\text{eff}}$  and  $M_X$  for  $\{\tan\beta, m_0, m_{1/2}, \mu\} = \{3, 1448.2 \text{ GeV}, 155.93 \text{ GeV}, 1 \text{ TeV}\}$  (corresponding to  $m_{\tilde{q}} = 1.5 \text{ TeV}$ ,  $m_{\tilde{W}} = 130 \text{ GeV}$ ), and  $\alpha_3(M_Z) = 0.1176$ . (a):  $r = 1/200$ . (b):  $r = 1/250$ . (c):  $r = 1/300$ . The vertical and horizontal dashed lines correspond to the experimentally allowed lowest values of  $M_X$  and  $M_{\text{eff}}$  which arise from limits on  $\Gamma^{-1}(p \rightarrow e^+\pi^0)$  and  $\Gamma^{-1}(p \rightarrow \bar{\nu}K^+)$  respectively, for central values of relevant parameters (see text).

Now, using expressions for proton decay rates (see below) one finds that the empirical lower limit on  $\Gamma^{-1}(p \rightarrow \bar{\nu}K^+)$  requires that  $M_{\text{eff}} \gtrsim 2.91 \cdot 10^{19} \text{ GeV}$  (for reasonable scenarios for the Yukawa couplings, see discussions in section 4.1), while that on  $\Gamma^{-1}(p \rightarrow e^+\pi^0)$  requires (owing to eq. (3.4))  $r \lesssim 1/150$ . Using the particular choice of SUSY parameters stated above, and the ranges for  $M_{\text{eff}}$  and  $r$  as given in eqs. (2.7) and (3.3), we therefore illustrate our results on the correlation between  $M_{\text{eff}}$  and  $M_X$  in figure 2 by confining to the ranges  $M_{\text{eff}} \simeq (2.91 - 6) \times 10^{19} \text{ GeV}$  and  $r \simeq (1/200 - 1/300)$ .<sup>3</sup>

Two crucial differences between our model and that of minimal SUSY SU(5) (see ref. [42, 43]–[46]) are: (i)  $M_{H_c}$  of SU(5) is replaced by  $M_{\text{eff}} \cos \gamma$  which can be significantly larger than  $M_{\text{GUT}}$  (for  $\tan \beta \geq 3$  say); and (ii) whereas the range for  $r$  is severely restricted by symmetry considerations for the SO(10) model (see eq. (3.3)), this is not so for the minimal SUSY SU(5) model. These two distinctions make the SO(10) model presented here more predictive for proton decay into  $e^+\pi^0$  on the one hand, and viable on the other. In particular, as discussed below, with  $\alpha_3(M_Z)$  being consistent with experiments, the  $d = 5$  proton decay rate is in full accord with the experimental limit.

<sup>3</sup>While such large values of  $M_{\text{eff}}$  lie in their natural ranges, they correspond via eq. (3.1) to rather small values of  $\hat{p}/p \sim 10^{-4}$ . Thus  $\hat{p}/p \sim 10^{-4}$  is the only parameter of the model whose smallness remains unexplained on grounds of naturalness.

## 4 Nucleon decay

There are two main mechanisms for proton decay corresponding to  $d = 5$  and  $d = 6$ , which respectively yield  $p \rightarrow \bar{\nu}K^+$  and  $p \rightarrow e^+\pi^0$  as the dominant decay modes. Although a priori these two modes are largely independent, owing to the correlation given in eq. (3.4) and figure 2, they get linked in our model such that the observed lower limit on the inverse decay rate of either mode implies an *upper limit* on that of the other. The latter is found to be especially constrained for the  $p \rightarrow e^+\pi^0$  mode. The rates for  $d = 6$  decay modes  $p \rightarrow e^+\pi^0$  and  $p \rightarrow \bar{\nu}\pi^+$ , which are largely independent of the details of Yukawa couplings and SUSY spectrum, are given by:

$$\Gamma(p \rightarrow e^+\pi^0) \simeq \frac{m_p}{64\pi f_\pi^2} (1+D+F)^2 \bar{\alpha}_H^2 \left( \frac{g_X^2 A_R}{M_X^2} \right)^2 f(p), \quad \Gamma(p \rightarrow \bar{\nu}\pi^+) \simeq 2\Gamma(p \rightarrow e^+\pi^0) \frac{f(p)-4}{f(p)}, \quad (4.1)$$

where  $f(p) = 4 + (1 + 1/(1+p^2))^2$ . Here  $\alpha_H$  denotes the hadronic matrix element. Recent lattice calculation yields  $\alpha_H \simeq 0.012 \text{ GeV}^3$  at  $\mu = 2 \text{ GeV}$  [47, 48].  $D$  and  $F$  are chiral lagrangian parameters with  $D \simeq 0.8$ ,  $F \simeq 0.47$ .  $g_X$  denotes the effective  $X, Y$  boson coupling at  $M_X$ . The correlation curves (see figure 2 for a representative case) restrict  $M_X$  in the range of about  $(6 - 10) \times 10^{15} \text{ GeV}$ . Taking an average of the three gauge couplings, which nearly unify at  $M_X$ , lying in the range as given above (see figure 1), we obtain  $\alpha_G(M_X) = g_X^2/(4\pi) \simeq (1/20)(1 \pm 0.1)$ , where the error reflects variations in the GUT scale spectrum or equivalently in the parameters of the superpotential lying in a natural range. The function  $f(p)$  varies between the limits 8 and 5 as  $p$  varies from 0 to  $\infty$ ; correspondingly one obtains  $\Gamma(p \rightarrow e^+\pi^0)/\Gamma(p \rightarrow \bar{\nu}\pi^+) \simeq (1, 1.4, 2.5)$  for  $(p \lesssim 1/3, p \approx 1, p \gg 1)$ . The case of  $p \rightarrow \infty$  represents the SU(5) limit. If this branching ratio is measured to be significantly smaller than 2.5, that would be strongly suggestive of SO(10) (as opposed to SU(5)) grand unification. The quantity  $A_R$  in eq. (4.1) denotes the net renormalization of the  $d = 6$  operator, including short ( $A_S^{d=6} \simeq 2.22$ ) and long distance effects ( $R_L \simeq 1.25$ ). In our model  $A_R \simeq 2.78$ . From eq. (4.1) we get:

$$\Gamma_{d=6}^{-1}(p \rightarrow e^+\pi^0) \simeq 1.0 \times 10^{34} \text{ yrs} \left( \frac{0.012 \text{ GeV}^3}{\alpha_H} \right)^2 \left( \frac{2.78}{A_R} \right)^2 \left( \frac{5.12}{f(p)} \right) \left( \frac{1/20}{\alpha_G(M_X)} \right)^2 \left( \frac{M_X}{6.24 \times 10^{15} \text{ GeV}} \right)^4. \quad (4.2)$$

Allowing for uncertainty in  $|\alpha_H|$  by  $\pm 25\%$  (see discussion in ref. [47, 48]) and that in  $\alpha_G(M_X)$  by  $\pm 10\%$ , and letting  $p$  vary in the theoretically favored range of  $p \simeq 1$  to  $p = 10$ , we see that the empirical lower limit on  $\Gamma^{-1}(p \rightarrow e^+\pi^0) \gtrsim 1.01 \times 10^{34} \text{ yrs}$  [49–52] (For latest limits see webpage of SuperK [53]) requires  $M_X \gtrsim (6.26 \cdot 10^{15} \text{ GeV}) \cdot (1 \pm 0.14)$ , where for the central value we have used  $p = 4$ , and the errors are added in quadratures.<sup>4</sup>

Without further theoretical constraint on  $M_X$ , given that  $\Gamma_{d=6}^{-1}(p \rightarrow e^+\pi^0) \propto M_X^4$ , there has been considerable uncertainty in the literature so far on  $\Gamma^{-1}(p \rightarrow e^+\pi^0)$ , which

---

<sup>4</sup>This is only to indicate a reasonable range for  $(M_X)_{\min}$ , which, however, is not used for our explicit predictions.

has been quoted to lie in the range  $\sim 10^{34} - 10^{38}$  yrs [45, 54]. Such a range corresponds to (a priori reasonable) guesses on  $M_X \sim M_{\text{GUT}} \times (1/3 - 3) \sim (0.7 - 6) \times 10^{16}$  GeV, the higher values of  $M_X$  being allowed by letting  $r$  be arbitrarily small ( $\lesssim 10^{-4}$  say). Lifetimes much exceeding  $(\text{few} - 10) \times 10^{35}$  yrs would, however, be inaccessible to next generation proton decay experiments. As discussed below, the correlation eq. (3.4) together with the restriction on  $r$  (see eq. (3.3)) would provide a much stronger constraint on  $\Gamma^{-1}(p \rightarrow e^+ \pi^0)_{\text{max}}$ , which is fully accessible. To obtain an upper limit on  $M_X$  and thus on  $\Gamma^{-1}(p \rightarrow e^+ \pi^0)$  we first need to discuss  $d = 5$  proton decay.

#### 4.1 Fermion masses and $p \rightarrow \bar{\nu} K^+$ decay rate

In order to investigate  $d = 5$  proton decay, the Yukawa sector needs to be specified. We have found a self-consistent picture for fermion masses and mixings in the present setup in the same spirit as in ref. [16, 55]. We introduce a non-Abelian flavor symmetry  $Q_4$  (the quaternionic group) and assign the matter fields  $16_{1,2}$  as a doublet of this group,  $\vec{16} = (16_1, 16_2)$ .  $16_3$  transforms trivially under  $Q_4$ . Two  $Q_4$  doublet flavon fields  $\vec{X}, \vec{Y}$  both with VEVs along  $(1, 0)$  direction are also utilized. The  $Q_4$  symmetry also enables us to successfully address the SUSY FCNC problem [60, 61]. With  $U(1)$  charge assignment  $Q(\vec{16}) = -(1/2 + 3/k)$ ,  $Q(\vec{X}) = 3/k$ ,  $Q(\vec{Y}) = 7/k$ , the operators generating Dirac Yukawa couplings are  $16_3 16_3 H$ ,  $\frac{\vec{X}}{M_*} \vec{16} 16_3 H$ ,  $\frac{S Z^2 A}{M_*^4} \vec{16} \vec{16} H$ ,  $\frac{Z^3 C}{M_*^4} \vec{16} \vec{16} C'$ ,  $\frac{AC\vec{Y}}{M_* \langle Z \rangle^2} (\vec{16} \cdot 16_3 + 16_3 \cdot \vec{16}) C'$ , and  $\frac{AC}{M_* \langle Z \rangle^2} (\vec{X} \vec{16})(\vec{Y} \vec{16}) C'$ . The resulting mass matrices for the quarks and charged leptons at the GUT scale are:

$$M_u = \begin{matrix} & u_1^c & u_2^c & u_3^c \\ \begin{matrix} u_1 \\ u_2 \\ u_3 \end{matrix} & \begin{pmatrix} 0 & \epsilon' & 0 \\ -\epsilon' & 0 & \sigma \\ 0 & \sigma & 1 \end{pmatrix} & m_U^0, \end{matrix}$$

$$M_{d(e)} = \begin{matrix} & d_1^c(e_1) & d_2^c(e_2) & d_3^c(e_3) \\ \begin{matrix} d_1^c(e_1) \\ d_2^c(e_2) \\ d_3^c(e_3) \end{matrix} & \begin{pmatrix} 0 & \kappa_{d(e)} \epsilon' + \eta' & 0 \\ -\kappa_{d(e)} \epsilon' - \eta' & \kappa_{d(e)} \xi_{22}^d & \sigma + \kappa_{d(e)} \epsilon \\ 0 & \sigma + \kappa_{d(e)} \bar{\epsilon} & 1 \end{pmatrix} & m_D^0, \end{matrix} \quad (4.3)$$

where  $m_D^0 = m_U^0 \cos \gamma / \tan \beta$ ,  $\kappa_d = 1$  and  $\kappa_e = 3$ . Eqs. (4.3) provide a constrained system with fewer parameters than observables. A consistent fit for all masses and mixing parameters as well as observed CP violation is obtained with the choice  $\sigma = 0.0508$ ,  $\epsilon = -0.0188 + 0.0333i$ ,  $\bar{\epsilon} = 0.106 + 0.0754i$ ,  $\epsilon' = 1.56 \cdot 10^{-4}$ ,  $\eta' = -0.00474 + 0.00177i$ ,  $\xi_{22}^d = 0.014e^{4.1i}$  at the GUT scale. Following renormalization in going down to low energies (with  $m_t(m_t) = 160$  GeV and  $\tan \beta = 3$ ), these values reproduce the central values of the charged lepton masses. In addition, for the quark masses we obtain

$$\begin{aligned} m_u(2 \text{ GeV}) &= 3.55 \text{ MeV}, & m_c(m_c) &= 1.15 \text{ GeV}, \\ m_d(2 \text{ GeV}) &= 6.45 \text{ MeV}, & m_s(2 \text{ GeV}) &= 137.6 \text{ MeV}, & m_b(m_b) &= 4.67 \text{ GeV}. \end{aligned} \quad (4.4)$$

For the CKM mixings we obtain at  $\mu = M_Z$ ,

$$|V_{us}| = 0.225, \quad |V_{cb}| = 0.0414, \quad |V_{ub}| = 0.0034, \quad |V_{td}| = 0.00878, \quad \bar{\eta} = 0.334, \quad \bar{\rho} = 0.12. \quad (4.5)$$

Thereby we get  $\sin 2\beta = 0.663$ . All these are in a good agreement with experiments.

Let us now briefly discuss the neutrino sector. The operators  $\frac{Z^{k-4}\bar{Y}^2}{M_*^{k-3}M_{N'}^2}\bar{16}^2\bar{C}^2$ ,  $\frac{Z^{k-2}}{M_*^{k-2}M_{N'}^2}\bar{Y} \cdot \bar{16}16_3\bar{C}^2$ , and  $\frac{Z^{k-1}S}{M_*^{k-1}M_N^2}16_3^2\bar{C}^2$  are responsible for Majorana masses of the heavy right-handed neutrinos. We assume that in the first of these couplings the  $Q_4$  contraction  $\bar{Y}^2\bar{16}^2$  is in the  $1'$  channel. The heavy Majorana mass matrix  $M_R$  is given by:

$$M_R = \begin{matrix} & \nu_1^c & \nu_2^c & \nu_3^c \\ \nu_1 & \left( \begin{array}{ccc} b & 0 & 0 \\ 0 & b & a \\ 0 & a & 1 \end{array} \right) & & \\ \nu_2 & & & \\ \nu_3 & & & \end{matrix} M_0, \quad (4.6)$$

The Dirac mass matrix  $M_{\nu D}$  at GUT scale can be obtained from  $M_u$  (see eq. (4.3)) by the replacement  $\epsilon' \rightarrow -3\epsilon'$ . We can take the two dimensionless parameters  $(a, b)$  and the mass parameter  $M_0$  as input to fix  $\sqrt{\Delta m_{\text{atm}}^2}$ ,  $\theta_{12}$  and  $\theta_{23}$ . Two observables, viz.,  $\sqrt{\Delta m_{\text{sol}}^2/\Delta m_{\text{atm}}^2} \simeq m_2/m_3$  and  $\theta_{13}$ , will then be predictions of the model. The structures given in eq. (4.6) are valid at GUT scale. Applying renormalization, including threshold effects due to the different  $\nu^c$  masses, with  $\theta_{12} \simeq 30^\circ$  and  $\theta_{23} \simeq 43^\circ$  as inputs, we obtain  $m_2/m_3 \simeq 0.13$  and  $\theta_{13} \simeq 3.6^\circ$  as predictions. Such a fit is realized by choosing  $a = 0.0252e^{-0.018i}$ ,  $b = 1.61 \cdot 10^{-6}e^{-1.592i}$ , and  $M_0 = 1.89 \cdot 10^{13}$  GeV. The corresponding  $\nu^c$  masses are  $(M_{R1}, M_{R2}, M_{R3}) = (3.04 \times 10^7, 1.2 \times 10^{10}, 1.79 \times 10^{13})$  GeV. These results include all the relevant RG running effects. One sees broad, although not precise, agreement with data. We consider this fit, which provides large neutrino oscillation angles, together with small quark mixing angles as well as observed CP violation as fairly successful and highly nontrivial, especially in a quark-lepton unified framework with a stabilized doublet-triplet splitting.

With the Yukawa couplings specified, the inverse of the sum of partial decay widths, in  $p \rightarrow \bar{\nu}K^+$ , is computed to be:

$$\Gamma_{d=5}^{-1}(p \rightarrow \bar{\nu}K^+) = 3.5 \times 10^{33} \text{ yrs} \left( \frac{0.012 \text{ GeV}^3}{|\beta_H|} \right)^2 \left( \frac{6.91}{\bar{A}_S^\alpha} \right)^2 \left( \frac{1.25}{R_L} \right)^2 \left( \frac{M_{\text{eff}}}{3.38 \times 10^{19} \text{ GeV}} \right)^2 \times \\ \times \left( \frac{m_{\bar{q}}}{1.5 \text{ TeV}} \right)^4 \left( \frac{130}{m_{\bar{W}}} \right)^2 \left( \frac{3.1}{K_{d=5}^\nu} \right). \quad (4.7)$$

Here  $K_{d=5}^\nu$  denotes a sum of contributions to the total decay rate from the three neutrino flavors, reflecting the dependence of the  $d = 5$  operator on the Yukawa couplings:  $K_{d=5}^\nu \equiv |A_{\nu_e}|^2 + |A_{\nu_\mu}|^2 + |A_{\nu_\tau}|^2$ . Each individual  $A_{\nu_i}$  receives contributions from three types of diagrams leading to the  $d = 5$  operator: (a) those with only the first two families in the external legs, (b) those having the quark doublet of the third family in just one external line, and (c) those having the same as in (b) in two external lines. The last two contributions

incorporate the short distance renormalization of the  $d = 5$  operator that arises through the running of the top quark Yukawa coupling, from the GUT scale to the weak scale. Contributions from all three diagrams are found to be important, especially for  $|A_{\nu_\mu}|$  and  $|A_{\nu_\tau}|$ . The net result is that  $|A_{\nu_e}| \sim \mathcal{O}(10^{-1})$ ,  $|A_{\nu_\mu}| \sim |A_{\nu_\tau}| \sim \mathcal{O}(1)$  and  $K_{d=5}^Y \simeq 3.1$  [39].  $\bar{A}_S^\alpha$  in eq. (4.7) denotes the short distance RGE factor for the  $d = 5$  operator, corresponding to the running from the GUT scale to the weak scale, that arises purely from the gauge interactions, without the effects of the top quark Yukawa coupling. Note that  $\bar{A}_S^\alpha$  defined here differs from the RGE factor  $A_S$  defined conventionally [42, 43] in that  $A_S$  includes the effect of the running of  $m_c m_s$  in going from low energies to the GUT scale,  $\bar{A}_S^\alpha$  does not. Thus,  $A_S = \bar{A}_S^\alpha J$ , where  $J = (m_c m_s)_{\text{GUT}} / (m_c m_s)_\mu \sim \mathcal{O}(10^{-1})$  for  $\mu = 2 \text{ GeV}$  (with low  $\tan \beta \sim 3$  to 10).

We can now discuss the derivation of an upper limit on  $M_X$  and thus on  $\Gamma^{-1}(p \rightarrow e^+ \pi^0)$ . Owing to eq. (3.4) this would correspond to the minimum allowed value of  $M_{\text{eff}}$ . Now, taking conservatively  $m_{\bar{q}} \lesssim 1.5 \text{ TeV}$  and the experimental lower limit  $m_{\bar{W}} \gtrsim 125 \text{ GeV}$ , the observed lower limit on  $\Gamma^{-1}(p \rightarrow \bar{\nu} K^+) \gtrsim 2.8 \times 10^{33} \text{ yrs}$  [49–53] yields (via eq. (4.7)):  $(M_{\text{eff}})_{\text{min}} \gtrsim 2.91 \times 10^{19} \text{ GeV}$ . This in turn yields by using eq. (3.4) (with  $|\beta_H| = 0.012 \text{ GeV}^3$ , the lowest value of  $r = 1/300$  and  $\tan \beta = 3$ ):  $(M_X)_{\text{max}} \lesssim (5.16, 7.02, 9.45) \times 10^{15} \text{ GeV}$  for  $\alpha_3(M_Z) = (0.1156, 0.1176, 0.1196)$ . Thus, if we use central values of  $\alpha_3(M_Z), \alpha_H, \beta_H$  and  $\alpha_G(M_X)$  with  $p = 4$ , the upper limit on  $\Gamma^{-1}(p \rightarrow e^+ \pi^0)$  (using eq. (4.1)) would be  $1.61 \times 10^{34} \text{ yrs}$ . If the uncertainties in all these parameters are stretched to their extremes, *each in a direction so as to prolong the lifetime*, the stated upper limit could increase by at most a factor of 10.8. Considering that all the uncertainties having such extreme values, and in the same direction, to be very unlikely, we would regard something like the geometric mean of the two upper limits corresponding to the central and extreme values of the parameters to be a more realistic, yet conservative, upper limit for the lifetime. We thus predict:

$$\Gamma_{d=6}^{-1}(p \rightarrow e^+ \pi^0) \lesssim 5.3 \times 10^{34} \text{ yrs} . \tag{4.8}$$

If  $m_{\bar{q}} < 1.5 \text{ TeV}$ , or  $m_{\bar{W}} > 125 \text{ GeV}$ , or  $r > 1/300$ , or  $\tan \beta > 3$ , the upper limit would of course decrease further significantly.<sup>5</sup> Thus, the upper limit shown above on  $\Gamma^{-1}(p \rightarrow e^+ \pi^0)$ , stemming from eq. (3.4), is a robust and novel feature of the model. The predicted lifetime is accessible to proposed megaton size water Cherenkov (or equivalent) detectors.

Reversing the procedure given above, we can derive an upper limit on  $M_{\text{eff}}$  and thereby on  $\Gamma^{-1}(p \rightarrow \bar{\nu} K^+)$ . Owing to eq. (3.4), this would correspond to the minimum allowed value of  $M_X$  and  $r$ . Using central values of  $|\alpha_H|$  and  $\alpha_G(M_X)$  with  $p \approx 4$  (for concreteness), the observed lower limit on  $\Gamma^{-1}(p \rightarrow e^+ \pi^0) \geq 1.01 \times 10^{34} \text{ yrs}$  [49] yields via eq. (4.2):  $(M_X)_{\text{min}} \geq 6.26 \times 10^{15} \text{ GeV}$ . This in turn yields by using eq. (3.4) (with  $m_{\bar{q}} = 1.5 \text{ TeV}$ ,  $m_{\bar{W}} = 130 \text{ GeV}$  and the lowest values of  $r \approx 1/300$ ):  $(M_{\text{eff}})_{\text{max}} \leq (1.627, 4.105, 10.02) \times 10^{19} \text{ GeV} (3/\tan \beta)$

---

<sup>5</sup>While  $\Gamma^{-1}(p \rightarrow e^+ \pi^0)$  given by eq. (4.1) does not explicitly depend on  $m_{\bar{q}}, m_{\bar{W}}, r$  and  $\tan \beta$ , the *upper limit* on  $M_X$  and thereby  $\Gamma^{-1}(p \rightarrow e^+ \pi^0)$  does depend on these parameters via the correlation eq. (3.4). The latter relates  $(M_X)_{\text{max}}$  to  $(M_{\text{eff}})_{\text{min}}$  and thereby to the empirical lower limit on  $\Gamma^{-1}(p \rightarrow \bar{\nu} K^+)$  which depends on  $m_{\bar{q}}$  and  $m_{\bar{W}}$ . Because of this, the upper limit given in eq. (4.8) should in fact be multiplied by an approximate factor  $(m_{\bar{q}}/1.5 \text{ TeV})^{8/3} (125 \text{ GeV}/m_{\bar{W}})^{4/3} [(1/300)/r]^{4/3} (3/\tan \beta)^{4/3}$ .

for  $\alpha_3(M_Z) = (0.1156, 0.1176, 0.1196)$ .<sup>6</sup> If we use central values of the parameters and the spectrum as noted above, with  $p \approx 4$  and  $\tan \beta \geq 3$ , the upper limit on  $\Gamma^{-1}(p \rightarrow \bar{\nu}K^+)$  (using eq. (4.7)) would be  $5.16 \times 10^{33}$  yrs. Allowing for uncertainties in the parameters in a combined manner, analogous to the case of  $d = 6$  lifetime, we thus obtain<sup>7</sup>

$$\Gamma^{-1}(p \rightarrow \bar{\nu}K^+) \lesssim (3.1 \times 10^{34} \text{ yrs}) \times \left(\frac{m_{\tilde{q}}}{1.5 \text{ TeV}}\right)^4 \left(\frac{130 \text{ GeV}}{m_{\tilde{W}}}\right)^2 (3/\tan \beta)^2. \quad (4.9)$$

In eq. (4.9) the mild dependence of the curly bracket of eq. (3.4) on  $m_{\tilde{q}}$  and  $m_{\tilde{W}}$  is not exhibited. The actual lifetime is likely to be significantly lower than  $\text{few} \times 10^{34}$  yrs if  $M_{\text{eff}}$  is not stretched to its upper limit (corresponding to e.g.,  $M_X > (M_X)_{\text{min}}$ , or  $r > 1/300$ , or  $\tan \beta > 3$  and/or  $\alpha_3(m_Z) < 0.1196$ ), or if  $m_{\tilde{q}} < 1.5 \text{ TeV}$ , or  $m_{\tilde{W}} > 130 \text{ GeV}$ . We thus find that not only the  $p \rightarrow e^+\pi^0$  mode, but very likely even the  $p \rightarrow \bar{\nu}K^+$  mode should be observable by improving the current experimental sensitivity by about a factor of five to ten.

Some important details concerning the present work, including those pertaining to the issues of fermion masses and mixings, and some variants as regards the cancelation of the  $\mathcal{U}(1)_A$  Fayet-Iliopoulos term, will be presented in a forthcoming longer paper [39].

In summary, we have presented a class of supersymmetric SO(10) models with low dimensional Higgs system that fully resolves all the naturalness issues of doublet-triplet splitting, including stability against higher order operators, generation of  $\mu$ -term of order  $m_{\text{susy}}$ , and proton stability. The threshold corrections in these models are found to depend only on a few effective parameters, making the scenario very predictive. An intriguing feature of these models is the correlation equation and the corresponding constrained upper limit on  $\Gamma^{-1}(p \rightarrow e^+\pi^0)$ . We find that in this class of models proton decays into both  $e^+\pi^0$  and very likely  $\bar{\nu}K^+$  as well should show with an improvement in the current sensitivity by about a factor of five to ten. The building of a megaton water Cherenkov detector (or equivalent) would thus be most welcome.

## Acknowledgments

We would like to thank Takaaki Kajita and Edward Kearns for helpful communications. K.S.B. and Z.T. are supported in part by US Department of Energy, Grant Numbers DE-FG02-04ER41306 and DE-FG02-ER46140. J.C.P. is supported in part by the US Department of Energy, contract Number DE-AC02-76SF00515 and by DOE grant No. DE-FG02-96ER41015. Z.T. is also partially supported by GNSF grant 07\_462\_4-270.

---

<sup>6</sup>Note that these values of  $(M_{\text{eff}})_{\text{max}}$  are quite consistent with the estimate given in eq. (2.7).

<sup>7</sup>We have checked [39] that the rate for  $d = 5$  proton decay generated by Planck scale operators are sufficiently suppressed, owing to symmetries present in the model, so as not to disturb the upper limit quoted in eq. (4.9).

**Open Access.** This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

## References

- [1] J.C. Pati and A. Salam, *Is Baryon Number Conserved?*, *Phys. Rev. Lett.* **31** (1973) 661 [[SPIRES](#)].
- [2] J.C. Pati and A. Salam, *Lepton Number as the Fourth Color*, *Phys. Rev. D* **10** (1974) 275 [[SPIRES](#)].
- [3] H. Georgi and S.L. Glashow, *Unity of All Elementary Particle Forces*, *Phys. Rev. Lett.* **32** (1974) 438 [[SPIRES](#)].
- [4] H. Georgi, H.R. Quinn and S. Weinberg, *Hierarchy of Interactions in Unified Gauge Theories*, *Phys. Rev. Lett.* **33** (1974) 451 [[SPIRES](#)].
- [5] S. Dimopoulos, S. Raby and F. Wilczek, *Supersymmetry and the Scale of Unification*, *Phys. Rev. D* **24** (1981) 1681 [[SPIRES](#)].
- [6] W.J. Marciano and G. Senjanović, *Predictions of Supersymmetric Grand Unified Theories*, *Phys. Rev. D* **25** (1982) 3092 [[SPIRES](#)].
- [7] C. Giunti, C.W. Kim and U.W. Lee, *Running coupling constants and grand unification models*, *Mod. Phys. Lett. A* **6** (1991) 1745 [[SPIRES](#)].
- [8] P. Langacker and M.-x. Luo, *Implications of precision electroweak experiments for  $M_t$ ,  $\rho_0$ ,  $\sin^2 \theta_W$  and grand unification*, *Phys. Rev. D* **44** (1991) 817 [[SPIRES](#)].
- [9] U. Amaldi, W. de Boer and H. Furstenuau, *Comparison of grand unified theories with electroweak and strong coupling constants measured at LEP*, *Phys. Lett. B* **260** (1991) 447 [[SPIRES](#)].
- [10] H. Georgi, *The State Of The Art — Gauge Theories*, in *Particles and Fields*, proceedings of the Meeting of the APS Division of Particles and Fields, Williamsburg Virginia U.S.A, 1974, C. Carlson Ed., AIP New York U.S.A. (1975) [[SPIRES](#)].
- [11] H. Fritzsch and P. Minkowski, *Unified Interactions of Leptons and Hadrons*, *Ann. Phys.* **93** (1975) 193 [[SPIRES](#)].
- [12] S. Dimopoulos and F. Wilczek, *Incomplete Multiplets In Supersymmetric Unified Models*, KITP, Print-81-0600, Santa Barbara U.S.A. (1981) [[SPIRES](#)].
- [13] K.S. Babu and S.M. Barr, *Natural suppression of Higgsino mediated proton decay in supersymmetric SO(10)*, *Phys. Rev. D* **48** (1993) 5354 [[hep-ph/9306242](#)] [[SPIRES](#)].
- [14] K.S. Babu and S.M. Barr, *Natural gauge hierarchy in SO(10)*, *Phys. Rev. D* **50** (1994) 3529 [[hep-ph/9402291](#)] [[SPIRES](#)];  
K.S. Babu and S.M. Barr, *Supersymmetric SO(10) simplified*, *Phys. Rev. D* **51** (1995) 2463 [[hep-ph/9409285](#)] [[SPIRES](#)].
- [15] Z. Chacko and R.N. Mohapatra, *Economical doublet-triplet splitting and strong suppression of proton decay in SO(10)*, *Phys. Rev. D* **59** (1999) 011702 [[hep-ph/9808458](#)] [[SPIRES](#)].
- [16] K.S. Babu, J.C. Pati and F. Wilczek, *Fermion masses, neutrino oscillations and proton decay in the light of SuperKamiokande*, *Nucl. Phys. B* **566** (2000) 33 [[hep-ph/9812538](#)] [[SPIRES](#)].



- [17] Z. Berezhiani and Z. Tavartkiladze, *More missing VEV mechanism in supersymmetric SO(10) model*, *Phys. Lett. B* **409** (1997) 220 [[hep-ph/9612232](#)] [[SPIRES](#)].
- [18] N. Maekawa, *Neutrino masses, anomalous U(1) gauge symmetry and doublet-triplet splitting*, *Prog. Theor. Phys.* **106** (2001) 401 [[hep-ph/0104200](#)] [[SPIRES](#)].
- [19] N. Maekawa and T. Yamashita, *2 loop analysis of gauge coupling unification with anomalous U(1) symmetry and proton decay*, *Prog. Theor. Phys.* **108** (2002) 719 [[hep-ph/0205185](#)] [[SPIRES](#)].
- [20] S.M. Barr and S. Raby, *Minimal SO(10) unification*, *Phys. Rev. Lett.* **79** (1997) 4748 [[hep-ph/9705366](#)] [[SPIRES](#)].
- [21] K.S. Babu and R.N. Mohapatra, *Predictive neutrino spectrum in minimal SO(10) grand unification*, *Phys. Rev. Lett.* **70** (1993) 2845 [[hep-ph/9209215](#)] [[SPIRES](#)].
- [22] B. Bajc, G. Senjanović and F. Vissani, *b- $\tau$  unification and large atmospheric mixing: A case for non-canonical see-saw*, *Phys. Rev. Lett.* **90** (2003) 051802 [[hep-ph/0210207](#)] [[SPIRES](#)].
- [23] C.S. Aulakh, B. Bajc, A. Melfo, G. Senjanović and F. Vissani, *The minimal supersymmetric grand unified theory*, *Phys. Lett. B* **588** (2004) 196 [[hep-ph/0306242](#)] [[SPIRES](#)].
- [24] T. Fukuyama, A. Ilakovac, T. Kikuchi, S. Meljanac and N. Okada, *Detailed analysis of proton decay rate in the minimal supersymmetric SO(10) model*, *JHEP* **09** (2004) 052 [[hep-ph/0406068](#)] [[SPIRES](#)].
- [25] S. Bertolini, M. Frigerio and M. Malinsky, *Fermion masses in SUSY SO(10) with type-II seesaw: A non-minimal predictive scenario*, *Phys. Rev. D* **70** (2004) 095002 [[hep-ph/0406117](#)] [[SPIRES](#)].
- [26] K.S. Babu and C. Macesanu, *Neutrino masses and mixings in a minimal SO(10) model*, *Phys. Rev. D* **72** (2005) 115003 [[hep-ph/0505200](#)] [[SPIRES](#)].
- [27] B. Dutta, Y. Mimura and R.N. Mohapatra, *Proton Decay and Flavor Violating Thresholds in SO(10) Models*, *Phys. Rev. Lett.* **100** (2008) 181801 [[arXiv:0712.1206](#)] [[SPIRES](#)].
- [28] M.-C. Chen and K.T. Mahanthappa, *From CKM matrix to MNS matrix: A model based on supersymmetric SO(10)  $\times$  U(2)<sub>F</sub> symmetry*, *Phys. Rev. D* **62** (2000) 113007 [[hep-ph/0005292](#)] [[SPIRES](#)].
- [29] K.S. Babu, I. Gogoladze, P. Nath and R.M. Syed, *A unified framework for symmetry breaking in SO(10)*, *Phys. Rev. D* **72** (2005) 095011 [[hep-ph/0506312](#)] [[SPIRES](#)].
- [30] C.S. Aulakh and S.K. Garg, *Correcting  $\alpha_3(M_Z)$  in the NMSGUT*, *Mod. Phys. Lett. A* **24** (2009) 1711 [[arXiv:0710.4018](#)] [[SPIRES](#)].
- [31] M. Dine, N. Seiberg and E. Witten, *Fayet-Iliopoulos Terms in String Theory*, *Nucl. Phys. B* **289** (1987) 589 [[SPIRES](#)].
- [32] J.J. Atick, L.J. Dixon and A. Sen, *String Calculation of Fayet-Iliopoulos d Terms in Arbitrary Supersymmetric Compactifications*, *Nucl. Phys. B* **292** (1987) 109 [[SPIRES](#)].
- [33] M. Dine, I. Ichinose and N. Seiberg, *F terms and d Terms in String Theory*, *Nucl. Phys. B* **293** (1987) 253 [[SPIRES](#)].
- [34] N. Maekawa and T. Yamashita, *2 loop analysis of gauge coupling unification with anomalous U(1) symmetry and proton decay*, *Prog. Theor. Phys.* **108** (2002) 719 [[hep-ph/0205185](#)] [[SPIRES](#)].

- [35] K.S. Babu, I. Gogoladze and Z. Tavartkiladze, *Missing partner mechanism in SO(10) grand unification*, *Phys. Lett. B* **650** (2007) 49 [[hep-ph/0612315](#)] [[SPIRES](#)].
- [36] I. Antoniadis, J.R. Ellis, J.S. Hagelin and D.V. Nanopoulos, *The Flipped SU(5)  $\times$  U(1) String Model Revamped*, *Phys. Lett. B* **231** (1989) 65 [[SPIRES](#)].
- [37] A.E. Faraggi, *A New standard-like model in the four-dimensional free fermionic string formulation*, *Phys. Lett. B* **278** (1992) 131 [[SPIRES](#)].
- [38] A.E. Faraggi and J.C. Pati, *A family-universal anomalous U(1) in string models as the origin of supersymmetry breaking and squark degeneracy*, *Nucl. Phys. B* **526** (1998) 21 [[hep-ph/9712516](#)] [[SPIRES](#)].
- [39] K.S. Babu, J.C. Pati and Z. Tavartkiladze, to appear.
- [40] A.H. Chamseddine, R.L. Arnowitt and P. Nath, *Locally Supersymmetric Grand Unification*, *Phys. Rev. Lett.* **49** (1982) 970 [[SPIRES](#)].
- [41] R. Barbieri, S. Ferrara and C.A. Savoy, *Gauge Models with Spontaneously Broken Local Supersymmetry*, *Phys. Lett. B* **119** (1982) 343 [[SPIRES](#)].
- [42] J. Hisano, H. Murayama and T. Yanagida, *Nucleon decay in the minimal supersymmetric SU(5) grand unification*, *Nucl. Phys. B* **402** (1993) 46 [[hep-ph/9207279](#)] [[SPIRES](#)].
- [43] J. Hisano, T. Moroi, K. Tobe and T. Yanagida, *Limit on the color triplet Higgs mass in the minimum supersymmetric SU(5) model*, *Mod. Phys. Lett. A* **10** (1995) 2267 [[hep-ph/9411298](#)] [[SPIRES](#)].
- [44] H. Murayama and A. Pierce, *Not even decoupling can save minimal supersymmetric SU(5)*, *Phys. Rev. D* **65** (2002) 055009 [[hep-ph/0108104](#)] [[SPIRES](#)].
- [45] B. Bajc, P. Fileviez Perez and G. Senjanović, *Minimal supersymmetric SU(5) theory and proton decay: Where do we stand?*, [hep-ph/0210374](#) [[SPIRES](#)].
- [46] P. Nath and P. Fileviez Perez, *Proton stability in grand unified theories, in strings and in branes*, *Phys. Rept.* **441** (2007) 191 [[hep-ph/0601023](#)] [[SPIRES](#)].
- [47] Y. Aoki, C. Dawson, J. Noaki and A. Soni, *Proton decay matrix elements with domain-wall fermions*, *Phys. Rev. D* **75** (2007) 014507 [[hep-lat/0607002](#)] [[SPIRES](#)].
- [48] RBC-UKQCD collaboration, C. Allton et al., *Physical Results from 2 + 1 Flavor Domain Wall QCD and SU(2) Chiral Perturbation Theory*, *Phys. Rev. D* **78** (2008) 114509 [[arXiv:0804.0473](#)] [[SPIRES](#)].
- [49] SUPER-KAMIOKANDE collaboration, K. Kobayashi et al., *Search for nucleon decay via modes favored by supersymmetric grand unification models in Super-Kamiokande-I*, *Phys. Rev. D* **72** (2005) 052007 [[hep-ex/0502026](#)] [[SPIRES](#)].
- [50] SUPER-KAMIOKANDE collaboration, H. Nishino et al., *Search for Proton Decay via  $p \rightarrow e^+ \pi^0$  and  $p \rightarrow \mu^+ \pi^0$  in a Large Water Cherenkov Detector*, *Phys. Rev. Lett.* **102** (2009) 141801 [[arXiv:0903.0676](#)] [[SPIRES](#)].
- [51] K. Kaneyuki, *Result of nucleon decay search in Super-Kamiokande*, presentation at XI International Conference TAUP, Italy 2009.
- [52] E. Kearns, *Experimental searches for baryon number violation*, presentation at LBV 2009, Madison, WI U.S.A. 2009.
- [53] <http://www-sk.icrr.u-tokyo.ac.jp>.

- [54] PARTICLE DATA GROUP collaboration, C. Amsler et al., *Review of particle physics*, *Phys. Lett. B* **667** (2008) 1 [[SPIRES](#)].
- [55] K.S. Babu, J.C. Pati and P. Rastogi, *Tying in CP and flavor violations with fermion masses and neutrino oscillations*, *Phys. Rev. D* **71** (2005) 015005 [[hep-ph/0410200](#)] [[SPIRES](#)].
- [56] V. Lucas and S. Raby, *Nucleon decay in a realistic SO(10) SUSY GUT*, *Phys. Rev. D* **55** (1997) 6986 [[hep-ph/9610293](#)] [[SPIRES](#)].
- [57] R. Dermisek, A. Mafi and S. Raby, *SUSY GUTs under siege: Proton decay*, *Phys. Rev. D* **63** (2001) 035001 [[hep-ph/0007213](#)] [[SPIRES](#)].
- [58] C.H. Albright and S.M. Barr, *Fermion masses in SO(10) with a single adjoint Higgs field*, *Phys. Rev. D* **58** (1998) 013002 [[hep-ph/9712488](#)] [[SPIRES](#)].
- [59] C.H. Albright, K.S. Babu and S.M. Barr, *A minimality condition and atmospheric neutrino oscillations*, *Phys. Rev. Lett.* **81** (1998) 1167 [[hep-ph/9802314](#)] [[SPIRES](#)].
- [60] P. Pouliot and N. Seiberg, *(S)quark masses and nonAbelian horizontal symmetries*, *Phys. Lett. B* **318** (1993) 169 [[hep-ph/9308363](#)] [[SPIRES](#)].
- [61] K.S. Babu and J. Kubo, *Dihedral families of quarks, leptons and Higgses*, *Phys. Rev. D* **71** (2005) 056006 [[hep-ph/0411226](#)] [[SPIRES](#)].