

## RESEARCH

## Open Access



# Sums of products of two reciprocal Fibonacci numbers

Runnan Liu<sup>1</sup> and Andrew YZ Wang<sup>2\*</sup>

\*Correspondence:

yzwang@uestc.edu.cn

<sup>2</sup>School of Mathematical Sciences,  
University of Electronic Science and  
Technology of China, Chengdu,  
611731, P.R. ChinaFull list of author information is  
available at the end of the article**Abstract**

In this paper, we employ elementary methods to investigate the reciprocal sums of the products of two Fibonacci numbers in several ways. First, we consider the sums of the reciprocals of the products of two Fibonacci numbers and establish five interesting families of identities. Then we extend such analysis to the alternating sums and obtain five analogous results.

**MSC:** 11B39**Keywords:** Fibonacci number; reciprocal sum; floor function

## 1 Introduction

For an integer  $n \geq 0$ , the *Fibonacci number*  $F_n$  is defined by

$$F_n = F_{n-1} + F_{n-2} \quad \text{for } n \geq 2,$$

with  $F_0 = 0$  and  $F_1 = 1$ . There exists a simple and nonobvious formula for the Fibonacci numbers:

$$F_n = \frac{1}{\sqrt{5}} \left( \frac{1 + \sqrt{5}}{2} \right)^n - \frac{1}{\sqrt{5}} \left( \frac{1 - \sqrt{5}}{2} \right)^n.$$

The Fibonacci numbers play an important role in the theory and applications of mathematics, and its various properties have been investigated by many authors; see [1–4].

In recent years, there has been an increasing interest in studying the reciprocal sums of the Fibonacci numbers. For example, Elsner, Shimomura, and Shiokawa [5–8] investigated algebraic relations for reciprocal sums of the Fibonacci numbers. Ohtsuka and Nakamura [9] studied the partial infinite sums of the reciprocal Fibonacci numbers. They established the following results, where  $\lfloor \cdot \rfloor$  denotes the floor function.

**Theorem 1.1** For all  $n \geq 2$ ,

$$\left\lfloor \left( \sum_{k=n}^{\infty} \frac{1}{F_k} \right)^{-1} \right\rfloor = \begin{cases} F_{n-2} & \text{if } n \text{ is even;} \\ F_{n-2} - 1 & \text{if } n \text{ is odd.} \end{cases}$$

**Theorem 1.2** For each  $n \geq 1$ ,

$$\left[ \left( \sum_{k=n}^{\infty} \frac{1}{F_k^2} \right)^{-1} \right] = \begin{cases} F_n F_{n-1} - 1 & \text{if } n \text{ is even;} \\ F_n F_{n-1} & \text{if } n \text{ is odd.} \end{cases}$$

Recently, Wang and Wen [10] considered the partial finite sums of the reciprocal Fibonacci numbers and strengthened Theorem 1.1 and Theorem 1.2 to the finite-sum case.

**Theorem 1.3**

(i) For all  $n \geq 4$ ,

$$\left[ \left( \sum_{k=n}^{2n} \frac{1}{F_k} \right)^{-1} \right] = F_{n-2}.$$

(ii) If  $m \geq 3$  and  $n \geq 2$ , then

$$\left[ \left( \sum_{k=n}^{mn} \frac{1}{F_k} \right)^{-1} \right] = \begin{cases} F_{n-2} & \text{if } n \text{ is even;} \\ F_{n-2} - 1 & \text{if } n \text{ is odd.} \end{cases}$$

**Theorem 1.4** For all  $m \geq 2$  and  $n \geq 1$ , we have

$$\left[ \left( \sum_{k=n}^{mn} \frac{1}{F_k^2} \right)^{-1} \right] = \begin{cases} F_n F_{n-1} - 1 & \text{if } n \text{ is even;} \\ F_n F_{n-1} & \text{if } n \text{ is odd.} \end{cases}$$

Furthermore, Wang and Zhang [11] studied the reciprocal sums of the Fibonacci numbers with even or odd indexes and obtained the following main results.

**Theorem 1.5** We have

$$\left[ \left( \sum_{k=n}^{mn} \frac{1}{F_{2k}} \right)^{-1} \right] = \begin{cases} F_{2n-1} & \text{if } m = 2 \text{ and } n \geq 3; \\ F_{2n-1} - 1 & \text{if } m \geq 3 \text{ and } n \geq 1. \end{cases}$$

**Theorem 1.6** For all  $n \geq 1$  and  $m \geq 2$ , we have

$$\left[ \left( \sum_{k=n}^{mn} \frac{1}{F_{2k-1}} \right)^{-1} \right] = F_{2n-2}.$$

**Theorem 1.7** If  $n \geq 1$  and  $m \geq 2$ , then

$$\left[ \left( \sum_{k=n}^{mn} \frac{1}{F_{2k}^2} \right)^{-1} \right] = F_{4n-2} - 1.$$

**Theorem 1.8** For all  $n \geq 1$  and  $m \geq 2$ , we have

$$\left[ \left( \sum_{k=n}^{mn} \frac{1}{F_{2k-1}^2} \right)^{-1} \right] = F_{4n-4}.$$

More recently, Wang and Zhang [12] proceeded with investigating the reciprocal sums of the Fibonacci numbers according to the subscripts modulo 3 and found many identities. Here are a few examples.

**Theorem 1.9**

(i) For all  $n \geq 2$ ,

$$\left\lfloor \left( \sum_{k=n}^{2n} \frac{1}{F_{3k}} \right)^{-1} \right\rfloor = 2F_{3n-2}.$$

(ii) If  $m \geq 3$  and  $n \geq 1$ , then

$$\left\lfloor \left( \sum_{k=n}^{mn} \frac{1}{F_{3k}} \right)^{-1} \right\rfloor = \begin{cases} 2F_{3n-2} & \text{if } n \text{ is even;} \\ 2F_{3n-2} - 1 & \text{if } n \text{ is odd.} \end{cases}$$

**Theorem 1.10** If  $n \geq 2$  and  $m \geq 2$ , we have

$$\left\lfloor \left( \sum_{k=n}^{mn} \frac{1}{F_{3k}^2} \right)^{-1} \right\rfloor = \begin{cases} F_{3n}^2 - F_{3n-3}^2 & \text{if } n \text{ is even;} \\ F_{3n}^2 - F_{3n-3}^2 - 1 & \text{if } n \text{ is odd.} \end{cases}$$

In this article, we focus ourselves on the sums and alternating sums of the products of two reciprocal Fibonacci numbers. By evaluating the integer parts of these sums, we obtain several interesting families of identities concerning the Fibonacci numbers.

**2 Main results I: reciprocal sums**

We first introduce several well-known results on the Fibonacci numbers, which will be used throughout the article. The detailed proofs can be found in, for example, [4] and [13].

**Lemma 2.1** For any positive integers  $m$  and  $n$ , we have

$$F_m F_n + F_{m+1} F_{n+1} = F_{m+n+1}. \tag{2.1}$$

**Lemma 2.2** If  $n \geq 1$ , then

$$F_{2n} = F_{n+1}^2 - F_{n-1}^2, \tag{2.2}$$

$$F_{2n+1} = F_{n+1} F_{n+2} - F_{n-1} F_n. \tag{2.3}$$

**Lemma 2.3** Let  $a, b, c, d$  be positive integers with  $a + b = c + d$  and  $b \geq \max\{c, d\}$ . Then

$$F_a F_b - F_c F_d = (-1)^{a+1} F_{b-c} F_{b-d}. \tag{2.4}$$

**2.1 Reciprocal sum of  $F_k F_{k+1}$**

**Lemma 2.4** For all  $n \geq 1$ , we have

$$F_{2n+1}^2 + 1 > (F_{n+1}^2 + 1)^2 > F_n F_{n+1} (F_{n+1}^2 + 1). \tag{2.5}$$

*Proof* It follows from (2.1) that  $F_{2n+1} = F_n^2 + F_{n+1}^2$ . Hence,

$$F_{2n+1}^2 + 1 = F_n^4 + 2F_n^2F_{n+1}^2 + F_{n+1}^4 + 1 > F_{n+1}^4 + 2F_{n+1}^2 + 1 = (F_{n+1}^2 + 1)^2.$$

It is clear that  $F_{n+1}^2 \geq F_nF_{n+1}$ ; therefore,  $F_{n+1}^2 + 1 > F_nF_{n+1}$ , which yields the second inequality.  $\square$

**Theorem 2.5** *If  $m \geq 2$  and  $n \geq 1$ , then*

$$\left[ \left( \sum_{k=n}^{mn} \frac{1}{F_k F_{k+1}} \right)^{-1} \right] = \begin{cases} F_n^2 & \text{if } n \text{ is even;} \\ F_n^2 - 1 & \text{if } n \text{ is odd.} \end{cases}$$

*Proof* We first consider the case where  $n$  is even. By elementary manipulations and setting  $a = k - 1$ ,  $b = k + 1$ , and  $c = d = k$  in (2.4), we obtain, for  $k \geq 1$ ,

$$\begin{aligned} \frac{1}{F_k^2} - \frac{1}{F_k F_{k+1}} - \frac{1}{F_{k+1}^2} &= \frac{F_{k+1}^2 - F_k F_{k+1} - F_k^2}{F_k^2 F_{k+1}^2} \\ &= \frac{F_{k-1} F_{k+1} - F_k^2}{F_k^2 F_{k+1}^2} \\ &= \frac{(-1)^k}{F_k^2 F_{k+1}^2}. \end{aligned} \tag{2.6}$$

Now we have

$$\sum_{k=n}^{mn} \frac{1}{F_k F_{k+1}} = \frac{1}{F_n^2} - \frac{1}{F_{mn+1}^2} + \sum_{k=n}^{mn} \frac{(-1)^{k-1}}{F_k^2 F_{k+1}^2}.$$

Since  $n$  is even, it is easy to see that

$$\sum_{k=n}^{mn} \frac{(-1)^{k-1}}{F_k^2 F_{k+1}^2} < 0,$$

which implies that

$$\sum_{k=n}^{mn} \frac{1}{F_k F_{k+1}} < \frac{1}{F_n^2}. \tag{2.7}$$

A direct calculation shows that, for  $k \geq 1$ ,

$$\begin{aligned} \frac{1}{F_k^2 + 1} - \frac{1}{F_k F_{k+1}} - \frac{1}{F_{k+1}^2 + 1} &= \frac{F_k F_{k+1} (F_{k+1}^2 - F_k^2) - (F_k^2 + 1)(F_{k+1}^2 + 1)}{(F_k^2 + 1) F_k F_{k+1} (F_{k+1}^2 + 1)} \\ &= \frac{F_k F_{k+1} (F_{k+1}^2 - F_k^2 - F_k F_{k+1}) - F_k^2 - F_{k+1}^2 - 1}{(F_k^2 + 1) F_k F_{k+1} (F_{k+1}^2 + 1)} \\ &= \frac{F_k F_{k+1} (F_{k-1} F_{k+1} - F_k^2) - F_k^2 - F_{k+1}^2 - 1}{(F_k^2 + 1) F_k F_{k+1} (F_{k+1}^2 + 1)} \end{aligned}$$

$$= \frac{(-1)^k F_k F_{k+1} - F_k^2 - F_{k+1}^2 - 1}{(F_k^2 + 1)F_k F_{k+1}(F_{k+1}^2 + 1)} < 0.$$

Therefore,

$$\begin{aligned} \sum_{k=n}^{mn} \frac{1}{F_k F_{k+1}} &= \frac{1}{F_n^2 + 1} - \frac{1}{F_{mn+1}^2 + 1} + \sum_{k=n}^{mn} \frac{(-1)^{k-1} F_k F_{k+1} + F_k^2 + F_{k+1}^2 + 1}{(F_k^2 + 1)F_k F_{k+1}(F_{k+1}^2 + 1)} \\ &> \frac{1}{F_n^2 + 1} - \frac{1}{F_{2n+1}^2 + 1} + \frac{F_n^2 + F_{n+1}^2 + 1 - F_n F_{n+1}}{(F_n^2 + 1)F_n F_{n+1}(F_{n+1}^2 + 1)} \\ &> \frac{1}{F_n^2 + 1} + \frac{1}{F_n F_{n+1}(F_{n+1}^2 + 1)} - \frac{1}{F_{2n+1}^2 + 1} \\ &> \frac{1}{F_n^2 + 1}, \end{aligned} \tag{2.8}$$

where the last inequality follows from (2.5).

Combining (2.7) and (2.8), we have

$$\frac{1}{F_n^2 + 1} < \sum_{k=n}^{mn} \frac{1}{F_k F_{k+1}} < \frac{1}{F_n^2},$$

which means that the statement is true when  $n$  is even.

We now concentrate ourselves on the case where  $n$  is odd. It is obviously true for  $n = 1$ .

Now we assume that  $n \geq 3$ . A similar calculation shows that, for  $k \geq 3$ ,

$$\frac{1}{F_k^2 - 1} - \frac{1}{F_k F_{k+1}} - \frac{1}{F_{k+1}^2 - 1} = \frac{(-1)^k F_k F_{k+1} + F_k^2 + F_{k+1}^2 + 1}{(F_k^2 - 1)F_k F_{k+1}(F_{k+1}^2 - 1)} > 0,$$

from which we get

$$\sum_{k=n}^{mn} \frac{1}{F_k F_{k+1}} < \frac{1}{F_n^2 - 1} - \frac{1}{F_{mn+1}^2 - 1} < \frac{1}{F_n^2 - 1}. \tag{2.9}$$

It follows from (2.1) that

$$F_{2n+1} = F_{n-1}F_{n+1} + F_n F_{n+2} = F_n^2 + F_{n+1}^2,$$

which implies that  $F_{2n+1} \geq F_n F_{n+2}$ , and  $F_{2n+2} > F_{2n+1} > F_{n+1}^2$ . Therefore,

$$F_{2n+1}^2 F_{2n+2} > F_n^2 F_{n+1}^2 F_{n+2}^2. \tag{2.10}$$

Invoking (2.6), (2.2), and the fact  $n$  is odd, we have

$$\begin{aligned} \sum_{k=n}^{mn} \frac{1}{F_k F_{k+1}} &= \frac{1}{F_n^2} - \frac{1}{F_{mn+1}^2} + \sum_{k=n}^{mn} \frac{(-1)^{k-1}}{F_k^2 F_{k+1}^2} \\ &> \frac{1}{F_n^2} - \frac{1}{F_{2n+1}^2} + \frac{1}{F_n^2 F_{n+1}^2} - \frac{1}{F_{n+1}^2 F_{n+2}^2} \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{F_n^2} + \frac{F_{n+2}^2 - F_n^2}{F_n^2 F_{n+1}^2 F_{n+2}^2} - \frac{1}{F_{2n+1}^2} \\
 &= \frac{1}{F_n^2} + \frac{F_{2n+2}}{F_n^2 F_{n+1}^2 F_{n+2}^2} - \frac{F_{2n+2}}{F_{2n+1}^2 F_{2n+2}} \\
 &> \frac{1}{F_n^2},
 \end{aligned} \tag{2.11}$$

where the last inequality follows from (2.10).

Combining (2.9) and (2.11) yields that

$$\frac{1}{F_n^2} < \sum_{k=n}^{mn} \frac{1}{F_k F_{k+1}} < \frac{1}{F_n - 1},$$

from which the desired result follows immediately. □

### 2.2 Reciprocal sum of $F_{2k-1}F_{2k}$

**Lemma 2.6** *For all  $n \geq 1$ , we have*

$$F_{8n+1} > F_{2n-1}F_{2n}(F_{4n+1} + 1).$$

*Proof* Applying (2.1) repeatedly, we have

$$F_{8n+1} = F_{4n+1}^2 + F_{4n}^2 > F_{4n+1}^2 + F_{4n+1} = F_{4n+1}(F_{4n+1} + 1) > F_{2n-1}F_{2n}(F_{4n+1} + 1),$$

which completes the proof. □

**Theorem 2.7** *For all  $n \geq 2$  and  $m \geq 2$ ,*

$$\left[ \left( \sum_{k=n}^{mn} \frac{1}{F_{2k-1}F_{2k}} \right)^{-1} \right] = F_{4n-3}.$$

*Proof* It follows from (2.3) that

$$F_{4n+1} = F_{2n+1}F_{2n+2} - F_{2n-1}F_{2n}, \tag{2.12}$$

$$F_{4n-3} = F_{2n-1}F_{2n} - F_{2n-3}F_{2n-2}. \tag{2.13}$$

Employing (2.4), we can easily get that

$$F_{2n-2}F_{2n+1} = F_{2n-1}F_{2n} - 1, \tag{2.14}$$

$$F_{2n-3}F_{2n+2} = F_{2n-1}F_{2n} + 2. \tag{2.15}$$

Applying (2.12), (2.13), (2.14), and (2.15), it is easy to see that, for all  $k \geq 2$ ,

$$\begin{aligned}
 \frac{1}{F_{4k-3}} - \frac{1}{F_{2k-1}F_{2k}} - \frac{1}{F_{4k+1}} &= \frac{F_{2k-1}F_{2k}(F_{4k+1} - F_{4k-3}) - F_{4k-3}F_{4k+1}}{F_{4k-3}F_{2k-1}F_{2k}F_{4k+1}} \\
 &= \frac{F_{2k-3}F_{2k-2}F_{2k+1}F_{2k+2} - F_{2k-1}^2 F_{2k}^2}{F_{4k-3}F_{2k-1}F_{2k}F_{4k+1}}
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{F_{2k-1}F_{2k} - 2}{F_{4k-3}F_{2k-1}F_{2k}F_{4k+1}} \\
 &> 0,
 \end{aligned}$$

which implies that

$$\sum_{k=n}^{mn} \frac{1}{F_{2k-1}F_{2k}} < \sum_{k=n}^{mn} \left( \frac{1}{F_{4k-3}} - \frac{1}{F_{4k+1}} \right) = \frac{1}{F_{4n-3}} - \frac{1}{F_{4mn+1}} < \frac{1}{F_{4n-3}}. \tag{2.16}$$

It follows from (2.1) that

$$F_{4n+1} > F_{4n} = F_{2n-1}F_{2n} + F_{2n}F_{2n+1} > F_{2n-1}F_{2n} + 1.$$

Therefore, by (2.13) we obtain

$$F_{4n+1} - F_{2n-3}F_{2n-2} > F_{2n-1}F_{2n} - F_{2n-3}F_{2n-2} + 1 = F_{4n-3} + 1. \tag{2.17}$$

Elementary manipulations and (2.17) yield, for  $k \geq 2$ ,

$$\begin{aligned}
 \frac{1}{F_{4k-3} + 1} - \frac{1}{F_{2k-1}F_{2k}} - \frac{1}{F_{4k+1} + 1} &= \frac{F_{2k-3}F_{2k-2} - F_{4k+1} - 3}{(F_{4k-3} + 1)F_{2k-1}F_{2k}(F_{4k+1} + 1)} \\
 &< \frac{-1}{F_{2k-1}F_{2k}(F_{4k+1} + 1)}.
 \end{aligned}$$

Now we can deduce that

$$\begin{aligned}
 \sum_{k=n}^{mn} \frac{1}{F_{2k-1}F_{2k}} &> \frac{1}{F_{4n-3} + 1} - \frac{1}{F_{4mn+1} + 1} + \sum_{k=n}^{mn} \frac{1}{F_{2k-1}F_{2k}(F_{4k+1} + 1)} \\
 &> \frac{1}{F_{4n-3} + 1} + \frac{1}{F_{2n-1}F_{2n}(F_{4n+1} + 1)} - \frac{1}{F_{4mn+1} + 1} \\
 &> \frac{1}{F_{4n-3} + 1} + \frac{1}{F_{2n-1}F_{2n}(F_{4n+1} + 1)} - \frac{1}{F_{8n+1} + 1} \\
 &> \frac{1}{F_{4n-3} + 1}, \tag{2.18}
 \end{aligned}$$

where the last inequality follows from Lemma 2.6.

Combining (2.16) and (2.18), we deduce

$$\frac{1}{F_{4n-3} + 1} < \sum_{k=n}^{mn} \frac{1}{F_{2k-1}F_{2k}} < \frac{1}{F_{4n-3}},$$

which yields the desired identity. □

Similarly, we can prove the following result, whose proof is left as an exercise to the readers.

**Theorem 2.8** *For all  $n \geq 2$  and  $m \geq 2$ ,*

$$\left[ \left( \sum_{k=n}^{mn} \frac{1}{F_{2k}F_{2k+1}} \right)^{-1} \right] = F_{4n-1} - 1.$$

### 2.3 Reciprocal sum of $F_{2k-1}F_{2k+1}$

**Lemma 2.9** For all  $n \geq 2$ , we have

$$F_{n+2}^2 - F_{n-2}^2 > 4F_{n-1}F_{n+1}.$$

*Proof* It is easy to check that

$$\begin{aligned} F_{n+2}^2 - F_{n-2}^2 &= (F_{n+2} - F_{n-2})(F_{n+2} + F_{n-2}) \\ &> (F_{n+1} + F_{n-1})(F_{n+1} + F_{n-1}) \\ &= F_{n+1}^2 + F_{n-1}^2 + 2F_{n-1}F_{n+1} \\ &= (F_{n+1} - F_{n-1})^2 + 4F_{n-1}F_{n+1} \\ &> 4F_{n-1}F_{n+1}. \end{aligned}$$

□

**Lemma 2.10** If  $n \geq 1$ , then

$$F_{8n+2} > (F_{4n-2} + 1)(F_{4n+2} + 1).$$

*Proof* It follows from (2.2) that

$$F_{8n+2} = F_{4n+2}^2 - F_{4n}^2 = (F_{4n+2} - F_{4n})(F_{4n+2} + F_{4n}) = F_{4n+1}(F_{4n+2} + F_{4n}).$$

It is obvious that  $F_{4n+1} > F_{4n-2} + 1$  and  $F_{4n} > 1$ , which completes the proof.

□

**Theorem 2.11** For all  $n \geq 1$  and  $m \geq 2$ ,

$$\left[ \left( \sum_{k=n}^{mn} \frac{1}{F_{2k-1}F_{2k+1}} \right)^{-1} \right] = F_{4n-2}.$$

*Proof* Employing (2.2), we can readily see that

$$\begin{aligned} F_{4n+2} &= F_{2n+2}^2 - F_{2n}^2 = F_{2n+1}(F_{2n+2} + F_{2n}) = F_{2n+1}^2 + 2F_{2n}F_{2n+1}, \\ F_{4n-2} &= F_{2n}^2 - F_{2n-2}^2 = F_{2n-1}(F_{2n} + F_{2n-2}) = 2F_{2n-1}F_{2n} - F_{2n-1}^2. \end{aligned}$$

Applying (2.4), we can establish the following identities:

$$\begin{aligned} F_{2n+1}^2 &= F_{2n}F_{2n+2} + 1, \\ F_{2n-1}^2 &= F_{2n-2}F_{2n} + 1, \\ F_{2n}^2 &= F_{2n-2}F_{2n+2} + 1. \end{aligned}$$

With the help of these identities, we now arrive at

$$\begin{aligned} F_{4n+2} - F_{4n-2} &= F_{2n+1}^2 + F_{2n-1}^2 + 2F_{2n}F_{2n+1} - 2F_{2n-1}F_{2n} \\ &= F_{2n+1}^2 + F_{2n-1}^2 + 2F_{2n}^2 \end{aligned}$$



$$\begin{aligned}
 &= (F_{2n}F_{2n+2} + 1) + (F_{2n-2}F_{2n} + 1) + (F_{2n-2}F_{2n+2} + 1) + F_{2n}^2 \\
 &= (F_{2n+2} + F_{2n})(F_{2n} + F_{2n-2}) + 3 \\
 &= \frac{F_{4n+2}F_{4n-2}}{F_{2n-1}F_{2n+1}} + 3.
 \end{aligned} \tag{2.19}$$

Elementary manipulations and (2.19) yield that, for  $k \geq 1$ ,

$$\begin{aligned}
 \frac{1}{F_{4k-2}} - \frac{1}{F_{2k-1}F_{2k+1}} - \frac{1}{F_{4k+2}} &= \frac{F_{2k-1}F_{2k+1}(F_{4k+2} - F_{4k-2}) - F_{4k-2}F_{4k+2}}{F_{4k-2}F_{2k-1}F_{2k+1}F_{4k+2}} \\
 &= \frac{3F_{2k-1}F_{2k+1}}{F_{4k-2}F_{2k-1}F_{2k+1}F_{4k+2}} \\
 &> 0,
 \end{aligned} \tag{2.20}$$

which implies that

$$\sum_{k=n}^{mn} \frac{1}{F_{2k-1}F_{2k+1}} < \sum_{k=n}^{mn} \left( \frac{1}{F_{4k-2}} - \frac{1}{F_{4k+2}} \right) = \frac{1}{F_{4n-2}} - \frac{1}{F_{4mn+2}} < \frac{1}{F_{4n-2}}. \tag{2.21}$$

Invoking (2.20) and Lemma 2.9, we have

$$\begin{aligned}
 &\frac{1}{F_{4k-2} + 1} - \frac{1}{F_{2k-1}F_{2k+1}} - \frac{1}{F_{4k+2} + 1} \\
 &= \frac{F_{2k-1}F_{2k+1}(F_{4k+2} - F_{4k-2}) - F_{4k-2}F_{4k+2}}{(F_{4k-2} + 1)F_{2k-1}F_{2k+1}(F_{4k+2} + 1)} \\
 &\quad - \frac{F_{4k-2} + F_{4k+2} + 1}{(F_{4k-2} + 1)F_{2k-1}F_{2k+1}(F_{4k+2} + 1)} \\
 &= \frac{3F_{2k-1}F_{2k+1} - (F_{2k+2}^2 - F_{2k-2}^2) - 1}{(F_{4k-2} + 1)F_{2k-1}F_{2k+1}(F_{4k+2} + 1)} \\
 &< \frac{3F_{2k-1}F_{2k+1} - 4F_{2k-1}F_{2k+1} - 1}{(F_{4k-2} + 1)F_{2k-1}F_{2k+1}(F_{4k+2} + 1)} \\
 &< -\frac{1}{(F_{4k-2} + 1)(F_{4k+2} + 1)},
 \end{aligned}$$

from which we deduce that

$$\begin{aligned}
 \sum_{k=n}^{mn} \frac{1}{F_{2k-1}F_{2k+1}} &> \sum_{k=n}^{mn} \left( \frac{1}{F_{4k-2} + 1} - \frac{1}{F_{4k+2} + 1} \right) + \sum_{k=n}^{mn} \frac{1}{(F_{4k-2} + 1)(F_{4k+2} + 1)} \\
 &= \frac{1}{F_{4n-2} + 1} - \frac{1}{F_{4mn+2} + 1} + \sum_{k=n}^{mn} \frac{1}{(F_{4k-2} + 1)(F_{4k+2} + 1)} \\
 &> \frac{1}{F_{4n-2} + 1} + \frac{1}{(F_{4n-2} + 1)(F_{4n+2} + 1)} - \frac{1}{F_{8n+2} + 1} \\
 &> \frac{1}{F_{4n-2} + 1},
 \end{aligned} \tag{2.22}$$

where the last inequality follows from Lemma 2.10.

Combining (2.21) and (2.22), we obtain

$$\frac{1}{F_{4n-2} + 1} < \sum_{k=n}^{mn} \frac{1}{F_{2k-1}F_{2k+1}} < \frac{1}{F_{4n-2}},$$

from which the desired result follows. □

Similarly, we can obtain the following result, whose proof is omitted here.

**Theorem 2.12** *For all  $n \geq 1$  and  $m \geq 2$ ,*

$$\left[ \left( \sum_{k=n}^{mn} \frac{1}{F_{2k}F_{2k+2}} \right)^{-1} \right] = F_{4n} - 1.$$

### 3 Main results II: alternating reciprocal sums

In this section, we extend the analysis of the sums of the products of two reciprocal Fibonacci numbers to alternating sums.

#### 3.1 Alternating reciprocal sum of $F_k F_{k+1}$

**Lemma 3.1** *For  $n \geq 1$ , we have*

$$\frac{F_{n+1}}{F_n} - \frac{1}{F_{2n}} = \frac{F_{2n+1} + (-1)^n - 1}{F_{2n}}, \tag{3.1}$$

$$\frac{F_{n+1}}{F_n} + \frac{1}{F_{2n}} = \frac{F_{2n+1} + (-1)^n + 1}{F_{2n}}. \tag{3.2}$$

*Proof* Applying (2.1) repeatedly and (2.4), we derive that

$$\begin{aligned} \frac{F_{n+1}}{F_n} - \frac{1}{F_{2n}} &= \frac{F_{n+1}F_{2n} - F_n}{F_n F_{2n}} \\ &= \frac{F_{n+1}(F_{n-1}F_n + F_n F_{n+1}) - F_n}{F_n F_{2n}} \\ &= \frac{F_{n+1}F_{n-1} + F_{n+1}^2 - 1}{F_{2n}} \\ &= \frac{(F_{n+1}F_{n-1} - F_n^2) + (F_n^2 + F_{n+1}^2) - 1}{F_{2n}} \\ &= \frac{F_{2n+1} + (-1)^n - 1}{F_{2n}}. \end{aligned}$$

Then (3.2) immediately follows from (3.1). □

**Lemma 3.2** *If  $n \geq 2$ , then*

$$\frac{1}{F_{2n} - 1} - \frac{F_{n+1}}{F_n F_{2n+1}} > 0, \tag{3.3}$$

$$\frac{1}{F_{2n} + 1} - \frac{F_{n+1}}{F_n F_{2n+1}} < 0. \tag{3.4}$$

*Proof* It follows from (2.4) that

$$\frac{1}{F_{2n}-1} - \frac{F_{n+1}}{F_n F_{2n+1}} = \frac{F_n F_{2n+1} - F_{n+1} F_{2n} + F_{n+1}}{(F_{2n}-1)F_n F_{2n+1}} = \frac{(-1)^{n+1} F_n + F_{n+1}}{(F_{2n}-1)F_n F_{2n+1}} > 0.$$

Similarly, we can prove (3.4). □

**Lemma 3.3** For  $n \geq 1$  and  $m \geq 2$ , we have

$$\frac{F_{2n+1}}{F_{2n}} - \frac{F_{mn+2}}{F_{mn+1}} > 0, \tag{3.5}$$

$$\frac{F_{2n+2}}{F_{2n+1}} - \frac{F_{mn+2}}{F_{mn+1}} \leq 0. \tag{3.6}$$

*Proof* With the help of (2.4), we see that

$$\frac{F_{2n+1}}{F_{2n}} - \frac{F_{mn+2}}{F_{mn+1}} = \frac{F_{2n+1} F_{mn+1} - F_{2n} F_{mn+2}}{F_{2n} F_{mn+1}} = (-1)^{2n+2} \frac{F_{(m-2)n+1}}{F_{2n} F_{mn+1}} > 0.$$

A similar analysis yields (3.6). □

**Theorem 3.4** If  $m \geq 2$  and  $n \geq 1$ , then

$$\left[ \left( \sum_{k=n}^{mn} \frac{(-1)^k}{F_k F_{k+1}} \right)^{-1} \right] = \begin{cases} F_{2n} - 1 & \text{if } n \text{ is even;} \\ -F_{2n} - 1 & \text{if } n \text{ is odd.} \end{cases}$$

*Proof* Employing (2.4), we derive that

$$\frac{(-1)^k}{F_k F_{k+1}} = \frac{F_{k+1}^2 - F_k F_{k+2}}{F_k F_{k+1}} = \frac{F_{k+1}}{F_k} - \frac{F_{k+2}}{F_{k+1}},$$

which implies that

$$\sum_{k=n}^{mn} \frac{(-1)^k}{F_k F_{k+1}} = \frac{F_{n+1}}{F_n} - \frac{F_{mn+2}}{F_{mn+1}}. \tag{3.7}$$

Furthermore, it follows from (3.6) and (2.4) that

$$\frac{F_{n+1}}{F_n} - \frac{F_{mn+2}}{F_{mn+1}} \leq \frac{F_{n+1}}{F_n} - \frac{F_{2n+2}}{F_{2n+1}} = \frac{F_{n+1} F_{2n+1} - F_n F_{2n+2}}{F_n F_{2n+1}} = (-1)^n \frac{F_{n+1}}{F_n F_{2n+1}}. \tag{3.8}$$

We first assume that  $n$  is even. Then combining (3.8) and (3.3), we obtain

$$\frac{F_{n+1}}{F_n} - \frac{F_{mn+2}}{F_{mn+1}} \leq \frac{F_{n+1}}{F_n F_{2n+1}} < \frac{1}{F_{2n}-1}. \tag{3.9}$$

On the other hand, it follows from (3.1) and (3.5) that

$$\frac{F_{n+1}}{F_n} - \frac{1}{F_{2n}} - \frac{F_{mn+2}}{F_{mn+1}} > 0,$$

which means that

$$\frac{F_{n+1}}{F_n} - \frac{F_{mn+2}}{F_{mn+1}} > \frac{1}{F_{2n}}. \tag{3.10}$$

Now combining (3.7), (3.9), and (3.10), we deduce that

$$\frac{1}{F_{2n}} < \sum_{k=n}^{mn} \frac{(-1)^k}{F_k F_{k+1}} < \frac{1}{F_{2n} - 1},$$

which shows that the statement is true when  $n$  is even.

We now consider the case where  $n$  is odd. It is clearly true for  $n = 1$ , so we assume that  $n \geq 3$ . Applying (3.8) and (3.4), we can see that

$$\frac{F_{n+1}}{F_n} - \frac{F_{mn+2}}{F_{mn+1}} \leq -\frac{F_{n+1}}{F_n F_{2n+1}} < -\frac{1}{F_{2n} + 1}. \tag{3.11}$$

It follows from (3.2) and (3.5) that

$$\frac{F_{n+1}}{F_n} + \frac{1}{F_{2n}} - \frac{F_{mn+2}}{F_{mn+1}} = \frac{F_{2n+1}}{F_{2n}} - \frac{F_{mn+2}}{F_{mn+1}} > 0,$$

which means that

$$\frac{F_{n+1}}{F_n} - \frac{F_{mn+2}}{F_{mn+1}} > -\frac{1}{F_{2n}}. \tag{3.12}$$

Combining (3.7), (3.11), and (3.12), we get that

$$-\frac{1}{F_{2n}} < \sum_{k=n}^{mn} \frac{(-1)^k}{F_k F_{k+1}} < -\frac{1}{F_{2n} + 1},$$

which yields the desired identity. □

### 3.2 Alternating reciprocal sums of $F_{2k-1}F_{2k}$

For  $n \geq 2$ , we define

$$\begin{aligned} f(n) &= \frac{1}{3F_{2n-2}F_{2n-1}} - \frac{(-1)^n}{F_{2n-1}F_{2n}} - \frac{1}{3F_{2n}F_{2n+1}}, \\ g(n) &= \frac{1}{3F_{2n-2}F_{2n-1} + 1} - \frac{(-1)^n}{F_{2n-1}F_{2n}} - \frac{1}{3F_{2n}F_{2n+1} + 1}, \\ s(n) &= \frac{-1}{3F_{2n-2}F_{2n-1} + 1} - \frac{(-1)^n}{F_{2n-1}F_{2n}} + \frac{1}{3F_{2n}F_{2n+1} + 1}, \\ t(n) &= \frac{-1}{3F_{2n-2}F_{2n-1}} - \frac{(-1)^n}{F_{2n-1}F_{2n}} + \frac{1}{3F_{2n}F_{2n+1}}. \end{aligned}$$

It is not hard to check that  $f(n), g(n), s(n)$ , and  $t(n)$  are all negative if  $n$  is even and positive otherwise.

**Lemma 3.5** For  $n \geq 2$ , we have

$$f(n) + f(n + 1) > 0. \tag{3.13}$$

*Proof* The statement is clearly true when  $n$  is odd, so we assume that  $n$  is even in the rest of the proof. Applying (2.3), we derive that

$$\begin{aligned} f(n) + f(n + 1) &= \frac{1}{3F_{2n-2}F_{2n-1}} - \frac{1}{F_{2n-1}F_{2n}} + \frac{1}{F_{2n+1}F_{2n+2}} - \frac{1}{3F_{2n+2}F_{2n+3}} \\ &= \frac{1}{3} \left( \frac{1}{F_{2n-2}F_{2n-1}} - \frac{1}{F_{2n+2}F_{2n+3}} \right) - \left( \frac{1}{F_{2n-1}F_{2n}} - \frac{1}{F_{2n+1}F_{2n+2}} \right) \\ &= \frac{1}{3} \frac{F_{2n+2}F_{2n+3} - F_{2n-2}F_{2n-1}}{F_{2n-2}F_{2n-1}F_{2n+2}F_{2n+3}} - \frac{F_{2n+1}F_{2n+2} - F_{2n-1}F_{2n}}{F_{2n-1}F_{2n}F_{2n+1}F_{2n+2}} \\ &= \frac{1}{3} \frac{F_{2n+2}F_{2n+3} - F_{2n}F_{2n+1} + F_{2n}F_{2n+1} - F_{2n-2}F_{2n-1}}{F_{2n-2}F_{2n-1}F_{2n+2}F_{2n+3}} \\ &\quad - \frac{F_{2n+1}F_{2n+2} - F_{2n-1}F_{2n}}{F_{2n-1}F_{2n}F_{2n+1}F_{2n+2}} \\ &= \frac{1}{3} \frac{F_{4n+3} + F_{4n-1}}{F_{2n-2}F_{2n-1}F_{2n+2}F_{2n+3}} - \frac{F_{4n+1}}{F_{2n-1}F_{2n}F_{2n+1}F_{2n+2}} \\ &= \frac{1}{3} \frac{3F_{4n+1}}{F_{2n-2}F_{2n-1}F_{2n+2}F_{2n+3}} - \frac{F_{4n+1}}{F_{2n-1}F_{2n}F_{2n+1}F_{2n+2}} \\ &= \frac{F_{4n+1}}{F_{2n-1}F_{2n+2}} \left( \frac{1}{F_{2n-2}F_{2n+3}} - \frac{1}{F_{2n}F_{2n+1}} \right) \\ &= \frac{F_{4n+1}}{F_{2n-1}F_{2n+2}} \cdot \frac{F_{2n}F_{2n+1} - F_{2n-2}F_{2n+3}}{F_{2n-2}F_{2n}F_{2n+1}F_{2n+3}} \\ &= \frac{F_{4n+1}}{F_{2n-1}F_{2n+2}} \cdot \frac{2}{F_{2n-2}F_{2n}F_{2n+1}F_{2n+3}} \\ &> 0, \end{aligned}$$

where the last equality follows from (2.4). □

**Remark** From the proof of Lemma 3.5 we can easily derive that if  $n$  is odd, then

$$\begin{aligned} f(n) + f(n + 1) &= \frac{F_{4n+1}}{F_{2n-1}F_{2n+2}} \left( \frac{1}{F_{2n-2}F_{2n+3}} + \frac{1}{F_{2n}F_{2n+1}} \right) \\ &> \frac{F_{4n+1}}{F_{2n-1}F_{2n+2}} \cdot \frac{2}{F_{2n-2}F_{2n}F_{2n+1}F_{2n+3}}. \end{aligned}$$

Therefore, whether  $n$  is even or odd, we always have

$$f(n) + f(n + 1) \geq \frac{F_{4n+1}}{F_{2n-1}F_{2n+2}} \cdot \frac{2}{F_{2n-2}F_{2n}F_{2n+1}F_{2n+3}} > \frac{2}{F_{2n-2}F_{2n}F_{2n+1}F_{2n+3}}. \tag{3.14}$$

**Lemma 3.6** If  $n \geq 2$  and  $m \geq 2$ , we have

$$f(n) + f(n + 1) + f(mn) > 0.$$

*Proof* If  $mn$  is odd, then the result follows from (3.13) and the fact  $f(mn) > 0$ . Now we consider the case where  $mn$  is even. It is straightforward to check that

$$\begin{aligned} f(mn) &= \frac{1}{3F_{2mn-2}F_{2mn-1}} - \frac{1}{F_{2mn-1}F_{2mn}} - \frac{1}{3F_{2mn}F_{2mn+1}} \\ &= \frac{F_{2mn}F_{2mn+1} - F_{2mn-2}F_{2mn-1} - 3F_{2mn-2}F_{2mn+1}}{3F_{2mn-2}F_{2mn-1}F_{2mn}F_{2mn+1}} \\ &= \frac{F_{2mn}(F_{2mn} + F_{2mn-1}) - (F_{2mn} - F_{2mn-1})F_{2mn-1} - 3F_{2mn-2}F_{2mn+1}}{3F_{2mn-2}F_{2mn-1}F_{2mn}F_{2mn+1}} \\ &= \frac{F_{2mn}^2 + F_{2mn-1}^2 - 3(F_{2mn} - F_{2mn-1})(F_{2mn} + F_{2mn-1})}{3F_{2mn-2}F_{2mn-1}F_{2mn}F_{2mn+1}} \\ &= \frac{4F_{2mn-1}^2 - 2F_{2mn}^2}{3F_{2mn-2}F_{2mn-1}F_{2mn}F_{2mn+1}}. \end{aligned}$$

Invoking (2.4), we get

$$\begin{aligned} F_{2mn}^2 - 2F_{2mn-1}^2 &= (F_{2mn-1} + F_{2mn-2})^2 - 2F_{2mn-1}^2 \\ &= 2F_{2mn-2}F_{2mn-1} + F_{2mn-2}^2 - F_{2mn-1}^2 \\ &= 2F_{2mn-2}F_{2mn-1} - F_{2mn-3}F_{2mn} \\ &= F_{2mn-2}F_{2mn-1} - (F_{2mn-3}F_{2mn} - F_{2mn-2}F_{2mn-1}) \\ &= F_{2mn-2}F_{2mn-1} - (-1)^{2mn-2} \\ &= F_{2mn-2}F_{2mn-1} - 1. \end{aligned}$$

Therefore, we have

$$f(mn) = \frac{-2(F_{2mn-2}F_{2mn-1} - 1)}{3F_{2mn-2}F_{2mn-1}F_{2mn}F_{2mn+1}} > \frac{-2}{3F_{2mn}F_{2mn+1}}. \tag{3.15}$$

Combining (3.14) and (3.15), we obtain

$$\begin{aligned} f(n) + f(n + 1) + f(mn) &> \frac{2}{F_{2n-2}F_{2n}F_{2n+1}F_{2n+3}} - \frac{2}{3F_{2mn}F_{2mn+1}} \\ &\geq \frac{2}{F_{2n}F_{2n+1}F_{2n-2}F_{2n+3}} - \frac{2}{3F_{4n}F_{4n+1}}. \end{aligned}$$

Since  $F_{4n+1} > F_{4n}$  and

$$F_{4n} = F_{2n-1}F_{2n} + F_{2n}F_{2n+1} = F_{2n-3}F_{2n+2} + F_{2n-2}F_{2n+3},$$

we determine that

$$f(n) + f(n + 1) + f(mn) > 0.$$

The proof is completed. □

**Lemma 3.7** *If  $n \geq 2$  is even, we have*

$$g(n) + g(n + 1) < 0.$$

*Proof* From the proof of Lemma 3.5 we know that

$$\begin{aligned} g(n) + g(n + 1) &= \frac{1}{3F_{2n-2}F_{2n-1} + 1} - \frac{1}{F_{2n-1}F_{2n}} + \frac{1}{F_{2n+1}F_{2n+2}} - \frac{1}{3F_{2n+2}F_{2n+3} + 1} \\ &= \frac{3(F_{2n+2}F_{2n+3} - F_{2n-2}F_{2n-1})}{(3F_{2n-2}F_{2n-1} + 1)(3F_{2n+2}F_{2n+3} + 1)} - \left( \frac{1}{F_{2n-1}F_{2n}} - \frac{1}{F_{2n+1}F_{2n+2}} \right) \\ &= \frac{9F_{4n+1}}{(3F_{2n-2}F_{2n-1} + 1)(3F_{2n+2}F_{2n+3} + 1)} - \frac{9F_{4n+1}}{9F_{2n-1}F_{2n}F_{2n+1}F_{2n+2}}. \end{aligned}$$

For  $n \geq 2$ , we have

$$\begin{aligned} &(3F_{2n-2}F_{2n-1} + 1)(3F_{2n+2}F_{2n+3} + 1) \\ &> 9F_{2n-2}F_{2n-1}F_{2n+2}F_{2n+3} + 3F_{2n+2}F_{2n+3} \\ &= 9F_{2n-1}F_{2n+2}(F_{2n-2}F_{2n+3} - F_{2n}F_{2n+1}) \\ &\quad + 3F_{2n+2}F_{2n+3} + 9F_{2n-1}F_{2n}F_{2n+1}F_{2n+2} \\ &= 9F_{2n-1}F_{2n}F_{2n+1}F_{2n+2} + 3F_{2n+2}F_{2n+3} - 18F_{2n-1}F_{2n+2} \\ &= 9F_{2n-1}F_{2n}F_{2n+1}F_{2n+2} + 3F_{2n+2}(F_{2n+3} - 6F_{2n-1}) \\ &= 9F_{2n-1}F_{2n}F_{2n+1}F_{2n+2} + 3F_{2n+2}(3F_{2n-2} - F_{2n-1}) \\ &> 9F_{2n-1}F_{2n}F_{2n+1}F_{2n+2}, \end{aligned}$$

which implies

$$g(n) + g(n + 1) < 0.$$

This completes the proof. □

**Lemma 3.8** *If  $n > 0$  is even, then*

$$g(n) + \frac{1}{3F_{2n}F_{2n+1} + 1} < 0.$$

*Proof* The result follows from the definition of  $g(n)$  and the fact  $3F_n > F_{n+2}$ . □

To introduce the property of  $s(n)$ , we need two preliminary results.

**Lemma 3.9** *If  $n \geq 5$ , then*

$$F_{2n}F_{2n+1} > 3F_{n-1}F_nF_{n+1}F_{n+2}.$$

*Proof* It is easy to see that  $2F_n > F_{n+1}$  for  $n \geq 3$ . We claim that  $5F_n \geq 3F_{n+1}$  if  $n \geq 3$ . First, the claim holds for  $n = 3$  and  $n = 4$ . Now we assume that  $n \geq 5$ . It is straightforward to

verify that

$$5F_n = 3F_n + 2F_n = 3F_n + 2F_{n-1} + 2F_{n-2} > 3F_n + 2F_{n-1} + F_{n-1} = 3F_{n+1}.$$

Since  $5F_n \geq 3F_{n+1}$  and  $2F_{n+1} > F_{n+2}$ , we have  $5F_n > F_{n+3}$  for  $n \geq 3$ .

It follows from (2.1) that  $F_{2n} = F_{n-1}F_n + F_nF_{n+1}$  and

$$F_{2n+1} = F_{n-2}F_{n+2} + F_{n-1}F_{n+3} = F_{n-2}F_{n+2} + F_{n-1}F_{n+2} + F_{n-1}F_{n+1},$$

from which we derive that

$$\begin{aligned} \frac{F_{2n}F_{2n+1}}{F_{n-1}F_nF_{n+1}F_{n+2}} &= \frac{F_{2n}}{F_nF_{n+1}} \cdot \frac{F_{2n+1}}{F_{n-1}F_{n+2}} \\ &= \left(1 + \frac{F_{n-1}}{F_{n+1}}\right) \left(1 + \frac{F_{n-2}}{F_{n-1}} + \frac{F_{n+1}}{F_{n+2}}\right) \\ &= 1 + \frac{F_{n-2}}{F_{n-1}} + \frac{F_{n+1}}{F_{n+2}} + \frac{F_{n-1}}{F_{n+1}} + \frac{F_{n-2}}{F_{n+1}} + \frac{F_{n-1}}{F_{n+2}} \\ &= 1 + \frac{F_{n-2}}{F_{n-1}} + \frac{F_{n+1}}{F_{n+2}} + \frac{F_n}{F_{n+1}} + \frac{F_{n-1}}{F_{n+2}} \\ &> 1 + \frac{3}{5} + \frac{3}{5} + \frac{3}{5} + \frac{1}{5} \\ &= 3, \end{aligned}$$

which completes the proof. □

**Lemma 3.10** *For  $n \geq 5$ , we have*

$$F_{2n+1}(F_{n+3} - 6F_{n-1}) > (F_{n-2}F_{n-1} + 1)F_{n+3}.$$

*Proof* It is easy to see that  $F_{n-2}^2F_{n+2} > F_{n+3}$  for  $n \geq 5$ , and thus we have

$$\begin{aligned} F_{2n+1}F_{n-2} &= (F_{n-2}F_{n+2} + F_{n-1}F_{n+3})F_{n-2} \\ &= F_{n-2}^2F_{n+2} + F_{n-2}F_{n-1}F_{n+3} \\ &> F_{n+3} + F_{n-2}F_{n-1}F_{n+3} \\ &= (F_{n-2}F_{n-1} + 1)F_{n+3}. \end{aligned}$$

It is straightforward to check that  $F_{n+3} = 5F_{n-1} + 3F_{n-2}$ , from which we get

$$F_{n+3} - 6F_{n-1} = 3F_{n-2} - F_{n-1} > F_{n-2}.$$

Combining the last two inequalities yields the desired result. □

**Lemma 3.11** *If  $n \geq 3$  is odd, then*

$$s(n) + s(n + 1) - \frac{1}{3F_{4n}F_{4n+1} + 1} > 0.$$



*Proof* Since  $n$  is odd, we have

$$\begin{aligned}
 s(n) + s(n + 1) &= \frac{-1}{3F_{2n-2}F_{2n-1} + 1} + \frac{1}{F_{2n-1}F_{2n}} - \frac{1}{F_{2n+1}F_{2n+2}} + \frac{1}{3F_{2n+2}F_{2n+3} + 1} \\
 &= \left( \frac{1}{F_{2n-1}F_{2n}} - \frac{1}{F_{2n+1}F_{2n+2}} \right) - \left( \frac{1}{3F_{2n-2}F_{2n-1} + 1} - \frac{1}{3F_{2n+2}F_{2n+3} + 1} \right) \\
 &= \frac{F_{4n+1}}{F_{2n-1}F_{2n}F_{2n+1}F_{2n+2}} - \frac{9F_{4n+1}}{(3F_{2n-2}F_{2n-1} + 1)(3F_{2n+2}F_{2n+3} + 1)} \\
 &> \frac{F_{4n+1}}{F_{2n-1}F_{2n}F_{2n+1}F_{2n+2}} - \frac{3F_{4n+1}}{(3F_{2n-2}F_{2n-1} + 1)F_{2n+2}F_{2n+3}} \\
 &= \frac{1}{F_{2n-1}F_{2n}F_{2n+1}F_{2n+2}} \cdot \frac{F_{4n+1}(F_{2n+3} - 6F_{2n-1})}{(3F_{2n-2}F_{2n-1} + 1)F_{2n+3}} \\
 &> \frac{1}{F_{2n-1}F_{2n}F_{2n+1}F_{2n+2}} \cdot \frac{F_{4n+1}(F_{2n+3} - 6F_{2n-1})}{(3F_{2n-2}F_{2n-1} + 3)F_{2n+3}} \\
 &> \frac{1}{3F_{2n-1}F_{2n}F_{2n+1}F_{2n+2}},
 \end{aligned}$$

where the last inequality follows from Lemma 3.10.

Applying Lemma 3.9, we have

$$3F_{4n}F_{4n+1} + 1 > 3F_{4n}F_{4n+1} > 9F_{2n-1}F_{2n}F_{2n+1}F_{2n+2},$$

which implies that

$$\begin{aligned}
 s(n) + s(n + 1) - \frac{1}{3F_{4n}F_{4n+1} + 1} &> s(n) + s(n + 1) - \frac{1}{9F_{2n-1}F_{2n}F_{2n+1}F_{2n+2}} \\
 &> \frac{1}{3F_{2n-1}F_{2n}F_{2n+1}F_{2n+2}} - \frac{1}{9F_{2n-1}F_{2n}F_{2n+1}F_{2n+2}} \\
 &> 0,
 \end{aligned}$$

which completes the proof. □

Applying a similar analysis of  $f(n)$ , we can obtain the following properties of  $t(n)$ , whose proofs are omitted here.

**Lemma 3.12** For  $n \geq 2$ ,

$$t(n) + t(n + 1) < 0.$$

**Lemma 3.13** If  $n \geq 2$  and  $m \geq 2$ , we have

$$t(n) + t(n + 1) + t(mn) < 0.$$

**Theorem 3.14** If  $m \geq 2$  and  $n \geq 3$ , then

$$\left[ \left( \sum_{k=n}^{mn} \frac{(-1)^k}{F_{2k-1}F_{2k}} \right)^{-1} \right] = \begin{cases} 3F_{2n-2}F_{2n-1} & \text{if } n \text{ is even;} \\ -3F_{2n-2}F_{2n-1} - 1 & \text{if } n \text{ is odd.} \end{cases}$$

*Proof* We first consider the case where  $n$  is even. With the help of  $f(n)$ , we have

$$\sum_{k=n}^{mn} \frac{(-1)^k}{F_{2k-1}F_{2k}} = \frac{1}{3F_{2n-2}F_{2n-1}} - \frac{1}{3F_{2mn}F_{2mn+1}} - \sum_{k=n}^{mn} f(k).$$

Lemma 3.5 implies that

$$\sum_{k=n+2}^{mn-1} f(k) > 0.$$

Furthermore, applying Lemma 3.6, we get

$$\sum_{k=n}^{mn} f(k) = f(n) + f(n+1) + f(mn) + \sum_{k=n+2}^{mn-1} f(k) > 0.$$

Hence, we obtain

$$\sum_{k=n}^{mn} \frac{(-1)^k}{F_{2k-1}F_{2k}} < \frac{1}{3F_{2n-2}F_{2n-1}}. \tag{3.16}$$

It follows from Lemma 3.7 and Lemma 3.8 that

$$\begin{aligned} \sum_{k=n}^{mn} \frac{(-1)^k}{F_{2k-1}F_{2k}} &= \frac{1}{3F_{2n-2}F_{2n-1} + 1} - \frac{1}{3F_{2mn}F_{2mn+1} + 1} - \sum_{k=n}^{mn} g(k) \\ &= \frac{1}{3F_{2n-2}F_{2n-1} + 1} - \sum_{k=n}^{mn-1} g(k) - \left( g(mn) + \frac{1}{3F_{2mn}F_{2mn+1} + 1} \right) \\ &> \frac{1}{3F_{2n-2}F_{2n-1} + 1}. \end{aligned} \tag{3.17}$$

Combining (3.16) and (3.17) yields

$$\frac{1}{3F_{2n-2}F_{2n-1} + 1} < \sum_{k=n}^{mn} \frac{(-1)^k}{F_{2k-1}F_{2k}} < \frac{1}{3F_{2n-2}F_{2n-1}},$$

which shows that the statement is true when  $n$  is even.

Next, we turn to the case where  $n$  is odd. It follows from Lemma 3.11 that

$$\begin{aligned} \sum_{k=n}^{mn} \frac{(-1)^k}{F_{2k-1}F_{2k}} &= \frac{-1}{3F_{2n-2}F_{2n-1} + 1} + \frac{1}{3F_{2mn}F_{2mn+1} + 1} - \sum_{k=n}^{mn} s(k) \\ &= \frac{-1}{3F_{2n-2}F_{2n-1} + 1} - \left( s(n) + s(n+1) - \frac{1}{3F_{2mn}F_{2mn+1} + 1} \right) - \sum_{k=n+2}^{mn} s(k) \\ &< \frac{-1}{3F_{2n-2}F_{2n-1} + 1} - \left( s(n) + s(n+1) - \frac{1}{3F_{4n}F_{4n+1} + 1} \right) - \sum_{k=n+2}^{mn} s(k) \\ &< \frac{-1}{3F_{2n-2}F_{2n-1} + 1}. \end{aligned} \tag{3.18}$$

If  $mn$  is even, then it follows from Lemma 3.12 that  $\sum_{k=n}^{mn} t(k) < 0$ , and hence

$$\begin{aligned} \sum_{k=n}^{mn} \frac{(-1)^k}{F_{2k-1}F_{2k}} &= \frac{-1}{3F_{2n-2}F_{2n-1}} + \frac{1}{3F_{2mn}F_{2mn+1}} - \sum_{k=n}^{mn} t(k) \\ &> \frac{-1}{3F_{2n-2}F_{2n-1}}. \end{aligned}$$

If  $mn$  is odd, then it follows from Lemma 3.13 that

$$\begin{aligned} \sum_{k=n}^{mn} \frac{(-1)^k}{F_{2k-1}F_{2k}} &= \frac{-1}{3F_{2n-2}F_{2n-1}} + \frac{1}{3F_{2mn}F_{2mn+1}} - \sum_{k=n}^{mn} t(k) \\ &= \frac{-1}{3F_{2n-2}F_{2n-1}} + \frac{1}{3F_{2mn}F_{2mn+1}} - \sum_{k=n+2}^{mn-1} t(k) \\ &\quad - (t(n) + t(n+1) + t(mn)) \\ &> \frac{-1}{3F_{2n-2}F_{2n-1}}. \end{aligned}$$

Therefore, if  $n$  is odd, then we always have

$$\sum_{k=n}^{mn} \frac{(-1)^k}{F_{2k-1}F_{2k}} > \frac{-1}{3F_{2n-2}F_{2n-1}}. \tag{3.19}$$

It follows from (3.18) and (3.19) that

$$\frac{-1}{3F_{2n-2}F_{2n-1}} < \sum_{k=n}^{mn} \frac{(-1)^k}{F_{2k-1}F_{2k}} < \frac{-1}{3F_{2n-2}F_{2n-1} + 1},$$

which shows that the assertion for odd  $n$  also holds. □

Similarly, we can consider the alternating reciprocal sums of  $F_{2k}F_{2k+1}$  and obtain the following result, whose proof is similar to that of Theorem 3.14 and is omitted here.

**Theorem 3.15** *If  $m \geq 2$  and  $n \geq 2$ , then*

$$\left[ \left( \sum_{k=n}^{mn} \frac{(-1)^k}{F_{2k}F_{2k+1}} \right)^{-1} \right] = \begin{cases} 3F_{2n-1}F_{2n} - 1 & \text{if } n \text{ is even;} \\ -3F_{2n-1}F_{2n}, & \text{if } n \text{ is odd.} \end{cases}$$

### 3.3 Alternating sums of $F_{2k-1}F_{2k+1}$

We first introduce the following notation:

$$\begin{aligned} \alpha(n) &= \frac{1}{3F_{2n-1}^2} - \frac{(-1)^n}{F_{2n-1}F_{2n+1}} - \frac{1}{3F_{2n+1}^2}, \\ \beta(n) &= \frac{1}{3F_{2n-1}^2 - 1} - \frac{(-1)^n}{F_{2n-1}F_{2n+1}} - \frac{1}{3F_{2n+1}^2 - 1}, \\ \gamma(n) &= \frac{-1}{3F_{2n-1}^2 - 1} - \frac{(-1)^n}{F_{2n-1}F_{2n+1}} + \frac{1}{3F_{2n+1}^2 - 1}, \end{aligned}$$

$$\delta(n) = \frac{-1}{3F_{2n-1}^2} - \frac{(-1)^n}{F_{2n-1}F_{2n+1}} + \frac{1}{3F_{2n+1}^2}.$$

It is not hard to check that  $\alpha(n)$ ,  $\beta(n)$ ,  $\gamma(n)$ , and  $\delta(n)$  are all negative if  $n$  is even and positive otherwise.

**Lemma 3.16** *If  $n > 0$  is even, then*

$$\alpha(n) + \alpha(n + 1) < 0.$$

*Proof* Since  $n$  is even, we have

$$\begin{aligned} \alpha(n) + \alpha(n + 1) &= \frac{1}{3F_{2n-1}^2} - \frac{1}{F_{2n-1}F_{2n+1}} + \frac{1}{F_{2n+1}F_{2n+3}} - \frac{1}{3F_{2n+3}^2} \\ &= \frac{1}{3} \left( \frac{1}{F_{2n-1}^2} - \frac{1}{F_{2n+3}^2} \right) - \frac{1}{F_{2n+1}} \left( \frac{1}{F_{2n-1}} - \frac{1}{F_{2n+3}} \right) \\ &= \frac{1}{3} \left( \frac{1}{F_{2n-1}} - \frac{1}{F_{2n+3}} \right) \left( \frac{1}{F_{2n-1}} + \frac{1}{F_{2n+3}} - \frac{3}{F_{2n+1}} \right) \\ &= \frac{1}{3} \left( \frac{1}{F_{2n-1}} - \frac{1}{F_{2n+3}} \right) \left( \frac{1}{F_{2n+3}} - \frac{F_{2n-3}}{F_{2n-1}F_{2n+1}} \right) \\ &= \frac{1}{3} \left( \frac{1}{F_{2n-1}} - \frac{1}{F_{2n+3}} \right) \left( \frac{F_{2n-1}F_{2n+1} - F_{2n-3}F_{2n+3}}{F_{2n-1}F_{2n+1}F_{2n+3}} \right) \\ &= - \left( \frac{1}{F_{2n-1}} - \frac{1}{F_{2n+3}} \right) \frac{1}{F_{2n-1}F_{2n+1}F_{2n+3}} \\ &< 0, \end{aligned}$$

where the last equality follows from (2.4). □

**Lemma 3.17** *For  $n > 0$ ,*

$$6F_{2n-1}F_{2n+1} > (3F_{n-1}^2 - 1)F_{n+3}^2.$$

*Proof* It is easy to see that the result holds when  $n < 5$ . Next we show that, for  $n \geq 5$ ,

$$2F_{2n-1}F_{2n+1} > F_{2n}F_{2n+1} > F_{n-1}^2F_{n+3}^2,$$

from which the desired result follows.

The first inequality is obvious. It follows from (2.1) that

$$\begin{aligned} F_{2n} &= F_{n-2}F_{n+3} + F_{n-3}F_{n+2}, \\ F_{2n+1} &= F_{n-1}F_{n+3} + F_{n-2}F_{n+2}, \end{aligned}$$

which implies that

$$\begin{aligned} F_{2n}F_{2n+1} &= F_{n-2}F_{n-1}F_{n+3}^2 + F_{n-2}^2F_{n+2}F_{n+3} + F_{n-3}F_{n-1}F_{n+2}F_{n+3} + F_{n-3}F_{n-2}F_{n+2}^2 \\ &= F_{n-2}F_{n-1}F_{n+3}^2 + (F_{n-3}F_{n-1} - (-1)^n)F_{n+2}F_{n+3} + F_{n-3}F_{n-1}F_{n+2}F_{n+3} \end{aligned}$$

$$\begin{aligned}
 &+ F_{n-3}F_{n-2}F_{n+2}^2 \\
 &= F_{n-2}F_{n-1}F_{n+3}^2 + 2F_{n-3}F_{n-1}F_{n+2}F_{n+3} + F_{n-3}F_{n-2}F_{n+2}^2 - (-1)^n F_{n+2}F_{n+3} \\
 &> F_{n-2}F_{n-1}F_{n+3}^2 + F_{n-3}F_{n-1}F_{n+3}^2 + F_{n-3}F_{n-2}F_{n+2}^2 - (-1)^n F_{n+2}F_{n+3} \\
 &= F_{n-1}^2F_{n+3}^2 + F_{n-3}F_{n-2}F_{n+2}^2 - (-1)^n F_{n+2}F_{n+3} \\
 &> F_{n-1}^2F_{n+3}^2,
 \end{aligned}$$

where the last inequality follows from the fact that, for  $n \geq 5$ ,

$$F_{n-3}F_{n-2}F_{n+2} \geq 2F_{n+2} > F_{n+3}.$$

This completes the proof. □

**Lemma 3.18** For  $n \geq 1$ ,

$$\beta(n) + \beta(n + 1) > 0.$$

*Proof* It is obviously true when  $n$  is odd, so we assume that  $n$  is even. Now we have

$$\begin{aligned}
 \beta(n) + \beta(n + 1) &= \left( \frac{1}{3F_{2n-1}^2 - 1} - \frac{1}{3F_{2n+3}^2 - 1} \right) - \left( \frac{1}{F_{2n-1}F_{2n+1}} - \frac{1}{F_{2n+1}F_{2n+3}} \right) \\
 &= \frac{3(F_{2n+3}^2 - F_{2n-1}^2)}{(3F_{2n-1}^2 - 1)(3F_{2n+3}^2 - 1)} - \frac{F_{2n+3} - F_{2n-1}}{F_{2n-1}F_{2n+1}F_{2n+3}} \\
 &= \frac{9F_{2n+1}(F_{2n+3} - F_{2n-1})}{(3F_{2n-1}^2 - 1)(3F_{2n+3}^2 - 1)} - \frac{F_{2n+3} - F_{2n-1}}{F_{2n-1}F_{2n+1}F_{2n+3}}.
 \end{aligned}$$

Since

$$\begin{aligned}
 &(3F_{2n-1}^2 - 1)(3F_{2n+3}^2 - 1) \\
 &= 9F_{2n-1}^2F_{2n+3}^2 - 3F_{2n-1}^2 - 3F_{2n+3}^2 + 1 \\
 &= 9F_{2n-1}F_{2n+3}(F_{2n+1}^2 + 1) - 3F_{2n-1}^2 - 3F_{2n+3}^2 + 1 \\
 &= 9F_{2n-1}F_{2n+1}F_{2n+3} + 9F_{2n-1}F_{2n+3} - 3F_{2n-1}^2 - 3F_{2n+3}^2 + 1 \\
 &< 9F_{2n-1}F_{2n+1}F_{2n+3} + 9F_{2n-1}F_{2n+3} - 3F_{2n+3}^2 \\
 &= 9F_{2n-1}F_{2n+1}F_{2n+3} + 3F_{2n+3}(3F_{2n-1} - F_{2n+3}) \\
 &= 9F_{2n-1}F_{2n+1}F_{2n+3} - 3F_{2n+3}(2F_{2n-1} + 3F_{2n-2}) \\
 &< 9F_{2n-1}F_{2n+1}F_{2n+3},
 \end{aligned}$$

we have

$$\beta(n) + \beta(n + 1) > 0,$$

which completes the proof. □

**Remark** From the proof of Lemma 3.18 we can derive that if  $n$  is odd, then

$$\begin{aligned}
 \beta(n) + \beta(n + 1) &= \left( \frac{1}{3F_{2n-1}^2 - 1} - \frac{1}{3F_{2n+3}^2 - 1} \right) + \left( \frac{1}{F_{2n-1}F_{2n+1}} - \frac{1}{F_{2n+1}F_{2n+3}} \right) \\
 &> \left( \frac{1}{3F_{2n-1}^2 - 1} - \frac{1}{3F_{2n+3}^2 - 1} \right) - \left( \frac{1}{F_{2n-1}F_{2n+1}} - \frac{1}{F_{2n+1}F_{2n+3}} \right) \\
 &= \frac{9F_{2n+1}(F_{2n+3} - F_{2n-1})}{(3F_{2n-1}^2 - 1)(3F_{2n+3}^2 - 1)} - \frac{F_{2n+3} - F_{2n-1}}{F_{2n-1}F_{2n+1}F_{2n+3}} \\
 &> \frac{9F_{2n+1}^2}{(3F_{2n-1}^2 - 1)(3F_{2n+3}^2 - 1)} - \frac{F_{2n+1}}{F_{2n-1}F_{2n+1}F_{2n+3}} \\
 &= \frac{9F_{2n+1}^2}{(3F_{2n-1}^2 - 1)(3F_{2n+3}^2 - 1)} - \frac{1}{F_{2n-1}F_{2n+3}} \\
 &> \frac{3F_{2n+1}^2}{(3F_{2n-1}^2 - 1)F_{2n+3}^2} - \frac{1}{F_{2n-1}F_{2n+3}} \\
 &= \frac{3F_{2n-1}(F_{2n+1}^2 - F_{2n-1}F_{2n+3}) + F_{2n+3}}{(3F_{2n-1}^2 - 1)F_{2n-1}F_{2n+3}^2} \\
 &= \frac{F_{2n+3} - 3F_{2n-1}}{(3F_{2n-1}^2 - 1)F_{2n-1}F_{2n+3}^2} \\
 &= \frac{2F_{2n} + F_{2n-2}}{(3F_{2n-1}^2 - 1)F_{2n-1}F_{2n+3}^2}.
 \end{aligned}$$

Thus, we have that, for all  $n > 0$ ,

$$\beta(n) + \beta(n + 1) > \frac{2F_{2n} + F_{2n-2}}{(3F_{2n-1}^2 - 1)F_{2n-1}F_{2n+3}^2} > \frac{2}{(3F_{2n-1}^2 - 1)F_{2n+3}^2}. \tag{3.20}$$

**Lemma 3.19** *If  $n \geq 1$  and  $m \geq 2$ , we have*

$$\beta(n) + \beta(n + 1) + \beta(mn) > 0.$$

*Proof* If  $mn$  is odd, then the result follows from the facts  $\beta(mn) > 0$  and  $\beta(n) + \beta(n + 1) > 0$ . Next, we focus ourselves on the case where  $mn$  is even. It is easy to see that

$$\begin{aligned}
 \beta(mn) &= \frac{1}{3F_{2mn-1}^2 - 1} - \frac{1}{F_{2mn-1}F_{2mn+1}} - \frac{1}{3F_{2mn+1}^2 - 1} \\
 &= \frac{3F_{2mn+1}^2 - 3F_{2mn-1}^2}{(3F_{2mn-1}^2 - 1)(3F_{2mn+1}^2 - 1)} - \frac{1}{F_{2mn-1}F_{2mn+1}} \\
 &> \frac{3F_{2mn+1}^2 - 3F_{2mn-1}^2}{3F_{2mn-1}^2 \cdot 3F_{2mn+1}^2} - \frac{1}{F_{2mn-1}F_{2mn+1}} \\
 &= \frac{F_{2mn+1}^2 - F_{2mn-1}^2}{3F_{2mn-1}^2 F_{2mn+1}^2} - \frac{1}{F_{2mn-1}F_{2mn+1}}.
 \end{aligned}$$

Furthermore, since

$$\begin{aligned}
 F_{2mn+1}^2 - F_{2mn-1}^2 &= F_{2mn+1}(F_{2mn-2} + 2F_{2mn-1}) - F_{2mn-1}^2 \\
 &= 2F_{2mn-1}F_{2mn+1} + F_{2mn-2}(F_{2mn-1} + F_{2mn}) - F_{2mn-1}^2
 \end{aligned}$$

$$\begin{aligned}
 &= 2F_{2mn-1}F_{2mn+1} + F_{2mn-2}F_{2mn-1} + (F_{2mn-2}F_{2mn} - F_{2mn-1}^2) \\
 &= 2F_{2mn-1}F_{2mn+1} + F_{2mn-2}F_{2mn-1} + (-1)^{2mn-1} \\
 &> 2F_{2mn-1}F_{2mn+1},
 \end{aligned}$$

we have

$$\beta(mn) > \frac{2F_{2mn-1}F_{2mn+1}}{3F_{2mn-1}^2F_{2mn+1}^2} - \frac{1}{F_{2mn-1}F_{2mn+1}} = -\frac{1}{3F_{2mn-1}F_{2mn+1}}.$$

From (3.20) we see that

$$\begin{aligned}
 \beta(n) + \beta(n + 1) + \beta(mn) &> \frac{2}{(3F_{2n-1}^2 - 1)F_{2n+3}^2} - \frac{1}{3F_{2mn-1}F_{2mn+1}} \\
 &\geq \frac{2}{(3F_{2n-1}^2 - 1)F_{2n+3}^2} - \frac{1}{3F_{4n-1}F_{4n+1}} \\
 &> 0,
 \end{aligned}$$

where the last inequality follows from Lemma 3.17. □

Applying a similar analysis of  $\beta(n)$ , we can obtain the following properties of  $\gamma(n)$ , and the details are left as an exercise.

**Lemma 3.20** *For  $n \geq 1$ , we have*

$$\gamma(n) + \gamma(n + 1) < 0.$$

**Lemma 3.21** *If  $n \geq 1$  and  $m \geq 2$ , then*

$$\gamma(n) + \gamma(n + 1) + \gamma(mn) < 0.$$

**Lemma 3.22** *If  $n \geq 1$  is odd, then we have*

$$\delta(n) + \delta(n + 1) - \frac{1}{3F_{4n+1}^2} > 0.$$

*Proof* Since  $n$  is odd, we have

$$\delta(n) + \delta(n + 1) = -\frac{1}{3F_{2n-1}^2} + \frac{1}{F_{2n-1}F_{2n+1}} - \frac{1}{F_{2n+1}F_{2n+3}} + \frac{1}{3F_{2n+3}^2}.$$

Applying the argument in the proof of Lemma 3.16, we obtain

$$\delta(n) + \delta(n + 1) = \frac{F_{2n+3} - F_{2n-1}}{F_{2n-1}F_{2n+1}F_{2n+3}^2} > \frac{1}{F_{2n-1}F_{2n+3}^2}.$$

Since  $F_{4n+1} = F_{2n-2}F_{2n+2} + F_{2n-1}F_{2n+3}$ , we have  $F_{4n+1} \geq F_{2n-1}F_{2n+3}$ . Thus,

$$3F_{4n+1}^2 > F_{2n-1}^2F_{2n+3}^2.$$

Combining the last two inequalities yields the desired result. □

**Theorem 3.23** *If  $m \geq 2$  and  $n \geq 2$ , then*

$$\left[ \left( \sum_{k=n}^{mn} \frac{(-1)^k}{F_{2k-1}F_{2k+1}} \right)^{-1} \right] = \begin{cases} 3F_{2n-1}^2 - 1 & \text{if } n \text{ is even;} \\ -3F_{2n-1}^2 & \text{if } n \text{ is odd.} \end{cases}$$

*Proof* We first consider the case where  $n$  is even. Now we have

$$\alpha(mn) + \frac{1}{3F_{2mn+1}^2} = \frac{1}{3F_{2mn-1}^2} - \frac{1}{F_{2mn-1}F_{2mn+1}} < 0,$$

where the inequality follows from the fact  $3F_{n-1} > F_{n+1}$ .

Combining Lemma 3.16 and the last inequality, we derive that

$$\begin{aligned} \sum_{k=n}^{mn} \frac{(-1)^k}{F_{2k-1}F_{2k+1}} &= \frac{1}{3F_{2n-1}^2} - \frac{1}{3F_{2mn+1}^2} - \sum_{k=n}^{mn} \alpha(k) \\ &= \frac{1}{3F_{2n-1}^2} - \sum_{k=n}^{mn-1} \alpha(k) - \left( \alpha(mn) + \frac{1}{3F_{2mn+1}^2} \right) \\ &> \frac{1}{3F_{2n-1}^2}. \end{aligned}$$

With the help of  $\beta(n)$ , Lemma 3.18, and Lemma 3.19, we get

$$\begin{aligned} \sum_{k=n}^{mn} \frac{(-1)^k}{F_{2k-1}F_{2k+1}} &= \frac{1}{3F_{2n-1}^2 - 1} - \frac{1}{3F_{2mn+1}^2 - 1} - \sum_{k=n}^{mn} \beta(k) \\ &< \frac{1}{3F_{2n-1}^2 - 1} - (\beta(n) + \beta(n+1) + \beta(mn)) - \sum_{k=n+2}^{mn-1} \beta(k) \\ &< \frac{1}{3F_{2n-1}^2 - 1}. \end{aligned}$$

Therefore, we arrive at

$$\frac{1}{3F_{2n-1}^2} < \sum_{k=n}^{mn} \frac{(-1)^k}{F_{2k-1}F_{2k+1}} < \frac{1}{3F_{2n-1}^2 - 1},$$

which shows that the statement is true when  $n$  is even.

We now turn to the case where  $n$  is odd. We have

$$\sum_{k=n}^{mn} \frac{(-1)^k}{F_{2k-1}F_{2k+1}} = \frac{-1}{3F_{2n-1}^2 - 1} + \frac{1}{3F_{2mn+1}^2 - 1} - \sum_{k=n}^{mn} \gamma(k).$$

If  $mn$  is even, we easily see that

$$\sum_{k=n}^{mn} \gamma(k) < 0$$



by Lemma 3.20. Therefore,

$$\sum_{k=n}^{mn} \frac{(-1)^k}{F_{2k-1}F_{2k+1}} > \frac{-1}{3F_{2n-1}^2 - 1}.$$

If  $mn$  is odd, then employing Lemma 3.20 and Lemma 3.21, we deduce

$$\begin{aligned} \sum_{k=n}^{mn} \frac{(-1)^k}{F_{2k-1}F_{2k+1}} &= \frac{-1}{3F_{2n-1}^2 - 1} + \frac{1}{3F_{2mn+1}^2 - 1} - \sum_{k=n+2}^{mn-1} \gamma(k) \\ &\quad - (\gamma(n) + \gamma(n+1) + \gamma(mn)) \\ &> \frac{-1}{3F_{2n-1}^2 - 1}. \end{aligned}$$

Thus, we always have

$$\sum_{k=n}^{mn} \frac{(-1)^k}{F_{2k-1}F_{2k+1}} > \frac{-1}{3F_{2n-1}^2 - 1},$$

provided that  $n$  is odd.

Since  $n$  is odd, it follows from Lemma 3.22 that

$$\sum_{k=n+2}^{mn} \delta(k) > 0.$$

Furthermore, applying the last inequality and Lemma 3.22, we derive that

$$\begin{aligned} \sum_{k=n}^{mn} \frac{(-1)^k}{F_{2k-1}F_{2k+1}} &= \frac{-1}{3F_{2n-1}^2} + \frac{1}{3F_{2mn+1}^2} - \sum_{k=n}^{mn} \delta(k) \\ &= \frac{-1}{3F_{2n-1}^2} - \sum_{k=n+2}^{mn} \delta(k) - \left( \delta(n) + \delta(n+1) - \frac{1}{3F_{2mn+1}^2} \right) \\ &< \frac{-1}{3F_{2n-1}^2} - \left( \delta(n) + \delta(n+1) - \frac{1}{3F_{4n+1}^2} \right) \\ &< \frac{-1}{3F_{2n-1}^2}. \end{aligned}$$

Therefore, when  $n$  is odd, we have

$$\frac{-1}{3F_{2n-1}^2 - 1} < \sum_{k=n}^{mn} \frac{(-1)^k}{F_{2k-1}F_{2k+1}} < \frac{-1}{3F_{2n-1}^2},$$

which yields the desired identity. □

Similarly, we can prove the following result, whose proof is omitted here.

**Theorem 3.24** *If  $m \geq 2$  and  $n \geq 2$ , then*

$$\left[ \left( \sum_{k=n}^{mn} \frac{(-1)^k}{F_{2k}F_{2k+2}} \right)^{-1} \right] = \begin{cases} 3F_{2n}^2 & \text{if } n \text{ is even;} \\ -3F_{2n}^2 - 1 & \text{if } n \text{ is odd.} \end{cases}$$

## 4 Conclusions

In this paper, we investigate the sums and alternating sums of the products of two reciprocal Fibonacci numbers in various ways. The results are new and interesting. In particular, the techniques for dealing with alternating sums can be applied to study other types of alternating sums, which will be presented in a future paper.

### Competing interests

The authors declare that they have no competing interests.

### Authors' contributions

Both authors contributed equally to deriving all the results of this article and read and approved the final manuscript.

### Author details

<sup>1</sup>School of Communication and Information Engineering, University of Electronic Science and Technology of China, Chengdu, 611731, P.R. China. <sup>2</sup>School of Mathematical Sciences, University of Electronic Science and Technology of China, Chengdu, 611731, P.R. China.

### Authors' information

R Liu is an undergraduate student supervised by A Wang in the Research Informed Teaching Activity for Discrete Mathematics. A Wang is an associate professor in Applied Mathematics, whose research interests include Combinatorics and Graph Theory.

### Acknowledgements

The authors would like to thank the anonymous reviewers for their helpful comments. This work was supported by the National Natural Science Foundation of China (No. 11401080).

Received: 4 February 2016 Accepted: 10 May 2016 Published online: 18 May 2016

### References

- Duncan, RL: Applications of uniform distribution to the Fibonacci numbers. *Fibonacci Q.* **5**, 137-140 (1967)
- Karaduman, E: An application of Fibonacci numbers in matrices. *Appl. Math. Comput.* **147**, 903-908 (2004)
- Koshy, T: *Fibonacci and Lucas Numbers with Applications*. Wiley, New York (2001)
- Vorobiev, NN: *Fibonacci Numbers*. Springer, Basel (2002)
- Elsner, C, Shimomura, S, Shiokawa, I: Algebraic relations for reciprocal sums of Fibonacci numbers. *Acta Arith.* **130**, 37-60 (2007)
- Elsner, C, Shimomura, S, Shiokawa, I: Algebraic relations for reciprocal sums of odd terms in Fibonacci numbers. *Ramanujan J.* **17**, 429-446 (2008)
- Elsner, C, Shimomura, S, Shiokawa, I: Algebraic independence results for reciprocal sums of Fibonacci numbers. *Acta Arith.* **148**, 205-223 (2011)
- Elsner, C, Shimomura, S, Shiokawa, I: Algebraic relations for reciprocal sums of even terms in Fibonacci numbers. *J. Math. Sci.* **180**, 650-671 (2012)
- Ohstuka, H, Nakamura, S: On the sum of reciprocal Fibonacci numbers. *Fibonacci Q.* **46/47**, 153-159 (2008/2009)
- Wang, AYZ, Wen, PB: On the partial finite sums of the reciprocals of the Fibonacci numbers. *J. Inequal. Appl.* **2015**, Article ID 73 (2015)
- Wang, AYZ, Zhang, F: The reciprocal sums of even and odd terms in the Fibonacci sequence. *J. Inequal. Appl.* **2015**, Article ID 376 (2015)
- Wang, AYZ, Zhang, F: The reciprocal sums of the Fibonacci 3-subsequences. *Adv. Differ. Equ.* **2016**, Article ID 27 (2016)
- Stone, RR: General identities for Fibonacci and Lucas numbers with polynomial subscripts in several variables. *Fibonacci Q.* **13**, 289-294 (1975)

Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](http://springeropen.com)