

Oxytocin induced cAMP-dependent protein kinase activation and urokinase-type plasminogen activator production in LLC-PK₁ renal epithelial cells is mediated by the vasopressin V₂-receptor

David A. Jans*, Imre Pavo** and Falk Fahrenholz

Max-Planck-Institut für Biophysik, Kennedyallee 70, D-6000 Frankfurt am Main 70, Germany

Received 20 October 1992; revised version received 16 November 1992

Using a variety of peptide analogues of oxytocin (OT) and Arg⁸-vasopressin (AVP), OT-mediated induction of urokinase-type plasminogen activator (uPA) was examined in LLC-PK₁ renal epithelial cells, which possess distinct high-affinity receptors of both the OT- and vasopressin renal (V₂-) types. OT or OT-receptor specific agonists induced concentration-dependent cAMP synthesis, activation of the cAMP-dependent protein kinase (cAMP-PK) and uPA production consistent with their respective binding affinities for the V₂- and not the OT-receptor. OT-mediated uPA induction could be inhibited in a concentration-dependent fashion by coincubation with a V₂/V₁-receptor specific antagonist, but not by an OT-receptor specific antagonist. Results implied that stimulation of cAMP- and uPA responses in LLC-PK₁ cells by OT was V₂-receptor-mediated.

Neurohypophyseal nonapeptide hormone; Specific antagonist; Vasopressin V₂-receptor; Urokinase-type plasminogen activator

1. INTRODUCTION

The specific binding of the neurohypophyseal nonapeptide oxytocin (OT) to high-affinity receptors in myometrium [1–3] leads to inositol-trisphosphate turnover and Ca²⁺ influx [4,5], responses which are similar to those induced by the binding of the other neurohypophyseal nonapeptide, vasopressin, to the hepatic vasopressin V₁-receptor [6,7]. Interestingly, OT can have either diuretic or antidiuretic osmoregulatory effects depending on the presence or absence of vasopressin [8–10], which may be explained by the ability of OT to bind to the adenylate cyclase (AC)-stimulating vasopressin renal V₂-type receptor present on distal tubules and collecting ducts [9,10]. OT has also been reported to enhance glomerular filtration rate [8] and to have a natriuretic effect [8,9], both of which are OT-

specific [8]. Stimulation of glomerular filtration is probably mediated by OT-receptors in the macula densa [11].

The LLC-PK₁ renal epithelial cell line has been reported to possess low density high-affinity OT-receptors [12,13], in addition to V₂-receptors (about 45,000/cell) [14], which couple to AC stimulation [14,15], activation of the cAMP-dependent protein kinase (cAMP-PK) [14–16], and production of the extracellularly secreted protease urokinase-type plasminogen activator (uPA) [14,15,17]. In addition to agents elevating intracellular cAMP, phorbol esters stimulate LLC-PK₁ cells to produce uPA by a cAMP-independent Ca²⁺/phospholipid-dependent protein kinase (PK-C)-mediated pathway [14,18–21].

We were interested in the second messenger pathway of OT-mediated uPA induction in LLC-PK₁ cells. We show here that OT is capable of stimulating cAMP synthesis, cAMP-PK activation and uPA induction in LLC-PK₁ cells. All of these responses, however, appear to be mediated by OT-binding to the V₂-receptor.

Correspondence address: F. Fahrenholz, Max-Planck-Institut für Biophysik, Kennedyallee 70, D-6000 Frankfurt am Main 70, Germany. Fax: (49) (69) 630 3244.

**Current address:* Institut für Medizinische Physik, Westfälische Wilhelms-Universität, Münster, Germany.

***Current address:* Endocrine Unit of the 1st Department of Medicine, Albert Szent-Györgyi University, Szeged, Hungary.

Abbreviations: cAMP-PK, cAMP-dependent protein kinase or ATP:protein phosphotransferase (EC 2.7.1.37); AC, adenylate cyclase or ATP pyrophosphatylase (cyclicizing, EC 4.6.1.1); AVP, Arg⁸-vasopressin; OT, oxytocin; uPA, urokinase-type plasminogen activator (EC 3.4.21.31); IBMX, 1-isobutyl-3-methylxanthine.

2. MATERIALS AND METHODS

2.1. Materials

5'-γ-[³²P]ATP, [³H]Arg⁸-vasopressin (AVP) and [³H]OT were from Amersham, and phosphocellulose paper (P-81) from Whatman. The OT and AVP analogues used, listed in Table I, were synthesized as described [3,22–25]. Other materials were from previously described sources [14,26].

2.2. Cell culture

Cells of the LLC-PK₁ pig kidney epithelial cell line [27] and the V₂-receptor deficient mutant M18 [14] were cultured as described previously [14,26].




Fig. 1. (A) Concentration dependence of uPA production by LLC-PK₁ cells in response to OT and AVP analogues. Cell monolayers were washed extensively and then incubated for 8 h with agonists in serum-free DMEM as indicated: O (oxytocin), sO (dSOT), V (AVP), sV (SAVP), na (no addition), and the numbers represent the molar concentrations in negative log form. Medium (5 μ l) was spotted onto the casein-plasminogen agar, and, after incubation at 37°C for 2–3 h, uPA activity became visible as zones of lysis in the agar. (B) uPA production by cells of the LLC-PK₁ and V₂-receptor deficient mutant M18 cell lines in response to OT and AVP analogues. Cell monolayers of the LLC-PK₁ and M18 (18) cell lines were treated as in Fig. 1A.

2.3. Enzyme assays

Extracts for the assay of cAMP-PK catalytic activity were prepared and assayed as previously, using Kemptide (L-R-R-A-S-A-G) as substrate [14,26]. The cAMP-PK activity ratio expresses the C-subunit activity present in cell extracts (assayed in the absence of cAMP) relative to the total stimutable activity (assayed in the presence of cAMP) [14,26]. The ratio estimates the extent of cAMP-PK activation induced by different agents elevating intracellular cAMP levels [26,28,29]. uPA activity was detected using the previously described spot test [30,31], with activity indicated by zones of proteolytic clearing in the casein, due to plasmin activation. No proteolytic activity was observed in the absence of plasminogen. Protein was estimated using the dye binding assay of Bradford [32] with BSA (fatty-acid-free) as standard.

2.4. cAMP determination

Cells were treated with hormones and other agonists for 8 h as previously described [33]. Medium samples were then treated as described [25,26], and cAMP determined using the competitive protein binding assay of Tovey et al. [34].

2.5. Receptor binding

Vasopressin and oxytocin binding by EDTA-suspended cells was measured at 30°C as described [14,35]. Dissociation constants (K_D), the concentration of hormone corresponding to 50% maximal binding, were determined from competition binding experiments [3,14,23].

3. RESULTS

Maximal specific [³H]AVP and [³H]OT binding (10⁻⁸ M ligand) by LLC-PK₁ cells was 223 ± 24 ($n = 4$) and

Table I

Dissociation constants of the neurohypophyseal structural analogues used in this study for [³H]AVP and [³H]OT in LLC-PK₁ cells

Peptide/description [reference]	K_D for binding sites* (nM) for:	
	[³ H]AVP	[³ H]OT
[Arg ⁸]vasopressin (AVP)	1.4 ± 0.6	4.6
Oxytocin (OT)	776	2.8 ± 0.4
[Mpa ¹ ,Sar ⁷]OT (dSOT)/OT-agonist [3,22]	820	4.5 ± 0.4
[Thr ⁴ ,Sar ⁷]OT (TSOT)/OT-agonist [3]	>103	ND*
[Sar ⁷]AVP (SAVP)/V ₂ -agonist [23]	3.6 ± 0.9	78
[Mpa ¹ ,Val ⁴ ,Sar ⁷]AVP (dVSAVP)/V ₂ -agonist [22]	ND*	ND*
[Mca ¹ ,MeAla ⁷]AVP (MMAVP)/V ₁ /OT-antagonist [23]	101	ND*
[Mca ¹ ,D-Phe ² ,Sar ⁷]AVP (MFSAVP)/V ₁ /OT-antagonist [23,24]	52 ± 8	4.4 ± 1.0
[Mca ¹ ,D-Phe ² ,Ile ⁴ ,Lys ⁸]AVP (MFIKAVP)/V ₂ /V ₁ -antagonist [25]	ND*	ND*
[Mca ¹ ,Sar ⁷ ,Orn ⁸]OT (MSOOT)/OT-antagonist ^a	6,510	40

* K_D was determined from competition binding studies with [³H]AVP and [³H]OT as described [23–25], using affinity constants (Scatchard analysis) for [³H]AVP and [³H]OT in LLC-PK₁ cells of 3.5 [14] and 1.9 [12], respectively. Results represent the mean from at least two separate experiments ($n = 3$ where S.D. is indicated).

*Not determined.

^aInhibitory activity in rat uterus in vitro: $pA_2 = 7.34$ (Mg²⁺-free medium); a detailed description of the synthesis and pharmacological properties of this OT-antagonist will be published elsewhere.




Fig. 2. uPA production by LLC-PK₁ cells stimulated by OT or dSOT in the absence (na) or presence of OT- (O), V₂/V₁- (V2) or V₁/OT- (V1) specific antagonists. Cell monolayers were treated as in the legend to Fig. 1A,B, in the presence of differing concentrations (designated as per Fig. 1A,B) of the antagonists MAAVP (V1), MFIKAVP (V2) or MSOOT (O).

34 ± 8.5 ($n = 4$) fmol/mg respectively. Binding was further characterized by competition with the various AVP and OT analogues listed in Table I. The dissociation constants (K_D) calculated from the displacement binding curves (not shown) are also listed. Reduced competition of [³H]AVP binding was exhibited by OT, OT-receptor and V₁-specific ligands (15- to 15,000-fold difference in K_D), compared to the V₂-specific agonists SAVP and dVSAVP (Table I, and not shown). This implied that [³H]AVP binds predominantly to the V₂-receptor of LLC-PK₁ cells.

[³H]OT binding could be competed quite well by OT- and V₁-specific ligands compared to the V₂-agonist SAVP (about a 20-fold difference in K_D , Table I). AVP, however, also competed reasonably well for [³H]OT-binding sites (K_D of 4.6 nM compared to 2.8 nM for OT itself), implying that the OT-receptor of LLC-PK₁ cells can bind AVP, consistent with similar observations for OT-receptors from other tissues [3,36,37]. In conclusion, the V₂-receptor accounted for about 95% of the

specific [³H]AVP binding at 10^{-8} M ligand (less than 5% of the binding likely to be due to binding to the OT-receptor); whilst the OT-receptor accounted for about 99% of the specific [³H]OT binding at 10^{-8} M OT (Table I, and not shown).

3.1. cAMP production and cAMP-PK activation in LLC-PK₁ cells in response to OT

We tested the analogues from Table I for the stimulation of cAMP production and cAMP-PK activation in LLC-PK₁ cells (Table IIA,B). Results for the non-receptor-mediated AC-activator forskolin are shown for comparison. Whilst even high concentrations of the OT-agonists dSOT and TSOT only very weakly stimulated cAMP synthesis (Table IIA, and not shown) and cAMP-PK activation (Table IIB, and not shown), high concentrations of OT induced marked responses. Whereas half-maximal responses were elicited at about 10^{-7} M OT (Table IIA,B, and not shown) [12], half-maximal cAMP production and cAMP-PK activation in response to AVP are at about 10^{-9} M (Table IIA,B) [12,37,38]. The V₂-specific agonists dVSAVP and SAVP showed similar K_a s (Table IIA,B, and not shown) [25]. Thus, although OT and to a minor extent OT-specific agonists were capable of inducing cAMP-mediated responses, the concentration-dependence of the responses was consistent with their respective affinities for the V₂-receptor of LLC-PK₁ cells. In contrast, half-maximal response in terms of Ca²⁺ influx occurs at about 5×10^{-9} M OT in LLC-PK₁ cells [12].

3.2. uPA induction in LLC-PK₁ cells by OT and OT-specific agonists

The above results were elaborated by testing the various analogues for the stimulation of uPA-production in LLC-PK₁ cells. uPA activity was detected by spotting 8h conditioned media onto a casein plasminogen agar, and incubating at 37°C [30,31], activity being indicated by zones of plasminogen-dependent proteolytic clearing in the agar (Fig. 1). Results (not shown) could be confirmed using a colorimetric assay with the synthetic peptide substrate S-2251 [14,19,20,26]. OT and dSOT stimulated uPA production (Fig. 1A) but only at concentrations about 100 times higher than those for maximal induction by AVP and SAVP, consistent with the affinity of the respective ligands for the V₂-receptor of LLC-PK₁ cells.

Treatment of the V₂-receptor deficient mutant M18 [14] resulted in no uPA production (Fig. 1B), supporting the conclusion that OT-stimulated induction of LLC-PK₁ cells was V₂-receptor mediated. Further evidence was provided by coincubating 10^{-9} M AVP or SAVP (not shown) or 10^{-7} M OT or dSOT (Fig. 2) with various concentrations of OT-, V₁/OT-, or V₂/V₁-specific antagonists. Whilst the OT-antagonist MSOOT failed to block OT- or dSOT-mediated uPA induction, the V₂/V₁-specific antagonist MFIKAVP inhibited uPA in-

duction in a concentration-dependent manner (Fig. 2). These results were consistent with OT- and OT-specific agonist uPA induction in LLC-PK₁ cells being V₂-mediated.

4. DISCUSSION

The results here show that OT as well as OT-specific agonists can stimulate cAMP synthesis, cAMP-PK activation and uPA induction in LLC-PK₁ cells. The con-

Table IIA

Extracellular cAMP production induced by V₂- and OT-specific agonists and antagonists in LLC-PK₁ and M18 cells

Treatment*	Extracellular cAMP production (pmol/10 ⁶ cells) ⁺			
	LLC-PK ₁		M18	
	-IBMX	+IBMX	-IBMX	+IBMX
No addition	21	116	15	59
50 μM forskolin	210	4,560	236	3,740
10 ⁻⁸ M AVP	380	8,830	14	141
10 ⁻⁸ M dVSAVP	282	6,010	14	112
10 ⁻⁷ M OT	38	555	11	52
10 ⁻⁷ M dSOT	23	112	13	50

*Cell monolayers were treated for 8 h in serum-free DMEM in the absence or presence of 500 μM IBMX as indicated.

⁺Medium cAMP was determined as described in section 2, and is expressed per 10⁶ cells. Data represent the mean from a single typical experiment where the S.D. was less than 22% the value of the mean.

Table IIB

cAMP-PK activation in LLC-PK₁ cells induced by forskolin, V₂- and OT-specific agonists and V₁/OT- and OT-specific antagonists

Treatment*	cAMP-PK activity ratio	
	-IBMX	+IBMX
No addition	0.02 ± 0.01	0.14 ± 0.04 (6)
50 μM forskolin	0.20 ± 0.04	0.63 ± 0.02
10 ⁻⁷ M AVP	0.18 ± 0.05	0.82 ± 0.07 (3)
10 ⁻⁸ M AVP	0.11 ± 0.01	0.58 ± 0.05 (3)
10 ⁻⁹ M AVP	0.07 ± 0.01	0.36 ± 0.06 (3)
10 ⁻⁶ M OT	0.07 ± 0.02	0.51 ± 0.05 (3)
10 ⁻⁷ M OT	0.05 ± 0.02	0.29 ± 0.12 (3)
10 ⁻⁸ M OT	0.03 ± 0.01	0.05 ± 0.01 (3)
10 ⁻⁶ M dSOT	0.03 ± 0.01	0.18 ± 0.02
10 ⁻⁷ M dSOT	0.03 ± 0.01	0.05 ± 0.01
10 ⁻⁶ M TSOT	0.06 ± 0.01	0.16 ± 0.02
10 ⁻⁷ M TSOT	0.02 ± 0.01	0.07 ± 0.01
10 ⁻⁸ M SAVP	0.08 ± 0.01	0.64 ± 0.06
10 ⁻⁶ M MAAVP	0.04 ± 0.01	0.08 ± 0.01

*Cell monolayers were treated for 30 min in serum-free DMEM in the absence or presence of 500 μM IBMX as indicated, prior to washing and preparation of cell extracts. Extracts were then assayed in the presence or absence of exogenously added cAMP (10 μM). Data represent the mean ± S.D. (*n* in parentheses where *n* > 2). Total cAMP-PK activity (assayed in the presence of 10 μM cAMP) was 2.8 ± 0.2 U/mg.

centration dependence of the responses, the lack of OT-response of the V₂-deficient LLC-PK₁ mutant M18, and the inhibition of OT-mediated uPA induction by an V₂/V₁-receptor specific, but not OT-receptor specific antagonist, strongly indicate that these responses are all mediated by binding of the respective ligands to the V₂-, and not to the OT-receptor of LLC-PK₁ cells. Although activation of the Ca²⁺-phospholipid dependent kinase (PK-C) by phorbol esters has been shown to stimulate uPA production in LLC-PK₁ cells independently of cAMP and cAMP-PK [14,18–21], binding of OT to the OT-receptor does not appear to lead to significant induction of uPA synthesis, at least within 8 h, as shown here. PK-C activation would accordingly not seem to couple to stimulation of the OT-receptor in LLC-PK₁ cells. Further investigations using specific OT-agonists and antagonists should assist in elaborating the precise events of signal transduction following OT-stimulation.

Acknowledgements: The authors gratefully acknowledge Patricia Jans for invaluable technical assistance, H. Multscher and A. Krafft for helpful discussions, and the Deutsche Forschungs Gemeinschaft (SFB 169) for financial support.

REFERENCES

- [1] Soloff, M.S. and Swatz, T.L. (1973) *J. Biol. Chem.* 248, 6471–6478.
- [2] Soloff, M.S. and Grzonka, Z. (1986) *Endocrinology* 119, 1564–1569.
- [3] Fahrenholz, F., Hackenberg, M. and Müller, M. (1988) *Eur. J. Biochem.* 174, 81–85.
- [4] Schrey, M.P., Read, A.M. and Steer, P.J. (1986) *Biosci. Rep.* 6, 613–619.
- [5] Batras, J.L. (1986) *Eur. J. Pharmacol.* 120, 57–61.
- [6] Bojanic, D. and Fain, J.N. (1986) *Biochem. J.* 240, 361–365.
- [7] Fitzgerald, T.J., Uhling, R.J. and Exton, J.H. (1986) *J. Biol. Chem.* 261, 16871–16877.
- [8] Conrad, K.P., Gellai, M., North, W.G. and Valtin, H. (1986) *Am. J. Physiol.* 251, F290–296.
- [9] Forsling, M.L. and Brimble, M.J. (1985) in: *Oxytocin: Clinical and Laboratory Studies* (J.A. Amico and A.G. Robinson, Eds.) Elsevier, Amsterdam, pp. 167–175.
- [10] Lyness, J., Robinson, A.G., Sheridan, M.N. and Gash, D.M. (1985) *Experientia* 41, 1444–1446.
- [11] Stoeckel, S. and Freund-Mercier, M.J. (1989) *Am. J. Physiol.* 257, F310–F314.
- [12] Stassen, F.L., Heckman, G., Schmidt, D., Papadopoulos, M.T., Nambi, P., Sarau, H., Aiyar, N., Gellai, M. and Kinter, L. (1988) *Mol. Pharmacol.* 33, 218–224.
- [13] Cantau, B., Barjon, J.N., Chicot, D., Baskevitch, P.P. and Jard, S. (1990) *Am. J. Physiol.* 258, F963–972.
- [14] Jans, D.A., Resink, T.J., Wilson, E.R., Reich, E. and Hemmings, B.A. (1986) *Eur. J. Biochem.* 160, 407–412.
- [15] Dayer, J.-M., Vassalli, J.-A., Bobbit, J.L., Hull, R.N., Reich, E.R. and Krane, S.H. (1981) *J. Cell Biol.* 91, 195–200.
- [16] Hemmings, B.A. (1985) *Curr. Topics in Cell Regulation* 27, 117–132.
- [17] Nagamine, Y., Sudol, M. and Reich, E. (1983) *Cell* 32, 1181–1190.
- [18] Degen, J.D., Estensen, R.D., Nagamine, Y. and Reich, E. (1985) *J. Biol. Chem.* 260, 12426–12433.
- [19] Jans, D.A. and Hemmings, B.A. (1986) *FEBS Lett.* 205, 127–131.
- [20] Jans, D.A., Dierks-Ventling, C. and Hemmings, B.A. (1987) *Exp. Cell Res.* 172, 76–83.

- [21] Pearson, D., Nigg, E.A., Nagamine, Y., Jans, D.A. and Hemmings, B.A. (1991) *Exp. Cell Res.* 192, 315-318.
- [22] Grzonka, Z., Kasprzykowski, F., Kojro, E., Darlak, K., Melin, P., Fahrenholz, F., Crause, P. and Boer, R. (1983) *J. Med. Chem.* 29, 96-99.
- [23] Fahrenholz, F., Boer, R., Crause, P., Fritzsche, G. and Grzonka, Z. (1984) *Eur. J. Pharmacol.* 100, 47-58.
- [24] Grzonka, Z., Lammek, B., Kasprzykowski, F., Gazis, D. and Schwartz, I.L. (1983) *J. Med. Chem.* 26, 555-559.
- [25] Jans, D.A., van Oost, B.A., Ropers, H.H. and Fahrenholz, F. (1990) *J. Biol. Chem.* 265, 15379-15382.
- [26] Jans, D.A., Resink, T.J. and Hemmings, B.A. (1987) *Biochem. J.* 243, 413-418.
- [27] Hull, R.N., Cherry, W.R. and Weaver, G.W. (1976) *In Vitro* 12, 670-677.
- [28] Corbin, J.D. (1983) *Methods Enzymol.* 99, 227-232.
- [29] Soderling, T.R., Corbin, J.D. and Park, C.R. (1974) *Methods Enzymol.* 38, 358-367.
- [30] Luzius, H., Jans, D.A. and Fahrenholz, F. (1990) *J. Receptor Res.* 10, 61-80.
- [31] Jans, D.A., Peters, R., Jans, P. and Fahrenholz, F. (1990) *Exp. Cell Res.* 191, 121-128.
- [32] Bradford, W.M. (1976) *Anal. Biochem.* 72, 248-255.
- [33] Steinberg, R.A., Steinberg, M.G. and van Daalen Wetters, T. (1979) *J. Cell Physiol.* 100, 579-588.
- [34] Tovey, K.C., Oldham, K.G. and Whelan, J.A.M. (1974) *Clin. Chem. Acta* 56, 221-234.
- [35] Jans, D.A. and Hemmings, B.A. (1991) *FEBS Lett.* 281, 267-271.
- [36] Müller, M., Soloff, M.S. and Fahrenholz, F. (1989) *FEBS Lett.* 242, 333-336.
- [37] Audigier, S. and Barberis, C. (1985) 4, 1407-1412.
- [38] Jans, D.A., Zsigo, J., Peters, R. and Fahrenholz, F. (1989) *EMBO J.* 8, 2431-2438.
- [39] Jans, D.A., Peters, R., Jans, P. and Fahrenholz, F. (1991) *J. Cell Biol.* 114, 53-60.