

Pairs of noncrossing free Dyck paths and noncrossing partitions

William Y.C. Chen^{a,*}, Sabrina X.M. Pang^a, Ellen X.Y. Qu^a, Richard P. Stanley^b

^a Center for Combinatorics, LPMC-TJKLC, Nankai University, Tianjin 300071, PR China

^b Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

ARTICLE INFO

Article history:

Received 18 April 2008

Received in revised form 20 June 2008

Accepted 20 June 2008

Available online 13 August 2008

Keywords:

Dyck path

Free Dyck path

Plane partition

Noncrossing partition

Vacillating tableau

ABSTRACT

Using the bijection between partitions and vacillating tableaux, we establish a correspondence between pairs of noncrossing free Dyck paths of length $2n$ and noncrossing partitions of $[2n + 1]$ with $n + 1$ blocks. In terms of the number of up steps at odd positions, we find a characterization of Dyck paths constructed from pairs of noncrossing free Dyck paths by using the Labelle merging algorithm.

© 2008 Elsevier B.V. All rights reserved.

1. Introduction

We use the bijection between vacillating tableaux and partitions to establish a correspondence between pairs of noncrossing free Dyck paths of length $2n$ and noncrossing partitions of $[2n + 1]$ with $n + 1$ blocks. Recall that a *Dyck path* is a lattice path from the origin to a point $(2n, 0)$ consisting of up steps $U = (1, 1)$ and down steps $D = (1, -1)$ that does not go below the x -axis. Moreover, a lattice path from the origin to $(2n, 0)$ using the steps U and D without the restriction on a Dyck path is called a *free Dyck path*. Usually, a (free) Dyck path of length $2n$ is represented as a sequence of n U 's and n D 's. A k -tuple (P_1, P_2, \dots, P_k) of (free) Dyck paths from $(0, 0)$ to $(2n, 0)$ is called *noncrossing* if each P_i never goes below P_{i+1} for $1 \leq i \leq k - 1$.

A *partition* of a finite set S is a collection $\pi = \{B_1, B_2, \dots, B_k\}$ of subsets of S such that: (i) $B_i \neq \emptyset$ for each i ; (ii) $B_i \cap B_j = \emptyset$ if $i \neq j$, and (iii) $B_1 \cup B_2 \cup \dots \cup B_k = S$. Each B_i is called a *block* of π . A *plane partition* is an array $\delta = (\delta_{ij})_{i,j \geq 1}$ of nonnegative integers such that δ has finitely many nonzero entries and is weakly decreasing in rows and columns. If $\sum \delta_{i,j} = n$, then we say that δ is a plane partition of n and write $|\delta| = n$. A *part* of a plane partition $\delta = (\delta_{ij})$ is a positive entry $\delta_{ij} > 0$. The shape of a plane partition δ is the integer partition λ for which δ has λ_i nonzero parts in the i th row.

Let $[n] = \{1, 2, \dots, n\}$. Given a partition P of $[n]$, the *standard representation* of the partition P is a graph G on $[n]$ such that a block $\{i_1, i_2, \dots, i_k\}$ of P written in the increasing order $i_1 < i_2 < \dots < i_k$ corresponds to a path (i_1, i_2, \dots, i_k) . For example, the standard representation of 1358–29–46–7 is illustrated in Fig. 1.1. Meanwhile, we may view the standard representation of P as a directed graph because each edge can always be considered as an arc (i, j) with $i < j$, and we say that i is the *left end point* and j is the *right end point*. In this paper, we use Π_n to denote the set of partitions of $[n]$.

Let $k \geq 2$ and $P \in \Pi_n$. Define a *k-crossing* (*k-nesting*) of P as a set of k arcs $(i_1, j_1), (i_2, j_2), \dots, (i_k, j_k)$ in the standard representation of P such that $i_1 < i_2 < \dots < i_k < j_1 < j_2 < \dots < j_k$ ($i_1 < i_2 < \dots < i_k < j_k < \dots < j_2 < j_1$).

* Corresponding author.

E-mail addresses: chen@nankai.edu.cn (W.Y.C. Chen), pang@cfc.nankai.edu.cn (S.X.M. Pang), xiaoying@cfc.nankai.edu.cn (E.X.Y. Qu), rstan@math.mit.edu (R.P. Stanley).




Fig. 1.1. The standard representation of 1358–29–46–7.




Fig. 2.1. A triple of noncrossing free Dyck paths and the corresponding plane partition.

We use $cr(P)$ ($ne(P)$) to denote the maximal k such that P has a k -crossing (k -nesting). In particular, a 2-crossing (2-nesting) is called a *crossing (nesting)* for short. The statistics cr and ne were studied in [1] via a bijection between vacillating tableaux and set-partitions. We will use this bijection to connect noncrossing free Dyck paths to noncrossing partitions.

The paper is organized as follows. In Section 2, we give a quick review of the correspondence between k -tuples of noncrossing free Dyck paths and plane partitions. As a consequence, we get the formula for the number of pairs of noncrossing free Dyck paths of length $2n$. Then we give an overview of the bijection between vacillating tableaux and partitions as shown in [1]. We find a correspondence between pairs of noncrossing free Dyck paths and vacillating tableaux such that there is at most one row in each shape. These vacillating tableaux allow us to construct the noncrossing partitions. In Section 3, we give a characterization of Dyck paths obtained from pairs of noncrossing free Dyck paths by applying the Labelle merging algorithm.

2. Pairs of noncrossing free Dyck paths

We begin with the enumeration of k -tuples of noncrossing free Dyck paths via the correspondence with plane partitions with bounded part size. Given a k -tuple of noncrossing free Dyck paths (P_1, P_2, \dots, P_k) with each P_i of length $2n$, they must lie in the region bounded by the paths P_0 consisting of n up steps followed by n down steps and the path P_{k+1} consisting of n down steps followed by n up steps. As illustrated in Fig. 2.1, we may obtain a plane partition by filling the areas with i in each square located in the region between the paths P_i and P_{i+1} for $1 \leq i \leq k$. Suppose that the resulting plane partition π is of shape λ . Then we see that $\lambda_1 \leq n$ and $\lambda'_1 \leq n$ and the largest part of π does not exceed k . Since this correspondence is one-to-one, the enumeration of k -tuples of noncrossing free Dyck paths can be converted into the enumeration of plane partitions with bounded part size.

Let $B(r, c, t)$ be the set of plane partitions with at most r rows and at most c columns, and with the largest part at most t . It is known that

$$\sum_{\pi \in B(n, n, k)} q^{|\pi|} = \frac{[k+1][k+2]^2 \cdots [k+n]^n [k+n+1]^{n-1} \cdots [k+2n-1]}{[1][2]^2 \cdots [n]^n [n+1]^{n-1} \cdots [2n-1]}, \tag{2.1}$$

where $[i] = 1 - q^i$; see, for example, [5, Theorem 7.21.7]. Let $F(n, k)$ denote the number of k -tuples of noncrossing free Dyck paths of length $2n$. Then one can deduce a formula for $F(n, k)$ by setting $q = 1$ in (2.1). In particular,

$$F(n, 2) = \frac{(2n)!(2n+1)!}{(n!(n+1)!)^2}. \tag{2.2}$$

It has been shown by Callan that the above number also equals the number of noncrossing partitions of $[2n+1]$ with $n+1$ blocks; see Sloane [4, Sequence A000891]. Hence we are led to find a bijection between the set of pairs of noncrossing free Dyck paths of length $2n$ and the set of noncrossing partitions of $[2n+1]$ with $n+1$ blocks.

Our bijection, denoted by ζ , consists of two steps. The first step is to transform a pair of noncrossing free Dyck paths into a vacillating tableau in which each shape has at most one row. Then we use the bijection of Chen, Deng, Du, Stanley and Yan [1] to construct the corresponding noncrossing partition. We now give a brief review of the construction in [1]. We assume that the reader is familiar with the RSK algorithm, and we will use row insertion as the basic operation.

Definition 2.1. A vacillating tableau V_λ^{2n} of shape λ and length $2n$ is a sequence $(\lambda^0, \lambda^1, \dots, \lambda^{2n})$ of partitions such that: (i) $\lambda^0 = \emptyset$, and $\lambda^{2n} = \lambda$, (ii) λ^{2i+1} is obtained from λ^{2i} by doing nothing (i.e., $\lambda^{2i+1} = \lambda^{2i}$) or deleting a square, and (iii) λ^{2i} is obtained from λ^{2i-1} by doing nothing or adding a square.

Given a partition P , let $E(P)$ denote the set of arcs in the standard representation of P . To construct a vacillating tableau, we will derive a sequence $(T_0, T_1, T_2, \dots, T_{2n})$ of standard Young tableaux. Then the vacillating tableau is just the sequence of the shapes of these tableaux. We work our way backwards from $T_{2n} = \emptyset$ by determining the tableau T_{i-1} from T_i . We can construct the tableaux T_{2k-1}, T_{2k-2} from T_{2k} ($k \leq n$) by the following rules:

1. Let $T_{2k-1} = T_{2k}$ if the integer k does not appear in T_{2k} . Otherwise, T_{2k-1} is obtained from T_{2k} by deleting the square occupied by the element k .
2. $T_{2k-2} = T_{2k-1}$ if $E(P)$ does not have any arc of the form (i, k) . Otherwise, there is a unique integer $i < k$ such that $(i, k) \in E(P)$. Then T_{2k-2} is obtained from T_{2k-1} by row inserting the element i into T_{2k-1} .

Let λ^i be the shape of T_i for $0 \leq i \leq 2n$. Then $(\lambda^0, \lambda^1, \lambda^2, \dots, \lambda^{2n})$ is the required vacillating tableau.

The inverse procedure can be described as follows. Given a vacillating tableau $V = (\emptyset = \lambda^0, \lambda^1, \dots, \lambda^{2n} = \emptyset)$, we will recursively generate a sequence $(T_0, T_1, \dots, T_{2n})$, where T_i is a SYT (standard Young tableau) of shape λ^i . Let T_0 be the empty SYT. Below are the rules to construct T_i from T_{i-1} :

1. If $\lambda^i = \lambda^{i-1}$, then $T_i = T_{i-1}$.
2. If $\lambda^i \supset \lambda^{i-1}$, then $i = 2k$ for some integer $k \in [n]$. Determine the tableau T_i such that T_i is obtained from T_{i-1} by adding the integer k in the position of $\lambda^i \setminus \lambda^{i-1}$.
3. If $\lambda^i \subset \lambda^{i-1}$, then $i = 2k - 1$ for some integer $k \in [n]$. Set T_i to be the unique SYT (on a suitable alphabet) of shape λ^i such that T_{i-1} is obtained from T_i by row inserting some element j . Moreover, we record the arc $A_i = (j, k)$.

After the completion of the above procedure we are led to a set of arcs generated in Step 3. These arcs form a standard representation of a partition P of $[n]$.

For example, given the vacillating tableau

$$(\emptyset, \emptyset, 1, 1, 11, 11, 111, 11, 11, 1, 2, 1, 1, \emptyset, \emptyset),$$

the sequence (A_i, T_i) is as follows:

i	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
T_i	\emptyset	\emptyset	1	1	1	1	1	2	2	3	35	3	3	\emptyset	\emptyset
A_i					2	2	2	3	3						
							3								
								(1, 4)		(2, 5)		(5, 6)		(3, 7)	

The corresponding partition is $P = 14-256-37$.

The following result can be derived from [1, Theorem 6].

Theorem 2.2. Let $P \in \Pi_n$ and $(\emptyset = \lambda^0, \lambda^1, \dots, \lambda^{2n} = \emptyset)$ be the corresponding vacillating tableau. Then $cr(P)$ is the largest number of rows among λ^i , and $ne(P)$ is the largest number of columns among λ^i .

We are now ready to describe the bijection ζ . Let (P, Q) be a pair of noncrossing free Dyck paths, and let $P = p_1 p_2 \dots p_{2n}$ and $Q = q_1 q_2 \dots q_{2n}$. Based on P and Q , we form the sequence (p_i, q_i) , where $i = 1, 2, \dots, 2n$. The bijection ζ consists of two phases. First, we transform (P, Q) into a vacillating tableau

$$\mathcal{V}_{\emptyset}^{4n+2} = (\lambda^0, \lambda^1, \dots, \lambda^{4n+2})$$

of empty shape, i.e., $\lambda^{4n+2} = \emptyset$, such that there is at most one row in each λ^i and there are a total number of n operations of adding a square in the process to obtain λ^{4n+2} from λ^0 . Once a vacillating tableau is constructed, we may turn the vacillating tableau into a partition by the bijection in [1].

For $1 \leq i \leq 2n$, we have the following procedure to determine λ^k for $0 \leq k \leq 4n + 2$. Keep in mind that all the involved tableaux have at most one row. Specifically, we have the rules:

1. $\lambda^0 = \lambda^1 = \lambda^{4n+2} = \emptyset$.
2. If $(p_i, q_i) = (U, U)$, then λ^{2i} is obtained from λ^{2i-1} by adding one square, and λ^{2i+1} is obtained from λ^{2i} by deleting one square.
3. If $(p_i, q_i) = (U, D)$, then λ^{2i} is obtained from λ^{2i-1} by adding one square, and $\lambda^{2i+1} = \lambda^{2i}$.
4. If $(p_i, q_i) = (D, U)$, then $\lambda^{2i} = \lambda^{2i-1}$, and λ^{2i+1} is obtained from λ^{2i} by deleting one square.
5. If $(p_i, q_i) = (D, D)$, then $\lambda^{2i-1} = \lambda^{2i} = \lambda^{2i+1}$.

Let $P_i = p_1 p_2 \dots p_i$ and $Q_i = q_1 q_2 \dots q_i$, and let $|(P_i, Q_i)|_{(U,U)}$ denote the number of the pairs $(p_j, q_j) = (U, U)$ in (P_i, Q_i) . Similarly, we can define $|(P_i, Q_i)|_{(U,D)}$, $|(P_i, Q_i)|_{(D,U)}$ and $|(P_i, Q_i)|_{(D,D)}$. Evidently, the number of U 's in P_i equals $|(P_i, Q_i)|_{(U,U)} + |(P_i, Q_i)|_{(U,D)}$, and the number of U 's in Q_i equals $|(P_i, Q_i)|_{(U,U)} + |(P_i, Q_i)|_{(D,U)}$. Since P and Q are noncrossing, the number of U 's in P_i is not less than that in Q_i . It follows that

$$|(P_i, Q_i)|_{(U,U)} + |(P_i, Q_i)|_{(U,D)} \geq |(P_i, Q_i)|_{(U,U)} + |(P_i, Q_i)|_{(D,U)}.$$




Fig. 2.2. A pair of noncrossing free Dyck paths and its corresponding partition.

Hence we find that

$$|(P_i, Q_i)|_{(U,D)} \geq |(P_i, Q_i)|_{(D,U)}. \tag{2.3}$$

To show that $(\lambda^0, \lambda^1, \lambda^2, \dots, \lambda^{4n+2})$ is a valid vacillating tableau, we should justify that the above constructions are feasible. Clearly, the items 1, 2, 3 and 5 are well defined. So we may restrict our attention to item 4, in which case $(p_i, q_i) = (D, U)$. We aim to show that $\lambda^{2i} = \lambda^{2i-1}$ is not an empty shape. Combining (2.3) and the relations $|(P_i, Q_i)|_{(D,U)} = |(P_{i-1}, Q_{i-1})|_{(D,U)} + 1$ and $|(P_i, Q_i)|_{(U,D)} = |(P_{i-1}, Q_{i-1})|_{(U,D)}$, we see that

$$|(P_{i-1}, Q_{i-1})|_{(U,D)} > |(P_{i-1}, Q_{i-1})|_{(D,U)}.$$

From the construction of $(\lambda^0, \lambda^1, \dots, \lambda^{2i-1})$ it can be seen that $\lambda^{2i-1} \neq \emptyset$. Thus we have reached the conclusion that $(\lambda^0, \lambda^1, \dots, \lambda^{4n+2})$ is a vacillating tableau.

It is easily seen from Theorem 2.2 that the partition corresponding to the above vacillating tableau, denoted by R , is noncrossing because λ^i contains at most one row for any i . It remains to show that the resulting partition R contains exactly $n + 1$ blocks. Since the free Dyck path P is of length $2n$, there are n left end points in the standard representation of R . This implies that there are n arcs in the standard representation of R . On the other hand, R contains $2n + 1$ elements, since the vacillating tableau is of length $4n + 2$. So we may deduce that there are $n + 1$ blocks in R . It is not difficult to see that the above procedure is reversible. Therefore, we have established the following result.

Theorem 2.3. *The above map ζ is a bijection between the set of pairs of noncrossing free Dyck paths of length $2n$ and the set of noncrossing partitions of $[2n + 1]$ with $n + 1$ blocks.*

Fig. 2.2 is an illustration of the bijection ζ .

We remark that the bijection ζ can be described in a simpler manner. For a given pair (P, Q) of noncrossing free Dyck paths, let $P = p_1 p_2 \cdots p_{2n}$ and $Q = q_1 q_2 \cdots q_{2n}$, and let l_i (resp. r_i) denote the left-degree (resp. the right-degree) of vertex i in the standard representation of the partition corresponding to (P, Q) , i.e., the number of vertices j with $j < i$ (resp. $j > i$) connected to i . First, set $l_1 = r_{2n+1} = 0$. Then the pair (r_i, l_{i+1}) is determined by the following rules for $1 \leq i \leq 2n$:

1. If $(p_i, q_i) = (U, U)$, then set $(r_i, l_{i+1}) = (1, 1)$.
2. If $(p_i, q_i) = (U, D)$, then set $(r_i, l_{i+1}) = (1, 0)$.
3. If $(p_i, q_i) = (D, U)$, then set $(r_i, l_{i+1}) = (0, 1)$.
4. If $(p_i, q_i) = (D, D)$, then set $(r_i, l_{i+1}) = (0, 0)$.

Observe that the degree sequences $(l_1, l_2, \dots, l_{2n+1})$ and $(r_1, r_2, \dots, r_{2n+1})$ consist of only zeros and ones. As shown in Fig. 2.3, we may use half arcs (intuitively called left half arcs and right half arcs) to represent the left and right degrees. We have the following unique way to pair up half arcs in order to form a noncrossing partition. At each step we always try to find the leftmost left half arc and pair it up with the nearest right half arc on its left. Iterate this procedure until all the half arcs are paired. Finally, we obtain the standard representation of the desired partition, say R .

It is necessary to show that the resulting partition R contains $2n + 1$ elements and $n + 1$ blocks and is noncrossing. First, from the sequence $(p_1, q_1), (p_2, q_2), \dots, (p_{2n}, q_{2n})$ we determine the degrees $(r_1, l_2), (r_2, l_3), \dots, (r_{2n}, l_{2n+1})$. Clearly, the underlying set of the partition R is $[2n + 1]$. From the definition of (r_i, l_{i+1}) , one sees that the total number of half arcs equals the number of U 's in P and Q . Since P and Q are free Dyck paths of length $2n$, the total number of U 's in P and Q equals $2n$. It follows that there are n arcs in the standard representation of R . Moreover, R is noncrossing since at each step the leftmost left half arc is paired up with the nearest right half arc on its left, and this operation does not cause any crossing. It is not hard to check that the above procedure is reversible. Fig. 2.3 gives an illustration of this procedure.




Fig. 2.3. A pair of noncrossing free Dyck paths and the corresponding partition.




Fig. 3.1. The Labelle merging algorithm.

3. The Labelle merging algorithm

In this section, we establish a correspondence between pairs of noncrossing free Dyck paths of length $2n$ and Dyck paths of length $4n + 2$ with $n + 1$ up steps at odd positions. According to a theorem of Sulanke [6], the number of Dyck paths of length $2n$ with k up steps at odd positions equals the Narayana number $N(n, k) = \frac{1}{n} \binom{n}{k-1} \binom{n}{k}$. This implies the formula (2.2).

Labelle [3] gives an algorithm to merge a pair of noncrossing free Dyck paths into a 2-Motzkin path. It is realized that one can further transform a 2-Motzkin path into a Dyck path by using a bijection due to Delest and Viennot [2]. We will present an equivalent algorithm that directly transforms a pair of noncrossing free Dyck paths into a single Dyck path, and will call it the Labelle merging algorithm.

Let $P = p_1 p_2 \cdots p_{2n}$ and $Q = q_1 q_2 \cdots q_{2n}$ be a pair of noncrossing free Dyck paths of length $2n$. Let $Q' = q'_1 q'_2 \cdots q'_{2n}$, where we define $U' = D$ and $D' = U$. Then we merge P and Q into a Dyck path

$$U p_1 q'_1 p_2 q'_2 \cdots p_{2n} q'_{2n} D.$$

The following theorem gives a characterization of the Dyck paths corresponding to pairs of noncrossing free Dyck paths.

Theorem 3.1. *The Labelle merging algorithm is a bijection between noncrossing free Dyck paths of length $2n$ and Dyck paths of length $4n + 2$ with $n + 1$ up steps at odd positions.*

The verification of the above statement is omitted. Fig. 3.1 is an illustration of the Labelle merging algorithm, where the thick lines represent up steps at odd positions.

Acknowledgments

The authors would like to thank Martin Rubey and the referee for valuable comments leading to an improvement of an earlier version. This work was supported by the 973 Project, the PCSIRT Project of the Ministry of Education, the Ministry of Science and Technology, the National Science Foundation of China, and the US NSF grants DMS-9988459 and DMS-0604423.

References

[1] W.Y.C. Chen, E.Y.P. Deng, R.R.X. Du, R.P. Stanley, C.H.F. Yan, Crossings and nestings of matchings and partitions, *Trans. Amer. Math. Soc.* 359 (4) (2007) 1555–1575.
 [2] M.P. Delest, G. Viennot, Algebraic languages and polyominoes enumeration, *Theor. Comput. Sci.* 34 (1984) 169–206.
 [3] J. Labelle, On pairs of non-crossing generalized Dyck paths, *J. Statist. Plann. Inference* 34 (1993) 209–217.
 [4] N.J.A. Sloane, The on-line encyclopedia of integer sequences, www.research.att.com/~njas/sequences/.
 [5] R.P. Stanley, *Enumerative Combinatorics*, vol. 2, Cambridge University Press, Cambridge, 1999.
 [6] R.A. Sulanke, Moments, Narayana numbers, and the cut and paste for lattice paths, *J. Statist. Plann. Inference* 135 (1) (2005) 229–244.