



CIRUGIA Y CIRUJANOS

Órgano de difusión científica de la Academia Mexicana de Cirugía
Fundada en 1933

www.amc.org.mx www.elsevier.es/circir



CLINICAL CASE

Hemangiolympangioma of the spermatic cord in a 17 year-old: A case report[☆]



José Fernando Rogel-Rodríguez^{a,*}, José Fernando Gil-García^b,
Pasquinelly Velasco-García^a, Fernando Romero-Espinoza^c,
Tahitiana Zaragoza-Salas^a, Gabriel Muñoz-Lumbreras^a

^a Departamento de Cirugía General, Centro Médico Adolfo López Mateos, Toluca, Estado de México, Mexico

^b Departamento de Urología, Centro Médico Adolfo López Mateos, Toluca, Estado de México, Mexico

^c Departamento de Cirugía Plástica y Reconstructiva, Centro Médico Adolfo López Mateos, Toluca, Estado de México, Mexico

Received 31 March 2014; accepted 19 November 2014

Available online 8 March 2016

KEYWORDS

Hemangiolympphan-
gioma;
Orchiectomy;
Spermatic cord

Abstract

Background: Hemangiolympangiomas are extremely rare tumours arising from blood and lymphatic vessels. It is a benign disorder, and 95% are of the neck and axilla.

Objective: To present a case of hemangiolympangioma of the spermatic cord with contralateral recurrence.

Clinical case: A 17-year-old patient with a progressively growing tumour in the right inguinoscrotal region. Examination revealed a painless, movable and soft right inguinoscrotal tumour, mobile and soft. Surgical resection showed a 25 cm × 25 cm tumour from spermatic cord, right testicle, and subcutaneous cellular tissue. Histopathological study reported a hemangiolympangioma.

Conclusions: The spermatic cord is an unusual location of hemangiolympangiomas with contralateral recurrence. Surgical treatment, with histopathological diagnosis, is associated with good prognosis.

© 2015 Academia Mexicana de Cirugía A.C. Published by Masson Doyma México S.A. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

[☆] Please cite this article as: Rogel-Rodríguez JF, Gil-García JF, Velasco-García P, Romero-Espinoza F, Zaragoza-Salas T, Muñoz-Lumbreras G. Hemangiolympangioma de cordón espermático en adolescente de 17 años, reporte de caso. Cirugía y Cirujanos. 2016;84:162–166.

* Corresponding author at: Departamento de Cirugía General, Centro Médico Adolfo López Mateos, Segundo piso, Av. Nicolás San Juan, s/n, Col Ex Hacienda las Magdalenas, C.P. 50100, Toluca, Estado de México, Mexico. Tel.: +52 (722) 276 0860.

E-mail address: fernandorogel6@gmail.com (J.F. Rogel-Rodríguez).

PALABRAS CLAVE

Hemangiolinfangioma;
Orquiectomía;
Cordón espermático

Hemangiolympangioma de cordón espermático en adolescente de 17 años, reporte de caso**Resumen**

Antecedentes: Los hemangiolympangiomas son tumores benignos raros, provenientes tanto de vasos sanguíneos como linfáticos. El 95% se encuentran en cuello y axila.

Objetivo: Reportar el caso de un hemangiolympangioma de cordón espermático con recurrencia contralateral.

Caso clínico: Paciente masculino de 17 años de edad, con tumor en región inguinoscrotal derecha, de crecimiento progresivo. A la exploración física se encontró tumor derecho inguinoscrotal doloroso, de consistencia blanda y móvil. Se resecó un tumor de 25 × 25 cm que tenía involucro de cordón espermático, testículo derecho y tejido celular subcutáneo. El estudio histopatológico confirmó el diagnóstico de hemangiolympangioma.

Conclusión: Es inusual la localización de hemangiolympangiomas de cordón espermático, que además tuvo recurrencia contralateral. Debe confirmarse su diagnóstico con estudio histopatológico. El tratamiento es quirúrgico y se obtiene buen pronóstico.

© 2015 Academia Mexicana de Cirugía A.C. Publicado por Masson Doyma México S.A. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Background

Hemangiolympangiomas are mixed lesion with a vascular and lymphatic component.¹ They are generally benign in nature leading to favourable prognosis. They can be congenital or appear in infancy.²

There is a prevalence in the male of 2.4:1. There is greater risk of the hemangiolympangiomas presenting in premature babies and in live newborns incidence is 1:12,000. 40–60% of these are discovered at birth (congenital), 80–90% during the first 2 years of life (the age with most frequent presentation) and this frequency decreases with age.³

In 95% of cases, presentation is more frequent in: head, neck and axilla; 5% may be located in: mediastinum, retroperitoneum and mesentery, and is extremely rare in the groin. It may be associated with Klippel–Trenaunay–Weber syndrome, which is a rare congenital disease which was characterised by arteriovenous fistulas and multiple hemangiomas.^{2–4}

They are benign tumours which may infiltrate adjacent organs and cause severe malformations in the cavity and extremities.

The aetiology is unknown. One theory considers that during embryogenesis lymphatic tissue was deposited in the wrong areas with the result that an appropriate connection with central lymphatic pathways was obstructed, and were joined together with blood vessels.⁴

In general the tumours are asymptomatic, and several reports describe that pain is rare, and that the main concern is over function and aesthetics. Complications will depend on the affected organ, and may include vascular compromise through obstruction, lymphedema, and compartment syndrome (which is extremely rare).

Aim

Our report is a case of hemangiolympangioma of the spermatic cord with contralateral recurrence and upon which no known reports exist on this location, according to our review of references.

Clinical case

A 17 year old male patient with a history of left scrotal and testicular hemangiolympangioma surgical resection at the age of 2, and tonic-clonic seizures treated with valproate and magnesium.

During the treatment and control of a tonic-clonic seizure by the Emergency Service of the Centro Médico Adolfo López Mateos (ISEM), a physical examination revealed the following in the right hypogastrium and inguinal and testicle region: increase in hypogastrium volume and a 25 cm × 25 cm tumour at pubis level spreading to the inguinal channel and right scrotum sac; it was painful to the touch, soft in consistency, moveable, with scrotal skin attached, enlargement of the spermatic cord and the right testicle; the left scrotal sac had no testicle. Adenomegalias were not found (Fig. 1). The patient stated that the increase in volume had been progressive, over 2 years and there was a piercing pain on touch, with discomfort even from contact with underclothes. An assessment was made by the Urology Unit, which requested computed tomography of the abdomen and pelvis. This showed amorphous heterogeneous imaging with densities of 114 to 42 Hounsfield Units in the right scrotal sac. When contrast imaging was used its spread to the inguinal canal and subcutaneous cellular tissues became clear, with an



Figure 1 25 cm × 25 cm increase in volume in the right inguinal scrotal region. Arrow: increase in volume in spermatic cord region.




Figure 2 Computed tomography of the abdomen and pelvis with heterogeneous increase in volume of 114 to 42 UH. Arrow on left side: origin of tumour of the spermatic cord. Arrow on the right side: great size of the tumour in the suprapubic region.

approximate dimension of 26.4 cm × 25.7 cm in antero posterior and transverse edges (Fig. 2). Tumour markers were found to be within normal limits. The laboratory report was as follows: 204 UI/l lactate dehydrogenase, 0.77 UI/ml alfa-fetoprotein, ≤ 1.2 mIU/ml chorionic gonadotropin fraction β hormone.

Surgical resection was performed with the following findings: 25 cm × 25 cm soft tumour, with celluloadipose and lymphatic macroscopic characteristics, well defined edges, with subcutaneous cellular tissue, spermatic cord, scrotum and the right testicle; with celluloadipose and lymphatic macroscopic characteristics. After orchiectomy, the surgical specimen was sent for testing to the pathology department (Fig. 3).

Histopathology

Two testicle specimens were received. The first measured 5 cm × 4 cm, with a scrotal region measuring 4 cm × 2 cm, a dark brown, rugged epidermis, with characteristic congestive hairs; the tunica albuginea was smooth and shiny, as was the paratesticular cystic lesion of 2 cm × 1 cm. The second specimen measured 28 cm × 10 cm × 7 cm, with 2 flaps of skin, the largest measuring 15 cm × 10 cm and the smallest 11 cm × 10 cm; the epidermal surface was rugged and a brownish greyish colour. The spermatic cord of 6 cm × 1 cm was identified, with an external reddish brownish colour and when dissected, contained dilated vascular structures in its 3 sections. Cavernous hemangiolympangioma of 15 cm of spermatic cord and paratesticular soft tissues (Fig. 4).



Figure 3 Surgical specimen: block resection which includes skin with subcutaneous cellular tissue. The arrow shows the spermatic cord from where the hemangiolympangioma is derived.




Figure 4 A benign lesion is identified with vascular channels of diverse calibre, hyalinized wall, flat endothelium and congestive lights. The arrows indicate the vascular component with erythrocytes.



Figure 5 Reconstruction of the exposed residual area, which is 30 cm × 20 cm with fasciocutaneous advancement flap. The arrow indicates the exposed site area.

A second operation was performed by the Plastic Surgery Service, where an abdominoplasty type abdominal advancement flap was performed (Fig. 5).

Discussion

Vascular malformations are localised or diffuse errors of angiogenesis or vasculogenesis (embryogenesis period) and

are classified as low-flow which are capillary, venous, or lymphatic malformations, or high-flow which are arteriovenous or arteriovenous fistula malformations.^{5,6}

The classification of lesions from lymphatic vessels are: *benign*: (a) lymphangiomas with their 3 variants: capillary, cystic and cavernous; (b) lymphoangioma; (c) mixed lymphangioma (hemangiolympangioma) which is our case. *Malignant*: lymphangiosarcomas.^{5,6}

The diagnosis of this type of lesion occurs through histopathological examination. Their differential diagnoses are hemangiopericytoma, angiosarcoma, pure hemangiomas, pure lymphatic malformations. The lymphatic malformations may mainly be classified in accordance with histopathological characteristics into cavernous, cystic or capillary.^{7,8}

They are clinically characterised by being grouped into blisters, forming raised plaques, being verrucous and pink or reddish blue in colour, with a content based on lymph or erythrocyte cells due to prominent and dilated lymphatic and vascular channels.⁹

Medical check-ups are recommended for patients with spontaneous regression. If tumours are superficial they may be treated with corticosteroids may be administered or photocoagulation with laser and good aesthetic and functional outcomes are obtained. For deep tumours, as in the case mentioned, the treatment of choice is block surgical resection.^{10,11}

Recurrence is variable and may occur in 10–100% of cases. It depends on the extension of the excision. Locations with main recurrence are mouth, larynx, hypopharynx and tongue. Mortality is reported to occur in 6–10% of cases.^{11,12}

Conclusions

Hemangiolympangiomas are rare lesions with a favourable prognosis. This was the case of our patient who despite this aesthetic deformity was treated satisfactorily. The patient is currently at the 6 month stage of check-ups and no recurrence has occurred. However, medical surveillance will be continued in order to identify new cases of recurrence at an early stage, identify early stage lesions and offer the right treatment with limited damage.

Conflict of interests

The authors have no conflict of interests to declare.

References

1. Yu Seob S, Ram AD, Myung Ki K. Cavernous hemangiolympangioma of the testis without cutaneous hemangiomatosis in an elderly patient. *Korean J Urol.* 2012;53(11):80–112. Available from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3502743/pdf/kju-53-810.pdf> [accessed 01.01.14].
2. Cabrerizo-Merino MC, Oñate Sánchez RE, Romero Maroto M. El hemangiolinfangioma en la infancia: localización inusual. *RCOE.* 2004;9(1):89–92. Available from <http://scielo.isciii.es/pdf/rcoe/v9n1/clinico2.PDF> [accessed 05.01.14].
3. Fay A, Nguyen J, Waner M. Conceptual approach to the management of infantile hemangiomas. *J Pediatr.* 2010;157(6),

- 881-888.e1-5. Available from <http://www.ncbi.nlm.nih.gov/pubmed/20884010> [accessed 05.01.14].
4. Navarrete FG, Vences CM, Vásquez RM. Hemangiolinfangioma localizado en la piel cabelluda. *Dermatología Rev Mex*. 2008;52(2):89-92. Available from <http://www.medigraphic.com/pdfs/derrevmex/rmd-2008/rmd082e.pdf> [accessed 05.01.14].
 5. Handra-Luca A, Montgomery E. Vascular malformations and hemangiolymphangiomas of the gastrointestinal tract: morphological features and clinical impact. *Int J Clin Exp Pathol*. 2011;4(5):430-43. Available from <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3127065/> [accessed 07.01.14].
 6. Pratap A, Tiwari A, Pandey S, Agrawal B, Paudel G, Adhikary S, et al. Giant cavernous hemangiolymphangioma of the bladder without cutaneous hemangiomas causing massive hematuria in a child. *J Pediatric Urol*. 2007;3(4):326-9. Available from <http://www.ncbi.nlm.nih.gov/pubmed/18947766> [accessed 07.01.14].
 7. Tseng JJ, Chou MM, Ho ES. Fetal axillary hemangiolymphangioma with secondary intralesional bleeding: serial ultrasound findings. *Ultrasound Obstet Gynecol*. 2002;19(4):403-6. Available from <http://www.ncbi.nlm.nih.gov/pubmed/11952973> [accessed 15.01.14].
 8. Dubois J, Rypens F. Vascular anomalies. *Ultrasound Clin*. 2009;4(4):471-95. Available from <http://www.mdconsult.com/das/article/body/4371116842/jorg=clinics&source=&sp=23056717&sid=0/N/740769/1.html?issn=1556-858X> [accessed 17.01.14].
 9. Pérez LH, Cisneros GLC, Barba RE. Planeación terapéutica en un hemangiolinfangioma gigante toracoabdominal. *Arch Inv Mat Inf*. 2003;5(3):113-7. Available from <http://www.medigraphic.com/pdfs/imi/imi-2013/imi133c.pdf> [accessed 17.01.14].
 10. Larry DH, Lisa B. Current management of infantile hemangiomas and their common associated conditions. *Otolaryngol Clin North Am*. 2012;45(3):545-56. Available from <http://www.ncbi.nlm.nih.gov/pubmed/22588036> [accessed 24.01.14].
 11. Grossgold E, Kusuda L. Scrotal lymphangioma in an adult case report. *Urology*. 2007;70(3), 590.e1-2. Available from <http://www.ncbi.nlm.nih.gov/pubmed/17905123> [accessed 28.01.14].
 12. Torres S, Navarrete G, Villagómez E. Hemangiolinfangioma. *Rev Cent Dermatol Pascua*. 2008;17(1):26-9. Available from <http://www.medigraphic.com/pdfs/derma/cd-2008/cd081f.pdf> [accessed 28.01.14].