

5-METHYL-2-THIOURIDINE: A NEW SULFUR-CONTAINING MINOR CONSTITUENT FROM RAT LIVER GLUTAMIC ACID AND LYSINE tRNAs

F. KIMURA-HARADA, M. SANEYOSHI and S. NISHIMURA

National Cancer Center Research Institute, Chuo-ku, Tokyo, Japan

Received 11 February 1971

1. Introduction

2-Thiouridine-5-acetic acid methyl ester and 5-methylaminomethyl-2-thiouridine were found in tRNA from yeast and *E. coli*, respectively [1, 2]. Recently, it was shown that these 2-thiouridine derivatives were possibly located in the first position of the anticodon of the glutamic acid tRNAs and had a specific role for codon recognition [3, 4]. Since a 2-thiouridine derivative is present in tRNA specific for a particular amino acid in both yeast and *E. coli*, it was of interest to investigate whether mammalian tRNAs also contain a 2-thiouridine derivative, and if so, what its structure is. This communication reports that rat liver glutamic acid and lysine tRNAs contain a 2-thiouridine derivative. It is neither 5-methylaminomethyl-2-thiouridine nor 2-thiouridine-5-acetic acid methyl ester. This new minor nucleoside found in the rat liver tRNAs was fully characterized as 5-methyl-2-thiouridine. This is the first report of the characterization of a sulfur-containing minor constituent from mammalian tRNA.

2. Materials and methods

Rat liver tRNA₃^{Glu} and tRNA₂^{Lys} were obtained by

Abbreviations:

m¹Ap: 1-methyladenosine 3'-phosphate
 m¹Gp: 1-methylguanosine 3'-phosphate
 m⁷Gp: N⁷-methylguanosine 3'-phosphate
 m⁵Cp: 5-methylcytidine 3'-phosphate
 N₁p, N₂p, N₃p, N₄p, N₅p: 3'-phosphate of unknown nucleosides

successive column chromatographies on DEAE-Sephadex A-50 [5, 6], and benzoylated DEAE-cellulose [7]. Details of the purification procedure will be published elsewhere [8]. A synthetic 2-thiouridine was prepared as described by Brown et al. [9]. Authentic 5-methyl-2-thiouridine was prepared from 2',3',5'-tri-*O*-benzoyl-5-methyl-2-thiouridine which was kindly supplied from Dr. M. Sano of Central Research Laboratory, Daiichi Seiyaku Co., Ltd. [10], by alkaline debenzoylation and was purified by paper chromatography. Synthetic 2-thiouridine-5-acetic acid methyl ester was a gift from Dr. R.H. Hall of McMaster University. RNase T₂ was purchased from Sankyo Co. *E. coli* alkaline phosphomonoesterase was obtained from Worthington Biochemical Co. Thin-layer glass plates (10 cm × 10 cm) coated with Avicel SF cellulose were products from Funakoshi Pharmaceutical Co., Tokyo.

To test for the presence of new minor constituents in rat liver tRNAs, 2 absorbance units of the dialyzed tRNAs were extensively hydrolyzed with RNase T₂, and subjected to two dimensional thin-layer chromatography using the solvent system described previously [11]. For isolation of 5-methyl-2-thiouridine 3'-phosphate from rat liver tRNAs on a relatively large scale, the RNase T₂ digest of the tRNA (30 absorbance units) was subjected to two dimensional paper chromatography (Toyo-Roshi No. 51A paper, 30 cm × 30 cm) using the same solvent system as for thin-layer chromatography. The spot containing 5-methyl-2-thiouridine 3'-phosphate was cut off and eluted with water. It was then converted to the corresponding nucleoside by treatment with *E. coli* alkaline phosphomonoesterase.




Fig. 1. Two dimensional thin-layer chromatography of RNase T₂ digests of rat liver tRNA₃^{Glu} and tRNA₂^{Lys}.




Fig. 2. Ultraviolet absorption spectra of (a) N₁, (b) N₂ and (c) 5-methyl-2-thiouridine. pH 7.0, —; pH 2.0, - - - - -; pH 12.0, - · - · - ·.

Desulfurization of 5-methyl-2-thiouridine and related compounds was carried out by a similar procedure to that described previously [4, 12] for desulfurization of 4-thiouridine and 5-methylaminomethyl-2-thiouridine. The sample was dissolved in 0.05 ml of 0.1 M sodium carbonate–sodium bicarbonate buffer (pH 8.9), and 0.005 ml of a 2% aqueous solution of cyanogen bromide was added. The mixture was kept at room temperature for 10 min and then 0.001 ml of 10 M NaOH was added, and the mixture was heated at

50° for 1 hr. The alkaline solution was neutralized with Dowex-50 resin, and lyophilized for analysis by thin-layer chromatography and electrophoresis.

3. Results

As shown in fig. 1, several minor components were detected in two dimensional chromatograms of RNase T₂ digests of rat liver tRNA₃^{Glu} and tRNA₂^{Lys}, respec-

Table 1

Relative chromatographic mobilities and electrophoretic mobilities of N_1 , N_2 , 5-methyl-2-thiouridine and related compounds.

| | Thin-layer chromatography R_f in solvent system | | | Electrophoresis $R 2', (3')$ UPM |
|--|---|------|------|----------------------------------|
| | A | B | C | |
| Nucleoside N_1 | 0.54 | 0.56 | 0.46 | 0.28 |
| Nucleoside N_2 | 0.54 | 0.56 | 0.46 | 0.28 |
| 5-Methyl-2-thiouridine | 0.54 | 0.56 | 0.46 | 0.28 |
| 2-Thiouridine-5-acetic acid methyl ester | 0.59 | 0.60 | 0.51 | 0.36 |
| 2-Thiouridine | 0.48 | 0.47 | | 0.41 |
| Uridine | 0.48 | 0.40 | | 0.00 |

The solvent systems used were: A) isobutyric acid-0.5 M NH_4OH (5:3, v/v); B) butanol-acetic acid- H_2O (5:3:2, v/v/v); C) 2-propanol-conc. NH_4OH-H_2O (7:1:2, v/v/v). Thin-layer electrophoresis was carried out using a glass plate coated with Avicel SF cellulose in 0.05 M triethylammonium bicarbonate buffer, pH 7.5.

tively. In addition to m^1Ap , m^1Gp , m^5Cp , m^7Gp and Ψp , two unknown minor nucleotides were detected in the digest of $tRNA_3^{Glu}$ and these were designated as N_1p and N_3p , respectively. In the case of $tRNA_2^{Lys}$, three unknown nucleotides, designated as N_2p , N_4p and N_5p , were detected together with m^5Cp , m^7Gp and Ψp . The present communication only deals with the characterization of N_1 and N_2 as 5-methyl-2-thiouridine. No ribothymidine 3'-phosphate was detected at all in these two tRNAs. This is the first observation of the absence of ribothymidine in a pure species of tRNA from eukaryotic cells, and its absence is of interest in understanding the function of ribothymidine in tRNA.

Fig. 2 shows a comparison of the ultraviolet (UV) absorption spectra of the nucleoside preparations, N_1 from $tRNA_3^{Glu}$ and N_2 from $tRNA_2^{Lys}$, with those of synthetic 5-methyl-2-thiouridine. The three spectra were identical in all respects at three different pH values, indicating that both N_1 and N_2 are either 2-thiouridine or a derivative of it. Table 1 shows the R_f values of N_1 and N_2 on thin-layer chromatography and their relative electrophoretic mobilities, as compared with those of 5-methyl-2-thiouridine and related compounds. The mobilities of the nucleoside preparations, N_1 and N_2 were the same as these of synthetic

Table 2

Relative chromatographic mobilities and electrophoretic mobilities of desulfurized nucleosides, ribothymidine and related compounds.

| | Thin-layer chromatography R_f in solvent system D | Electrophoresis $R 2', (3')$ UMP |
|---|---|----------------------------------|
| Desulfurized N_1 | 0.55 | 0.00 |
| Desulfurized N_2 | 0.55 | 0.00 |
| Ribothymidine | 0.55 | 0.00 |
| Desulfurized 5-methyl 2-thiouridine | 0.55 | 0.00 |
| Desulfurized 2-thiouridine 5-acetic acid methyl ester | 0.29 | 0.53 |
| 5-Methyl-2-thiouridine | | 0.24 |
| 2-Thiouridine 5-acetic acid methyl ester | | 0.31 |

The solvent system used was: D) 95% ethanol- H_2O (4:1, v/v). Paper electrophoresis was carried out as described in table 1.

5-methyl-2-thiouridine in all systems tested, but different from these of 2-thiouridine and 2-thiouridine-5-acetic acid methyl ester. These results strongly suggest that N_1 and N_2 are 5-methyl-2-thiouridine. The lack of identity of N_1 and N_2 with 2-thiouridine-5-acetic acid methyl ester was confirmed by chromatography and electrophoresis of N_1 and N_2 after alkaline treatment [1]. Treatment of N_1 and N_2 with 0.2 M NaOH for 1 hr at 100° did not cause a change in their chromatographic and electrophoretic mobilities.

Further proof that N_1 and N_2 are identical with 5-methyl-2-thiouridine was obtained on desulfurization of N_1 and N_2 to ribothymidine. As shown in table 2, the chromatographic and electrophoretic mobilities of desulfurized N_1 and N_2 were the same as those of authentic ribothymidine and desulfurized 5-methyl-2-thiouridine. Although not shown in the figure, the UV absorption spectra of desulfurized N_1 and N_2 were identical with those of a specimen of ribothymidine.

4. Discussion

A new minor nucleoside isolated from rat liver $tRNA_3^{Glu}$ and $tRNA_2^{Lys}$ was thoroughly characterized

as 5-methyl-2-thiouridine. 5-Methyl-2-thiouridine has not previously been isolated from any type of nucleic acid or tRNA. Moreover, there is no previous report of the isolation and characterization of a sulfur-containing minor nucleoside from mammalian tRNA. Eliceiri recently reported that ^{35}S from ^{35}S -cysteine was incorporated into the 4S fraction of mouse lymphoma cells [13]. He concluded that ^{35}S was incorporated into a fraction other than the 4-thiouridine moiety. Moreover, Carbon et al. showed that the chromatographic profile of one of the isoaccepting lysine tRNAs from rabbit liver is reversibly altered by iodine treatment [14]. These indirect pieces of evidence suggesting the presence of a sulfur containing nucleoside can now be explained by the existence of 5-methyl-2-thiouridine in these mammalian tRNAs.

Based on the assumption that 1 absorbance unit of tRNA is equal to 1.66 μmole [15], the amounts of 5-methyl-2-thiouridine in $\text{tRNA}_3^{\text{Glu}}$ and $\text{tRNA}_2^{\text{Lys}}$ were estimated to be approximately 1 mole per tRNA. As reported in a separate paper, $\text{tRNA}_3^{\text{Glu}}$ and $\text{tRNA}_2^{\text{Lys}}$ were preferentially recognized by GAA and AAA, respectively [8]. It was previously shown that there is a correlation between the existence of a 2-thiouridine derivative and preferential recognition of A in the third position of the anticodon in both yeast and *E. coli* tRNA^{Glu} [3, 4]. From the above results and those reported here, it seems likely that 5-methyl-2-thiouridine is also located in the first position of the anticodon of rat liver tRNA^{Glu} and tRNA^{Lys}. The existence of 2-thiouridine derivatives in particular tRNAs of such widely different types as mammalian and bacterial cells suggests that the 2-thiouridine moiety in tRNA is essential for the fundamental process of protein synthesis in cells. It is possibly important for precise codon recognition of tRNA, as discussed previously [4].

Acknowledgements

We wish to thank Dr. W. Nakahara, Director of this Institute for his encouragement. We are indebted to Mr. M. Fujimoto for large scale isolation of rat liver, to Dr. M. Sano for a gift of 2', 3', 5'-tri-*O*-benzoyl-5-methyl-2-thiouridine, and to Dr. R.H. Hall for a gift of 2-thiouridine-5-acetic acid methyl ester.

References

- [1] L. Baczynskyj, K. Biemann and R. Hall, *Science* 159 (1968) 1481.
- [2] J. Carbon, H. David and M.H. Studier, *Science* 161 (1968) 1146.
- [3] M. Yoshida, K. Takeishi and T. Ukita, *Biochem. Biophys. Res. Commun.* 39 (1970) 852.
- [4] Z. Ohashi, M. Saneyoshi, F. Harada, H. Hara and S. Nishimura, *Biochem. Biophys. Res. Commun.* 40 (1970) 866.
- [5] S. Nishimura, F. Harada, U. Narushima and T. Seno, *Biochim. Biophys. Acta* 142 (1967) 133.
- [6] S. Nishimura and I.B. Weinstein, *Biochemistry* 8 (1969) 832.
- [7] I. Gillam, S. Millward, D. Blew, M. von Tigerstrom, E. Wimmer and G.M. Tener, *Biochemistry* 6 (1967) 3043.
- [8] F. Ishii, F. Kimura-Harada and S. Nishimura, in preparation.
- [9] D.M. Brown, D.B. Parihar, Sir Alexander Todd and S. Varadarajan, *J. Chem. Soc.* (1958) 3028.
- [10] T. Naito and M. Sano, *Chem. Pharm. Bull. (Tokyo)* 9 (1961) 709.
- [11] T. Seno, M. Kobayashi and S. Nishimura, *Biochim. Biophys. Acta* 169 (1968) 80.
- [12] M. Saneyoshi and S. Nishimura, *Biochim. Biophys. Acta* 204 (1970) 389.
- [13] G.L. Eliceiri, *Biochim. Biophys. Acta* 209 (1970) 387.
- [14] J.A. Carbon, L. Hung and D.S. Jones, *Proc. Natl. Acad. Sci. U.S.A.* 53 (1965) 979.
- [15] R.M. Hoskinson and H.G. Khorana, *J. Biol. Chem.* 240 (1965) 2129.