



HOSTED BY



ELSEVIER

Contents lists available at [ScienceDirect](http://ScienceDirect.com)

Asian Pacific Journal of Tropical Biomedicine

journal homepage: www.elsevier.com/locate/apjtbOriginal article <http://dx.doi.org/10.1016/j.apjtb.2015.04.005>

Antibacterial activities of ethanol extracts of Philippine medicinal plants against multidrug-resistant bacteria

Demetrio L. Valle Jr.^{1,2}, Jeannie I. Andrade², Juliana Janet M. Puzon^{1,3}, Esperanza C. Cabrera⁴, Windell L. Rivera^{1,3*}¹Institute of Biology, College of Science, University of the Philippines, Diliman, Quezon City 1101, Philippines²Makati Medical Center, Makati City 1229, Philippines³Natural Sciences Research Institute, University of the Philippines, Diliman, Quezon City 1101, Philippines⁴Biology Department, De La Salle University, Taft Avenue, Manila City 1004, Philippines

ARTICLE INFO

Article history:

Received 23 Mar 2015

Received in revised form 22 Apr 2015

Accepted 30 Apr 2015

Available online 20 June 2015

Keywords:

Philippine herbal medicine

Multidrug-resistant bacteria

*Piper betle**Psidium guajava**Phyllanthus niruri**Ehretia microphylla*

ABSTRACT

Objective: To investigate the antibacterial activities of crude ethanol extracts of 12 Philippine medicinal plants.**Methods:** Crude ethanol extracts from 12 Philippine medicinal plants were evaluated for their antibacterial activity against methicillin-resistant *Staphylococcus aureus*, vancomycin-resistant *Enterococcus*, extended spectrum β -lactamase-producing, carbapenem-resistant Enterobacteriaceae and metallo- β -lactamase-producing *Pseudomonas aeruginosa* and *Acinetobacter baumannii*.**Results:** The leaf extracts of *Psidium guajava*, *Phyllanthus niruri*, *Ehretia microphylla* and *Piper betle* (*P. betle*) showed antibacterial activity against the Gram-positive methicillin-resistant *Staphylococcus aureus* and vancomycin-resistant *Enterococcus*. *P. betle* showed the highest antibacterial activity for these bacteria in the disk diffusion (16–33 mm inhibition diameter), minimum inhibitory concentration (19–156 μ g/mL) and minimum bactericidal concentration (312 μ g/mL) assays. *P. betle* leaf extracts only showed remarkable antibacterial activity for all the Gram-negative multidrug-resistant bacteria (extended spectrum β -lactamase-producing, carbapenem-resistant Enterobacteriaceae and metallo- β -lactamase-producing) in the disk diffusion (17–21 mm inhibition diameter), minimum inhibitory concentration (312–625 μ g/mL) and minimum bactericidal concentration (312–625 μ g/mL) assays.**Conclusions:** *P. betle* had the greatest potential value against both Gram-negative and Gram-positive multidrug-resistant bacteria. Favorable antagonistic activities were also exhibited by the ethanol extracts of *Psidium guajava*, *Phyllanthus niruri* and *Ehretia microphylla*.

1. Introduction

Antibiotic resistance is a problem that continues to challenge the healthcare sector in a large part of the world in both developing and developed countries. The spread of multidrug-

resistant (MDR) bacteria in hospital and community settings remains a widely unresolved problem and a heavy burden to health services [1]. Despite advances in antibiotic therapy, infectious complications remain an important cause of mortality and morbidity among hospitalized patients. Although medical practitioners can resort to second- or third-choice drugs for treating these patients, the use of these synthetic drugs may subject the patient to a higher risk, due to the possibility of the drugs producing more harmful side effects. To address this challenge, actions must be taken to reduce this problem, such as controlling the use of antibiotics, understanding the genetic mechanisms of resistance and developing new

*Corresponding author: Windell L. Rivera, Institute of Biology, College of Science, University of the Philippines, Diliman, Quezon City 1101, Philippines.

Tel: +63 29205471

E-mail: wrivera@science.upd.edu.ph

Peer review under responsibility of Hainan Medical University.

Foundation Project: Supported by Philippine Council for Health Research and Development of the Department of Science and Technology (Grant No. 2015PHD1).

antibiotics and new therapeutic strategies. Advances in identifying new sources of natural products with antimicrobial activities and expanding antibiotic chemical diversity are providing chemical leads for new drugs [2].

The vast majority of modern medications were derived originally from ancient herbal traditions. The practices of plant-based traditional medicine are founded on hundreds of years of belief and observations, which predate the development of modern medicine. Medicinal plants have been used for centuries as remedies for human diseases as they contain components of therapeutic value. There are numerous plant natural products which have antifungal, antibacterial and antiprotozoal activities that could be used either systemically or locally [3]. Several plants containing volatile oils, polyphenols and alkaloids as active constituents are utilized as popular folk medicines, while others gained popularity in the form of finished products collectively named phytomedicines [4]. During the second half of the 20th century, the acceptance of traditional medicine as an alternative form of health care and the development of microbial resistance to the classical antibiotics led researchers to investigate the antimicrobial activities of medicinal plants. Antimicrobials of plant origin have enormous therapeutic potential [3]. They are effective in the treatment of infectious diseases while simultaneously mitigating many of the side effects that are often associated with synthetic antimicrobials.

Traditionally used medicinal plants produce a variety of compounds of known therapeutic properties. One of the vital activities possessed by these medicinal plants is antimicrobial. The scarcity of infective diseases in plants is in itself an indication of the successful defense mechanisms developed by them [5]. The substances that can either inhibit the growth of bacteria or kill them, with no toxicity or minimum toxicity to host cells are considered candidates for developing new antimicrobial drugs [6]. Some of the bioactive compounds could hinder the life processes of disease-causing bacteria, either by itself or in combination with other therapeutic agents [7]. In recent years, antimicrobial properties of medicinal plants are being increasingly reported from different parts of the world [8–10]. It is expected that plant extracts showing target sites other than those used by antibiotics will be active against drug resistant microbial pathogens. However, very little information is available on such activity of medicinal plants [6]. There have been numerous documentations found in the medical literature concerning the significance of traditional medicinal plants as alternatives to synthetic antibacterial and antifungal medications [11–17]. Most of these published works come from many countries that are still practicing the use of herbal medicine for the treatment of various diseases for practical and economic reasons. These studies are valuable resources for local medical scientists who seek to explore and substantiate the antibacterial and antifungal activities of Philippine medicinal herbs, particularly against MDR bacteria. Knowledge on the different antimicrobial assays and the plants' bioactive compounds are vital for the design of future studies [2].

The Philippines is one of the Asian countries with a diverse flora, and numerous species are believed to possess curative properties. However, most of these claims lack scientific validation. The use of herbal medicines in the pre-Spanish era had been widespread, although its documentation was only initiated during the Spanish occupation, with the main practitioners being called the *mediquillos* (herbal scientists) or *herbolarios* [18]. In

1997, the Philippine Government passed Republic Act 8423, also known as the “Traditional and Alternative Medicine Act of 1997,” an act creating the Philippine Institute of Traditional and Alternative Health Care to accelerate the development of traditional and alternative health care in the Philippines [19]. Most noteworthy objectives include “the encouragement of scientific research on traditional and alternative health care systems that have direct impact on public health care, and the promotion of the use of traditional, alternative, preventive and curative health care modalities that have been proven safe, effective, cost-effective, and they are consistent with government standards on medical practice.”

The Infectious Disease Society of America has considered the following bacteria as especially challenging in terms of management: methicillin-resistant *Staphylococcus aureus* (MRSA), vancomycin-resistant *Enterococcus* (VRE), extended spectrum β -lactamase (ES β L)-producing and carbapenem-resistant Enterobacteriaceae (CRE), metallo- β -lactamase (M β L)-producing *Pseudomonas aeruginosa* (*P. aeruginosa*) and *Acinetobacter baumannii* (*A. baumannii*) [20]. In Asia, the same emerging MDR bacteria have turned into an immense threat. There have also been distressing reports regarding the occurrence of ES β L-producing Enterobacteriaceae in fresh imported culinary herbs from Thailand, Vietnam and Malaysia, which are usually consumed without appropriate heating [21]. Even *A. baumannii*, over the last 10 years, has emerged as one of the most problematic bacteria since it exhibits a wide spectrum of antimicrobial resistance mechanisms, such that treatment has been limited to a few antibiotics [22]. VRE likewise has emerged as one of the difficult-to-treat bacteria that cause several life-threatening infections, and for which a definitive and successful therapy remains elusive from the reach of an ever-growing number of patients [23]. The advent of these MDR bacteria has solicited some inquiries with regard to the future of therapeutic drugs in chemotherapy. In line with the aforementioned findings, it is crucial that therapeutic antimicrobial studies should be more focused on the inhibition of these medically significant bacteria [24].

This study aimed to investigate the antibacterial activities of crude ethanol extracts of 12 Philippine medicinal plants, namely, *Cassia alata* L. (locally known as “akapulko”) (*C. alata*), *Psidium guajava* L. (locally known as “bayabas”) (*P. guajava*), *Piper betle* L. (locally known as “ikmo”) (*P. betle*), *Vitex negundo* L. (locally known as “lagundi”) (*V. negundo*), *Mitrephora lanotan* (locally known as “lanotan”) (*M. lanotan*), *Zingiber officinale* Rosc. (locally known as “luya”) (*Z. officinale*), *Curcuma longa* (locally known as “luyang dilaw”) (*C. longa*), *Tinospora rumphii* Boerl (locally known as “makabuhay”) (*T. rumphii*), *Moringa oleifera* (locally known as “malunggay”) (*M. oleifera*), *Phyllanthus niruri* (locally known as “sampa-sampalukan”) (*P. niruri*), *Centella asiatica* (locally known as “takip kuhol”) (*C. asiatica*), and *Ehretia microphylla* Lam. (locally known as “tsaang gubat”) (*E. microphylla*) against different MDR bacteria isolated from a tertiary medical center in the Philippines. Selection of the plants was based on their documented medicinal values. All 12 plants have traditionally known antibacterial and antifungal activities, with no extensive bioassay studies performed locally. In order to identify the plants with potential bioactive molecules of pharmaceutical importance in the management of MDR bacteria, *in vitro* methods of evaluation on test organisms were performed using the plant extracts.

2. Materials and methods

2.1. Plant materials

The 12 medicinal plant samples tested in this study are shown in Table 1. These were collected from different regions of the Philippines. All the plant materials were identified and compared to voucher specimens at the Herbarium of the Institute of Biology, University of the Philippines Diliman for authentication. The different plant parts were washed with water to remove all unwanted materials, air-dried, pulverized in a mill and stored in a sterile air-tight container until further use.

2.2. Plant extract preparation

Plant extracts were prepared in accordance to the methods of Basri and Fan with minor modifications [25]. Briefly, 150 g of each powdered plant material was soaked in 500 mL of ethanol for 24 h with shaking. The resultant extracts were centrifuged at 3000 r/min for 5 min at 4 °C. The supernatant was filtered through Whatman filter paper No. 1 (Whatman, UK), while the residues were used for a second extraction with 300 mL of ethanol. After the second extraction, the filtrates were concentrated under reduced pressure using a rotary evaporator at 50 °C. The crude extracts were collected and allowed to dry at room temperature.

2.3. Test microorganisms

The panel of test organisms for primary *in vitro* antibacterial screening in this study is summarized in Table 2. About 10 MDR bacteria were isolated from patients of the Makati Medical Center, Makati City, Philippines. All isolates were identified by automated biochemical tests using Vitek[®]MS (bioMérieux, Marcy l'Etoile, France) Gram-positive colorimetric identification card. The susceptibility patterns were obtained using Vitek[®]MS aspartate aminotransferase (bioMérieux, Marcy l'Etoile, France) following minimum inhibitory concentration (MIC) interpretive standards from the Clinical Laboratory Standard Institute M100-S25 [26].

2.4. Antibacterial assay of plant extracts

The plant extracts were tested for antibacterial activity using the disk diffusion method. The test organisms were subcultured in 5% sheep blood agar plate (BAP) for 24 h at (35 ± 2) °C. The colonies were inoculated in normal saline solution. The turbidity was then adjusted to equal the turbidity of 0.5 McFarland standard giving a final inoculum of 1.5 × 10⁸ CFU/mL. About 100 µL of inoculum of test organism was spread on Mueller-Hinton agar plate (Remel™, Thermo Fisher Scientific, USA). Sterile 6 mm paper disks (Becton Dickinson and Company, USA) with the plant extracts (200 µg) or solvent blank (dimethyl

Table 1

Philippine medicinal plants assayed against different MDR bacteria.

Scientific name	Family	Local name	Plant part used	Place of collection
<i>P. guajava</i>	Myrtaceae	Bayabas	Leaves	Tacloban, Leyte
<i>M. oleifera</i>	Moringaceae	Malunggay	Leaves	Cainta, Rizal
<i>T. rumphii</i>	Menispermaceae	Makabuhay	Stem	Tacloban, Leyte
<i>P. niruri</i>	Euphorbiaceae	Sampa-sampalukan	Aerial parts	Tacloban, Leyte
<i>C. asiatica</i>	Apiaceae	Takip kuhol	Leaves	Luisiana, Laguna
<i>P. betle</i>	Piperaceae	Ikmo	Leaves	General Nakar, Quezon
<i>C. longa</i>	Zingiberaceae	Luyang dilaw	Rhizome	Luisiana, Laguna
<i>Z. officinale</i>	Zingiberaceae	Luya	Rhizome	Luisiana, Laguna
<i>V. negundo</i>	Verbenaceae	Lagundi	Leaves	Roxas, Palawan
<i>M. lanotan</i>	Annonaceae	Lanotan	Leaves	Mt. Isarog, Camarines Sur
<i>E. microphylla</i>	Boraginaceae	Tsaang gubat	Leaves	Tanay, Rizal
<i>C. alata</i>	Fabaceae	Akapulko	Leaves	Angono, Rizal

Table 2

Panel of test organisms for primary antibacterial *in vitro* screening.

Characteristic	Species	Source	Antibiotic resistance pattern
Gram-positive coccus	<i>S. aureus</i>	ATCC 29223	Susceptible
Enterobacteriaceae	<i>E. coli</i>	ATCC 25922	Susceptible
Enterobacteriaceae, encapsulated	<i>K. pneumoniae</i>	ATCC BAA-1705	Susceptible
Non-Enterobacteriaceae	<i>P. aeruginosa</i>	ATCC 27853	Susceptible
Gram-positive coccus, MDR, #1	MRSA	12/Male, wound	AM, FOX, OX, P
Gram-positive coccus, MDR, #2	MRSA	69/Male, wound	AM, FOX, OX, P
Gram-positive coccus, MDR, #3	MRSA	42/Male, blood	AM, FOX, OX, P
Gram-positive coccus, MDR, #4	MRSA	35/Female, sputum	AM, FOX, OX, P
Enterobacteriaceae, MDR	ESβL, <i>E. coli</i>	55/Female, blood	AM, FEP, CTX, CTZ, CRO
Enterobacteriaceae, encapsulated, MDR	ESβL, <i>K. pneumoniae</i>	25/Female, urine	AM, FEP, CTX, CTZ, CRP
Non-Enterobacteriaceae, MDR	MβL, <i>P. aeruginosa</i>	64/Male, blood	FEP, CRZ, IPM, MEM
Non-Enterobacteriaceae, MDR	MβL, <i>A. baumannii</i>	53/Female, blood	FEP, CRZ, IPM, MEM
Enterobacteriaceae, MDR	CRE, <i>E. coli</i>	48/Male, urine	AM, FEP, CTX, CTZ, CRO, IPM, MEM
Enterobacteriaceae, encapsulated, MDR	CRE, <i>K. pneumoniae</i>	58/Female, sputum	AM, FEP, CTZ, CTZ, CRO, IPM, MEM
VRE	VRE	45/Male, urine	P, VA

AM: ampicillin; FEP: cefepime; CTX: cefotaxime; FOX: ceftazidime; CTZ: ceftazidime; CRO: ceftriaxone; IPM: imipenem; MEM: meropenem; OX: oxacillin; P: penicillin; VA: vancomycin; *S. aureus*: *Staphylococcus aureus*; *E. coli*: *Escherichia coli*; *K. pneumoniae*: *Klebsiella pneumoniae*.

sulfoxide) were then placed on the inoculated plates. The plates were incubated at (35 ± 2) °C for 16–24 h. Representative antibiotics of the test isolates based on susceptibility patterns were used as positive controls (Table 2). Antibacterial activities were evaluated by measuring the diameters of zones of inhibition in mm against the test organism.

2.5. Determination of MIC and minimum bactericidal concentration (MBC) of the plant extracts

The MICs of the plant extracts were determined in sterile 96-well microplates using the broth microdilution method of the Clinical and Laboratory Standards Institute, M07-A8 [26]. Each test was done in triplicate. The plant extracts were serially diluted to produce final concentrations of 19 µg/mL to 10 000 µg/mL. Cation-adjusted Mueller-Hinton broth (Becton Dickinson and Company, USA) was used as diluent. The set-up included bacterial growth controls in wells containing 10 µL of the test inoculum and negative controls without bacterial inoculum. Reference drug controls were likewise included in the set-up.

The inoculum was prepared by direct saline suspension of isolated bacterial colonies selected from an 18–24 h 5% sheep BAP culture. Suspension was adjusted to achieve a turbidity equivalent to 0.5 McFarland turbidity standard, which approximated 1.5×10^8 CFU/mL. Within 15 min after standardization, 10 µL of the adjusted inoculum was added into each well containing 100 µL plant extract in the dilution series, and mixed. The sealed microdilution trays were incubated at (35 ± 2) °C for 16–20 h in an ambient air incubator.

The MIC was determined by selecting the lowest concentration of plant extract that completely inhibited the growth of the organism in the well as detected by the unaided eye. To determine the growth end points, the amount of growth in the wells containing the plant extracts was compared with the amount of growth in the growth-control well (no plant extracts) used in each set of tests. For a test to be considered valid, acceptable growth (2 mm button or definite turbidity) must occur in the growth-control well.

The MBC was determined following the methods described by Irobi and Daramola with slight modifications [27]. Wells with no visible growth in MIC assays were subcultured using a 10 µL inoculating loop onto a 5% sheep BAP at (35 ± 2) °C for 16–20 h incubation. The MBC was defined as the lowest concentration of the extract that did not permit any growth.

3. Results

3.1. Disk diffusion method

The disk diffusion method for antimicrobial susceptibility testing was initially performed to determine the antibacterial activities of the 12 medicinal plants against MDR bacteria, namely, MRSA, VRE, MβL *P. aeruginosa* and *A. baumannii*, ESβL *E. coli*, ESβL *K. pneumoniae* and CRE *K. pneumoniae*. ATCC control strains of *S. aureus*, *E. coli*, *K. pneumoniae* and *P. aeruginosa* were also included. Of the 12 medicinal plants, the ethanolic extracts of *E. microphylla*, *P. niruri*, *P. guajava* and *P. betle* showed inhibition of bacterial growth against some or all of the test organisms (Table 3). The plant extracts

Table 3

Diameters of zones of inhibition of ethanol extracts from 12 medicinal plants against clinical isolates of MDR bacteria and ATCC reference strains (mm)^a.

Plant extracts	MRSA	MRSA	MRSA	MRSA	VRE	MβL	MβL	MβL	ESβL	ESβL	ESβL	ESβL	CRE	<i>S. aureus</i> (ATCC 25923)	<i>E. coli</i> (ATCC 25922)	<i>K. pneumoniae</i> (ATCC BAA1705)	<i>P. aeruginosa</i> (ATCC 27853)	
	1	2	3	4		<i>P. aeruginosa</i>	<i>A. baumannii</i>	<i>E. coli</i>	<i>K. pneumoniae</i>	<i>E. coli</i>	<i>K. pneumoniae</i>	<i>P. aeruginosa</i>						
<i>C. alata</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>C. asiatica</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>C. longa</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>E. microphylla</i>	19	14	12	9	12	-	12	-	-	-	-	-	-	12	-	-	10	-
<i>M. lanotan</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>M. oleifera</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>P. niruri</i>	9	11	9	9	13	-	-	-	-	-	-	-	-	14	-	-	-	-
<i>P. betle</i>	32	34	28	34	28	17	23	20	20	21	21	16	16	30	16	17	17	17
<i>P. guajava</i>	18	16	18	13	12	-	-	-	-	-	-	-	-	14	-	-	-	-
<i>T. rumphii</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>V. negundo</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Z. officinale</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

^a Zone size includes 6-mm disk; -: No zone of inhibition.

exhibited inhibition zones ranging from 9 mm to 34 mm diameter, with the most noteworthy results shown by *P. betle* (Figure 1). Cefoxitin 30 µg disk (14 mm) showed a resistant result. The diameters included the 6 mm filter paper disk.

The ethanolic extract prepared from the *P. betle* leaf demonstrated inhibition zones greater than 15 mm in diameter, as shown in Table 3, with the greatest zones (28–34 mm) produced against the clinical strains of VRE and MRSA and for the *S. aureus* ATCC control strain. The leaf extract likewise showed growth inhibition against all Gram-negative clinical MDR bacterial strains and their respective ATCC controls tested, with zones ranging from 16 mm to 23 mm. On the other hand, *P. guajava* and *P. niruri* produced zones of growth inhibition against both clinical MRSA and VRE and ATCC strain of *S. aureus*, but failed to inhibit the growth of Gram-negative organisms tested. *E. microphylla* likewise exhibited inhibition zones for clinical strains of MRSA, VRE and *S. aureus* ATCC 25923. In addition, it inhibited the Gram-negative MβL *A. baumannii* and *P. aeruginosa* ATCC 27853.

Results of the disk diffusion assay showed that the most commonly inhibited bacteria by the ethanolic plant extracts were the clinical MRSA and VRE strains, and the ATCC control *S. aureus*, followed by MβL *A. baumannii* and *P. aeruginosa* ATCC 27853.

3.2. MIC

The MIC assay of the *P. betle* leaf extract consistently showed strong antagonistic activities against all microorganisms tested, with MIC values ranging from 19 to 1250 µg/mL (Table 4). The strongest activity was exhibited against the clinical strains of Gram-positive bacteria: VRE (19 µg/mL), MRSA 4 (78 µg/mL), MRSA 1–3 (each with MIC of 156 µg/mL) and *S. aureus* ATCC 25923 (312 µg/mL). For the Gram-negative bacteria, the most common MIC value was 625 µg/mL for MβL *A. baumannii*, ESβL *K. pneumoniae*, *E. coli* ATCC 25922 and *P. aeruginosa* ATCC 27853, followed by 312 µg/mL for MβL *P. aeruginosa*, ESβL *E. coli* and CRE




Figure 1. Zones of inhibition of *P. betle* (32 mm), *E. microphylla* (19 mm), *V. negundo* (0 mm) and *M. lanotan* (0 mm) against clinical isolate MRSA 1.

Table 4

MIC of ethanol extracts from 12 medicinal plants against clinical isolates of MDR bacteria and ATCC reference isolates (µg/mL).

Plant extracts	MRSA	MRSA	MRSA	MRSA	VRE	MβL	MβL	MβL	MβL	ESβL	ESβL	ESβL	ESβL	CRE	S. aureus (ATCC 25923)	E. coli (ATCC 25922)	K. pneumoniae (ATCC BAA1705)	P. aeruginosa (ATCC 27853)
<i>C. alata</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>C. asiatica</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>C. longa</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>E. microphylla</i>	1250	1250	2500	1250	1250	5000	-	-	-	-	-	-	-	1250	625	1250	2500	625
<i>M. lanotan</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>M. oleifera</i>	5000	2500	2500	5000	1250	-	-	-	-	-	-	-	-	2500	-	-	-	-
<i>P. niruri</i>	156	156	156	78	19	625	625	312	312	625	625	625	312	312	625	1250	625	625
<i>P. betle</i>	1250	625	1250	625	5000	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>P. guajava</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>T. rumphii</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>V. negundo</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Z. officinale</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

-: No antibacterial activity.

Table 5
MBC of ethanol extracts from 12 medicinal plants against clinical isolates of MDR bacteria and ATCC reference isolates ($\mu\text{g/mL}$).

Plant extracts	MRSA 1	MRSA 2	MRSA 3	MRSA 4	VRE	M β L <i>P. aeruginosa</i>	M β L <i>A. baumannii</i>	ES β L <i>E. coli</i>	ES β L <i>K. pneumoniae</i>	CRE	<i>S. aureus</i> (ATCC 25923)	<i>E. coli</i> (ATCC 25922)	<i>K. pneumoniae</i> (ATCC BAA1705)	<i>P. aeruginosa</i> (ATCC 27853)
<i>C. alata</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>C. asiatica</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>C. longa</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>E. microphylla</i>	1250	2500	2500	1250	1250	-	5000	-	-	-	2500	-	-	2500
<i>M. lanotan</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>M. oleifera</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>P. niruri</i>	5000	2500	2500	5000	1250	-	-	-	-	-	2500	-	-	-
<i>P. betle</i>	312	156	156	78	19	312	625	625	625	625	312	625	1250	625
<i>P. guajava</i>	2500	1250	1250	625	5000	-	-	-	-	-	1250	-	-	-
<i>T. rumphii</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>V. negundo</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Z. officinale</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-

-: No antibacterial activity.

K. pneumoniae. The *K. pneumoniae* ATCC BAA-1705 only showed MIC greater than 1000 $\mu\text{g/mL}$ (1250 $\mu\text{g/mL}$).

P. guajava and *P. niruri* similarly showed good patterns of inhibition against all Gram-positive bacteria tested, although the MIC values were much higher than those for *P. betle*. The MIC values ranged from 625 to 5000 $\mu\text{g/mL}$ and 1250–5000 $\mu\text{g/mL}$, respectively (Table 4). Similar to *P. niruri*, the ethanol extract from *E. microphylla* showed MIC values of 1250–5000 $\mu\text{g/mL}$, with the lowest MICs for MRSA, VRE and *S. aureus* ATCC 25923 and the highest MIC for M β L *A. baumannii*.

3.3. MBC

The MBCs were established by subculturing the samples with no visible growth in the MIC assays. Most of the MBC and MIC values for the ethanol plant extracts were closely similar, with only *P. niruri* showing exactly identical values for both assays (Table 5). Slightly higher MBC values for MRSA 1 (312 $\mu\text{g/mL}$; MIC 156 $\mu\text{g/mL}$), ES β L *E. coli* (625 $\mu\text{g/mL}$; MIC 312 $\mu\text{g/mL}$) and CRE *K. pneumoniae* (625 $\mu\text{g/mL}$; MIC 312 $\mu\text{g/mL}$) were obtained with *P. betle*. For *P. guajava*, the MBC values slightly differed for MRSA 1 (2500 $\mu\text{g/mL}$; MIC 250 $\mu\text{g/mL}$) and MRSA 2 (1250 $\mu\text{g/mL}$; MIC 625 $\mu\text{g/mL}$) strains, whereas *E. microphylla* had slightly higher MBC values for MRSA 2 (2500 $\mu\text{g/mL}$; MIC 1250 $\mu\text{g/mL}$) and ATCC *S. aureus* control strain (2500 $\mu\text{g/mL}$; MIC 1250 $\mu\text{g/mL}$).

4. Discussion

The current problem associated with emerging MDR bacteria presents a serious global medical crisis, requiring constant surveillance, which continuously challenges the scientific community [24]. The diminishing efficacy and increasing toxicity of synthetic drugs further aggravate this problem, thus, scientists are directed to seek more natural or organic materials for solutions. Traditional medicine has been practiced worldwide for centuries, particularly the application of herbal plants for therapeutic purposes. Philippines possess a rich source of medicinal plants, although not as extensive as India or China, enough to provide us with alternative remedies [18]. The 12 plants have been used locally for their traditional medicinal properties, all of which have documented antimicrobial activities. However, their efficacies against MDR bacteria have not been studied. In developing countries, it is imperative that effective but less expensive antibacterials should be developed to accommodate all patients, regardless of financial status, in order to eliminate some of the human factors that can cause MDR.

In recent years, there are increasing published reports showing successful antimicrobial activities of various traditional medicinal plants against MDR bacteria, such as MRSA, ES β L, VRE, and MDR *P. aeruginosa* [9,28–38]. In the present study, favorable antagonistic activities against various MDR bacteria were exhibited by the ethanol extracts of four Philippine medicinal plants, namely, *P. betle*, *P. guajava*, *P. niruri* and *E. microphylla*.

As previously mentioned, the medicinal plants have known antimicrobial properties, but their efficacies against MDR

bacteria have not been well-documented in the local medical literature or in other countries. *P. guajava* has been separately documented to be inhibitory against MRSA [39], MDR *V. cholera* [40] and MDR *S. aureus* [41]. In relation to the extraction method, it was reported that only ethanol extracts showed antagonistic activities against Gram-negative bacteria [10,42], while Mishra and Babele documented that the highest antimicrobial activity was obtained with the methanol extract of *P. guajava* leaf against drug resistant and susceptible *E. coli* and *P. aeruginosa* [43]. In the present study, the ethanol extract from the *P. guajava* leaf demonstrated significant inhibitory activities against MRSA and VRE and no activity against the Gram-negative bacteria tested. MICs and MBCs obtained in the current study are almost similar to the values obtained from another study by Esimone against MRSA, which ranged from 1250–5000 µg/mL for the water extracts, and 625–2500 µg/mL for the methanol extracts [39].

P. niruri is traditionally used for the treatment of kidney stones, gallstones, wound healing, genito-urinary tract infections and sexually transmitted diseases [18,44,45]. Bokarey et al. determined that different extracts from the *P. niruri* plant were effective against common human bacteria such as *S. aureus*, *E. coli*, *Proteus vulgaris* and *P. aeruginosa* [46]. Our present study is the first to show the antibacterial activity of *P. niruri* against MRSA and VRE.

P. betle is a multifunctional medicinal plant, which is cultivated throughout the Philippines and can also be found in other countries, such as Malaysia and India. The leaves have been traditionally used to control caries, periodontal disease and prevention of halitosis and is proven to have significant gastrointestinal and hepatoprotective effects [47,48]. The 80% methanol and 70% ethanol leaf extracts have been reported to be effective against strains of *S. aureus*, *E. coli* and *P. aeruginosa*, with the methanolic extract exhibiting wider zones of inhibition in the modified agar well diffusion method by Khan and Kumar [49]. The zones of inhibition for the methanolic extract ranged from 15 to 29.5 mm, whereas the ethanol extracts showed inhibition zones ranging from 14 to 17 mm. MIC for both extracts ranged from 2 to 8196 µg/mL. A similar study against the three bacteria has been performed but with the addition of two extraction methods: cold aqueous and ethyl acetate extracts of several Indian varieties of *P. betle* leaves [50]. Another study showed the efficacy of the ethanolic leaf extract against *K. pneumoniae* and *P. vulgaris*, wherein the MIC value obtained for the two bacteria was 25 µg/mL [51]. In a more recent study related to the activities of 90% methanol and 90% ethanol leaf extracts of *P. betle* against clinical oral isolates (*S. aureus*, *Staphylococcus epidermidis*, *Bacillus cereus*, *Bacillus subtilis*, *Listeria monocytogenes*, *E. coli*, *Salmonella typhimurium*, *Salmonella enteritidis*, *K. pneumoniae* and *P. aeruginosa*), the extracts demonstrated good antimicrobial activities against the Gram-positive and Gram-negative bacteria, except for *P. aeruginosa* [52]. At present, studies on the antimicrobial activities of *P. betle* against MRSA, ESβL, CRE, VRE, or MβL-producing *P. aeruginosa* and *A. baumannii* have been noted in scientific literature. The results of the present study showed that *P. betle* had the greatest potential value against both Gram-negative and Gram-positive MDR bacteria, exhibiting wide zones of growth inhibition in the disk diffusion assay and with the lowest concentrations of the extract required to inhibit the growth and

affect death of the pathogens, as supported by the MIC and MBC assays, respectively.

E. microphylla, *P. guajava* and *P. niruri* also showed significant antagonistic activities against the Gram-positive bacteria, but the limited or lack of activity against Gram-negative bacteria has reduced their value as an ideal or broad spectrum antibacterial for MDR microorganisms. Nevertheless, they can still be developed as second or third choices for the treatment of susceptible MDR bacteria. In addition, *E. microphylla* could provide us with an alternative treatment for MβL-producing *A. baumannii* and *P. aeruginosa*. *In vitro* antimicrobial studies regarding *E. microphylla* are rarely encountered. It has been surmised that the antibacterial activities of herbal plant extracts and essential oils are focused on the structures and cellular membranes and due to the presence of various bioactive compounds and extensive chemical profiles, it is likely that the antimicrobial potency is not just caused by one solitary mechanism but rather by several events at a cellular level [53].

The outer membrane found in the Gram-negative cell wall is composed of structural lipopolysaccharides which render the cell wall impermeable to lipophilic solutes, unlike Gram-positive bacteria which do not have this outer membrane. This morphologic difference influences their reaction to antibacterial agents. In addition, Gram-negative bacteria also have inherent overexpressed or multiple efflux pumps that prevent the intracellular accumulation of antibacterial agents. Thus, there is a need to discover and develop antibacterial agents that are capable of bypassing or suppressing the efflux pumps and that could also re-establish the antibacterial potency of older antibiotics [54]. Although previous studies on *P. guajava* [10,42] and *P. niruri* [46] reported significant antagonistic activities against Gram-negative bacteria, the results of the present study did not show the efficacies of these two plants on the Gram-negative bacteria tested. The variation in sensitivities could be attributed to the reinforced defense mechanisms acquired by the MDR bacteria.

However, this study resolutely established the efficacy of the ethanolic leaf extract of *P. betle* against known Gram-positive and Gram-negative MDR bacteria, indicating that the plant produces compounds that are able to combat the bacterial defenses. Henceforth, the results justify that further investigation should be performed on *P. betle*. Additional or complementary studies are necessary concerning the application of different extraction methods, phytochemical screening, phytophysical analysis, bioassay-guided isolation, purification and quantification of the bioactive components, including the detection of toxicity, side effects and pharmacokinetic properties in preparation for its *in vivo* assessment. Completion and favorable outcomes obtained from succeeding studies would definitely strengthen its potential as a novel and cost-effective agent against MDR bacteria.

Conflict of interest statement

We declare that we have no conflict of interest.

Acknowledgments

This study was supported by a research grant from the Philippine Council for Health Research and Development of the Department of Science and Technology (Grant No. 2015PHD1).

References

- [1] Cosgrove SE, Carmeli Y. The impact of antimicrobial resistance on health and economic outcomes. *Clin Infect Dis* 2003; **36**(11): 1433-7.
- [2] Nascimento GGF, Locatelli J, Freitas PC, Silva GL. Antibacterial activity of plant extracts and phytochemicals on antibiotic-resistant bacteria. *Braz J Microbiol* 2000; **31**(4): 247-56.
- [3] Cowan MM. Plant products as antimicrobial agents. *Clin Microbiol Rev* 1999; **12**(4): 564-82.
- [4] Kaushik P, Goyal P. *In vitro* evaluation of *Datura innoxia* (thorn-apple) for potential antibacterial activity. *Indian J Microbiol* 2008; **48**(3): 353-7.
- [5] Bolla JM, Alibert-Franco S, Handzlik J, Chevalier J, Mahamoud A, Boyer G, et al. Strategies for bypassing the membrane barrier in multidrug resistant Gram-negative bacteria. *FEBS Lett* 2011; **585**(11): 1682-90.
- [6] Lee CK, Kim H, Moon KH, Shin KH. Screening and isolation of antibiotic resistance inhibitors from herb materials-resistance inhibition of volatile components of Korean aromatic herbs. *Arch Pharm Res* 1998; **21**(1): 62-6.
- [7] Sivananthan M. Antibacterial activity of 50 medicinal plants used in folk medicine. *Int J Biosci* 2013; **3**(4): 104-21.
- [8] Grosvenor PW, Supriono A, Gray DO. Medicinal plants from Riau Province, Sumatra, Indonesia. Part 2: antibacterial and antifungal activity. *J Ethnopharmacol* 1995; **45**(2): 97-111.
- [9] Ahmad I, Beg AZ. Antimicrobial and phytochemical studies on 45 Indian medicinal plants against multi-drug resistant human pathogens. *J Ethnopharmacol* 2001; **74**(2): 113-23.
- [10] Abu-Shanab BA, Adwan G, Adwan KM, Abu-Shanab F. Efficacy of aqueous and ethanol extracts of some Palestinian medicinal plants for potential antibacterial activity. *Islam Univ J Gaza* 2008; **16**(2): 77-86.
- [11] Panda SK. Ethno-medicinal uses and screening of plants for antibacterial activity from Similipal Biosphere Reserve, Odisha. *India. J Ethnopharmacol* 2014; **151**(1): 158-75.
- [12] Raut JS, Karuppayil SM. A status review on the medicinal properties of essential oils. *Ind Crops Prod* 2014; **62**: 250-64.
- [13] Elfahmi, Woerdenbag HJ, Kayser O. *Jamu*: Indonesian traditional herbal medicine toward rational phytopharmacological use. *J Herb Med* 2014; **4**(2): 51-73.
- [14] Tan YP, Chan EWC. Antioxidant, antityrosinase and antibacterial properties of fresh and processed leaves of *Anacardium occidentale* and *Piper betle*. *Food Biosci* 2014; **6**: 17-23.
- [15] Aumeeruddy-Elalfi Z, Gurib-Fakim A, Mahomoodally F. Antimicrobial, antibiotic potentiating activity and phytochemical profile of essential oils from exotic and endemic medicinal plants of Mauritius. *Ind Crops Prod* 2015; **71**: 197-204.
- [16] Akhtar N, Ihsan-ul-Haq, Mirza B. Phytochemical analysis and comprehensive evaluation of antimicrobial and antioxidant properties of 61 medicinal plant species. *Arab J Chem* 2015; <http://dx.doi.org/10.1016/j.arabjc.2015.01.013>.
- [17] Marasini B, Baral P, Aryal P, Ghimire KR, Neupane S, Dahal N, et al. Evaluation of antibacterial activity of some traditionally used medicinal plants against human pathogenic bacteria. *Biomed Res Int* 2015; <http://dx.doi.org/10.1155/2015/265425>.
- [18] Quisumbing E. *Medicinal plants of the Philippines*. Quezon: Katha Publishing Co., Inc; 1978.
- [19] Arellano Law Foundation. An act creating the Philippine Institute of Traditional and Alternative Health Care (Pitahc) to accelerate the development of traditional and alternative health care in the Philippines, providing for a traditional and alternative health care development fund and for other purposes. Philippines: Arellano Law Foundation. 1997. [Online] Available from: http://www.lawphil.net/statutes/repacts/ra1997/ra_8423_1997.html [Accessed on 21st March, 2015]
- [20] Boucher HW, Talbot GH, Bradley JS, Edwards JE, Gilbert D, Rice LB, et al. Bad bugs, no drugs: no ESKAPE! An update from the Infectious Diseases Society of America. *Clin Infect Dis* 2009; **48**(1): 1-12.
- [21] Veldman K, Kant A, Dierikx C, van Essen-Zandbergen A, Wit B, Mevius D. Enterobacteriaceae resistant to third-generation cephalosporins and quinolones in fresh culinary herbs imported from Southeast Asia. *Int J Food Microbiol* 2014; **177**: 72-7.
- [22] Neonakis IK, Spandidos DA, Petinaki E. Confronting multidrug-resistant *Acinetobacter baumannii*: a review. *Int J Antimicrob Agents* 2011; **37**(2): 102-9.
- [23] Ranotkar S, Kumar P, Zutshi S, Prashanth KS, Bezbaruah B, Anand J, et al. Vancomycin-resistant enterococci: troublemaker of the 21st century. *J Glob Antimicrob Resist* 2014; **2**(4): 205-12.
- [24] Lai CC, Lee K, Xiao YH, Ahmad N, Veeraraghavan B, Thamlikitkul V, et al. High burden of antimicrobial drug resistance in Asia. *J Glob Antimicrob Resist* 2014; **2**(3): 141-7.
- [25] Basri D, Fan SH. The potential of aqueous and acetone extracts of galls of *Quercus infectoria* as antibacterial agents. *Indian J Pharmacol* 2005; **37**(1): 26-9.
- [26] Clinical and Laboratory Standards Institute. Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically. Wayne: Clinical Laboratory Standards Institute. 2009. [Online] Available from: <http://antimicrobianos.com.ar/ATB/wp-content/uploads/2012/11/03-CLSI-M07-A9-2012.pdf> [Accessed on 21st March, 2014]
- [27] Irobi ON, Daramola SO. Bactericidal properties of crude extracts of *Mitracarpus villosus*. *J Ethnopharmacol* 1994; **42**(1): 39-43.
- [28] Tsuchiya H, Sato M, Miyazaki T, Fujiwara S, Tanigaki S, Ohyama M, et al. Comparative study on the antibacterial activity of phytochemical flavanones against methicillin-resistant *Staphylococcus aureus*. *J Ethnopharmacol* 1996; **50**(1): 27-34.
- [29] Aqil F, Khan MS, Owais M, Ahmad I. Effect of certain bioactive plant extracts on clinical isolates of beta-lactamase producing methicillin-resistant *Staphylococcus aureus*. *J Basic Microbiol* 2005; **45**(2): 106-14.
- [30] Sato M, Tanaka H, Oh-Uchi T, Fukai T, Etoh H, Yamaguchi R. Antibacterial activity of phytochemicals isolated from *Erythrina zeyheri* against vancomycin-resistant enterococci and their combinations with vancomycin. *Phytother Res* 2004; **18**(11): 906-10.
- [31] Ahmad I, Aqil F. *In vitro* efficacy of bioactive extracts of 15 medicinal plants against ES β L-producing multidrug-resistant enteric bacteria. *Microbiol Res* 2007; **162**(3): 264-75.
- [32] Jesonbabu J, Spandana N, Aruna Lakshmi K. *In vitro* antimicrobial potentialities of chloroform extracts of ethnomedicinal plant against clinically isolated human pathogens. *Int J Pharm Pharm Sci* 2012; **4**(3): 624-6.
- [33] Chowdhury AN, Ashrafuzzaman M, Ali H, Liza LN, Zinnah KMA. Antimicrobial activity of some medicinal plants against multidrug resistant human pathogens. *Adv Biosci Bioeng* 2013; **1**(1): 1-24.
- [34] Fankam A, Kuate JR, Kuete V. Antibacterial activities of *Beilschmiedia obscura* and six other Cameroonian medicinal plants against multidrug-resistant Gram-negative phenotypes. *BMC Complement Altern Med* 2014; **14**: 241.
- [35] Buru AS, Pichika MR, Neela V, Mohandas K. *In vitro* antibacterial effects of *Cinnamomum* extracts on common bacteria found in wound infections with emphasis on methicillin-resistant *Staphylococcus aureus*. *J Ethnopharmacol* 2014; **153**(3): 587-95.
- [36] Chakraborty B, Nath A, Saikia H, Sengupta M. Bactericidal activity of selected medicinal plants against multidrug-resistant bacterial strains from clinical isolates. *Asian Pac J Trop Med* 2014; [http://dx.doi.org/10.1016/S1995-7645\(14\)60271-6](http://dx.doi.org/10.1016/S1995-7645(14)60271-6).
- [37] Shari SP, Vincy M, Brilliant R. *In vitro* antibacterial efficacy of medicinal plants against multidrug-resistant bacteria a comparative analysis. *Asian Acad Res J Multidiscip* 2014; **1**(27): 356-69.
- [38] Rath S, Padhy RN. Antibacterial efficacy of five medicinal plants against multidrug-resistant enteropathogenic bacteria infecting under-5 hospitalized children. *J Integr Med* 2015; **13**(1): 45-57.
- [39] Esimone CO, Attama AA, Mundi KW, Ibekwe NN, Chah KF. Antimicrobial activity of *Psidium guajava* Linn. stem extracts against methicillin-resistant *Staphylococcus aureus*. *Afr J Biotechnol* 2012; **11**(89): 15556-9.
- [40] Rahim N, Gomes DJ, Watanabe H, Rahman SR, Chomvarin C, Endtz HP, et al. Antibacterial activity of *Psidium guajava* leaf and bark against multidrug-resistant *Vibrio cholerae*: implication for cholera control. *Jpn J Infect Dis* 2010; **63**(4): 271-4.

- [41] Anas K, Jayasree PR, Vijayakumar T, Manish Kumar PR. *In vitro* antibacterial activity of *Psidium guajava* Linn. leaf extract on clinical isolates of multidrug-resistant *Staphylococcus aureus*. *Indian J Exp Biol* 2008; **46**(1): 41-6.
- [42] Pesewu GA, Cutler RR, Humber DP. Antibacterial activity of plants used in traditional medicines of Ghana with particular reference to MRSA. *J Ethnopharmacol* 2008; **116**(1): 102-11.
- [43] Mishra A, Babele P. A comparative study of antibacterial activity of crude extracts preparations of five medicinal plants: *Ocimum sanctum*, *Ajadirachta indica*, *Punica granatum*, *Psidium guajava*, and *Syzygium aromaticum* against multidrug-resistant bacteria of clinical origin. *World J Pharm Res* 2014; **3**(5): 1174-87.
- [44] Campos AH, Schor N. *Phyllanthus niruri* inhibits calcium oxalate endocytosis by renal tubular cells: its role in urolithiasis. *Nephron* 1999; **81**(4): 393-7.
- [45] Shanbhag T, Amuthan A, Shenoy S, Sudhakar. Effect of *Phyllanthus niruri* Linn. on burn wound in rats. *Asian Pac J Trop Med* 2010; **3**(2): 105-8.
- [46] Bokarey MR, Thakre PP, Wadhai VS. Antibacterial activity and phytochemical analysis of *Phyllanthus niruri* against common human pathogenic bacteria. *Rev Res* 2012; **1**(6): 30.
- [47] Ramji N, Ramji N, Iyer R, Chandrasekaran S. Phenolic antibacterials from *Piper betle* in the prevention of halitosis. *J Ethnopharmacol* 2002; **83**(1-2): 149-52.
- [48] Baliga MS, Fazal F, Priya MR, Ratnu VS, Rai MP. Betel leaf (*Piper betel* Linn.): the wrongly maligned medicinal and recreational plant possesses potent gastrointestinal and hepatoprotective effects. In: Watson RR, Preedy VR, editors. *Bioactive food as dietary interventions for liver and gastrointestinal disease*. California: Academic Press; 2013.
- [49] Khan JA, Kumar N. Evaluation of antibacterial properties of extracts of *Piper betle* leaf. *J Pharm Biomed Sci* 2011; **11**(11): 1-3.
- [50] Agarwal T, Singh R, Shukla AD, Waris I, Gujrati A. Comparative analysis of antibacterial activity of four *Piper betel* varieties. *Adv Appl Sci Res* 2012; **3**(2): 698-705.
- [51] Datta A, Ghoshdastidar S, Singh M. Antimicrobial property of *Piper betle* leaf against clinical isolates of bacteria. *Int J Pharm Sci Res* 2011; **2**: 104-9.
- [52] Nouri L, Nafchi AM, Karim AA. Phytochemical, antioxidant, antibacterial, and α -amylase inhibitory properties of different extracts from betel leaves. *Ind Crops Prod* 2014; **62**: 47-52.
- [53] Farzaneh V, Carvalho IS. A review of the health benefit potentials of herbal plant infusions and their mechanisms of actions. *Ind Crops Prod* 2015; **65**: 247-58.
- [54] Piddock LJ. Clinically relevant chromosomally encoded multidrug resistance efflux pumps in bacteria. *Clin Microbiol Rev* 2006; **19**(2): 382-402.