

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

Biochimica et Biophysica Acta 1757 (2006) 206–214

<http://www.elsevier.com/locate/bba>

A functionally inactive, cold-stabilized form of the *Escherichia coli* F₁F_o ATP synthase

Mikhail A. Galkin¹, Robert R. Ishmukhametov, Steven B. Vik*

Department of Biological Sciences, Southern Methodist University, Dallas, TX 75275-0376, USA

Received 22 December 2005; received in revised form 15 February 2006; accepted 20 February 2006

Available online 20 March 2006

Abstract

An unusual effect of temperature on the ATPase activity of *E. coli* F₁F_o ATP synthase has been investigated. The rate of ATP hydrolysis by the isolated enzyme, previously kept on ice, showed a lag phase when measured at 15 °C, but not at 37 °C. A pre-incubation of the enzyme at room temperature for 5 min completely eliminated the lag phase, and resulted in a higher steady-state rate. Similar results were obtained using the isolated enzyme after incorporation into liposomes. The initial rates of ATP-dependent proton translocation, as measured by 9-amino-6-chloro-2-methoxyacridine (ACMA) fluorescence quenching, at 15 °C also varied according to the pre-incubation temperature. The relationship between this temperature-dependent pattern of enzyme activity, termed thermohysteresis, and pre-incubation with other agents was examined. Pre-incubation of membrane vesicles with azide and Mg²⁺, without exogenous ADP, resulted in almost complete inhibition of the initial rate of ATPase when assayed at 10 °C, but had little effect at 37 °C. Rates of ATP synthesis following this pre-incubation were not affected at any temperature. Azide inhibition of ATP hydrolysis by the isolated enzyme was reduced when an ATP-regenerating system was used. A gradual reactivation of azide-blocked enzyme was slowed down by the presence of phosphate in the reaction medium. The well-known Mg²⁺ inhibition of ATP hydrolysis was shown to be greatly enhanced at 15 °C relative to at 37 °C. The results suggest that thermohysteresis is a consequence of an inactive form of the enzyme that is stabilized by the binding of inhibitory Mg-ADP.

© 2006 Elsevier B.V. All rights reserved.

Keywords: F₁F_o; ATP synthase; Thermohysteresis; Azide; Mg-ADP inhibition

1. Introduction

F₁F_o-ATP synthases, present in mitochondria, chloroplasts and bacteria, catalyze ATP synthesis driven by proton or sodium translocation across the membrane. In some bacteria, this enzyme can also function as an ion translocator driven by ATP hydrolysis. The enzyme consists of two distinct parts: F₁, containing six nucleotide-binding sites and three catalytic centers, and membrane-embedded F_o responsible for proton translocation [1]. In the *E. coli* ATP synthase F₁ is composed of five subunits in a stoichiometry of α₃β₃γδε [2]. F_o consists of

three subunits in a stoichiometry of ab₂c₁₀ [3,4]. It is well established that the coupling between the ATP hydrolytic reaction and proton translocation is mediated by rotation of the central stalk γεc₁₀ (rotor) relative to the stator (α₃β₃ab₂δ) [5–11]. F₁ can be separated physically from F_o and it catalyses energy-uncoupled ATP hydrolysis at a high rate. The isolated F_o can act as a passive proton pore. Despite great success in studying this nano-motor, many mechanistic aspects of its function remain unknown. The *E. coli* ATP synthase is among the simplest of those that have been studied. Studies of the homologous enzymes from *Bacillus PS3*, *P. modestum* and *I. tartaricus* have also contributed to our understanding of ATP synthesis [9,12,13]. These enzymes share many, but certainly not all, features with the more complex enzymes found in mitochondria.

A common feature of ATP synthases is that the soluble F₁-ATPases, but not the whole complex F₁F_o, undergoes cold inactivation accompanied by dissociation of the enzyme into

Abbreviations: DCCD, N,N'-dicyclohexylcarbodiimide; FCCP, carbonyl cyanide *p*-(trifluoromethoxy)phenylhydrazine; ACMA, 9-amino-6-chloro-2-methoxyacridine; LDAO, lauryldimethylamine oxide

* Corresponding author. Tel.: +1 214 768 4228; fax: +1 214 768 3955.

E-mail address: svik@smu.edu (S.B. Vik).¹ Present address: Department of Molecular Physiology and Biological Physics, University of Virginia School of Medicine, Charlottesville, VA 22908, USA.

sub-complexes [14–16]. The intact F_1F_o complex is less sensitive to incubation at low temperature, and its sensitivity to the cold has been interpreted as mediated by dissociation of F_1 from the membrane and subsequent cold inactivation of the soluble F_1 [14]. A different type of temperature effect on the rate of ATP hydrolysis by the *E. coli* F_1 -ATPase was described by Laget over 25 years ago [17]. The rate of ATP hydrolysis as measured at 10 °C was found to be dependent upon the incubation temperature of the enzyme prior to the assay. This was interpreted as a slow conversion between two conformational states of the enzyme, termed a high activity state and a low activity state. It was called a hysteretic property, according to the definition of Frieden [18,19], in which such an enzyme responds slowly to a rapid change in ligand concentration. In this case, it is a slow response to a change in temperature, and so can be referred to as thermohysteresis.

Laget also examined the effects of ATP and ADP on the hysteretic properties of the *E. coli* F_1 -ATPase [20]. Pre-incubation of the enzyme with ATP or with ADP affected both the equilibrium between the two states, and the kinetic properties of the two states. Both nucleotides shifted the equilibrium towards the high activity state, but they had different effects on kinetic properties. The results were interpreted in terms of a high activity state that is stabilized by binding ATP, and a low activity state that is stabilized by binding ADP. The ratio of ATP/Mg²⁺ was kept constant at 2.0 during assays, and thus the role of Mg²⁺ was not investigated.

An intriguing feature of mitochondrial and chloroplast F_1 -ATPases is unidirectional inhibition by Mg-ADP in the direction of hydrolysis of ATP [21–24]. Hysteretic behavior with respect to ADP binding has been described in a variety of systems [20,21,25,26]. Mg-ADP inhibition has been studied extensively in the F_1 -ATPase from the thermophilic *Bacillus* PS3 [27–29], but less so in the enzyme from *E. coli* [30,31]. In the case of the bacterial F_1 -ATPase, studies of inhibition can be complicated by the epsilon subunit, an endogenous inhibitor that can partially dissociate during assay conditions [32,33].

Recent studies in this laboratory have resulted in a rapid procedure for isolation of the ATP synthase from *E. coli* in a highly active form [34]. Procedures for the measurement of ATP hydrolysis and ATP-dependent proton translocation in continuous assay systems were optimized. During the course of these studies an unusual temperature dependence of ATP hydrolysis by the isolated F_1F_o was observed, which resembled prior results obtained by Laget using F_1 -ATPase [15,18]. Therefore, the sensitivity of F_1F_o to low temperature was explored systematically, and its possible relationship to Mg-ADP inhibition was examined.

2. Materials and methods

2.1. Purification and preparations

F_1F_o was purified from *E. coli* strain DK8 harboring plasmid pFV2, as previously described [34]. Growth of cultures, preparation of membrane vesicles, purification of F_1F_o , and reconstitution into liposomes were also carried out as described previously [34]. Azide-treated membrane vesicles, termed N-particles, were prepared by two-fold dilution with medium containing 250 mM

sucrose, 10 mM HEPES, pH 8.0, 5 mM MgCl₂, 100 mM KCl, 0.1 mM EDTA, 2 mM malonate (potassium salt), 1 mg/ml bovine serum albumin, and 2 mM NaN₃. Phosphate-treated membrane vesicles, termed P-particles, were prepared by two-fold dilution with medium containing 250 mM sucrose, 10 mM HEPES, pH 8.0, 10 mM phosphate (sodium salt), 100 mM KCl, 0.1 mM EDTA, 2 mM malonate, 1 mg/ml bovine serum albumin. Pretreatment was carried out for 1 h at room temperature.

2.2. Functional assays

ATP hydrolysis activity was measured either with an ATP regenerating system or with the pH-indicator phenol red, using the following conditions, unless otherwise specified. For measurements with an ATP regenerating system the medium (1 ml) contained 10 mM HEPES/KOH, pH 8.0, 100 mM KCl, 2.5 mM MgCl₂, 0.1 mM EDTA, 1 mM ATP, 200 μM NADH, 2 mM phosphoenolpyruvate, lactate dehydrogenase (5 units/ml), pyruvate kinase (5 units/ml), and 5 μM FCCP (in the case of reconstituted F_1F_o). Activity was measured by following the decrease in optical density at 340 nm. Using the pH-indicator phenol red, ATP hydrolysis activity was measured at 557 nm by following scalar proton release, essentially as described [34]. The medium (2 ml) contained 10 mM HEPES/KOH, pH 8.0, 100 mM KCl, 10 mM ATP, 4 mM MgCl₂, 0.1 mM EDTA, 60 μM phenol red, and 5 μM FCCP (in the case of membrane vesicles). The phenol red assay was always checked at pH 8.0 to ensure that the ratio of protons released to phosphate released during ATP hydrolysis was approximately 1.0. Measurements of ATP-dependent ACMA-fluorescence quenching were performed as described previously [34]. In all assays, volumes of enzyme added were 5–20 μl per reaction. ATP synthesis by preparations of membrane vesicles was monitored using a modified PTI Delta-Scan 1 fluorimeter, which allowed real-time measurement of ATP synthesis using luciferase. Prior to measurement of ATP synthesis, membrane vesicles were resuspended in 1 ml of French press buffer (200 mM Tricine/HCl, pH 7.8, 100 mM KCl, 5 mM MgCl₂, 0.1 mM EDTA, and 2.5% glycerol), passed through a 10-ml Sephadex column, equilibrated with the same buffer, ultracentrifuged as before, and resuspended in French press buffer. This step eliminated any energization-independent ATP synthase activity. The membranes were diluted twice with buffer used for preparation of P-particles. The reaction was initiated by addition of 20 mM succinate to 2 ml of a medium containing 200 μg membrane protein, 10 mM Tricine/KOH, pH 8.0, 100 mM KCl, 5 mM MgCl₂, 1 mM P_i, 0.1 mM ADP, 125 μM luciferin, and 100 ng luciferase. 20 nmol ATP was added for calibration after each reaction was finished. No ATP synthesis was observed without ADP or P_i, or with uncoupler FCCP.

2.3. Analytical methods

Native electrophoresis and determination of protein concentration were performed as described earlier [34].

3. Results

3.1. Temperature dependence of ATP hydrolysis

Fig. 1A shows the time course of ATP hydrolysis by isolated *E. coli* F_1F_o using an ATP-regenerating assay at 15 °C. When enzyme was taken directly from ice (“cold” enzyme), its activity demonstrated a pronounced lag phase (trace b). Conversely, the activity of the enzyme after pre-incubation for 5 min at room temperature (“hot” enzyme) showed no lag phase, and was higher than in the first case (trace a). If the “hot” enzyme was returned to ice for 5 min before activity measurement, the time course became similar to that of the “cold” one (trace c). When the enzyme was subjected to additional cycles of warming and cooling, it exhibited the behavior of the “hot” or “cold” enzyme, according to its last incubation (data not shown). When the activity measurements were made at 37 °C, the duration of the




Fig. 1. Time-course of ATP hydrolysis by isolated F_1F_o (A, B) and by F_1F_o incorporated into proteoliposomes (C, D) at 15 °C (A, C) and at 37 °C (B, D). ATPase activity was measured with an ATP regenerating system as described in Materials and methods. The reactions were initiated by addition of 2.5 μg (A), 0.7 μg (B), 1.2 μg (C) or 0.4 μg (D) of protein. In all panels trace a represents “hot” enzyme, in which the sample was taken from ice and pre-incubated at room temperature for 5 min before measurement. All traces b represent “cold” enzyme, in which the sample was taken directly from ice for measurement. In panels A and C trace c represents enzyme that was taken from ice, then incubated at room temperature for 5 min, and returned to ice for 5 min, before measurement, and the lines marked d represent the initial rates of “cold” enzyme (trace b). Numbers shown near the curves indicate the rates of ATP hydrolysis expressed in μmol ATP per min per mg of protein.

lag phase for the “cold” enzyme (Fig. 1B, trace b) was much shorter, and there was no significant difference in the steady-state rates of the two forms of enzyme. In panel A the initial rate of the “cold” enzyme is shown by the straight line marked d. In five different experiments, the average ratio of the initial rates (traces d and a) was 0.53 ± 0.06 for isolated F_1F_o .

The same set of experiments were conducted for the case of F_1F_o reconstituted into liposomes (Fig. 1C, D). Similar patterns of activity were observed: a lag phase for the “cold” enzyme when assayed at 15 °C with a reversible transition between the “hot” and “cold” enzyme (Fig. 1C), and no significant difference between the two forms of enzyme when assayed at 37 °C (Fig. 1D). In panel C, the initial rate of the “cold” enzyme is shown by the straight line marked d. In five different experiments, the average ratio of the initial rates (traces d and a) was 0.49 ± 0.06 for proteoliposomes.

The two forms of enzyme were also tested for ATP-dependent proton translocation by the ACMA fluorescence quenching assay. At 15 °C, the initial rate of ATP-dependent ACMA fluorescence quenching by the “cold” enzyme, previously reconstituted into liposomes (Fig. 2A, trace b),

was about one half that of the “hot” enzyme (trace a). In eight experiments, the average ratio of initial rates was 0.54 ± 0.03 . As before, the rate of quenching by the “hot” reconstituted enzyme, returned to ice for 5 min, was the same as that of the “cold” enzyme (trace c). There was no difference in the initial rate of quenching between the two forms of enzyme when the assay was carried out at 37 °C (Fig. 2B). One difference between the ATP hydrolysis assay, shown in Fig. 1, and the proton translocation assay, shown in Fig. 2, is the length of time from addition of the enzyme until the start of recording. For technical reasons, this interval is about 30 s for ATP hydrolysis, and about 1 min for proton translocation, and this might affect the estimation of the initial rates.

3.2. Dissociation of the enzyme

As was first shown about 40 years ago, soluble F_1 -ATPase promptly loses its activity at low temperature [15]. This cold inactivation is accompanied by dissociation of F_1 into sub-complexes. In contrast, the thermohysteretic behavior of *E. coli* F_1 -ATPase as discovered by Laget [17], is not




Fig. 2. ATP-dependent ACMA fluorescence quenching by reconstituted F_1F_o at 15 °C (A) and at 37 °C (B). The assay was carried out as described in Materials and methods using 1.2 μ g of protein. The reaction was initiated by the addition of 0.2 mM ATP. In both panels trace a represents “hot” enzyme, in which the sample was taken from ice and pre-incubated at room temperature for 5 min before measurement; trace b represents “cold” enzyme, in which the sample was taken directly from ice for measurement. In panel A trace c represents enzyme that was taken from ice, then incubated at room temperature for 5 min, and returned to ice for 5 min before measurement.

due to cold dissociation of the subunits. To check for the possibility of subunit dissociation by F_1F_o , two approaches were used: estimation of ATPase activity after multiple cycles of temperature variation, and analysis of treated samples by native gel electrophoresis.

In the first approach, the “cold” enzyme was frozen at -20 °C for 1 h and subsequently thawed at 0 °C for 40 min. The thawed enzyme was divided into two portions. One portion was incubated on ice, and the other was incubated at room temperature for 5 min before measurement of activity. The time course of ATP hydrolysis by these enzymes was similar to

the unfrozen “hot” and “cold” ones. Recovery of the ATPase activity of the “hot” enzyme after freeze–thawing was 80% (data not shown).

In a second approach, F_1F_o was first pre-incubated either on ice or at room temperature for 5 min, and then frozen rapidly in liquid nitrogen (<5 s). The frozen suspensions were thawed at 37 °C until reaching 0 °C (about 5 s) and immediately thereafter the ATPase activity was measured. Even this severe freeze–thaw cycle did not considerably change the time courses (or activities) of ATP hydrolysis observed, relative to the unfrozen “hot” and “cold” enzymes, shown previously.

Finally, the possibility of subunit dissociation was investigated directly by native gel electrophoresis. F_1F_o was taken up either from ice or after a slow cycle of freezing–thawing and was run on native gels at 4 °C, and at room temperature, for 50 min, as described previously [34]. At both temperatures only a single band was observed, indicating that F_1F_o remained intact under all conditions and did not undergo cold dissociation (data not shown).

3.3. Investigations into the mechanism of thermohysteretic behavior

F_1 -ATPases are well known to be inhibited generally by ADP, and the *E. coli* enzyme in particular is inhibited by Mg^{2+} ions [34,35]. Furthermore, a Mg-ADP inhibited form of F_1 has been identified in several different species, leading to hysteretic behavior [36,37]. Azide is thought to stabilize the Mg-ADP inhibited form of F_1 , [31,37,38] although in *E. coli* this is controversial [30,39]. The *E. coli* F_1 is inhibited by azide, with a K_i of 25 μ M at 37 °C and 7 μ M at 25 °C [40]. Therefore, the relationship of Mg^{2+} ions, ADP and azide to the thermohysteretic properties described above was investigated.

In the previous experiments (Fig. 1), ATP hydrolysis was measured using an ATP regenerating system, in which little or no ADP accumulates during the assay. To investigate the consequences of the accumulation of ADP as ATP is hydrolyzed, a phenol red system was also used. The difference between the two assay systems can be demonstrated by comparing the rates of ATP hydrolysis by the isolated F_1F_o ATP synthase, previously incubated for 5 min at room temperature (“hot” enzyme), at both 37 °C and at 15 °C. At 37 °C the rate found in the phenol red assay system is 95% of that using the ATP regenerating system. At 15 °C the rate in the phenol red system is only 31% of that in the ATP regenerating system. This suggests that the isolated F_1F_o is especially sensitive to inhibition by ADP at low temperatures. Furthermore, in the presence of 100 μ M azide, while the enzyme is seen to be inhibited using both assay systems, the rate is essentially zero using the phenol red assay, and is less than 5% of the rate in the ATP regenerating system.

In contrast to the isolated enzyme (Fig. 1), the ATPase activity of membrane vesicles did not depend upon the temperature of pre-incubation when measured at 15 °C as

shown in Fig. 3A. In trace b, the membranes were taken directly from ice, and in trace a, the membrane were pre-incubated for 5 min at room temperature. There was also no effect of pre-incubation when assayed at 37 °C (data not shown). These measurements were made using the phenol red assay, to avoid complications due to NADH dehydrogenases present in the membranes. For comparison, the results from isolated F_1F_0 assayed under the same conditions of Fig. 3A are shown in panel B, in which a lag phase for the “cold” enzyme is apparent. Pre-incubation of the enzyme (isolated, reconstituted F_1F_0 , or in membrane vesicles) on ice for more than 5 min did not change the pattern of activities seen in Figs. 1, 2, or 3. The traces shown are typical ones from several experiments.

The addition of up to 200 μ M ADP to the pre-incubation medium had no effect on ATPase activity under all conditions tested. In contrast, pre-incubation with 1 mM azide, even in the




Fig. 3. Time-courses of ATP hydrolysis by membrane vesicles (A) and by isolated F_1F_0 (B) without an ATP-regenerating assay system. In both panels, ATPase activity was measured with phenol red at 15 °C as described in Materials and methods, using 220 μ g (A) or 7 μ g (B) of protein. In both panels trace a represents “hot” enzyme, in which the sample was taken from ice and pre-incubated at room temperature for 5 min before measurement; trace b represents “cold” enzyme, in which the sample was taken directly from ice for measurement. Numbers shown near the curves indicate the rates of ATP hydrolysis expressed in μ mol ATP per min per mg of protein.

absence of exogenous ADP, had profound effects on the initial rates of ATPase activity, as shown by the following experiments. Membrane vesicles that were pre-incubated with 1 mM azide for 1 h at room temperature (called N-particles) were compared with those that were pre-incubated without azide (control). The results, as shown in Table 1, indicate that azide is a potent inhibitor of the initial rate of ATPase activity, in a way that depends upon both temperature and the concentrations of ATP and Mg^{2+} . When assayed at 10 °C, membrane vesicles had initial rates of ATP hydrolysis that varied with the ATP concentration (Mg^{2+} was kept at 4 mM). Under the same assay conditions, membranes pre-incubated with azide (N-particles) had initial rates of zero. After 7 min, the rates increased somewhat, if assayed at higher concentrations of ATP (Table 1 and Fig. 4). If the assay medium also included 10 mM P_i , the lag phase was extended, and the rate decreased, as shown in Fig. 4 (comparing traces b and c). Inhibition of initial rates due to pre-incubation with azide was completely eliminated if the assay medium included 0.3% LDAO (Table 1, assay 4). Under assay conditions of 10 mM ATP/4 mM Mg^{2+} , if the assay temperature was increased from 10 °C (assay 3) to 25 °C (assay 5) or to 37 °C (assay 6) the initial rates increased dramatically at the higher temperatures. In addition, the effect of pre-incubation with azide decreased from 100% inhibition at 10 °C to only 6% at 37 °C.

The effects of azide seen here are due primarily to the pre-incubation, since if the equivalent amount of azide is diluted into the assay medium the inhibition is only about 7%, if measured at 10 °C. Furthermore, the effects of azide are limited to ATP hydrolysis, since pre-incubation of membrane vesicles with azide had no effect on the rates of ATP synthesis when assayed at 5 °C to 22 °C. For example, at 10 °C, the measured rates of ATP synthesis were between 11 and 13 nmol ATP/min/mg protein for N-particles and control particles, with or without a pre-incubation with 2 mM P_i . No lag phase was apparent at any temperature. Likewise, the rates at 22 °C were between 28 and 30 nmol ATP/min/mg protein in each case.

The effect of Mg^{2+} concentration on the thermohysteretic properties of F_1F_0 was also investigated, and the results are presented in Fig. 5. The ATP concentration was fixed at 10 mM and the Mg^{2+} concentration was varied from 1 to 20 mM. Rates of ATP hydrolysis by isolated F_1F_0 (panel A) and by membrane vesicles (panel B) were measured using the phenol red method. Samples were pre-incubated for 5 min at room temperature, and assayed at 15 °C (filled circles) or 37 °C (open circles). The rates of ATP hydrolysis by isolated F_1F_0 at both temperatures show a similar dependence upon Mg^{2+} concentration (A), but by plotting the ratio of the rate at 15 °C over the rate at 37 °C it can be seen that ATP hydrolysis by isolated F_1F_0 at 15 °C is increasingly inhibited by Mg^{2+} (panel C, squares). This indicates that Mg^{2+} is an essential component of the thermohysteretic properties of the “cold” enzyme. In contrast, the ratio of the rates of ATP hydrolysis by membrane vesicles at the two temperatures does not depend upon Mg^{2+} concentration (panel 6C, triangles). This is consistent with the prior observation (Fig. 4) that F_1F_0 in membrane vesicles does not exhibit thermohysteretic properties. A plot of the ratio of rates at the two

Table 1
Inhibition of ATP hydrolysis by pre-incubation with azide at 23 °C^a

Assay number	Assay conditions ^b (mM ATP/mM Mg ²⁺)	Initial rates of ATP hydrolysis ($\mu\text{mol}/\text{min}/\text{mg}$ protein)		Ratio of rates (N-particles/Control)	Rates of ATP hydrolysis by N-particles after 7 min in assay medium
		Control	N-particles		
1	10 °C (0.5/4)	0.16	0	0	0
2	10 °C (2/4)	0.27	0	0	0.045
3	10 °C (10/4)	0.31	0	0	0.12
4	10 °C (0.5/4) 0.3% LDAO	0.2	0.2	1.0	ND ^c
5	25 °C (10/4)	0.75	0.56	0.74	ND
6	37 °C (10/4)	1.54	1.44	0.94	ND

^a Rates shown are a typical set of values from at least three different experiments with similar results. N-particles were prepared by two-fold dilution of membrane vesicles with medium containing 250 mM sucrose, 10 mM HEPES, pH 8.0, 5 mM MgCl₂, 100 mM KCl, 0.1 mM EDTA, 2 mM malonate (potassium salt), 2 mM NaN₃, and incubated at room temperature for 1 h. Control particles were treated similarly, without the NaN₃.

^b Reactions were initiated by addition of 415 μg membrane protein (up to 20 μl) to 2 ml of medium, using the phenol red system.

^c ND, Not determined.

temperatures by F₁F_o reconstituted into liposomes falls between the other two plots (data not shown).

4. Discussion

The thermohysteretic behavior described previously by Laget [17,20] for hydrolysis of ATP by the soluble F₁-ATPase from *E. coli* has been shown to occur in the isolated F₁F_o ATP

synthase. Laget showed that the rate of ATP hydrolysis at 10 °C depends upon the temperature of pre-incubation of the enzyme. He found that F₁-ATPase incubated at low temperature had lower rates than did identical samples pre-incubated at higher temperatures. We found a similar dependence upon the rate of ATP hydrolysis by the isolated F₁F_o ATP synthase when measured at 15 °C. These results appear to extend to both ATP hydrolysis and ATP-driven proton translocation by the isolated enzyme, reconstituted in liposomes. Remarkably, ATP hydrolysis by membrane vesicles at 15 °C did not exhibit a lag phase when pre-incubated on ice, suggesting that the transition was too rapid to be observed. This might be a consequence of the native lipid environment, but further studies will be necessary to understand this.

Hysteretic behavior was not a consequence of the well-known cold denaturation of F₁-ATPase [14–16]. Native gel electrophoresis showed that subunit dissociation did not occur as a result of the incubation on ice, and multiple cycles of pre-incubation temperatures showed that this transition was readily reversible. As was concluded by Laget [17,20] with respect to F₁, this behavior is consistent with two states of the enzyme in equilibrium: a high activity state and a low activity state. At temperatures above 25 °C, the states are in rapid equilibrium, and the high activity state is favored. At temperatures below 15 °C, the low activity state becomes more prevalent, and the rate of inter-conversion between the two states is reduced.

Further experiments showed that the low temperature state of the F₁F_o ATP synthase is more sensitive to inhibition by the presence of ADP or Mg²⁺ in the assay medium, and to azide in the pre-incubation medium. Therefore, it is proposed that the low activity state of the isolated F₁F_o can be stabilized by tightly-bound Mg-ADP. Both mitochondrial and chloroplast F₁-ATPases are thought to be subject to inhibition by tightly-bound Mg-ADP [21,26,37,38], leading to hysteretic behavior. This inhibition is gradually lost after the addition of sufficient ATP, such as occurs during an assay of ATP hydrolysis, and can be seen as a lag phase in ATP hydrolysis. The F₁-ATPase from the thermophilic *Bacillus* PS3 has been extensively studied with respect to such inhibition [28,29,41]. For example, the loss of




Fig. 4. Time courses of ATP hydrolysis by membrane vesicles pre-incubated with azide. Rates of ATP hydrolysis were measured using the phenol red assay system at 10 °C in the presence of 4 mM Mg²⁺. Trace *a* shows the rate of control membrane vesicles using 10 mM ATP. Traces *b*–*e* are from azide-treated membranes (N-particles). Trace *b* shows the rate of ATP hydrolysis by N-particle in the presence of 10 mM ATP. Trace *c* is the same as trace *b*, except that the medium contains 10 mM Pi. Trace *d* shows the rate of ATP hydrolysis by N-particles using 2 mM ATP, and trace *e* shows the rate with 0.5 mM ATP. All samples were kept at room temperature prior to the assay. The traces shown were typical ones from at least three sets of measurements.




Fig. 5. The effect of Mg²⁺ concentration on the thermohysteretic properties of F₁F₀. In panel A, the rates of ATP hydrolysis by "hot" isolated F₁F₀ were measured using the phenol red system at 37 °C (open circles) and at 15 °C (filled circles). In panel B, the same measurements were made using membrane vesicles at 37 °C (open circles) and at 15 °C (filled circles). In panel C, the ratios of the rates at 15 °C and at 37 °C are plotted for the isolated F₁F₀ (squares) and for membrane vesicles (triangles).

ATP binding due to mutations at non-catalytic sites was shown to greatly enhance the inhibition of ATP hydrolysis by tightly-bound Mg-ADP. LDAO was found to eliminate Mg-ADP inhibition when included in the assay medium at a concentration of 0.1%, and similar results were shown here in Table 1.

Prior studies of the *E. coli* F₁-ATPase have failed to detect Mg-ADP inhibition of ATP hydrolysis under assay conditions at room temperature or above [30]. Hyndman et al. [31] demonstrated a lag phase in ATP hydrolysis by incubating the F₁-ATPase, free of its epsilon subunit, in Mg-ADP, then diluting into 50 nM ATP, 50 μM Mg²⁺. Since the studies reported here

used intact F₁F₀, there were no complications from dissociation of the inhibitory epsilon subunit that can occur by dilution of the F₁-ATPase during the assay. The low temperatures used in this study appear to have been essential for observation of the lag phase in ATP hydrolysis by isolated F₁F₀.

The apparent initial rate of ATP hydrolysis depends primarily upon temperature, but is also influenced by the method of assay. The thermohysteretic effect is enhanced when using an assay that does not regenerate ATP. It appears that at low temperatures inhibitory ADP binds with high affinity to the isolated F₁F₀. This ADP could be from commonly found endogenous ADP [30,42], from ATP hydrolysis, or from ADP contamination of commercial ATP. Recent results by Turina et al. [43] showed that at 34 °C, the rate of ATP hydrolysis as measured by the phenol red system were virtually identical to the rates measured by an ATP regenerating system, using membrane vesicles from *Rhodobacter capsulatus*. At 23 °C, the rate measured by phenol red was only about 70% of the maximal rate measured using a regenerating system, and it exhibited a slight lag phase. Those authors did not invoke temperature as an explanation of the difference in rates, but our results would support such an interpretation.

Results presented in Fig. 4 suggest that inorganic phosphate might play a similar inhibitory role to azide at low temperature. Fischer et al. [44] showed that ATP hydrolysis by the *E. coli* F₁F₀ ATP synthase in liposomes was inhibited sharply by a combination of inorganic phosphate and ADP in the assay medium. Our results indicate that such inhibition is enhanced at low temperature, and when using a phenol red assay.

The essential role of Mg²⁺ in the thermohysteretic properties of F₁F₀ is also demonstrated by results presented here. The inhibition of ATP hydrolysis by the *E. coli* enzyme due to high concentrations of Mg²⁺ is well known [34,35]. The optimal ratio of Mg²⁺/ATP for ATP hydrolysis is approximately 0.3–0.5, and at higher concentrations Mg²⁺ is inhibitory. Our results show that the response to Mg²⁺ of ATP hydrolysis by isolated F₁F₀ is temperature dependent. Above the optimal concentration of Mg²⁺, the isolated enzyme is increasingly inhibited at 15 °C relative to at 37 °C, as shown in Fig. 5C.

Inhibition of the mitochondrial and chloroplast F₁-ATPase by azide is thought to occur by stabilization of the Mg-ADP inhibited state. Hyndman et al. [31] have shown a correlation between inhibition of ATP hydrolysis of the *E. coli* F₁-ATPase by azide, and the amount of tightly bound ADP present after passage through a centrifuge column. Their results were seemingly contradicted by studies [39] that showed that 3 molecules of ATP are bound at catalytic sites in the presence of inhibitory azide, by monitoring the fluorescence of reporter tryptophan residues at the catalytic sites. Those results show that ADP binding at catalytic sites is not necessary for inhibition by azide, although ATP binding did occur. Other studies, using the *Bacillus PS3* enzyme, have shown that it may not always be possible to distinguish between the binding of ADP and ATP [45] through such fluorescence measurements. Results presented here show that at low temperatures, and in the presence of Mg-ADP, azide is a potent inhibitor of ATP hydrolysis, but not of ATP synthesis.

The epsilon and gamma subunits are likely to play a role in the conformation of the low activity state, and in the binding of inhibitory Mg-ADP. For example, Yoshida and colleagues [10,46] have discovered a partially-rotated state of the gamma subunit during assays of ATP hydrolysis using video microscopy. Gamma appears to pause after a rotation of about 85°, before it continues to the fully rotated position of 120°. It has been previously suggested that the position of gamma is related to the binding of inhibitory Mg-ADP [47]. The position of the C-terminal alpha-helices of the epsilon subunit have also been implicated in the inhibition of ATP hydrolysis, in a nucleotide and inorganic phosphate dependent way [48,49]. Recently, Yoshida and colleagues [50] have shown that the C-terminus of epsilon and the N-terminus of gamma can form a cross-link by a disulfide bond during conditions that promote ATP synthesis, but not during ATP hydrolysis. LDAO is known to increase the rate of ATP hydrolysis by the *E. coli* F₁ and F₁F_o, and also to prevent a cross-link between the C-terminus of the epsilon and beta subunits [51,52]. We propose that the low activity form of the *E. coli* F₁F_o seen at low temperatures represents a low energy state, in which the gamma subunit is at an intermediate position, and the C-terminal alpha helix of the epsilon subunit is docked near the N-terminus of gamma. At sufficiently low temperatures the enzyme can entrap inhibitory Mg-ADP and show hysteretic behavior when assayed. This might be related to strengthening electrostatic interactions between the C-terminal alpha-helix of epsilon and the DEL-SEED region of the beta subunit. At higher temperatures, the enzyme more easily escapes from the low activity state. Azide is proposed to stabilize the low activity state, increasing the affinity of Mg-ADP.

Studies by Fischer et al. [44] have shown the importance of $\Delta\tilde{\mu}_{H^+}$ in stimulating the rate of ATP hydrolysis by the *E. coli* F₁F_o reconstituted into liposomes and measured at 23 °C. Activation was seen only if ADP or inorganic phosphate, which were inhibitory, were included in the assay. In the absence of ADP and inorganic phosphate the rate of ATP hydrolysis was equal to the activated rate. It can be imagined that this transient energization is able to rotate the gamma subunit in the direction of ATP synthesis, allowing escape from the deactivated state. Lack of a lag phase in ATP synthesis supports this view.

The results presented here also have important implications for interpretations of the temperature dependence of rates of ATP hydrolysis by the *E. coli* ATP synthase. Cold temperature deactivation of an enzyme could lead to breaks in an Arrhenius plot, as has been discussed by Silvius et al. [53,54]. Membrane-dependent effects on the temperature dependence of ATP hydrolysis have been reported with the *E. coli* enzyme [55]. Other studies [56], using the bovine F₁-ATPase have demonstrated breaks in the Arrhenius plot of ATP hydrolysis without a membrane environment. The activation energy at temperatures below the transition temperature, about 18 °C, was diminished if the enzyme was depleted of endogenous nucleotides. Their results were entirely consistent with our findings that the *E. coli* ATP synthase undergoes cold deactivation that depends upon Mg²⁺ and ADP.

Acknowledgments

This work was supported by grant GM40508 from the National Institutes of Health, and grant N-1378 from the Welch Foundation. We thank Dr. Pia Vogel for helpful discussions, Dr. Larry Ruben for assistance in modifying the PTI fluorimeter for luminescence, and Dr. Ivan Charamisinau and the machine shop of the Department of Mechanical Engineering at SMU for assistance in construction of the luminometer cell chamber.

References

- [1] P.D. Boyer, The ATP synthase-A splendid molecular machine, *Annu. Rev. Biochem.* 66 (1997) 717–749.
- [2] P.D. Bragg, C. Hou, Subunit composition, function, and spatial arrangement in the Ca²⁺- and Mg²⁺-activated adenosine triphosphatases of *Escherichia coli* and *Salmonella typhimurium*, *Arch. Biochem. Biophys.* 167 (1975) 311–321.
- [3] D.L. Foster, R.H. Fillingame, Stoichiometry of subunits in the H⁺-ATPase complex of *Escherichia coli*, *J. Biol. Chem.* 257 (1982) 2009–2015.
- [4] W. Jiang, J. Hermolin, R.H. Fillingame, The preferred stoichiometry of c subunits in the rotary motor sector of *Escherichia coli* ATP synthase is 10, *Proc. Natl. Acad. Sci. U. S. A.* 98 (2001) 4966–4971.
- [5] T.M. Duncan, V.V. Bulygin, Y. Zhou, M.L. Hutcheon, R.L. Cross, Rotation of subunits during catalysis by *Escherichia coli* F₁-ATPase, *Proc. Natl. Acad. Sci. U. S. A.* 92 (1995) 10964–10968.
- [6] D. Sabbit, S. Engelbrecht, W. Junge, Intersubunit rotation in active F-ATPase, *Nature* 381 (1996) 623–625.
- [7] Y. Sambongi, Y. Iko, M. Tanabe, H. Omote, A. Iwamoto-Kihara, I. Ueda, T. Yanagida, Y. Wada, M. Futai, Mechanical rotation of the c subunit oligomer in ATP synthase (F₀F₁): direct observation, *Science* 286 (1999) 1722–1724.
- [8] H. Itoh, A. Takahashi, K. Adachi, H. Noji, R. Yasuda, M. Yoshida, K. Kinoshita, Mechanically driven ATP synthesis by F₁-ATPase, *Nature* 427 (2004) 465–468.
- [9] H. Noji, R. Yasuda, M. Yoshida, K. Kinoshita Jr., Direct observation of the rotation of F₁-ATPase, *Nature* 386 (1997) 299–302.
- [10] R. Yasuda, H. Noji, M. Yoshida, K. Kinoshita Jr., H. Itoh, Resolution of distinct rotational substeps by submillisecond kinetic analysis of F₁-ATPase, *Nature* 410 (2001) 898–904.
- [11] S. Hong, Y.H. Ko, P.L. Pedersen, Rotary catalysis within ATP synthases: a bioinformatic approach provides novel insight into how large pH-dependent movements of the C-terminal helix of subunit c may be accommodated, *Arch. Biochem. Biophys.* 394 (2001) 275–279.
- [12] J. Xing, H. Wang, C. von Ballmoos, P. Dimroth, G. Oster, Torque generation by the Fo motor of the sodium ATPase, *Biophys. J.* 87 (2004) 2148–2163.
- [13] T. Meier, P. Polzer, K. Diederichs, W. Welte, P. Dimroth, Structure of the rotor ring of F-Type Na⁺-ATPase from *Ilyobacter tartaricus*, *Science* 308 (2005) 659–662.
- [14] A. Bennun, E. Racker, Partial resolution of the enzymes catalyzing photophosphorylation. IV. Interaction of coupling factor 1 from chloroplasts with components of the chloroplast membrane, *J. Biol. Chem.* 244 (1969) 1325–1331.
- [15] H.S. Penefsky, R.C. Warner, Partial resolution of the enzymes catalyzing oxidative phosphorylation. VI. Studies on the mechanism of cold inactivation of mitochondrial adenosine triphosphatase, *J. Biol. Chem.* 240 (1965) 4694–4702.
- [16] G. Vogel, R. Steinhart, ATPase of *Escherichia coli*: purification, dissociation, and reconstitution of the active complex from the isolated subunits, *Biochemistry* 15 (1976) 208–216.
- [17] P.P. Laget, Hysteretic properties of soluble F₁ ATPase from *Escherichia coli*. Evidence for a slow transition between two conformational states of the enzyme, *Arch. Biochem. Biophys.* 189 (1978) 122–131.

- [18] C. Frieden, Kinetic aspects of regulation of metabolic processes. The hysteretic enzyme concept, *J. Biol. Chem.* 245 (1970) 5788–5799.
- [19] C. Frieden, Slow transitions and hysteretic behavior in enzymes, *Annu. Rev. Biochem.* 48 (1979) 471–489.
- [20] P.P. Laget, Hysteretic properties of soluble F_1 ATPase from *Escherichia coli*. II. Nucleotide effects on the slow changes of the enzyme kinetic behaviour, *Arch. Biochem. Biophys.* 192 (1979) 474–481.
- [21] J.M. Jault, W.S. Allison, Hysteretic inhibition of the bovine heart mitochondrial F_1 -ATPase is due to saturation of noncatalytic sites with ADP which blocks activation of the enzyme by ATP, *J. Biol. Chem.* 269 (1994) 319–325.
- [22] A.F. Fitin, E.A. Vasilyeva, A.D. Vinogradov, An inhibitory high affinity binding site for ADP in the oligomycin-sensitive ATPase of beef heart submitochondrial particles, *Biochem. Biophys. Res. Commun.* 86 (1979) 434–439.
- [23] V. Shoshan, H. Strotmann, The effect of phosphate on light-induced exchange of ADP at the tight nucleotide binding site of CF_1 , *J. Biol. Chem.* 255 (1980) 996–999.
- [24] D. Recktenwald, B. Hess, A slow conformation change in the transient state kinetics of soluble ATPase of yeast mitochondria, *FEBS Lett.* 80 (1977) 187–189.
- [25] B. Schobert, J.K. Lanyi, Hysteretic behavior of an ATPase from the archaeobacterium, *Halobacterium saccharovorum*, *J. Biol. Chem.* 264 (1989) 12805–12812.
- [26] A. Di Pietro, F. Penin, C. Godinot, D.C. Gautheron, “Hysteric” behavior and nucleotide binding sites of pig heart mitochondrial F_1 adenosine 5'-triphosphatase, *Biochemistry* 19 (1980) 5671–5678.
- [27] S.R. Paik, J.M. Jault, W.S. Allison, Inhibition and inactivation of the F_1 adenosinetriphosphatase from *Bacillus* PS3 by dequalinium and activation of the enzyme by lauryl dimethylamine oxide, *Biochemistry* 33 (1994) 126–133.
- [28] J.M. Jault, T. Matsui, F.M. Jault, C. Kaibara, E. Muneyuki, M. Yoshida, Y. Kagawa, W.S. Allison, The $\alpha_3\beta_3\gamma$ complex of the F_1 -ATPase from thermophilic *Bacillus* PS3 containing the α D261N substitution fails to dissociate inhibitory MgADP from a catalytic site when ATP binds to noncatalytic sites, *Biochemistry* 34 (1995) 16412–16418.
- [29] J.M. Jault, C. Dou, N.B. Grodsky, T. Matsui, M. Yoshida, W.S. Allison, The $\alpha_3\beta_3\gamma$ subcomplex of the F_1 -ATPase from the thermophilic *Bacillus* PS3 with the β T165S substitution does not entrap inhibitory MgADP in a catalytic site during turnover, *J. Biol. Chem.* 271 (1996) 28818–28824.
- [30] A.E. Senior, R.S. Lee, M.K. al-Shawi, J. Weber, Catalytic properties of *Escherichia coli* F_1 -ATPase depleted of endogenous nucleotides, *Arch. Biochem. Biophys.* 297 (1992) 340–344.
- [31] D.J. Hyndman, Y.M. Milgrom, E.A. Bramhall, R.L. Cross, Nucleotide-binding sites on *Escherichia coli* F_1 -ATPase. Specificity of noncatalytic sites and inhibition at catalytic sites by MgADP, *J. Biol. Chem.* 269 (1994) 28871–28877.
- [32] J.B. Smith, P.C. Sternweiss, Purification of membrane attachment and inhibitory subunits of the proton translocating adenosine triphosphatase from *Escherichia coli*, *Biochemistry* 16 (1977) 306–311.
- [33] P.C. Sternweiss, J.B. Smith, Characterization of the inhibitory (ϵ) subunit of the proton-translocating adenosine triphosphatase from *Escherichia coli*, *Biochemistry* 19 (1980) 526–531.
- [34] R.R. Ishmukhametov, M.A. Galkin, S.B. Vik, Ultrafast purification and reconstitution of His-tagged cysteine-less *Escherichia coli* F_1F_0 ATP synthase, *Biochim. Biophys. Acta* 1706 (2005) 110–116.
- [35] J.G. Wise, T.M. Duncan, L.R. Latchney, D.N. Cox, A.E. Senior, Properties of F_1 -ATPase from the *uncD412* mutant of *Escherichia coli*, *Biochem. J.* 215 (1983) 343–350.
- [36] Y.M. Milgrom, L.L. Ehler, P.D. Boyer, The characteristics and effect on catalysis of nucleotide binding to noncatalytic sites of chloroplast F_1 -ATPase, *J. Biol. Chem.* 266 (1991) 11551–11558.
- [37] M.B. Murataliev, Y.M. Milgrom, P.D. Boyer, Characteristics of the combination of inhibitory Mg^{2+} and azide with the F_1 ATPase from chloroplasts, *Biochemistry* 30 (1991) 8305–8310.
- [38] E.A. Vasilyeva, I.B. Minkov, A.F. Fitin, A.D. Vinogradov, Kinetic mechanism of mitochondrial adenosine triphosphatase. Inhibition by azide and activation by sulphite, *Biochem. J.* 202 (1982) 15–23.
- [39] J. Weber, A.E. Senior, Effects of the inhibitors azide, dicyclohexylcarbodiimide, and aurovertin on nucleotide binding to the three F_1 -ATPase catalytic sites measured using specific tryptophan probes, *J. Biol. Chem.* 273 (1998) 33210–33215.
- [40] T. Noumi, M. Maeda, M. Futai, Mode of inhibition of sodium azide on H^+ -ATPase of *Escherichia coli*, *FEBS Lett.* 213 (1987) 381–384.
- [41] D. Bald, E. Muneyuki, T. Amano, J. Kruij, T. Hisabori, M. Yoshida, The noncatalytic site-deficient $\alpha_3\beta_3\gamma$ subcomplex and F_0F_1 -ATP synthase can continuously catalyze ATP hydrolysis when P_i is present, *Eur. J. Biochem.* 262 (1999) 563–568.
- [42] H. Hanada, T. Noumi, M. Maeda, M. Futai, Uni-site catalysis by *Escherichia coli* F_1 -ATPase with different numbers of bound nucleotides, *FEBS Lett.* 257 (1989) 465–467.
- [43] P. Turina, D. Giovannini, F. Gubellini, B.A. Melandri, Physiological ligands ADP and P_i modulate the degree of intrinsic coupling in the ATP synthase of the photosynthetic bacterium *Rhodobacter capsulatus*, *Biochemistry* 43 (2004) 11126–11134.
- [44] S. Fischer, P. Gräber, P. Turina, The activity of the ATP synthase from *Escherichia coli* is regulated by the transmembrane proton motive force, *J. Biol. Chem.* 275 (2000) 30157–30162.
- [45] K. Dong, H. Ren, W.S. Allison, The fluorescence spectrum of the introduced tryptophans in the $\alpha_3(\beta F155W)_3\gamma$ subcomplex of the F_1 -ATPase from the thermophilic *Bacillus* PS3 cannot be used to distinguish between the number of nucleoside di- and triphosphates bound to catalytic sites, *J. Biol. Chem.* 277 (2002) 9540–9547.
- [46] R. Yasuda, T. Masaike, K. Adachi, H. Noji, H. Itoh, K. Kinoshita Jr., The ATP-waiting conformation of rotating F_1 -ATPase revealed by single-pair fluorescence resonance energy transfer, *Proc. Natl. Acad. Sci. U. S. A.* 100 (2003) 9314–9318.
- [47] W.S. Allison, J.M. Jault, C. Dou, N.B. Grodsky, Does the gamma subunit move to an abortive position of ATP hydrolysis when the F_1 .ADP.Mg complex isomerizes to the inactive F_1 .ADP.Mg complex? *J. Bioenerg. Biomembr.* 28 (1996) 433–438.
- [48] J. Mendel-Hartvig, R.A. Capaldi, Catalytic site nucleotide and inorganic phosphate dependence of the conformation of the epsilon subunit in *Escherichia coli* adenosinetriphosphatase, *Biochemistry* 30 (1991) 1278–1284.
- [49] J. Mendel-Hartvig, R.A. Capaldi, Nucleotide-dependent and dicyclohexylcarbodiimide-sensitive conformational changes in the epsilon subunit of *Escherichia coli* ATP synthase, *Biochemistry* 30 (1991) 10987–10991.
- [50] T. Suzuki, T. Murakami, R. Iino, J. Suzuki, S. Ono, Y. Shirakihara, M. Yoshida, F_0F_1 -ATPase/synthase is geared to the synthesis mode by conformational rearrangement of epsilon subunit in response to proton motive force and ADP/ATP balance, *J. Biol. Chem.* 278 (2003) 46840–46846.
- [51] H.R. Löttscher, C. deJong, R.A. Capaldi, Inhibition of the adenosinetriphosphatase activity of *Escherichia coli* F_1 by the water-soluble carbodiimide 1-ethyl-3-[3-dimethylamino]propyl carbodiimide is due to modification of several carboxyls in the β subunit, *Biochemistry* 23 (1984) 4134–4140.
- [52] H.R. Löttscher, C. deJong, R.A. Capaldi, Interconversion of high and low adenosinetriphosphatase activity forms of *Escherichia coli* F_1 by the detergent lauryldimethylamine oxide, *Biochemistry* 23 (1984) 4140–4143.
- [53] J.R. Silvius, R.N. McElhaney, Non-linear Arrhenius plots and the analysis of reaction and motional rates in biological membranes, *J. Theor. Biol.* 88 (1981) 135–152.
- [54] J.R. Silvius, B.D. Read, R.N. McElhaney, Membrane enzymes: artifacts in Arrhenius plots due to temperature dependence of substrate-binding affinity, *Science* 199 (1978) 902–904.
- [55] M.K. Al-Shawi, C.J. Ketchum, R.K. Nakamoto, Energy coupling, turnover, and stability of the F_0F_1 ATP synthase are dependent on the energy of interaction between gamma and beta subunits, *J. Biol. Chem.* 272 (1997) 2300–2306.
- [56] A. Baracca, E. Amler, G. Solaini, G. Parenti Castelli, G. Lenaz, J. Housteck, Temperature-induced states of isolated F_1 -ATPase affect catalysis, enzyme conformation and high-affinity nucleotide binding sites, *Biochim. Biophys. Acta* 976 (1989) 77–84.