

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

Physics Letters B 639 (2006) 456–461

PHYSICS LETTERS B

www.elsevier.com/locate/physletb

Four-nucleon force in chiral effective field theory

E. Epelbaum

Jefferson Laboratory, Theory Division, Newport News, VA 23606, USA

Received 18 April 2006; accepted 6 June 2006

Available online 5 July 2006

Editor: W. Haxton

Abstract

We derive the leading contribution to the four-nucleon force within the framework of chiral effective field theory. It is governed by the exchange of pions and the lowest-order nucleon–nucleon contact interaction and includes effects due to the nonlinear pion–nucleon couplings and the pion self interactions constrained by the chiral symmetry of QCD. The resulting four-nucleon force does not contain any unknown parameters and can be tested in few- and many-nucleon studies.

© 2006 Elsevier B.V. Open access under [CC BY license](http://creativecommons.org/licenses/by/4.0/).

PACS: 21.45.+v; 21.30.-x; 25.10.+s

1. Chiral effective field theory (EFT) is a powerful tool for analyzing the properties of hadronic systems at low energies in a systematic and model-independent way. It exploits the approximate and spontaneously broken chiral symmetry of QCD which governs low-energy hadron structure and dynamics. Over the past years, considerable progress has been achieved in understanding the structure of the nuclear force in this framework [1–5], see also recent review articles [6–8] and references therein. In particular, two- (2NFs) and three-nucleon forces (3NFs) have been worked out and applied in few-nucleon studies upto next-to-next-to-next-to-leading (N³LO) [9,10] and next-to-next-to-leading orders (N²LO) [11] in the chiral expansion, respectively. Parallel to these developments, significant progress has been achieved in the microscopic description of few- and many-nucleon systems. Advances in the development of methods, such as the Faddeev–Yakubovsky scheme, Green’s function Monte Carlo and hyperspherical harmonics methods as well as the no-core shell model, coupled with a significant increase in computational resources allow one to obtain the binding energies of light nuclei with high numerical accuracy reaching 0.2% for the α -particle [12] and 1–2% for heavier nuclei [13,14]. Three- and four-nucleon continuum observables are also being investigated. These studies open the door for precise and non-trivial tests of the underlying dynamics. In this context, three-nucleon forces have deserved much attention and were shown to be important e.g. for understanding the spectra of light nuclei [14]. Four-nucleon forces (4NFs) have not yet been explored in few-body calculations. Although the existing studies leave little room for the contribution of the possible 4NFs, see e.g. [12], this needs to be verified via explicit calculations. In this Letter we make a step in this direction and derive the leading 4NF in the framework of chiral EFT which appears at N³LO in the chiral expansion. The resulting 4NF is local and parameter-free. Its application in future few-body studies will provide a challenging test of our understanding of the nature of nuclear forces.

2. Let us first recall the structure of the effective chiral Lagrangian for pions and nucleons, which at lowest order reads:

$$\mathcal{L}_{\pi\pi}^{(2)} = \frac{F_\pi^2}{4} \text{tr}[\partial_\mu U \partial^\mu U^\dagger + M_\pi^2 (U + U^\dagger)],$$

$$\mathcal{L}_{\pi N}^{(1)} = N^\dagger \left(i D_0 - \frac{g_A}{2} \vec{\sigma} \cdot \vec{u} \right) N,$$

E-mail address: epelbaum@jlab.org (E. Epelbaum).

$$\mathcal{L}_{NN}^{(0)} = -\frac{1}{2}C_S(N^\dagger N)(N^\dagger N) - \frac{1}{2}C_T(N^\dagger \vec{\sigma} N) \cdot (N^\dagger \vec{\sigma} N). \quad (1)$$

Here the superscripts refer to the number of derivatives and/or M_π -insertions, $F_\pi = 92.4$ MeV is the pion decay constant and $g_A = 1.267$ is the nucleon axial-vector coupling. The low-energy constants (LECs) C_S and C_T determine the strength of the leading NN short-range interaction [1,2]. The Lagrangians $\mathcal{L}_{\pi N}^{(1)}$ and $\mathcal{L}_{NN}^{(0)}$ are given in the heavy-baryon formulation with N representing a non-relativistic nucleon field and $\vec{\sigma}$ denoting the Pauli spin matrices. Further, the SU(2) matrix $U = u^2$ collects the pion fields, $D^\mu = \partial^\mu + \frac{1}{2}[u^\dagger, \partial^\mu u]$ denotes the covariant derivative of the nucleon field and $u_\mu = iu^\dagger \partial_\mu U u^\dagger$. The first terms in the expansion of the matrix $U(\pi)$ in powers of the pion fields take the form, see e.g. [15]:

$$U(\pi) = 1 + \frac{i}{F_\pi} \boldsymbol{\tau} \cdot \boldsymbol{\pi} - \frac{1}{2F_\pi^2} \boldsymbol{\pi}^2 - \frac{i\alpha}{F_\pi^3} (\boldsymbol{\tau} \cdot \boldsymbol{\pi})^3 + \frac{8\alpha - 1}{8F_\pi^4} \boldsymbol{\pi}^4 + \dots, \quad (2)$$

where $\boldsymbol{\tau}$ denote the Pauli isospin matrices and α is an arbitrary constant. Notice that only the coefficients in front of the linear and quadratic terms in the pion field are fixed uniquely from the unitarity condition $U^\dagger U = 1$ and the proper normalization of the pion kinetic energy. The explicit α -dependence of the matrix U represents the freedom in the definition of the pion field. Clearly, all measurable quantities are α -independent.

3. The structure of the nuclear forces can be visualized utilizing a diagrammatic language. Given the non-uniqueness of the nuclear potential, the precise meaning of diagrams is scheme-dependent. Typical examples include Feynman graphs with subtracted iterative contributions, see e.g. [15], irreducible time-ordered diagrams [1,2] or time-ordered-like graphs in the method of unitary transformation [8]. Within the power counting scheme of Refs. [1,2] adopted in this work,¹ the importance of a particular contribution to the nuclear force is determined by the power ν of the expansion parameter Q/Λ . Here, Q and Λ refer to the generic low-momentum scale associated with external nucleon momenta or M_π and the pertinent hard scale, respectively. For a N -nucleon diagram with L loops, C separately connected pieces and V_i vertices of type i , the low-momentum dimension is given by [1,2]

$$\nu = -2 + 2N - 2C + 2L + \sum_i V_i \Delta_i \quad \text{with } \Delta_i = d_i + \frac{1}{2}n_i - 2. \quad (3)$$

Here, n_i is the number of nucleon field operators and d_i the number of derivatives and/or insertions of M_π . The interaction terms of lowest possible chiral dimension $\Delta_i = 0$ are listed in Eq. (1). The power counting in Eq. (3) naturally explains the dominance of the two-nucleon force (2NF) which starts to contribute at order $\nu = 0$ [1,2]. The leading 3NF formally arises at order $\nu = 2$. The corresponding contributions are known to vanish provided one uses an energy-independent formulation such as the method of unitary transformation [18–20], see also [4,21] for a similar conclusion based on the framework of time-ordered perturbation theory. The first non-vanishing 3NFs appear at order $\nu = 3$ [4,11]. At present, the 2NFs and 3NFs have been worked out and applied in few-body studies upto orders $\nu = 4$ [9,10] and $\nu = 3$ [11], respectively, see [8] for more details. Derivation of the leading 3NF corrections at order $\nu = 4$ is in progress.

According to Eq. (3), the leading 4NF is expected to arise at order $\nu = 2$ from disconnected tree diagrams with the lowest-order vertices, see Fig. 1. Using the method of unitary transformation, it is easy to see that both graphs in Fig. 1 yield vanishing contributions, see also [4] for an earlier discussion based on time-ordered perturbation theory. The first non-vanishing 4NFs appear at order $\nu = 4$ where one has to take into account disconnected tree diagrams with one insertion of a $\Delta_i = 2$ vertex as well as disconnected loop and connected tree graphs with the lowest-order vertices depicted in Figs. 2 and 3, respectively. In addition, one needs to consider relativistic $1/m$ -corrections to disconnected diagrams shown in Fig. 1. We have verified that *all* disconnected graphs at this order yield vanishing contributions to the 4NF. One is, therefore, left with connected tree diagrams shown in Fig. 3. We have used the method of unitary transformation [20] to evaluate the corresponding 4NF contributions. To that aim we first switch from the Lagrangian in Eq. (1) to the effective πN Hamilton operator using the canonical formalism and then apply a suitable unitary transformation in Fock space in order to decouple purely nucleonic states from the ones which contain pions. The resulting




Fig. 1. Disconnected diagrams at order $\nu = 2$ that lead to vanishing contributions to the four-nucleon force. Solid and dashed lines represent nucleons and pions, respectively.

¹ Notice that different counting schemes have been suggested in Refs. [16,17] for contact interactions. In particular, according to [17], certain contact two-nucleon terms have to be promoted to the lowest order in the low-momentum expansion. These findings do not affect our results for the long-range part of the 4NF but would lead to additional contributions involving contact interactions.




Fig. 2. Examples of disconnected diagrams at order $\nu = 4$ that lead to vanishing contributions to the four-nucleon force.




Fig. 3. The leading contributions to the four-nucleon force. Graphs resulting by the interchange of the vertex ordering and/or nucleon lines are not shown.

Hamilton operator acts on the nucleon subspace of the Fock space and provides a definition of the nuclear forces. A rather tedious calculation along the lines of Ref. [20] yields the following result for the contributions of individual diagrams in Fig. 3:

$$\begin{aligned}
 V^a &= -\frac{2g_A^6}{(2F_\pi)^6} \frac{\vec{\sigma}_1 \cdot \vec{q}_1 \vec{\sigma}_4 \cdot \vec{q}_4}{[\vec{q}_1^2 + M_\pi^2][\vec{q}_{12}^2 + M_\pi^2]^2[\vec{q}_4^2 + M_\pi^2]} \\
 &\quad \times [(\boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_4 \boldsymbol{\tau}_2 \cdot \boldsymbol{\tau}_3 - \boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_3 \boldsymbol{\tau}_2 \cdot \boldsymbol{\tau}_4) \vec{q}_1 \cdot \vec{q}_{12} \vec{q}_4 \cdot \vec{q}_{12} + \boldsymbol{\tau}_1 \times \boldsymbol{\tau}_2 \cdot \boldsymbol{\tau}_4 \vec{q}_1 \cdot \vec{q}_{12} \vec{q}_{12} \times \vec{q}_4 \cdot \vec{\sigma}_3 \\
 &\quad + \boldsymbol{\tau}_1 \times \boldsymbol{\tau}_3 \cdot \boldsymbol{\tau}_4 \vec{q}_4 \cdot \vec{q}_{12} \vec{q}_1 \times \vec{q}_{12} \cdot \vec{\sigma}_2 + \boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_4 \vec{q}_{12} \times \vec{q}_1 \cdot \vec{\sigma}_2 \vec{q}_{12} \times \vec{q}_4 \cdot \vec{\sigma}_3] + \text{all perm.}, \\
 V^c &= -\frac{2g_A^4}{(2F_\pi)^6} \frac{\vec{\sigma}_1 \cdot \vec{q}_1 \vec{\sigma}_4 \cdot \vec{q}_4}{[\vec{q}_1^2 + M_\pi^2][\vec{q}_{12}^2 + M_\pi^2][\vec{q}_4^2 + M_\pi^2]} \\
 &\quad \times [(\boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_4 \boldsymbol{\tau}_2 \cdot \boldsymbol{\tau}_3 - \boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_3 \boldsymbol{\tau}_2 \cdot \boldsymbol{\tau}_4) \vec{q}_{12} \cdot \vec{q}_4 + \boldsymbol{\tau}_1 \times \boldsymbol{\tau}_2 \cdot \boldsymbol{\tau}_4 \vec{q}_{12} \times \vec{q}_4 \cdot \vec{\sigma}_3] + \text{all perm.}, \\
 V^e &= \frac{g_A^4}{(2F_\pi)^6} \frac{\vec{\sigma}_2 \cdot \vec{q}_2 \vec{\sigma}_3 \cdot \vec{q}_3 \vec{\sigma}_4 \cdot \vec{q}_4}{[\vec{q}_2^2 + M_\pi^2][\vec{q}_3^2 + M_\pi^2][\vec{q}_4^2 + M_\pi^2]} \boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_2 \boldsymbol{\tau}_3 \cdot \boldsymbol{\tau}_4 \vec{\sigma}_1 \cdot (\vec{q}_3 + \vec{q}_4) + \text{all perm.}, \\
 V^f &= \frac{g_A^4}{2(2F_\pi)^6} [(\vec{q}_1 + \vec{q}_2)^2 + M_\pi^2] \frac{\vec{\sigma}_1 \cdot \vec{q}_1 \vec{\sigma}_2 \cdot \vec{q}_2 \vec{\sigma}_3 \cdot \vec{q}_3 \vec{\sigma}_4 \cdot \vec{q}_4}{[\vec{q}_1^2 + M_\pi^2][\vec{q}_2^2 + M_\pi^2][\vec{q}_3^2 + M_\pi^2][\vec{q}_4^2 + M_\pi^2]} \boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_2 \boldsymbol{\tau}_3 \cdot \boldsymbol{\tau}_4 + \text{all perm.}, \\
 V^k &= 4C_T \frac{g_A^4}{(2F_\pi)^4} \frac{\vec{\sigma}_1 \cdot \vec{q}_1 \vec{\sigma}_3 \times \vec{\sigma}_4 \cdot \vec{q}_{12}}{[\vec{q}_1^2 + M_\pi^2][\vec{q}_{12}^2 + M_\pi^2]^2} [\boldsymbol{\tau}_1 \cdot \boldsymbol{\tau}_3 \vec{q}_1 \times \vec{q}_{12} \cdot \vec{\sigma}_2 - \boldsymbol{\tau}_1 \times \boldsymbol{\tau}_2 \cdot \boldsymbol{\tau}_3 \vec{q}_1 \cdot \vec{q}_{12}] + \text{all perm.},
 \end{aligned}$$

$$\begin{aligned}
V^l &= -2C_T \frac{g_A^2}{(2F_\pi)^4} \frac{\vec{\sigma}_1 \cdot \vec{q}_1 \vec{\sigma}_3 \times \vec{\sigma}_4 \cdot \vec{q}_{12}}{[\vec{q}_1^2 + M_\pi^2][\vec{q}_{12}^2 + M_\pi^2]} \boldsymbol{\tau}_1 \times \boldsymbol{\tau}_2 \cdot \boldsymbol{\tau}_3 + \text{all perm.}, \\
V^n &= 2C_T^2 \frac{g_A^2}{(2F_\pi)^2} \frac{\vec{\sigma}_1 \times \vec{\sigma}_2 \cdot \vec{q}_{12} \vec{\sigma}_3 \times \vec{\sigma}_4 \cdot \vec{q}_{12}}{[\vec{q}_{12}^2 + M_\pi^2]^2} \boldsymbol{\tau}_2 \cdot \boldsymbol{\tau}_3 + \text{all perm.}
\end{aligned} \tag{4}$$

Here, the subscripts refer to the nucleon labels and $\vec{q}_i = \vec{p}'_i - \vec{p}_i$ with \vec{p}'_i and \vec{p}_i being the final and initial momenta of the nucleon i . Further, $\vec{q}_{12} = \vec{q}_1 + \vec{q}_2 = -\vec{q}_3 - \vec{q}_4 = -\vec{q}_{34}$ is the momentum transfer between the nucleon pairs 12 and 34. Notice that some terms in the sum over all possible 24 permutations of the nucleon labels are the same. Diagrams (b), (d), (g), (h), (i), (j), (m), (o) and (p) lead to vanishing contributions to the 4NF. Notice further that the short-range 4NFs only depend on the LEC C_T which can be traced back to the fact that the C_S -vertex commutes with the other lowest-order vertices. Details of the derivation will be published elsewhere.

The results given in Eq. (4) deserve some special comments. The easiest way to evaluate the contributions from graphs (e) and (f), which do not include reducible topologies (i.e. time-ordered diagrams with purely nucleonic intermediate states), is by calculating the corresponding Feynman diagrams. The resulting 4NFs depend individually on the parameter α which enters the effective Lagrangian in Eq. (1). When they are added together, this α -dependence cancels out, see also [22,23] for an earlier discussion of these contributions. We only show explicitly the remaining, α -independent terms in Eq. (4). The 3π -exchange 4NF proportional to g_A^2 can be obtained by evaluating the Feynman diagram (g) in Fig. 3. Due to the four-momentum conservation at each vertex and the fact that the Weinberg–Tomozawa vertex contains a time derivative of the pion field, this contribution is suppressed by Q^2/m^2 and, therefore, shifted to higher orders. In the method of unitary transformation, one needs to consider not only graph (g) but also graph (h), where the $\pi\pi NN$ vertex of dimension $\Delta_i = 0$,

$$\mathcal{H}_{\text{int}} = \frac{1}{2(2F_\pi)^4} (N^\dagger \boldsymbol{\tau} \times \boldsymbol{\pi} N) \cdot (N^\dagger \boldsymbol{\tau} \times \boldsymbol{\pi} N), \tag{5}$$

arises in the Hamiltonian through the application of the canonical formalism to the Lagrangian in Eq. (1), see e.g. [24]. We have verified that the 4NF contributions from diagrams (g) and (h) in this framework cancel against each other, which is consistent with the result based on the Feynman graph technique.

Before discussing the numerical size of the derived 4NF, another important comment is in order. Nuclear forces are not unique and can, in general, be changed via unitary transformations. We have explored a large class of unitary transformations which act on the nucleon subspace of the Fock space and affect nuclear forces at order $\nu = 4$ (and higher). We found that the associated ambiguity in the definition of the nuclear forces is resolved if one requires renormalizability of the 1-loop 3NF contributions. Consider, for example, 3π -exchange 3NF corresponding to diagrams (a)–(c) in Fig. 4. Ultraviolet-divergent loop integrals resulting from 2π -exchange between the nucleons 1 and 2 have to be renormalized by a redefinition of the LEC D which enters the expression for the 3NF at order $\nu = 3$ [4,11], see diagram (d) in Fig. 4:

$$V_{3N} = -D \frac{g_A}{8F_\pi^2} \frac{\vec{\sigma}_2 \cdot \vec{q}_3 \vec{\sigma}_3 \cdot \vec{q}_3}{[\vec{q}_3^2 + M_\pi^2]} \boldsymbol{\tau}_2 \cdot \boldsymbol{\tau}_3 + \text{all perm.} \tag{6}$$

For this to be possible, ultraviolet-divergent pieces in the 3π -exchange 3NF must have the same form as V_{3N} . As a consequence, no terms proportional to $(\vec{q}_3^2 + M_\pi^2)^{-2}$, $(\vec{q}_3^2 + M_\pi^2)^{-3/2}$ or $(\sqrt{\vec{q}_3^2 + M_\pi^2} + \sqrt{\vec{l}_1^2 + M_\pi^2})^{-1}$ with \vec{l}_1 being the momentum of one of the two pions exchanged between the nucleons 1 and 2 should appear in the 3NF. This restriction imposes non-trivial constraints on the choice of the unitary transformations and, therefore, also on the form of the 4NFs corresponding to graphs (a) and (b) in Fig. 3. These considerations may remind one of the recent findings in the context of large N_c QCD [25]. One should, of course, keep in mind that S-matrix elements, which involve iterations of the potential, are not affected by unitary transformations. Based on the above arguments, we have not found any ambiguity in the form of the 4NFs (for the considered class of unitary transformations). We postpone a detailed discussion of this issue to a forthcoming publication.

In order to test the effects of the 4NFs in few-nucleon systems, explicit calculations will need to be performed. To get a rough idea about the size of the 4NF contributions to e.g. the α -particle binding energy, one can look at the strength of the corresponding




Fig. 4. Selected contributions to the three-nucleon force.

r -space potentials. For V^e , for example, it is given by

$$\frac{1}{(4\pi)^3} \frac{2g_A^4}{(2F_\pi)^6} M_\pi^7 \sim 80 \text{ keV}, \quad (7)$$

see also [23]. Here, the factor $(4\pi)^{-3}$ arises from integrations over the relative momenta. Other 3π -exchange 4NFs have similar strengths. The short-range terms can be expected to be less important due to numerically small values of the LEC C_T found in [26] and (for certain cut-off combinations) in [10]. We remind the reader that in EFT without or with perturbative pions, one has $C_T = 0$ in the limit when both NN S-wave scattering lengths go to infinity [27]. To justify the validity of the above numerical estimation and to get an idea about its accuracy, we perform a similar estimation for the strength of the 3NF at N²LO and compare it with the findings of Ref. [11]. For the 2π -exchange 3NF proportional to LEC c_3 , for example, one finds

$$\frac{1}{(4\pi)^2} \frac{g_A^2}{(2F_\pi)^2} \frac{|c_3|}{F_\pi^2} M_\pi^6 \sim 300 \text{ keV}, \quad (8)$$

where we use the same value for c_3 as in [11]. The 2π -exchange 3NF terms proportional to the LECs c_1 and c_4 have similar strengths. The above estimation agrees reasonably well with the expectation values of the 2π -exchange 3NF in ^3H which are found to be -390 and -730 keV for the two considered cutoff choices [11]. A detailed discussion on the size of various 2NF and 3NF contributions can be found in [28].

Finally, we would like to emphasize that one should expect sizable corrections to the 4NF due to intermediate Δ -excitations [23], see also [29] for a recent discussion on the role of the Δ -isobar in the 3NF. In the EFT without explicit Δ 's, these effects are taken into account implicitly, i.e. through the values of the corresponding LECs, and first show up at higher orders. We postpone a systematic investigation of the corrections due to intermediate Δ -excitations to a later publication.

4. To summarize, we have derived the leading 4NF contributions based on chiral EFT that appear at N³LO in the low-momentum expansion. Chiral symmetry of QCD fixes the structure and strength of the pion couplings and plays a crucial role in the derivation. The strength of the lowest-order NN contact interaction entering the expressions for the 4NF is determined in the two-nucleon system [9,10], so that the resulting 4NF is parameter-free. Applications of this 4NF to study the properties of the α -particle and the spectra of other light nuclei will provide a challenging test of our understanding of the nuclear force.

Acknowledgements

The author would like to thank Sid Coon, Walter Glöckle, Ulf-G. Meißner, Vladimir Pascalutsa and Rocco Schiavilla for sharing their insights and stimulating discussions. This work has been supported by the US Department of Energy Contract No. DE-AC05-84ER40150 under which the Southeastern Universities Research Association (SURA) operates the Thomas Jefferson National Accelerator Facility.

References

- [1] S. Weinberg, Phys. Lett. B 251 (1990) 288.
- [2] S. Weinberg, Nucl. Phys. B 363 (1991) 3.
- [3] C. Ordóñez, L. Ray, U. van Kolck, Phys. Rev. Lett. 72 (1994) 1982.
- [4] U. van Kolck, Phys. Rev. C 49 (1994) 2932.
- [5] C. Ordóñez, L. Ray, U. van Kolck, Phys. Rev. C 53 (1996) 2086, hep-ph/9511380.
- [6] S.R. Beane, P.F. Bedaque, W.C. Haxton, D.R. Phillips, M.J. Savage, nucl-th/0008064.
- [7] P.F. Bedaque, U. van Kolck, Annu. Rev. Nucl. Part. Sci. 52 (2002) 339, nucl-th/0203055.
- [8] E. Epelbaum, Prog. Part. Nucl. Phys. 57 (2006) 654, nucl-th/0509032.
- [9] D.R. Entem, R. Machleidt, Phys. Rev. C 68 (2003) 041001, nucl-th/0304018.
- [10] E. Epelbaum, W. Glöckle, U.-G. Meißner, Nucl. Phys. A 747 (2005) 362, nucl-th/0405048.
- [11] E. Epelbaum, et al., Phys. Rev. C 66 (2002) 064001, nucl-th/0208023.
- [12] A. Nogga, H. Kamada, W. Glöckle, B.R. Barrett, Phys. Rev. C 65 (2002) 054003, nucl-th/0112026.
- [13] P. Navrátil, W.E. Ormand, Phys. Rev. C 68 (2003) 034305, nucl-th/0305090.
- [14] S.C. Pieper, K. Varga, R.B. Wiringa, Phys. Rev. C 66 (2002) 044310, nucl-th/0206061.
- [15] N. Kaiser, Phys. Rev. C 61 (2000) 014003, nucl-th/9910044.
- [16] S.R. Beane, P.F. Bedaque, M.J. Savage, U. van Kolck, Nucl. Phys. A 700 (2002) 377, nucl-th/0104030.
- [17] A. Nogga, R.G.E. Timmermans, U. van Kolck, Phys. Rev. C 72 (2005) 054006, nucl-th/0506005.
- [18] S.A. Coon, J.L. Friar, Phys. Rev. C 34 (1986) 1060.
- [19] J.A. Eden, M.F. Gari, Phys. Rev. C 53 (1996) 1510, nucl-th/9601025.
- [20] E. Epelbaum, W. Glöckle, U.-G. Meißner, Nucl. Phys. A 637 (1998) 107, nucl-th/9801064.
- [21] S.N. Yang, W. Glöckle, Phys. Rev. C 33 (1986) 1774.
- [22] H. McManus, D.O. Riska, Phys. Lett. B 92 (1980) 29.
- [23] M.R. Robilotta, Phys. Rev. C 31 (1985) 974.

- [24] S. Weinberg, *Phys. Lett. B* 295 (1992) 114, hep-ph/9209257.
- [25] T.D. Cohen, *Phys. Rev. C* 66 (2002) 064003, nucl-th/0209072.
- [26] E. Epelbaum, U.-G. Meißner, W. Glöckle, C. Elster, *Phys. Rev. C* 65 (2002) 044001, nucl-th/0106007.
- [27] T. Mehen, I.W. Stewart, M.B. Wise, *Phys. Rev. Lett.* 83 (1999) 931, hep-ph/9902370.
- [28] J.L. Friar, *Few Body Syst.* 22 (1997) 161, nucl-th/9607020.
- [29] V.R. Pandharipande, D.R. Phillips, U. van Kolck, *Phys. Rev. C* 71 (2005) 064002, nucl-th/0501061.