

Fhit proteins can also recognize substrates other than dinucleoside polyphosphates

Andrzej Guranowski^{a,*}, Anna M. Wojdyła^a, Małgorzata Pietrowska-Borek^b, Paweł Bieganowski^c, Elena N. Khurs^d, Matthew J. Cliff^e, G. Michael Blackburn^f, Damian Błaziak^g, Wojciech J. Stec^g

^a Department of Biochemistry and Biotechnology, The University of Life Sciences, 35 Wołyńska Street, 60-637 Poznań, Poland

^b Department of Plant Physiology, The University of Life Sciences, 60-637 Poznań, Poland

^c International Institute of Molecular and Cell Biology, Warsaw, Poland

^d Engelhardt Institute of Molecular Biology, Russian Academy of Sciences, Moscow, Russia

^e Department of Molecular Biology and Biotechnology, Krebs Institute, University of Sheffield, Sheffield, UK

^f Department of Chemistry, Krebs Institute, University of Sheffield, Sheffield, UK

^g Center of Molecular and Macromolecular Studies, Polish Academy of Sciences, Łódź, Poland

Received 1 July 2008; revised 17 July 2008; accepted 31 July 2008

Available online 9 August 2008

Edited by Hans Eklund

This work is dedicated to Professor Antonio Sillero on the occasion of his 70th birthday and to Professor Maria Antonia Günther Sillero.


Abstract We show here that Fhit proteins, in addition to their function as dinucleoside triphosphate hydrolases, act similarly to adenylylsulfatases and nucleoside phosphoramidases, liberating nucleoside 5'-monophosphates from such natural metabolites as adenosine 5'-phosphosulfate and adenosine 5'-phosphoramidate. Moreover, Fhits recognize synthetic nucleotides, such as adenosine 5'-O-phosphorfluoridate and adenosine 5'-O-(γ -fluorotriphosphate), and release AMP from them. With respect to the former, Fhits behave like a phosphodiesterase I concomitant with cleavage of the P–F bond. Some kinetic parameters and implications of the novel reactions catalyzed by the human and plant (*Arabidopsis thaliana*) Fhit proteins are presented. © 2008 Published by Elsevier B.V. on behalf of the Federation of European Biochemical Societies.

Keywords: Fhit protein; Dinucleoside triphosphatase activity; Nucleoside phosphoramidase activity; Adenylylsulfatase activity; Phosphodiesterase I activity; P–F bond cleavage

1. Introduction

Cells contain various minor nucleotides. Among these are the dinucleoside 5',5'''-P¹,Pⁿ-polyphosphates (Np_nN's, where N and N' are 5'-O-nucleosides and *n* represents the number of phosphate residues in the polyphosphate chain that esterifies N and N' at their 5' position) [1]. Np_nN's accumulate as a result of the activity of certain ligases and transferases that catalyze transfer of a nucleotidyl moiety onto various acceptors containing a pyrophosphate residue, including NTPs (ppNs) and NDPs (ppNs), from a variety of donors. For reviews see [2,3]. Np_nN's play different intracellular and extracellular functions [4–6]. The cellular level of Np_nN's can be controlled by various hydrolases or phosphorylases [7]. Among specific hydrolases is the dinucleoside triphosphatase (EC 3.6.1.29) that preferentially hydrolyzes NpppN' to an NMP and N'DP (see Reaction 1):

This enzyme was first discovered in Silleros' laboratory in extracts of rat liver [8] and subsequently in extracts of yellow



*Corresponding author. Fax: +48 61 8487146.

E-mail address: guranow@au.poznan.pl (A. Guranowski).

Abbreviations: APS or S-pA, adenosine 5'-phosphosulfate; NH₂-pA, adenosine 5'-phosphoramidate; ATP-F or F-pppA, adenosine 5'-O-(γ -fluorotriphosphate); HPLC, high performance liquid chromatography; TLC, thin layer chromatography

lupin seeds [9], *Saccharomyces cerevisiae* [10], *Artemia* [11], and green algae [12]. The intriguing finding that *FHIT* (from fragile histidine triad), a putative human tumor suppressor gene, encodes a typical dinucleoside triphosphatase [13] focused much attention on this particular enzyme. *FHIT* genes occur in different eukaryotic organisms [14] and homogeneous Fhit proteins have been obtained by overexpression of *FHIT* from human [15], yeast [16] and *Arabidopsis thaliana* (present study).

Work described here stemmed from the investigations of the regiospecificity of the hydrolysis of N_pN_n 's and in particular from a study on the mechanism of action of Fhit/ N_pN_n ' hydrolases. Preference for N_pN_3 's and a unique mode of substrate cleavage are features that distinguish N_pN_3 ' hydrolases from other specific hydrolases acting on N_pN_n 's. Using $H_2^{18}O$ and mass spectrometry of the isolated reaction products, it was first shown that the N_pN_3 ' hydrolase from yellow lupin exclusively hydrolyzes the anhydride bond of its substrates (ApppA, AppppA, AppCH₂pA or AppCHFPa) between P^α and P^β, incorporating ^{18}O only into AMP [17]. It was subsequently shown by Abend et al. that human Fhit/ N_pN_3 ' hydrolase acts in the same fashion [18]. Moreover, they demonstrated that cleavage proceeds with overall retention of configuration at phosphorus, implying a double inversion mechanism, and they postulated that all three histidines of the histidine triad (His94, His96 and His98) are involved in the formation of an intermediate in which His96 first undergoes adenylation and then deadenylation by water. Pursuing that issue, Frey et al. showed that Fhit and specifically mutated Fhit variants catalyzed the hydrolysis of adenosine 5'-phosphoimidazolide (APS or S-pA) [19] and, poorly, of *p*-nitrophenyl-AMP [20]. Thus, Fhits behave like a nucleoside phosphoramidase and phosphodiesterase, respectively.

We demonstrated that two naturally occurring AMP derivatives: sulfoadenylate (S-pA), a key metabolite on the sulfate assimilation pathway, and adenosine 5'-phosphoramidate (NH₂-pA, aminoadenylate), a less known natural metabolite [21] originating from APS by enzymatic displacement of its sulfate moiety by ammonia [22], are Fhits' substrates. We also showed that both human and plant (*Arabidopsis*) Fhits liberate AMP from such synthetic AMP congeners as fluoro-adenylate (F-pA) and fluoro-ATP (ATP-F).

2. Materials and methods

2.1. Materials

Adenosine 5'-*O*-fluorophosphate (F-pA) was synthesized according to [23] and adenosine 5'-*O*-(γ -fluorophosphate) (Fpp-pA) according to procedure developed for the synthesis of GTP γ F [24]. Other adenine (di)nucleotides were from Sigma, St. Louis, MO, USA. NH₂-pA was custom-labeled with tritium at its C-8 by Moravsek Biochemicals, Brea, CA, USA. Amino-[8-³H]inosylate (NH₂-p[8-³H]I) was obtained by deamination of NH₂-p[8-³H]A catalyzed by adenosine phosphate deaminase (EC 3.5.4.17) from *Helix pomatia* [25].

Primers 5'-CGACGCATATGTCGTCTACTTGTCTTCG and 5'-CGGCTCGAGCTAGCAATCGAAAAGAGATCTG were used to amplify the coding sequence of *A. thaliana* FHIT. The PCR product obtained with *A. thaliana* cDNA was cloned using NdeI and XhoI restriction sites included in the primer sequences into pSG02 vector [26]. Plasmid for the human Fhit expression was described earlier [27]. *A. thaliana* and human Fhit proteins were expressed in *Escherichia coli* strain BL21. Cells were lysed by sonication in buffer A, containing 100 mM NaCl, 20 mM Tris-HCl, pH 7.5 and 2 mM dithiothreitol. Nucleic acids were precipitated using polyethyleneimine at 0.1% concentration. Insoluble debris was removed by centrifugation and

remaining proteins were precipitated with ammonium sulfate added to 70% saturation. Precipitated proteins were resuspended in buffer A and ammonium sulfate precipitation was repeated. The resulting protein pellet was resuspended in buffer A and loaded onto an AMP-agarose column. Unbound proteins were washed out with buffer A and Fhit proteins were eluted with buffer A supplemented with 1 mM adenosine. This procedure yielded proteins that were about 90% pure by SDS-PAGE gel. Molecular mass of the human Fhit monomer is 16800 Da and *Arabidopsis* Fhit monomer 18120 Da. Both Fhit samples were then dialyzed against 20 mM Hepes/NaOH, pH 7.5 containing 100 mM NaCl and 5% glycerol, concentrated by ultrafiltration on Microcone filters from Millipore to 1.3 mg/ml in case of the human Fhit and to 1.5 mg/ml in case of the *Arabidopsis* Fhit, and stored at -20 °C.

2.2. Enzyme assays

Hydrolytic activities of the Fhit proteins were assayed in reaction mixture (0.1 ml) containing 50 mM Mes/KOH (pH 6.5), 5 mM MgCl₂, 1 mM substrate and rate-limiting amounts of either human or plant Fhit protein. The reactions were carried out at 30 °C. At time intervals (0, 5, 10 and 30 min), 20 μ l aliquots were withdrawn and the reaction stopped by heating the samples for 5 min at 96 °C. The samples were chilled, diluted three-fold with 50 mM TEAB (triethylamine buffer, pH 7.4), filtered through ultrafree-MC filters (from Millipore) and 10 μ l aliquots subjected to high performance liquid chromatography (HPLC) on a Discovery C18 column (4.6 \times 250 mm, 5 μ m); Supelco at a flow rate 1 ml/min. The column was eluted with a linear gradient of 50 mM TEAB (pH 7.4) (solvent A) and solvent A:acetonitrile (60:40, v/v) (solvent B); 0–19 min, 40% B. The retention times of the nucleotides are presented in Table 1. At the aforementioned experimental conditions there was a linear dependence between time and AMP, the reaction product, peak areas. This allowed to calculate and compare the rates of the hydrolysis of investigated substrates.

The nucleoside phosphoramidase activity of the Fhits was estimated in a reaction mixture containing 50 mM Mes/KOH (pH 6.5) and appropriate concentration of NH₂-p[8-³H]A. When the K_m values were estimated the radiolabeled substrate concentration varied between 1 and 15 μ M. At time intervals, 5 μ l aliquots of the reaction mixture were spotted onto thin layer chromatography (TLC) aluminum plates pre-coated with silica gel containing fluorescent indicator (from Merck), standards of NH₂-pA and AMP applied at the origin and the plates developed for 60 min in dioxane/ammonia/water (6:1:4, by vol.). Spots of the nucleotides were visualized under short-wave UV light and those of the reaction product (AMP) cut out, immersed in a scintillation cocktail, and radioactivity counted.

K_i values of different nucleotides used in competition with NH₂-p[8-³H]A in the nucleoside phosphoramidase reaction were determined in a reaction mixture (50 μ l) containing 50 mM Mes/KOH (pH 6.5), 5 mM MgCl₂, 2.75 μ M NH₂-p[8-³H]A (580000 c.p.m.) and varied concentrations of one of the following nucleotides: AMP, ADP, ATP, Ap₃A, S-pA, F-pA or adenosine-5'-*O*-(γ -fluoridetriphosphate) (ATP-F or F-pppA). Reaction rates were estimated as described above. K_i values were calculated according [28].

Table 1
Various substrates of human and *Arabidopsis* Fhit proteins

Nucleotide	Retention time (min) ^a	Relative velocities of hydrolysis (%) ^b	
		Human Fhit	<i>Arabidopsis</i> Fhit
NH ₂ -pA	9.7	100	100
S-pA	10.3	65	75
ApppA	11.8	58	51
F-pppA	10.2	43	90
F-pA	13.1	25	62

^aThe conditions of the used HPLC are in Section 2. The retention times for the reactions' product were: 8.8 min for AMP (pA) and 10.3 for ADP (ppA).

^bThe reaction mixture contained 50 mM Mes/KOH (pH 6.5), 5 mM MgCl₂, 1 mM substrate and rate-limiting amounts of either human or *Arabidopsis* Fhit protein. The k_{cat} values for the preferred substrate NH₂-pA were 1.26 and 1.27 s⁻¹, respectively.

2.3. NMR spectroscopy

^{31}P NMR spectra were recorded on a Bruker DRX 500 spectrometer with a 5 mm BB probe. ^{19}F NMR spectra were recorded on the same spectrometer using a 5 mm $^1\text{H}/^{19}\text{F}$ dual probe. Data were processed using Felix (Felix-NMR Inc., San Diego). Standard Bruker referencing was used (1 M phosphoric acid for ^{31}P , and trichlorofluoromethane for ^{19}F).

3. Results

3.1. Identification and cloning of the *A. thaliana* FHIT gene

BLAST algorithm was used to search for the *A. thaliana* proteins similar to known Fhit proteins. The search identified product of the At5g58240 gene as the most likely candidate. The At5g58240 protein was 50% identical to human Fhit and 46% identical to *S. cerevisiae* Hnt2, whereas its similarity to Hint nucleoside phosphoramidases from the same organisms was much lower (27% and 23% identity to *S. cerevisiae* Hnt1 and human Hint1, respectively). Based on these results and on the catalytic properties we decided that At5g58240 gene encoded genuine Fhit protein. The complete *A. thaliana* FHIT-coding sequence was PCR-amplified, cloned and sequenced. The obtained sequence was identical to the mRNA sequence deposited in GENE BANK under accession number AK228164.

3.2. Fhit proteins behave as nucleoside phosphoramidases

Earlier observations that human Fhit protein effectively hydrolyzed the P–N bond in adenosine 5'-phosphoimidazolide [19] led us to check whether $\text{NH}_2\text{-pA}$, the simplest and naturally occurring nucleoside phosphoramidate, is also recognized as substrate by the human and plant Fhits. Analysis of reaction mixtures in which $\text{NH}_2\text{-pA}$ was the only potential substrate of the Fhits was performed first by TLC (Fig. 1) and then by HPLC (Table 1). It clearly showed that Fhits act as adenosine phosphoramidases and catalyze the Reaction 2:






Fig. 1. Hydrolysis of different substrates by the human Fhit analyzed by TLC. The reaction mixture (50 μl final volume) contained 50 mM Mes/KOH (pH 6.5), 5 mM MgCl_2 , 1 mM indicated substrate and 0.2 μg of recombinant human Fhit. Incubation was carried out at 30 $^\circ\text{C}$. At times indicated, 3 μl aliquots were spotted on a TLC plate (aluminium precoated with silica containing fluorescent indicator, from Merck). The chromatogram was developed in dioxane:ammonia:water (6;1:4, by vol.) and photographed under short-wave UV light.

In order to estimate the kinetic parameters of this reaction by the most direct and sensitive method, we used tritium-labeled $\text{NH}_2\text{-pA}$ as a substrate to determine pH optimum, metal ion requirements, K_m and k_{cat} values, and also K_i s for different nucleotides that act in the phosphoramidase reaction as substrate competitors. The two Fhit proteins investigated catalyzed cleavage of the P–N bond in $\text{NH}_2\text{-pA}$ most effectively at neutral pH in a reaction independent of Mg^{2+} . K_m values estimated in 50 mM Mes/KOH buffer (pH 6.8) were $3 \pm 0.7 \mu\text{M}$ both for the human and *Arabidopsis* Fhits and k_{cat} values were calculated to be 1.26 and 1.27 s^{-1} , respectively. Using non-specific adenosine phosphate deaminase [25] we converted $\text{NH}_2\text{-p}[8\text{-}^3\text{H}]\text{A}$ into $\text{NH}_2\text{-p}[8\text{-}^3\text{H}]\text{I}$ and demonstrated that the latter was also deaminated to IMP by the Fhits. Cleavage of the P–N bonds in those two substrates (1 mM) proceeded at the same rate. Finally, we estimated the K_i values for the human and *Arabidopsis* Fhits from the effects exerted by nucleotides used by the enzymes either as substrates for other reactions (see below) or as products (Table 2). Generally, human Fhit recognized fluoroadenylates more poorly than did its plant counterpart. Tested as an adenosine phosphoramidase, human Fhit was practically not inhibited by F-pA and F-pppA, whereas *Arabidopsis* Fhit was; with the K_i values approximately one (for F-pA) and two (for F-pppA) orders of magnitude higher than the K_m for $\text{NH}_2\text{-pA}$. Both Fhits were inhibited with comparable effectiveness by AMP, one of the reaction products. ADP and ATP, poorly inhibited the plant enzyme and were practically without effect when tested in the reaction catalyzed by the human one. The nucleoside-phosphoramidase reaction catalyzed by Fhits, i.e., the hydrolysis of $\text{NH}_2\text{-pA}$ or $\text{NH}_2\text{-pI}$, was inhibited neither by Ap_3A nor by F-pppA in the absence of Mg^{2+} , which is a co-substrate of the reactions of hydrolysis of Ap_3A or F-pppA (see below). Analysis of the data (Tables 1 and 2) shows that the more effective is the nucleotide as a substrate of Fhit, the stronger is its inhibitory effect on the conversion of $\text{NH}_2\text{-pA}$ into AMP and NH_3 .

3.3. Fhit proteins act as adenylyl sulfate sulfohydrolase


We further tested the promiscuity of Fhits in catalyzing the hydrolysis of different AMP-containing compounds. We found that the mixed anhydride, sulfoadenylate (S-pA), was also recognized by the Fhits as a substrate (Reaction 3):



This liberation of AMP from S-pA did not require Mg^{2+} . At 1 mM substrate concentration the catalytic release of AMP from S-pA proceeded at only slightly lower rates than the release of AMP from NH_2 -pA: 1.5-fold lower with the human Fhit and 1.3-fold lower with *Arabidopsis* Fhit (Table 1).

3.4. Fhit proteins catalyze the hydrolysis of P–F bond


Adenylyl fluoride (F-pA), was also tested as a potential substrate of human and plant Fhits. We have found that this non-natural nucleotide is also recognized as substrate for which P–F cleavage does not depend on Mg^{2+} (Fig. 1 and Table 1, see Reaction 4):



The rates of release of AMP from F-pA were 4- and 1.6-fold lower for the human and plant Fhits, respectively, than those from NH_2 -pA.

3.5. Liberation of AMP from F-pppA

We have also examined the substrate behavior of F-pppA. This ATP analogue is recognized as a substrate of Fhits only in the presence of Mg^{2+} and was cleaved to AMP (Fig. 1) and fluoropyrophosphate (Reaction 5):



The hydrolysis products were identified by ^{31}P NMR. Three major peaks are present in the 1D spectrum of the reaction products (Fig. 2). These correspond to AMP (3.6 ppm, [29]), and fluoropyrophosphate. The $^2J_{PP}$ coupling (17 Hz) and $^1J_{PF}$ coupling (920 Hz) are as expected for fluoropyrophosphate, while fluoromonophosphate would give rise to a doublet at -3 ppm [30,31].

We also confirmed earlier observations [13] that Fhit proteins do not hydrolyze ATP, ADP and adenosine 5'-tetraphosphate. It can be added here that ADP-ribose was also not cleaved to AMP and that diadenosine pyrophosphate (AppA), a good substrate for phosphodiesterase I, was hydrolyzed at an

extremely low rate; more than 10000-fold slower than the rate of the hydrolysis of NH_2 -pA.

4. Discussion

Fhit proteins comprise one of three branches of proteins within the HIT superfamily [14]. Whereas Fhits have been recognized primarily as typical dinucleoside triphosphatases [10,16], their adenosine phosphoramidase activity is known as a feature of a different branch called Hint [14,32,33]. In their elucidation of the mechanism of human Fhit action, Huang

et al. showed that the enzyme cleaves adenosine phosphoimidazole, an analog of the reaction intermediate having a P–N bond. We have used the simplest nucleoside phosphoramidate, NH_2 -pA, and established that both human and *Arabidopsis* Fhits function as effective nucleoside phosphoramidases. It has already been shown that two enzymatic activities capable of deaminating NH_2 -pA and yielding AMP exist in a higher plant (*Lupinus luteus*). One of them was a feature of the yellow lupin Ap_3A hydrolase and the other a typical nucleoside

phosphoramidase that did not exhibit the Ap_3A -ase activity [34]. Those and the present observations concerning the *Arabidopsis* Fhit strongly suggest that the yellow lupin enzyme characterized previously as a dinucleoside triphosphatase [35] is in fact a Fhit protein.

Our current study also shows that Fhit proteins catalyze cleavage of S-pA to AMP and sulfate. Till now that reaction

Table 2
Inhibition of the adenosine phosphoramidase activity of human and *Arabidopsis* Fhit proteins by various nucleotides

Inhibitor	K_i (μM)	
	Human Fhit	<i>Arabidopsis</i> Fhit
S-pA	13.6 \pm 2.2	7.1 \pm 1.6
ApppA/Mg ²⁺	18.0 \pm 3.0	2.1 \pm 0.2
F-pppA/Mg ²⁺	>1000	235 \pm 35
F-pA	>1000	18.2 \pm 0.7
pA	110 \pm 15	113 \pm 18
ppA	>1500	140 \pm 20
pppA	>2000	145 \pm 22

K_i values are means of three independent determinations. For details see Section 2.2. The K_m values estimated for adenosine 5'-phosphoramidate for the both Fhits were 3 μM .

has been assigned to an adenylyl sulfate sulfohydrolase (EC 3.6.2.1). Such an enzymatic activity was identified in rat liver extracts and it did not hydrolyze ATP [36]; as is the case with Ap₃A hydrolases/Fhits [33, and this study]. Neither ApppA nor NH₂-pA was then tested as potential substrates of that sulfohydrolase.

Because of the nucleoside phosphoramidase and the adenylyl sulfate sulfohydrolase activities identified in this work, Fhits must now be considered as enzymes involved in the metabolism of many AMP-containing compounds, a list minimally including diadenosine triphosphate, naturally occurring nucleoside 5'-phosphoramidates and adenosine 5'-phosphosulfate. The influence of Fhits on the metabolism of adenosine 5'-phosphosulfate may be even more important for cells than Fhits regulation of the metabolism of dinucleoside triphosphates. The newly revealed activities should be taken into account particularly in the study of the anti-oncogenic function of Fhit and extend it beyond the binding and/or hydrolysis of dinucleoside polyphosphates [37,38].

It has been known for some time that mutations or deletions of the Fhit genes results in an increased incidence of spontaneous tumor formation in humans [37]. Recently, a similar correlation has been observed in mice having a deletion of the Hint gene [39]. The results of our study suggest that in vivo Fhit

may recognize Hint substrates, i.e., at least nucleoside phosphoramidates. The loss of Hint may thereby be alleviated by Fhit enzymatic activity. Therefore, potential defects caused by the loss of HINT may be visible only when FHIT is also inactivated.

Their capacity to liberate AMP from F-pA broadens the catalytic promiscuity of Fhits and shows that, with respect to cleavage of the P–F bond, Fhits act like a phosphodiesterase I (EC 3.1.4.1), as first reported by Wittmann [23]. Subsequently, the latter enzyme was demonstrated to split the P–F bond in uridine 5'-*O*-phosphorofluoridate [40], inosine 5'-*O*-phosphorofluoridate [41] and thymidine 5'-*O*-phosphorofluoridate [42].

Liberation of AMP from F-pppA sheds new light on the substrate requirements of Fhits. F-pppA appears to mimic one of the natural substrates of Fhits, ApppA, both nucleotides having three phosphate negative charges. In Fhit-catalyzed reactions, the nucleophilic histidine attacks P ^{α} in both nucleotides to displace ADP in case of ApppA but fluoropyrophosphate in the case of F-pppA. In the past, F-pppA has been studied in various enzymatic systems. Haley and Yount reported that the compound was cleaved to AMP and fluoropyrophosphate by snake venom phosphodiesterase [43]. Thus, the latter behaves as the Fhits.

The newly identified properties of Fhits can also be used in studies on the delivery of pronucleotides to target cells or organisms [44]. It seems viable that nucleoside phosphoramidates can be employed as nucleotide prodrugs. The same may be true for nucleoside phosphofluoridates that are also less polar than nucleoside monophosphates and therefore should penetrate the cell membrane better. As mentioned above, the plant-specific phosphoramidase did not hydrolyze ApppA [34]. Whether such specific phosphoramidases can cleave nucleoside phosphofluoridates thereby liberating nucleoside monophosphates from prodrugs is worthy of further study. Our work shows that both the specific nucleoside phosphoramidases (Hint proteins) and Fhits may be important in the metabolism of such prodrugs.

Acknowledgements: This work was supported by the Polish Ministry of Science and Higher Education (Grant PBZ-MNiSW-07/1/2007 to A.G., A.M.W. and M.P.-B. and Grant 2 P04A 05029 to P.B.).




Fig. 2. ³¹P NMR spectrum of the products of the reaction of ATP- γ F with human Fhit. The reaction mixture (0.1 ml total volume) contained 50 mM HEPES/KOH (pH 7.8), 5 mM MgCl₂, 1 mM F-pppA and 1 μg of the human Fhit. The reaction was carried out at 25 °C and its completion confirmed by TLC. Peaks are assigned as follows: 3.6 ppm (singlet) AMP; -5.8 ppm (d, ²J_{PP} = 17 Hz), fluoropyrophosphate (FPP); -16 ppm (dd, ¹J_{FP} = 920 Hz, ²J_{PP} = 17 Hz), fluoropyrophosphate (FPP). Spectra were recorded with 32 K scans using a spectral width of 12 136 Hz and 4 K points resulting in 338 ms acquisition time. Couplings are accurate to 3 Hz. Assignments made in comparison to those published [30,45].

References

- [1] Garrison, P.N. and Barnes, L.D. (1992) Determination of dinucleoside polyphosphates in: Ap₄A and Other Dinucleoside polyphosphates (McLennan, A.G., Ed.), pp. 29–61, CRC Press, Boca Raton, FL.
- [2] Guranowski, A. (2004) Metabolism of diadenosine tetraphosphate (Ap₄A) and related nucleotides in plants; review with historical and general perspective.
- [3] Guranowski, A., deDiego, A., Sillero, A. and Günther Sillero, M.A. (2004) Uridine 5'-polyphosphates (p₄U) and (p₅U) and uridine(5')polyphospho(5')nucleosides (Up_nNs) can be synthesized by UTP:glucose-1-phosphate uridylyltransferase from *Saccharomyces cerevisiae*. FEBS Lett. 561, 83–88.
- [4] McLennan, A.G. (2000) Dinucleoside polyphosphates – friend or foe? Pharmacol. Ther. 87, 73–89.
- [5] McLennan, A.G., Barnes, L.D., Blackburn, G.M., Brenner, Ch., Guranowski, A., Miller, A.D., Rovira, J.M., Rotllán, P., Soria, B., Tanner, J.A. and Sillero, A. (2001) Recent progress in the study of the intracellular functions of diadenosine polyphosphates. Drug Dev. Res. 52, 249–259.
- [6] Hoyle, C.H.V., Hilderman, R.H., Pintor, J.J., Schlüter, H. and King, B.F. (2001) Diadenosine polyphosphates as extracellular signal molecules. Drug Dev. Res. 52, 260–273.
- [7] Guranowski, A. (2000) Specific and nonspecific enzymes involved in the catabolism of mononucleoside and dinucleoside polyphosphates. Pharmacol. Ther. 87, 117–139.
- [8] Lobatón, C.D., Sillero, M.A.G. and Sillero, A. (1975) Diadenosine triphosphate splitting by rat liver extracts. Biochem. Biophys. Res. Commun. 67, 279–286.
- [9] Jakubowski, H. and Guranowski, A. (1983) Enzymes hydrolyzing ApppA and/or AppppA in higher plants; purification and some properties of diadenosine triphosphatase, diadenosine tetraphosphatase, and phosphodiesterase from yellow lupin (*Lupinus luteus*) seeds. J. Biol. Chem. 258, 9982–9989.
- [10] Brevet, A., Chen, J., Fromant, M., Blanquet, S. and Plateau, P. (1991) Isolation and characterization of a dinucleoside triphosphatase from *Saccharomyces cerevisiae*. J. Bacteriol. 173, 5275–5279.
- [11] Prescott, M., Thorne, N.M.H., Milne, A.D. and McLennan, A.G. (1992) Characterisation of a bis(5'-nucleosidyl)triphosphate pyrophosphohydrolase from encysted embryos of the brine shrimp *Artemia*. Int. J. Biochem. 24, 565–571.
- [12] McLennan, A.G., Mayers, E., Hankin, S., Thorne, N.M.H., Prescott, M. and Powls, R. (1994) The green alga *Scenedesmus obliquus* contains both diadenosine 5',5'''-P¹,P⁴-tetraphosphate (asymmetrical) pyrophosphohydrolase and phosphorylase activities. Biochem. J. 300, 183–189.
- [13] Barnes, L.D., Garrison, P.N., Siprashvili, Z., Guranowski, A., Robinson, A.K., Ingram, S.W., Croce, C.M., Ohta, M. and Huebner, K. (1996) Fhit, a putative tumor suppressor in humans, is a dinucleoside 5',5'''-P¹,P³-triphosphate hydrolase. Biochemistry 35, 11529–11535.
- [14] Brenner, Ch. (2002) Hint, Fhit, and GalT: function, structure, evolution, and mechanism of three branches of the histidine triad superfamily of nucleotide hydrolases and transferases. Biochemistry 41, 9003–9014.
- [15] Pawelczyk, T., Kowara, R., Gołębowski, F. and Matecki, A. (2000) Expression in *Escherichia coli* and simple purification of human Fhit protein. Protein Express. Purif. 18, 320–326.
- [16] Chen, J., Brevet, A., Blanquet, S. and Plateau, P. (1998) Control of 5',5'-dinucleoside triphosphate catabolism by APHI, a *Saccharomyces cerevisiae* analog of human FHIT. J. Bacteriol. 180, 2345–2349.
- [17] Guranowski, A., Brown, P., Ashton, P.A. and Blackburn, G.M. (1994) Regiospecificity of the hydrolysis of diadenosine polyphosphates catalyzed by three specific pyrophosphohydrolases. Biochemistry 33, 235–240.
- [18] Abend, A., Garrison, P.N., Barnes, L.D. and Frey, P.A. (1999) Stereochemical retention of the configuration in the action of Fhit on phosphorus-chiral substrates. Biochemistry 38, 3668–3676.
- [19] Huang, K., Arabshahi, A., Wei, Y. and Frey, P.A. (2004) The mechanism of action of the fragile histidine triad, Fhit: isolation of a covalent adenylyl enzyme and chemical rescue of H96G-Fhit. Biochemistry 43, 7637–7642.
- [20] Huang, K., Arabshahi, A. and Frey, P.A. (2005) pH-dependence in the hydrolytic action of the human fragile histidine triad. Eur. J. Org. Chem., 5198–5206.
- [21] Frankhauser, H., Berkowitz, G.A. and Schiff, J.A. (1981) A nucleotide with the properties of adenosine 5' phosphoramidate from *Chlorella* cells. Biochem. Biophys. Res. Commun. 101, 524–532.
- [22] Frankhauser, H., Schiff, J.A. and Garber, L.J. (1981) Purification and properties of adenylyl sulphate:ammonia adenylyltransferase from *Chlorella* catalyzing the formation of adenosine 5'-phosphoramidate from adenosine 5'-phosphosulphate and ammonia. Biochem. J. 195, 545–560.
- [23] Wittmann, R. (1963) Die reaktion der phosphorsäuren mit 2,4-dinitro-fluorobenzol. I: Eine neue synthese von monofluorophosphorsäuremonoestern. Chem. Ber. 96, 771–779.
- [24] Stumber, M., Herrmann, Ch., Wohlgemuth, S., Kalbitzer, H.R., Jahn, W. and Geyer, M. (2002) Synthesis, characterization and application of two nucleoside triphosphate analogues, GTPγNH₂ and GTPγF. Eur. J. Biochem. 269, 3270–3278.
- [25] Guranowski, A., Starzyńska, E., Günther Sillero, M.A. and Sillero, A. (1995) Conversion of adenosine (5')oligophospho(5')adenosines into inosine(5')oligophospho(5')inosines by non-specific adenylylase from the snail *Helix pomatia*. Biochim. Biophys. Acta 1243, 78–84.
- [26] Ghosh, S. and Lowenstein, J.M. (1996) A multifunctional vector system for heterologous expression of proteins in *Escherichia coli*. Expression of native and hexahistidyl fusion proteins, rapid purification of the fusion protein, and removal of fusion peptide by Kex2 protease. Gene 176, 249–255.
- [27] Brenner, Ch., Pace, H.C., Garrison, P.N., Rösler, A., Liu, X.H., Blackburn, G.M., Croce, C.M., Huebner, K. and Barnes, L.D. (1997) Purification and crystallization of complexes modeling the active site of the fragile histidine triad protein. Protein Eng. 10, 1461–1463.
- [28] Dixon, M. and Webb, E.C. (1964) Enzymes, second ed, Academic Press, New York.
- [29] Gradwell, M.J., Fan, T.W.M. and Lane, A.N. (1998) Analysis of phosphorylated metabolites in crayfish extracts by two-dimensional ¹H–³¹P NMR heteronuclear total correlation spectroscopy (heteroTOCSY). Anal. Biochem. 263, 139–149.
- [30] Iulucci, R.J. and Meier, B.H. (1998) A characterization of the linear P–O–P bonds in M4 + (P2O7) compounds: bond-angle determination by solid-state NMR. J. Am. Chem. Soc. 120, 9059–9062.
- [31] Yoza, N., Nakashima, S., Ueda, N., Miyajima, T., Nakamura, T. and Vast, P. (1994) High performance liquid chromatographic characterization of monofluorophosphate, difluorophosphate and hexafluorophosphate. J. Chromatogr. A 664, 111–116.
- [32] Guranowski, P., Garrison, P.N., Hodawedekar, S.C., Faye, G., Barnes, L.D. and Brenner, Ch. (2002) Adenosine monophosphoramidase activity of Hint and Hint1 supports function of Kin28, Cc11, and Tfb3. J. Biol. Chem. 277, 10852–10860.
- [33] Krakowiak, A., Pace, H.C., Blackburn, G.M., Adams, M., Mekhalfia, A., Kaczmarek, R., Baraniak, J., Stec, W.J. and Brenner, Ch. (2004) Biochemical, crystallographic, and mutagenic characterization of Hint, the AMP-lysine hydrolase, with novel substrates and inhibitors. J. Biol. Chem. 279, 18711–18716.
- [34] Guranowski, A., Bieganski, P., Baraniak, J., Rydzik, A., Stepiński, J. and Jemielity, J. (2006) Catabolism of nucleoside phosphoramidates in higher plants can be controlled by nucleoside phosphoramidase (Hint protein) and dinucleoside triphosphatase (Fhit protein). Acta Biochim. Pol. 53 S1, 183–184.
- [35] Guranowski, A., Starzyńska, E., Bojarska, E., Stepiński, J. and Darzynkiewicz, E. (1996) Dinucleoside 5',5'''-P¹,P³-triphosphate hydrolase from yellow lupin (*Lupinus luteus*) seeds; purification to homogeneity and hydrolysis of mRNA 5'-cap analogs. Protein Express. Purif. 8, 416–422.
- [36] Bailey-Wood, R., Dodgson, K.S. and Rose, F.A. (1969) A rat liver sulphohydrolase enzyme acting on adenylyl sulphate. Biochem. J. 112, 257–258.
- [37] Siprashvili, Z., Sozzi, G., Barnes, L.D., McCue, P., Robinson, A.K., Eryomin, V., Sard, L., Tagliabue, E., Greco, A., Fusetti, L., Schwartz, G., Pierotti, M.A., Croce, C.M. and Huebner, K. (1997) Replacement of Fhit in cancer cells suppresses tumorigenicity. Proc. Natl. Acad. Sci. USA 94, 13771–13776.

- [38] Fisher, D.I. and McLennan, A.G. (2008) Correlation of intracellular diadenosine triphosphate (Ap₃A) with apoptosis in Fhit-positive HEK293 cells. *Cancer Lett.* 259, 186–191.
- [39] Su, T., Suzui, M., Wang, L., Lin, Ch.-Sh., Xing, W.-Q. and Weinstein, I.B. (2003) Deletion of histidine triad nucleotide-binding protein 1/PKC-interacting protein in mice enhances cell growth and carcinogenesis. *Proc. Natl. Acad. Sci. USA* 100, 7824–7829.
- [40] Kučerová, Z. and Škoda, J. (1971) Resistance of uridine 5'-fluorophosphate to alkaline phosphatase and its sensitivity to 5'-nucleotidase. *Biochim. Biophys. Acta* 247, 194–196.
- [41] Nichol, A.W., Nomura, A. and Hampton, A. (1967) Studies on phosphate binding sites of inosinic acid dehydrogenase and adenylosuccinate synthetase. *Biochemistry* 6, 1008–1015.
- [42] Misiura, K., Szymanowicz, D. and Stec, W.J. (1998) Synthesis and chemical and enzymatic reactivity of thymidine 3'-*O*- and 5'-*O*-phosphorofluoridothioates. *Chem. Commun.*, 515–516.
- [43] Haley, B. and Yount, R.G. (1972) γ -Fluoroadenosine triphosphate. Synthesis, properties, and interaction with myosin and heavy meromyosin. *Biochemistry* 11, 2863–2871.
- [44] Chou, T.F., Baraniak, J., Kaczmarek, R., Zhou, X., Cheng, J., Ghosh, B. and Wagner, C.R. (2007) Phosphoramidate pronucleotides: a comparison of the phosphoramidase substrate specificity of human and *Escherichia coli* histidine triad nucleotide binding proteins. *Mol. Pharm.* 4, 208–217.
- [45] Falius, H. (1968) Synthesis of dipotassium difluorodiphosphate by thermal reaction of tetrachlorophosphorus decaoxide with potassium fluoride. *Angew. Chem. Int. Ed.* 7, 622.