



ELSEVIER

Biochimica et Biophysica Acta 1412 (1999) 295–300

BIOCHIMICA ET BIOPHYSICA ACTA

BBAwww.elsevier.com/locate/bba

Short sequence-paper

Characterization of the human *SDHD* gene encoding the small subunit of cytochrome *b* (cybS) in mitochondrial succinate–ubiquinone oxidoreductase

Hiroko Hirawake ^a, Masafumi Taniwaki ^b, Akiko Tamura ^b, Hisako Amino ^a, Eriko Tomitsuka ^a, Kiyoshi Kita ^{a,*}^a Department of Biomedical Chemistry, Graduate School of Medicine, The University of Tokyo, 7-3-1, Hongo, Bunkyo-ku, Tokyo 113-0033, Japan^b The Third Department of Internal Medicine, Kyoto Prefectural University of Medicine, Kyoto, Japan

Received 25 May 1999; accepted 17 June 1999

Abstract

We have mapped large (cybL) and small (cybS) subunits of cytochrome *b* in the succinate–ubiquinone oxidoreductase (complex II) of human mitochondria to chromosome 1q21 and 11q23, respectively (H. Hirawake et al., *Cytogenet. Cell Genet.* 79 (1997) 132–138). In the present study, the human *SDHD* gene encoding cybS was cloned and characterized. The gene comprises four exons and three introns extending over 19 kb. Sequence analysis of the 5' promoter region showed several motifs for the binding of transcription factors including nuclear respiratory factors NRF-1 and NRF-2 at positions –137 and –104, respectively. In addition to this gene, six pseudogenes of cybS were isolated and mapped on the chromosome. © 1999 Elsevier Science B.V. All rights reserved.

Keywords: Mitochondria; Succinate dehydrogenase; Cytochrome *b*; Gene structure; *SDHD*; Nuclear respiratory factor (NRF); (Human)

Succinate–ubiquinone oxidoreductase (complex II) is an important enzyme complex in both the tricarboxylic acid cycle and the aerobic respiratory chains of eukaryotic cell mitochondria and prokaryotic cells. Complex II catalyzes the oxidation of succinate to fumarate (succinate dehydrogenase: SDH) and transfers its reducing equivalent to ubiquinone [1,2]. Complex II also catalyzes the reduction of fumarate (fumarate reductase: FRD), which is the reverse of the reaction catalyzed by SDH, in the respiratory

chain of anaerobic bacteria and in the mitochondria of facultative anaerobic animals such as adult *Ascaris suum* [3,4]. Complex II is generally composed of four polypeptides with apparent molecular masses of 70, 30, 15, and 13 kDa, and contains five prosthetic groups, one covalently linked FAD, three iron–sulfur clusters (2Fe–2S, 4Fe–4S, and 3Fe–4S), and heme *b*. The larger two subunits, the flavoprotein subunit (Fp) and the iron–sulfur protein subunit (Ip), comprise the catalytic portion of the enzyme complex and catalyze electron transfer from succinate to artificial electron acceptors such as phenazine methosulfate (PMS). The amino acid sequence of Fp and Ip are highly conserved and cDNAs and genes have

* Corresponding author. Fax: +81-3-5841-3444;
E-mail: kitak@m.u-tokyo.ac.jp



Fig. 1. Genomic structure of the human *SDHD* gene. The four exons of the gene (E1–4) are boxed. Restriction sites for *SacI* are shown. Numbers indicate positions from the possible transcription start site (+1) determined by 5'-end analysis of the cDNA.

been cloned from various species using homology probing strategies (see for reviews [1,2]). The presence of a two-subunit cytochrome *b* composed of large (cybL, also referred to as QPs-1 or C_{II-3}) and small (cybS, also referred to as QPs-3 or C_{II-4}) subunits acting as hydrophobic membrane anchor peptides is a general feature of mitochondrial complex II [5,6]. In addition, the cytochrome *b* in complex II is essential for the interaction between the complex and quinone species. The ubiquinone-binding domains in cybL and cybS of bovine heart complex II have been suggested to be residues 113–140 and 20–37, respectively, by matching the sequences of azide-ubiquinone linked peptides to their respective protein sequences [7,8]. However, the function and structure of cytochrome *b* in complex II is poorly understood as compared with the well-characterized Fp and Ip because of a lack of sequence conservation at the nucleotide and/or amino acid sequence level.

To elucidate the molecular basis of a mitochondrial disease caused by a deficiency in the SDH activity of complex II ([9] and see [10] for review), we cloned the cDNAs for all four subunits of human liver complex II and found their unique features [11–13]. All the genes for human complex II are en-

coded on nuclear DNA, and the genes for Fp (*SDHA*) and Ip (*SDHB*) have been mapped on chromosomes 5 [14] and 1 [15]. Recently, we mapped the genes for cybL and cybS, *SDHC* and *SDHD*, to chromosomes 1q21 and 11q23, respectively [13]. Among these genes, the structures of *SDHB* and *SDHC* have been reported [15,16]. In this study, the entire nucleotide sequence of human *SDHD* was determined and characteristics of the promoter region were analyzed.

In the previous study of mapping the *SDHD* gene, we obtained two independent genomic clones from a human EMBL-3 library (Clontech, Lot 1125), S-3 and S-36, and used both in the chromosome assignment by FISH [13]. We determined the complete nucleotide sequence of the DNA fragment from clone S-36, since restriction enzyme analysis and Southern hybridization using probes specific to the cDNA for liver cybS showed that the S-3 clone (11.7 kb) contains three exons from the 5'-terminal, and the S-36 clone (18.6 kb) contains the entire coding sequence of four exons, the entire 3'-untranslated region, and at least part of the 5'-untranslated region. Fig. 1 shows a schematic representation of the genomic structure of the human *SDHD* gene,

Table 1
Potential sites for transcription factors in 5' upstream region of the *SDHD* gene

Transcription factor	Sequence	Position
CdxA	AATAATAATA	-5560, -4813, -4490, -2915, -1043
GATA-1	GCCAATCAGA	-1881
GATA-2	ACCTATCCCC	-1857
Lyf-1	TTTGGGAGG	-5512
Nkx-2	CACTTGA	-5662, -5354, -2830, -2800
NRF-1	TTGCGCATGCGCG	-137
NRF-2	GACTTCCGGT	-428, -104
Sox-5	TTAACAATAC	-4949
SRY	AAACAAA	-4597, -3995, -3916
USF	GCACGTGG	-1130, -596
YY1	GATAGTAAATGGTTT	-3263, -3227, -1857, -945

Consensus sequences as targets for transcription factors were analyzed by TFSEARCH (Y. Akiyama, Kyoto University).




Fig. 2. Nucleotide sequence of the human SDHD gene. Uppercase letters indicate exon sequences and lowercase letters indicate promoter and intron sequences. Amino acid sequences of the coding region are also shown. Splice donor and acceptor sites are underlined for each intron. Consensus sequences as targets for transcription factors were analyzed by TFSEARCH (Y. Akiyama, Kyoto University). Potential nuclear respiratory factor binding sites, NRF-1 and NRF-2, are also underlined. This sequence is available in the DDBJ, EMBL, and NCBI databases under accession number AB026906.




Fig. 3. Localization of introns in the human *SDHD* gene. Sites are indicated by arrows. ▼ indicates conserved histidine residues. ▽ indicates the amino termini of mature peptides. Amino acids identical to those in human *cybS* are boxed. *S. cerevisiae SDH4* [26]; *Homo sapiens* [13]; *A. suum* [27]; *Escherichia coli sdhd* [28].

and the sequence, including the entire coding region, is shown in Fig. 2. The nucleotide sequence of the human *SDHD* gene appears in the DDBJ, EMBL, and Gen Bank nucleotide sequence databases with the accession number AB026906. All exons are quite small, 52, 117, 145, and 163 nucleotides in exons 1, 2, 3, and 4, respectively, and the nucleotide sequence of the coding regions is completely identical to that of the liver cDNA, which we reported previously [13]. Donor and acceptor sites for RNA splicing were confirmed except 'gc' in the third intron. The localization of introns in the amino acid sequence of liver *cybS* is shown in Fig. 3. The first intron was found in the mitochondrial presequence that is essential for the import of mitochondrial proteins encoded by nuclear DNA [17]. The second intron is located near the cleavage site of the mitochondria presequence. This site corresponds to the putative amino-terminal of bovine heart *cybS* (Ser-Asp-Ser-Lys-Ala-Ala-Ser) determined from the peptide [18]. The third intron divides the region containing two conserved histidine residues in the *cybS* peptide of mitochondria from multicellular organisms. These two histidine residues are found in the putative second transmembrane segment and appear to be axial ligands of heme *b* in complex II [19–21].

An analysis of the approximately 6 kb upstream of exon 1 was carried out to identify potential consensus sequences as targets for transcription factors. The *SDHD* gene is typical of mammalian housekeeping

genes with TATA-less promoters. Interestingly, there are two distinct potential nuclear respiratory factor binding sites, NRF-1 and NRF-2 [22], at positions -137 and -104, respectively (Figs. 2 and 4). These sites have been found upstream of *SDHB* [23] and *SDHC* [16], indicating that two factors and sites are required for the coordinated control of gene expression for the stoichiometric assembly of a functional complex II. In addition to these sites, several potential sites for other transcription factors were found (Table 1). Many experiments, including gel shift assay, will be necessary to determine the requirements for these factors and their binding sites in the expression of *SDHD*.

Genetic heterogeneity and the presence of tissue specific isoforms of human complex II have been suggested from the study of the variable clinical expression of complex II deficiency including tissue-to-tissue variation [24]. Recently, we found two different stage-specific isoforms of complex II containing distinct cytochrome *bs* in *A. suum* mitochondria [25]. In this regard, it is of interest to note the high positive background obtained during the screening of the




Fig. 4. Schematic representation of the putative promoter region of human *SDHD*.

Table 2
Chromosomal localization of pseudogenes for human *SDHD*

Clone	Localization	Mutation	Comment
#6	1p36.1	No	poly A, 6 amino acid substitutions
#8	2q32.1	Stop, deletion	
#11	3p21.3	Stop, frame shift	
#12	3q26.3	Deletion, frame shift	
#10	7q32.1	Stop	
#5	18q11.2	Stop, frame shift	

No introns were found in these pseudogenes.

SDHD gene using a cDNA probe for cybS. Eleven clones in addition to S-3 and S-36 were isolated, sequenced, and mapped. These clones were divided into six groups and appear to be for pseudogenes of *SDHD* because they contain stop codons, deletions, and frame shifts in their sequences (Table 2). One exception is clone S-1, which was mapped to chromosome 1p36 by FISH. This clone has an open reading frame almost identical to the liver cDNA with only six amino acid substitutions, L4P, L14Q, L77P, M91T, L120S, and L131F. Since it would be of interest to prove the presence of two or more active genes of cybS, the expression of the clone S-1 gene and screening of another candidate for a second gene are now in progress.

This work was supported by a Grant-in-Aid for Scientific Research on Priority Areas from the Ministry of Education, Science, Culture and Sports of Japan (08281105, 10129202, and 11470065). This work was also performed partly under the management of Research Institute of Innovative Technology for the Earth (RITE) as a part of Development of a High-Performance Bioreactor for the Production of Biochemicals supported by New Energy and Industrial Technology Development Organization (NEDO).

References

- [1] C. Hägerhäll, Succinate:quinone oxidoreductase; variations on a conserved theme, *Biochim. Biophys. Acta* 1320 (1997) 107–141.
- [2] I.E. Scheffler, Molecular genetics of succinate:quinone oxidoreductase in eukaryotes, *Prog. Nucleic Acid Res. Mol. Biol.* 60 (1998) 267–315.
- [3] K. Kita, H. Hirawake, S. Takamiya, Cytochromes in the respiratory chain of helminth mitochondria, *Int. J. Parasitol.* 27 (1997) 617–630.
- [4] A.G.M. Tielens, J.J. Van Hellemond, The electron transport chain in anaerobically functioning eukaryotes, *Biochim. Biophys. Acta* 1365 (1998) 71–78.
- [5] C. Hägerhäll, L. Hederstedt, A structural model for the membrane-integral domain of succinate–quinone oxidoreductases, *FEBS Lett.* 389 (1996) 25–31.
- [6] K.S. Oyedotun, B.D. Lemire, The *Saccharomyces cerevisiae* succinate–ubiquinone reductase contains a stoichiometric amount of cytochrome *b*₅₆₂, *FEBS Lett.* 442 (1999) 203–207.
- [7] G.Y. Lee, D.Y. He, L. Yu, C.A. Yu, Identification of the ubiquinone-binding domain in QPsl of succinate–ubiquinone reductase, *J. Biol. Chem.* 270 (1995) 6193–6198.
- [8] S.K. Shenoy, L. Yu, C.A. Yu, The smallest membrane anchoring subunit (QPs3) of bovine heart mitochondrial succinate–ubiquinone reductase: cloning, sequencing, topology, and Q-binding domain, *J. Biol. Chem.* 272 (1997) 17867–17872.
- [9] K. Isobe, S. Kishino, K. Inoue, D. Takai, H. Hirawake, K. Kita, S. Miyabayashi, J. Hayashi, Identification of inheritance modes of mitochondrial diseases by introduction of pure nuclei from mtDNA-less HeLa cells to patient-derived fibroblasts, *J. Biol. Chem.* 272 (1997) 12606–12610.
- [10] D.C. Wallace, Mitochondrial diseases in man and mouse, *Science* 283 (1999) 1482–1488.
- [11] K. Kita, H. Oya, R.B. Gennis, B.A.C. Ackrell, M. Kasahara, Human complex II (succinate–ubiquinone oxidoreductase): cDNA cloning of iron–sulfur (Ip) subunit of liver mitochondria, *Biochem. Biophys. Res. Commun.* 166 (1990) 101–108.
- [12] H. Hirawake, H. Wang, T. Kuramochi, S. Kojima, K. Kita, Human complex II (succinate–ubiquinone oxidoreductase): cDNA cloning of the flavoprotein (Fp) subunit of liver mitochondria, *J. Biochem.* 116 (1994) 221–227.
- [13] H. Hirawake, M. Taniwaki, A. Tamura, S. Kojima, K. Kita, Cytochrome *b* in human complex II (succinate–ubiquinone oxidoreductase): cDNA cloning of the components in liver mitochondria and chromosome assignment of the genes for the large (SDHC) and small (SDHD) subunits to 1q21 and 11q23, *Cytogenet. Cell Genet.* 79 (1997) 132–138.
- [14] T. Bourgeron, P. Rustin, D. Chretien, M. Birch-Machin, M. Bourgeois, E. Viegas-Péquignot, A. Munnich, A. Rötig, Mutation of a nuclear succinate dehydrogenase gene results in mitochondrial respiratory chain deficiency, *Nat. Genet.* 11 (1995) 144–149.
- [15] H.C. Au, L.A. Bellew, P.L.E. Broomfield, M. Saghbini, I.E. Scheffler, Structural organization of the gene encoding the human iron–sulfur subunit of succinate dehydrogenase, *Gene* 159 (1995) 249–253.
- [16] A. Elbehti-Green, H.C. Au, J.T. Mascarello, D. Ream-Robinson, I.E. Scheffler, Characterization of the human *SDHC* gene encoding one of the integral membrane proteins of succinate–quinone oxidoreductase in mitochondria, *Gene* 213 (1998) 133–140.

- [17] E.C. Hurt, A.P. vanLoon, How proteins find mitochondria and intramitochondrial compartments, *Trends Biochem. Sci.* 11 (1986) 204–207.
- [18] B. Cochran, R.A. Capaldi, B.A.C. Ackrell, The cDNA sequence of beef heart C_{II-3}, a membrane-intrinsic subunit of succinate-ubiquinone oxidoreductase, *Biochim. Biophys. Acta* 1188 (1994) 162–166.
- [19] K. Nakamura, M. Yamaki, M. Sarada, S. Nakayama, C.R.T. Vibat, R.B. Gennis, T. Nakayashiki, H. Inokuchi, S. Kojima, K. Kita, Two hydrophobic subunits are essential for the heme b ligation and functional assembly of complex II (succinate-ubiquinone oxidoreductase) from *Escherichia coli*, *J. Biol. Chem.* 271 (1996) 521–527.
- [20] C.R.T. Vibat, G. Cecchini, K. Nakamura, K. Kita, R.B. Gennis, Localization of histidine residues responsible for heme axial ligation in cytochrome *b556* of complex II (succinate:ubiquinone oxidoreductase) in *Escherichia coli*, *Biochemistry* 37 (1998) 4148–4159.
- [21] S.K. Shenoy, L. Yu, C.A. Yu, Identification of quinone-binding and heme-ligating residues of the smallest membrane-anchoring subunit (QPs3) of bovine heart mitochondrial succinate:ubiquinone reductase, *J. Biol. Chem.* 274 (1999) 8717–8722.
- [22] R.C. Scarpulla, Nuclear respiratory factors and the pathways of nuclear-mitochondrial interaction, *Trends Cardiovasc. Med.* 6 (1996) 39–45.
- [23] H.C. Au, I.E. Scheffler, Analysis of the promoter of the succinate dehydrogenase iron-protein gene: both nuclear respiratory factors NRF-1 and NRF-2 are required, *Eur. J. Biochem.* 251 (1998) 164–174.
- [24] M.A. Birch-Machin, C. Marsac, G. Ponsot, B. Parfait, R.W. Taylor, P. Rustin, A. Munnich, Biochemical investigations and immunoblot analyses of two unrelated patients with an isolated deficiency in complex II of the mitochondrial respiratory chain, *Biochem. Biophys. Res. Commun.* 220 (1996) 57–62.
- [25] F. Saruta, T. Kuramochi, K. Nakamura, S. Takamiya, Y. Yu, T. Aoki, K. Sekimizu, S. Kojima, K. Kita, Stage-specific isoforms of complex II (succinate-ubiquinone oxidoreductase) in mitochondria from the parasitic nematode, *Ascaris suum*, *J. Biol. Chem.* 270 (1995) 928–932.
- [26] B.L. Bullis, B.D. Lemire, Isolation and characterization of the *Saccharomyces cerevisiae SDH4* gene encoding a membrane anchor subunit of succinate dehydrogenase, *J. Biol. Chem.* 269 (1994) 6543–6549.
- [27] F. Saruta, H. Hirawake, S. Takamiya, Y. Ma, T. Aoki, K. Sekimizu, S. Kojima, K. Kita, Cloning of a cDNA encoding the small subunit of cytochrome *b558* (cybS) of mitochondrial fumarate reductase (complex II) from adult *Ascaris suum*, *Biochim. Biophys. Acta* 1270 (1996) 1–5.
- [28] D. Wood, M.G. Darlison, R.J. Wilde, J.R. Guest, Nucleotide sequence encoding the flavoprotein and hydrophobic subunits of the succinate dehydrogenase of *Escherichia coli*, *Biochem. J.* 222 (1984) 519–534.