

The Oncogenic MicroRNA miR-22 Targets the TET2 Tumor Suppressor to Promote Hematopoietic Stem Cell Self-Renewal and Transformation

Su Jung Song,^{1,2,3,4,9} Keisuke Ito,^{1,2,3,4,9,10} Ugo Ala,^{1,2,3,4} Lev Kats,^{1,2,3,4} Kaitlyn Webster,^{1,2,3,4} Su Ming Sun,⁶ Mojca Jongen-Lavrencic,⁶ Katia Manova-Todorova,⁷ Julie Teruya-Feldstein,⁸ David E. Avigan,⁵ Ruud Delwel,⁶ and Pier Paolo Pandolfi^{1,2,3,4,*}

¹Cancer Genetics Program, Beth Israel Deaconess Cancer Center

²Department of Medicine, Division of Genetics, Beth Israel Deaconess Medical Center

³Department of Pathology, Beth Israel Deaconess Medical Center

⁴Division of Genetics, Beth Israel Deaconess Medical Center

⁵Division of Hematology and Oncology, Beth Israel Deaconess Medical Center

Harvard Medical School, Boston, MA 02215, USA

⁶The Institute for Hematology, Erasmus Medical Center, Rotterdam 5201, Netherlands

⁷Molecular Cytology

⁸Department of Pathology

Sloan-Kettering Institute, Memorial Sloan-Kettering Cancer Center, New York, NY 10065, USA

⁹These authors contributed equally to this work

¹⁰Present address: Ruth L. and David S. Gottesman Institute for Stem Cell and Regenerative Medicine Research, Departments of Cell Biology/Stem Cell Institute and Medicine, Albert Einstein Cancer Center, Albert Einstein College of Medicine, Bronx, NY 10461, USA

*Correspondence: ppandolf@bidmc.harvard.edu

<http://dx.doi.org/10.1016/j.stem.2013.06.003>

SUMMARY

MicroRNAs are frequently deregulated in cancer. Here we show that miR-22 is upregulated in myelodysplastic syndrome (MDS) and leukemia and its aberrant expression correlates with poor survival. To explore its role in hematopoietic stem cell function and malignancy, we generated transgenic mice conditionally expressing miR-22 in the hematopoietic compartment. These mice displayed reduced levels of global 5-hydroxymethylcytosine (5-hmC) and increased hematopoietic stem cell self-renewal accompanied by defective differentiation. Conversely, miR-22 inhibition blocked proliferation in both mouse and human leukemic cells. Over time, miR-22 transgenic mice developed MDS and hematological malignancies. We also identify TET2 as a key target of miR-22 in this context. Ectopic expression of TET2 suppressed the miR-22-induced phenotypes. Downregulation of TET2 protein also correlated with poor clinical outcomes and miR-22 overexpression in MDS patients. Our results therefore identify miR-22 as a potent proto-oncogene and suggest that aberrations in the miR-22/TET2 regulatory network are common in hematopoietic malignancies.

INTRODUCTION

Epigenetic mechanisms such as DNA methylation and histone modification drive stable and clonally propagated changes in

gene expression and have recently been shown to serve as molecular mediators of pathway dysfunction in neoplasia (Esteller, 2008; Rodríguez-Paredes and Esteller, 2011). Indeed, both myelodysplastic syndrome (MDS) and leukemias are characterized by frequent epigenetic abnormalities and aberrant DNA hypermethylation has been linked to poor prognosis in cases of MDS, as it is associated with more rapid progression to acute myeloid leukemia (AML) (Shen et al., 2010; Shih et al., 2012). Importantly, genome-wide discovery efforts in patients with myeloid malignancies have led to the identification of a number of causal genetic abnormalities that impact the epigenetic landscape. These genes include *TET2*, *IDH1*, *IDH2*, *DNMT3a*, and *EZH2*, all of which affect DNA and/or histone lysine methylation (Shih et al., 2012).

Recent studies have examined in detail one such gene, *TET2* (ten-eleven-translocation gene 2), which is located in 4q24 and whose mutation or deletion is extremely frequent in hematological malignancies, affecting 19% of patients with MDS, 12% of patients with myeloproliferative neoplasm (MPN), and 24% of patients with secondary AML (Abdel-Wahab et al., 2009; Delhommeau et al., 2009; Jankowska et al., 2009; Tefferi et al., 2009). *TET2* mutations are observed in CD34⁺ progenitor cells and are detected most frequently after MPN and MDS progression to AML (Abdel-Wahab et al., 2010).

TET2 catalyzes the oxidation of 5-methylcytosine (5-mC) to 5-hydroxymethylcytosine (5-hmC) and is known to control hematopoiesis, presumably by regulating gene expression through its effect on DNA methylation (Branco et al., 2012; Ito et al., 2011; Ko et al., 2010; Shih et al., 2012). *Tet2* deficiency in mice impairs 5-mC hydroxylation and leads to skewed differentiation, as well as enhancement of the self-renewal and repopulating capacity of hematopoietic stem cells (HSCs). This in turn promotes malignant transformation in mice, causing

disorders resembling MDS and MPN, as well as chronic myelomonocytic leukemia (CMML) (Moran-Crusio et al., 2011; Quivoron et al., 2011). Recent studies have likewise shown that leukemia-associated Isocitrate Dehydrogenase (IDH) 1 or 2 mutant enzymes can induce carboxylation of glutamine-derived α -ketoglutarate (α -KG), with a concomitant increase in synthesis of 2-hydroxyglutarate (2-HG), an inhibitory metabolite of TET2 (Figueroa et al., 2010; Sasaki et al., 2012). Collectively, these studies implicate functional and mutational losses of TET2 as key events underlying aberrant hematopoiesis and leukemogenesis, in turn suggesting that the inactivation of TET2 function by additional upstream cues could play a critical role in hematological malignancies.

MicroRNA (miRNA) deregulation is also known to contribute to hematological malignancies, including MDS and AML (Bartel, 2004; Chen et al., 2004; Gangaraju and Lin, 2009). A combined approach involving assessment of miRNA expression (using microarrays) and bioinformatic prediction of mRNA targets has revealed that distinct miRNA signatures fine-tune each step of hematopoiesis, including the reconstitution potential of HSCs (Arnold et al., 2011; Gangaraju and Lin, 2009). In turn, miRNAs that target critical leukemia suppressors could act as powerful proto-oncogenes in the hematopoietic compartment (Calin and Croce, 2006).

In this study, we demonstrate that miR-22 controls hematopoiesis and HSC maintenance by negatively regulating TET2 protein levels. Importantly, we identify miR-22 as an epigenetic modifier and the key oncogenic determinant for the pathogenesis of MDS and hematological malignancies in vivo, findings with prognostic and critical therapeutic implications for a large number of hematological disorders.

RESULTS

miR-22 Is Highly Expressed in Human Hematological Malignancies and Increases the Replating Ability of HSCs

In order to build on the insights of miRNA biology into epigenetic alterations in myeloid malignancies, we attempted to identify those miRNAs playing a critical role in the pathogenesis of the stem cell disorder MDS and its progression to overt leukemia. We first analyzed a previously reported miRNA microarray profiling database and found that miR-22 is markedly upregulated in patients with MDS (Figures S1A and S1B available online; Dostalova Merkerova et al., 2011). These results prompted us to perform expression analysis by in situ hybridization in a large set of patient samples, which included bone marrow with no sign of disease ($n = 37$), early stage MDS (refractory anemia [RA], 5q syndrome [5q-], refractory cytopenia with multilineage dysplasia [RCMD], and RA with ringed sideroblasts [RARS]) ($n = 40$), and RA with excess blasts (RAEB) ($n = 67$). Our comprehensive in situ hybridization analysis revealed that miR-22 is highly expressed in patients with MDS (Figure 1A and Figure S1C). We next estimated the survival rates of the sets of MDS patients by using a Kaplan-Meier analysis and applied a log-rank test to compare the survival rates of MDS patients on the basis of miR-22 expression levels. Remarkably, we found that aberrant expression of miR-22 directly correlates with poor survival rates in patients (Figure 1B). It is noteworthy that

the distribution of blasts into each World Health Organization (WHO) category of MDS stages was not directly associated with the expression level of miR-22, but a significant correlation between miR-22 levels and poor survival rates was observed within each WHO classification, as well as among MDS patients harboring a normal karyotype (Figures S1D–S1F). This indicates that the poorer survival expectancy linked to miR-22 overexpression does not simply reflect confounding factors such as blast count and cytogenetic karyotype. Taken together, these findings led us to investigate the role of miR-22 in the hematopoietic system and the genesis of hematological malignancies.

To explore the function of miR-22 in hematopoiesis and leukemia, we first transduced mouse hematopoietic stem/progenitor cells (HSPCs; c-Kit^{pos}Sca-1^{pos}Lin^{neg}, hereafter referred to as KSL cells) isolated from wild-type donor mice with either a retroviral vector encoding miR-22 and a green fluorescent protein (GFP) or a control vector encoding GFP only. The GFP-expressing KSL cells were then resorted by flow cytometry and plated in methylcellulose for an in vitro colony-forming unit (CFU) assay (Figure S1G). We observed in the first plating that the cells expressing miR-22 produced fewer colonies compared to control cells; however, when GFP⁺KSL cells were resorted and replated, miR-22-transduced cells produced more colonies in the second plating (Figure S1H). Strikingly, these cells also maintained this ability throughout serial replatings: whereas control cells only formed colonies in the first two platings, miR-22-expressing KSL cells were able to produce colonies after being replated at least seven times (Figure S1H). Microscopic analysis of the cells from these colonies revealed a homogenous blast-like morphology (Figure S1I, top). Consistent with these results, miR-22-expressing KSL cells maintained their characteristics and c-Kit expression (a marker of undifferentiated cells) during their replatings (Figure S1I, bottom and Figures S1J and S1K). To ensure that miR-22-transduced and control cells were subjected to the same experimental treatment, we also performed an in vitro competitive colony assay by mixing GFP⁺ KSL cells expressing miR-22 and GFP⁻ control KSL cells (Figure S1L, top). This analysis confirmed that miR-22-expressing GFP⁺, but not GFP⁻, KSL cells remained mostly in colonies through consecutive replatings (Figure S1L, bottom). Taken together, these data suggest that miR-22 can drive aberrant replating capacity of HSPCs and may contribute to malignant transformation.

miR-22 Affects HSC Biology In Vivo

We next sought to examine the biological effects of miR-22 in vivo and generated a conditional transgenic mouse model overexpressing miR-22 in the hematopoietic compartment (Figure S2A). miR-22 expression was induced by *Mx1-Cre*-mediated excision of a *LoxP* flanked transcriptional STOP cassette by administration of seven doses of polyinosine-polycytidine (pIpC) over 14 days (Figures S2B and S2C). Two weeks after pIpC administration, we isolated KSL cells from *miR-22^{F/+};Mx1-Cre* mice and *Mx1-Cre* littermate controls and analyzed their properties by using CFU and long-term culture-initiating cell (LTC-IC) assays (Figures 1C–1G and Figure S2D). In agreement with our previous retroviral transduction experiments, we found that KSL cells from *miR-22^{F/+};Mx1-Cre* transgenic mice displayed an enhanced replating potential




Figure 1. miR-22 Is Upregulated in MDS Patients and Leads to Increased Replating Capacity In Vitro

(A) miR-22 is highly expressed in MDS patient samples. Expression levels of miR-22 in blasts in subtypes of MDS were measured by in situ hybridization analysis. Normal (n = 37), RA (n = 30), 5q- (n = 5), RCMD (n = 4), RARS (n = 1), RAEB-1 (n = 32), and RAEB-2 (n = 35) are shown. Representative images of in situ hybridization for miR-22 in MDS patients and normal bone marrow are also shown (left). Insets of panels show miR-22 staining in blasts (arrow with B) and in differentiated myeloid cells (e.g., neutrophil, arrow with M) with a high magnification. The expression levels of miR-22 were also scored as described in the Supplemental Experimental Procedures (right). The data were analyzed by Chi-square test. *r*, Pearson's *r*. Scale bars, 60 μ m.

(B) miR-22 overexpression correlates with poor survival rates of human MDS patients. MDS patients were divided into two groups; low-miR-22-expressing patients' group (miR-22 Score 1 or 2, n = 49, blue) and high-miR-22-expressing patients' group (miR-22 Score 3, n = 58, red). Overall survival of these patients is demonstrated. *p* value was generated by a log-rank test.

(C) Overview of the experimental design for the conditional miR-22 expression in the hematopoietic compartment. Two-month-old miR-22^{F/+};Mx1-Cre mice or Mx1-Cre littermate controls were treated with plpC for two weeks. Two weeks after plpC administration, c-Kit^{pos} Sca-1^{pos}Lin^{neg} (KSL) cells were sorted to evaluate the characteristics of miR-22-expressing hematopoietic stem/progenitor cells. LTC-IC, long-term culture -initiating cells; BMT, bone marrow transplantation.

(D) Colony-forming capacity of miR-22-expressing progenitor cells. Two weeks after plpC administration, sorted KSL cells from miR-22^{F/+};Mx1-Cre mice or littermate controls were cultured in semisolid medium. Counting and classification of colonies were performed in independent littermate pairs (n = 3). GEMM, Colony-forming unit-granulocyte, erythroid, macrophage, and

megakaryocyte; GM, Colony-forming unit-granulocyte and macrophage; M, Colony-forming unit-macrophage; E, Burst-forming unit-erythroid.

(E) miR-22 expression retains the ability to serially replating and generate colonies. In vitro colony replating assay was performed as shown in Figure S2D and colony counts in the indicated replatings were scored (n = 3). N.D., nondetectable.

(F) Ectopic expression of miR-22 results in an increase of HSPCs in in vitro long-term culture with stromal cells. KSL cells isolated as described in (C) were cocultured with OP-9 stromal cells for 2 weeks. Maintained Lin^{neg} cells and KSL cells were evaluated (n = 3). Representative flow cytometry data of Lin^{neg} and the KSL cells (left) are shown. Mean percentage \pm SD of KSL cells in Lin^{neg} cells (middle) and mean absolute numbers \pm SD of CD45^{pos}KSL cells (right) are also shown.

(G) miR-22 increases the myeloid compartment. The percentages of CD45^{pos} cells (left), CD11b^{pos} cells (middle), or Gr-1^{pos} cells (right) in total mononuclear cells (MNCs) were investigated 2 weeks after the coculture with stromal cells from (F) (n = 3).

All error bars represent \pm SD. See also Figures S1, S2A–S2D, and S7.

(Figures 1D and 1E). Furthermore, ectopic expression of miR-22 caused a robust increase in the number of both KSL and mature myeloid cells, as determined by LTC-IC assay, indicating that miR-22 enhances HSPC maintenance and predisposes those cells to differentiate into the myeloid lineage (Figures 1F and 1G).

Next, we examined the ability of miR-22-overexpressing HSPCs to compete with wild-type counterparts in vivo by utilizing an in vivo competitive transplantation assay. KSL cells from miR-22^{F/+};Mx1-Cre and Mx1-Cre mice (Ly45.2) were purified 2 weeks after plpC administration, mixed with Ly45.1/Ly45.2

competitor bone marrow mononuclear cells (BM MNCs), and transplanted into lethally irradiated Ly45.1 congenic recipient mice. The ability of donor cells to contribute to the reconstitution was then determined by flow cytometric analysis of peripheral blood stained for Ly45.1 and Ly45.2. Within 3 weeks of transplantation, donor cells from miR-22^{F/+};Mx1-Cre mice exhibited a slight increase in chimerism compared to those from control animals (Figure S2E). Over time this difference became more pronounced as miR-22^{F/+};Mx1-Cre KSL cells outcompeted their wild-type counterparts (Figures 2A–2C and Figure S2F). At




Figure 2. miR-22 Leads to an Enhanced Repopulating Capacity of HSPCs In Vivo

(A and B) Ly45.1 recipient mice were transplanted with 1,500 KSL cells from *miR-22^{F/+};Mx1-Cre* mice (n = 15) or littermate controls (n = 10) after plpC administration together with 4 × 10⁵ Ly45.1/Ly45.2 competitor BM MNCs. Donor-derived chimerism in peripheral blood was analyzed 6 weeks after the transplantation. Representative flow data of the CD45.1/CD45.2 positivity (A) and the percentages of both donor-derived cells in myeloid (CD11b^{pos} and/or Gr-1^{pos}) (B, left) and donor-derived myeloid cells in MNCs (B, right) of recipient mice are shown.

(C–E) Donor contribution in hematopoiesis of recipient mice 9 weeks after the transplantation. Representative flow data of the CD45.1/CD45.2 positivity with the percentages of donor-derived CD45.1^{neg}CD45.2^{pos} cells (C, left) are shown. Mean percentages ± SD of donor-derived cells in MNCs (C, right), myeloid cells (D, left), B cells (E, left), and T cells (E, right) are shown (n = 10 for littermate controls and n = 15 for *miR-22^{F/+};Mx1-Cre*).

(F) c-Kit^{pos} cells are observed in peripheral blood of recipient mice transplanted with KSL cells from *miR-22^{F/+};Mx1-Cre* mice. Representative flow cytometry data of c-Kit positivity in donor-derived cells (left) and the percentages of c-Kit^{pos} cells in donor-derived cells (right) are shown. (n = 10 for littermate controls and n = 15 for *miR-22^{F/+};Mx1-Cre*).

All error bars represent ± SD. See also Figures S2E, S2F, and S7.

6 weeks posttransplantation, an increased chimerism was observed in all BM MNCs as well as within the myeloid fraction, and by 9 weeks this effect was also evident in the c-Kit^{pos} fraction (Figures 2A–2F). Taken together, these data strongly suggest that miR-22 enhances the self-renewal capacity of HSPCs in vivo.

miR-22 Triggers MDS-like Syndromes and Hematological Malignancies In Vivo

To further test whether increased aberrant hematopoiesis caused by miR-22 overexpression ultimately progresses to malignant hematological disease, we continued to follow up primarily transplanted recipients. Notably, at 12 weeks posttransplantation, recipient mice transplanted with KSL cells from *miR-22^{F/+};Mx1-Cre* donors, but not from *Mx1-Cre* donors, developed an illness that closely resembled human MDS, as characterized by defective erythroid maturation and anemia, splenomegaly with myeloid infiltration, low white blood cell (WBC) counts, and dysplastic myeloid cells in the peripheral

circulation (Figure 3 and Figures S2G and S2H). To determine whether this disorder is transplantable, the donor-derived KSL cells resorted from primary recipient mice were subjected to a secondary transplantation using fresh competitor cells. In agreement with our earlier results, all secondary recipients developed a lethal hematological disease within 4 weeks after secondary transplantation (Figure S2I).

Furthermore, we also monitored disease onset and progression to malignancy in miR-22 transgenic mice. The development of disease similar to that found in our chimeric transplantation model was observed in *miR-22^{F/+};Mx1-Cre* mice but not *Mx1-Cre* littermate controls at 8 months after plpC administration: the hallmarks of this distress included dysplasia in many lineages including erythroid, myeloid, and platelet; splenomegaly; and an increased number of c-Kit^{pos} undifferentiated blastic cells in the peripheral blood (Figures 4A–4C). By 16 months after plpC administration, all miR-22 transgenic mice had developed MDS, affecting multilineage hematopoiesis. Importantly, ~13% of mice succumbed to myeloid leukemia as well as T cell lymphoma by 11 months, and all miR-22 transgenic mice suffered from hematological abnormality by 16 months. Finally, ~70% of miR-22 transgenic mice developed myeloid leukemia within 2 years (Figures 4D–4G).




Figure 3. miR-22 Overexpression Develops Hematological Syndromes

(A and B) miR-22 overexpression leads to human MDS-like phenotypes characterized by defective erythropoiesis (A) and cytopenia (B) in miR-22 transgenic mice. After plpC administration, 1,500 KSL cells from *miR-22^{F/+};Mx1-Cre* mice or littermate controls were transplanted into lethally irradiated Ly45.1 recipient mice with 4×10^5 Ly45.1/Ly45.2 competitor BM MNCs ($n = 10$ for littermate controls and $n = 15$ for *miR-22^{F/+};Mx1-Cre*). Twelve weeks after transplantation, peripheral blood of recipient mice was evaluated. Representative flow data of the Ter119/CD71 positivity (A, left) and the percentages of R1 (Ter119^{pos}CD71^{pos}) (A, middle) and R2 (Ter119^{pos}CD71^{pos}) (A, right) are shown. White blood cell (WBC) (B, left) and platelet (Plt) counts (B, right) are also shown. (C and D) Splenomegaly and myeloid infiltration into spleen were observed in recipient mice transplanted with KSL cells from *miR-22^{F/+};Mx1-Cre* mice. Representative images of spleens (C, left), spleen weight ($n = 3$) (C, right), and representative flow data of the positivity of c-Kit and CD11b and/or Gr-1 in spleen (D) are shown. (E) miR-22 expression leads to a differentiation defect in erythroid compartment. Representative flow data of the Ter119/CD71 positivity in spleen are shown. All error bars indicate \pm SD. See also Figures S2G–S2L.

We also confirmed that the disease was triggered by miR-22 from KSL cells sorted from miR-22-expressing colonies. To this end, miR-22-expressing cells were subjected to consecutive colony replating and at the sixth replating, the residual c-Kit^{pos}Lin^{neg} (KL) cells were transplanted into lethally irradiated recipient mice (Figure S3A). Consistent with our in vivo miR-22 transgenic mice data, by 14 weeks after transplantation, recipient mice transplanted with miR-22-expressing cells developed myeloid leukemia, as characterized by increased myeloid blasts in the peripheral blood, defective erythroid maturation and anemia, splenomegaly with the infiltration of myeloid blasts, and increased c-Kit^{pos} blasts in bone marrow (Figures S3B–S3H). Taken together, these data strongly suggest that miR-22 can enhance HSPC function and contribute to leukemia development in vivo.

miR-22 Directly Targets TET2 and Affects the Epigenetic Landscape of the Hematopoietic Compartment In Vivo

To identify relevant molecular targets of miR-22, we next undertook a bioinformatic analysis using TargetScan 6.0 (<http://targetscan.org>) (Lewis et al., 2005), microRNA.org (<http://www.microrna.org/microrna>) (Betel et al., 2008), and miRBase (<http://www.mirbase.org>) (Griffiths-Jones et al., 2006). Among various potential targets, we focused on ten eleven translocation 2 (TET2) as it possesses critically conserved nucleotides indicative of a legitimate target (Figure S4A). We then confirmed a direct interaction of miR-22 with the 3'UTR

region of the *TET2* gene by luciferase reporter assay (Figure S4B and Song et al., 2013). TET2 is a member of the TET methylcytosine dioxygenase family, which catalyzes the conversion of 5-methylcytosine (5-mC) to 5-hydroxymethylcytosine (5-hmC). TET proteins have been implicated in epigenetic reprogramming, embryonic stem cell maintenance, and early development (Ficz et al., 2011; Ito et al., 2010). Importantly, recurrent *TET2* mutations have been identified in a variety of hematological malignancies including MDS, MPN, CMMoL, and AML (Abdel-Wahab et al., 2009; Delhommeau et al., 2009; Jankowska et al., 2009; Tefferi et al., 2009). It is noteworthy that the phenotypes elicited by miR-22 overexpression closely phenocopy the inactivation of *Tet2* in the hematopoietic system both in vitro and in vivo (Li et al., 2011; Moran-Crusio et al., 2011; Quivoron et al., 2011). It has also been documented that *Tet2* expression is increased during hematopoietic differentiation, and conversely, that miR-22 is highly expressed in HSPCs while its expression declines in mature cells (Figure S4C, Moran-Crusio et al., 2011). We therefore tested whether miR-22 could act as an authentic *TET2*-targeting miRNA in the hematopoietic compartment by utilizing our conditional transgenic mouse model. To this end, we measured *Tet2* expression in the peripheral blood and bone marrow of *miR-22^{F/+};Mx1-Cre* or *Mx1-Cre* littermate mice at 2–3 weeks after plpC administration. Importantly, real-time qPCR and western blot analyses revealed a marked reduction in levels of *Tet2* mRNA and protein, respectively, in *miR-22^{F/+};Mx1-Cre* mice compared to littermate controls (Figures 5A, 5B, 5E–5G and Figure S4D).




Figure 4. miR-22 Transgenic Mice Develop Primary Hematological Diseases

(A and B) miR-22 transgenic mice develop MDS-like hematological syndromes. Representative smears of peripheral blood of 8-month-old mice after plpC administration are shown (A). Representative images of dysplastic erythroid cells (poikilocytosis, A; polychromasia, B-i, arrowhead), dysplastic platelets (giant platelet, B-ii, arrow, B-vii), dysplastic neutrophils (hypersegmented neutrophils, B-ii-iv, arrows; a pseudo-Pelger-Huet anomaly, B-v and vi) and dysplastic blasts (B-viii arrows) in miR-22 transgenic mice are also shown. Scale bars, 50 μ m (A) and 10 μ m (B).

(C) c-Kit^{pos} immature blasts are increased in miR-22 transgenic mice. Representative flow cytometry data (left) and mean percentages \pm SD of c-Kit^{pos} cells in Lin^{neg} compartment (right) are shown (n = 3).

(D) Disease-free survival of miR-22^{F/+};Mx1-Cre mice (n = 26) and littermate controls (n = 13).

(legend continued on next page)

On the basis of these observations, we then sought to determine whether miR-22-mediated TET2 inactivation is sufficient to impact the global epigenetic landscape in blood cells. We therefore analyzed 5-hmC and 5-mC levels in the genomic DNA of *miR-22^{F/+};Mx1-Cre* mice. Immunofluorescence and dot blot analyses revealed a drastic (more than 2-fold) reduction in 5-hmC levels and a concomitant increase in 5-mC levels in bone marrow of these mice (Figures 5C and 5D), suggesting that changes in an important epigenetic mark (e.g., 5-hmC) may contribute to the ability of miR-22 to affect hematopoiesis and leukemogenesis.

To further understand the consequences of repression of TET2 by miR-22 in hematopoiesis, we next examined the effects of miR-22 overexpression on putative targets of TET2 (Kallin et al., 2012; Ko et al., 2010; Yamazaki et al., 2012). A comprehensive real-time qPCR analysis revealed that the expression levels of a substantial number of putative Tet2 target genes, including *Aim2*, *Hal*, *Igfb2*, and *Sp140*, were markedly decreased in the HSC compartment (CD150^{pos}CD48^{neg}Flt3^{neg}CD34^{neg}KSL cells) purified from *miR-22^{F/+};Mx1-Cre* mice (Figure 5E and Figure S4E). Interestingly, absence in myeloma 2 (AIM2) has been reported to have a role in the reduction of cell proliferation by cell cycle arrest and to also be highly methylated in MDS, especially in CMMoL patients with *TET2* mutations (Figuerola et al., 2010; Patsos et al., 2010; Yamazaki et al., 2012). Therefore we hypothesized that its silencing via methylation of the promoter might provide a growth advantage to cancer cells. Additionally, polymorphisms of *SP140* (SP140 nuclear body protein) have been found to correlate with chronic lymphoid leukemia, and its hypermethylation has been also demonstrated in CMMoL patients with *TET2* mutations, and hence its hypermethylation may influence the risk of leukemogenesis (Di Bernardo et al., 2008; Ko et al., 2010; Yamazaki et al., 2012). Indeed, we confirmed that the expression of *Aim2* and *Sp140* was drastically reduced at the levels of mRNA and protein in the bone marrow of *miR-22^{F/+};Mx1-Cre* mice compared to littermate controls, and this reduction was associated with an increase in promoter methylation (Figures 5F and 5G and Figure S4F). Taken together, these results suggest that miR-22 represses TET2 expression and thereby can remodel the epigenetic landscape with global changes in 5-hmC levels and alteration in the expression of putative TET2 target genes including *AIM2* and *SP140*.

miR-22 Enhances Stem Cell Function and Triggers Hematological Transformation through Repression of TET2

Next, we explored the relevance of the crosstalk between miR-22 and TET2 in hematopoiesis and hematological malignancies. To test whether TET2 directly contributes to the function of miR-22 in the hematopoietic compartment, we first infected KSL cells isolated from wild-type mice with a retroviral vector encoding miR-22 and GFP. After the resorting, GFP⁺KSL cells were additionally infected with a *TET2*-expressing lentiviral

particle. In vitro colony replating and LTC-IC assays demonstrated that the phenotypes caused by miR-22 overexpression in hematopoietic compartment are potently suppressed by ectopic expression of TET2 (Figures S4G–S4J). Notably, an in vivo transplantation assay revealed that ectopic expression of TET2 leads to survival advantages for recipient mice transplanted with miR-22-expressing KSL cells (Figure S4K).

To further examine this interaction, we purified KSL cells from miR-22 transgenic mice and infected them with a retroviral vector encoding Cre recombinase and GFP. After the resorting of GFP⁺KSL cells, cells were additionally infected with a TET2-expressing lentiviral particle and the resulting phenotypes were analyzed. In vitro colony replating and LTC-IC analyses demonstrated that the oncogenic function of miR-22 was antagonized by ectopic expression of TET2 (Figures 6A and 6B). Notably, when KSL cells of *miR-22^{F/+}* mice were simultaneously coinfecting with a Cre recombinase construct and a TET2-expressing construct, ectopic expression of TET2 more strikingly impaired the replating potential of GFP⁺KSL cells (Figures S5A and S5B). In line with these observations, an in vivo transplantation assay revealed significant survival advantages in recipient mice transplanted with *miR-22^{F/+}* KSL cells coinfecting with Cre and TET2 compared to those with *miR-22^{F/+}* KSL cells coinfecting with Cre and control vector (Figure 6C). Importantly, the positive survival effect drawn by ectopic expression of TET2 was more pronounced in secondary transplantation (Figure S5C).

Finally, we purified KSL cells from miR-22 transgenic mice after plpC administration, infected them with a TET2-expressing lentiviral particle, and evaluated the resulting phenotypes (Figure 6D). We found that all the phenotypes elicited by an elevation of miR-22 in vitro and in vivo in hematopoietic compartment, including repression of the expression of *Aim2* and *Sp140* in bone marrow, were also significantly suppressed by ectopic expression of TET2 (Figures 6E–6H and Figure S5D).

Importantly, when we purified KSL cells from primary miR-22 transgenic mice and infected them with TET2 followed by in vivo transplantation together with competitor cells, ectopic expression of TET2 delayed the onset of hematological disorder and was accompanied by significant advantages in disease-free survival (Figure S5E). Taken together, these data suggest that TET2 is one of the major targets responsible for the miR-22 proto-oncogenic function in hematopoiesis and the development of hematological malignancies.

miR-22 Decoying Offers Therapeutic Opportunities and Its Overexpression Correlates with the Silencing of TET2 Protein in MDS and AML Patients

Next, we investigated whether the miR-22/TET2 interaction may offer a therapeutic opportunity for the treatment of hematological malignancies. We therefore tested whether inhibition of miR-22 affects TET2 expression and activity, as well as the biology of

(E and F) Representative lethal hematological syndromes observed in miR-22 transgenic mice. Representative images of spleens (E, left) and H&E staining (E, right) are shown. Scale bars, 100 μ m. Representative smears of peripheral blood of miR-22 transgenic mice (6 months old) with increased myeloid blasts (F) are also shown. Scale bars, 20 μ m.

(G) Pie charts representing the disease spectrum in *miR-22^{F/+};Mx1-Cre* mice at the indicated ages (n = 26). MPN, myeloproliferative neoplasm. All error bars indicate \pm SD. See also Figure S3.




Figure 5. miR-22 Acts as an Epigenetic Modifier by Directly Targeting TET2

(A and B) A reduction in the levels of *Tet2* mRNA in miR-22 transgenic mice. Expression levels of miR-22 (left) and *Tet2* mRNA (right) in mononuclear cells from peripheral blood (A) and bone marrow (BM) (B) of *Mx1-Cre* control or *miR-22^{F/+};Mx1-Cre* mice 2–3 weeks after plpC administration were determined by real-time qPCR (n = 6).

(C) miR-22 expression results in a significant reduction in the levels of 5-hmC in the hematopoietic compartment. 5-hmC and 5-mC were evaluated by immunofluorescence analysis in BM cells isolated from *miR-22^{F/+};Mx1-Cre* mice and littermate controls 3 weeks after plpC administration (left). Cells expressing high levels of 5-hmC show low levels of 5-mC (arrow) and conversely, cells expressing low levels of 5-hmC exhibit high levels of 5-mC (arrowhead). Bar graph depicts mean percentages \pm SD of cells expressing high levels of 5-hmC (n = 3) (right). Scale bars, 10 μ m.

(legend continued on next page)

leukemias. To this end, we inhibited miR-22 in human leukemia cell lines, K562 and U937, both of which express relatively high levels of miR-22 (Figure S5F, Wang et al., 2012), by using a miR-22 retroviral sponge or a custom-made locked nucleic acid (LNA; Petersen and Wengel, 2003) miR-22 decoy as the miR-22 sponge. Critically, inhibition of miR-22 in human leukemic cells resulted in a significant reduction in cell proliferation and an elevation of the expression of TET2 and its targets, AIM2 and SP140 (Figures 6I and 6J and Figures S5G–S5I). It is worth noting that the U937 cell line harbors phosphatase and tensin homolog (*PTEN*) mutant with premature termination of coding sequence (Liu et al., 2000). Hence in this setting the effects observed upon miR-22 decoying are not mediated by the ability of miR-22 to downregulate *PTEN* expression.

In full agreement with these observations, an *in vitro* colony replating assay revealed that the inhibition of miR-22 by a miR-22 retroviral sponge in KSL cells purified from miR-22 transgenic mice led to a reduction in their colony-forming capability (Figure S5J). Taken together, these data provide a rationale for the therapeutic potential of targeting miR-22 with LNA miR-22 inhibitors in the treatment of hematological malignancies.

The pathological roles of TET2 in myeloid malignancies have been previously highlighted as its mutations are widely observed in myeloid malignancies and MDS patients. Most of these earlier studies, however, were limited to mutation analysis and found little relationship between the progression of MDS to AML and TET2 expression levels, although the phenotypes of *Tet2* knockout mice gave indications to the contrary (Moran-Crusio et al., 2011; Quivoron et al., 2011). Likewise, emerging evidence demonstrated no significant prognostic association between TET2 mutations in MDS patients and WHO subtypes, the International Prognostic Scoring System (IPSS) score, cytogenetic status, or transformation to AML (Kosmider et al., 2009; Smith et al., 2010).

However, on the basis of our data we hypothesized that a miR-22-dependent reduction in the levels of TET2 might have been overlooked and could play a key role in the development of myeloid malignancies. To test this hypothesis, we investigated a large cohort of patients with MDS by an immunohistochemical analysis. Strikingly, this analysis revealed that ~60% of 107 MDS patients analyzed displayed a reduction in the levels of TET2 (36% of 107 MDS patients show Score 1 and 24% show Score 2) and that this reduction was directly correlated with poor survival rates of patients (Figures 7A and 7B and Figures S6A–S6D). Importantly, the percentage of blasts distributed in each WHO category of MDS stage was not directly associated with expression level of TET2, but a significant correlation was found between TET2 expression and poor survival rates within each WHO classification, as well as among MDS patients harboring a normal karyotype, suggesting that the poorer survival expect-

tancy linked to TET2 downregulation does not simply reflect confounding factors, such as blast count and cytogenetic karyotype (Figure 7C and Figures S6E and S6F). Together, these results suggest that the contribution of TET2 to the pathogenesis of hematological malignancies extends far beyond the initial impact of its mutations.

Importantly, combined *in situ* hybridization and immunohistochemical analyses revealed that miR-22 expression was directly anticorrelated with the levels of TET2 in a large-cohort data set of MDS patients ($n = 107$) (Pearson's $r = -0.47$, $p = 2.4 \times 10^{-7}$): 28% of MDS patients that showed the reduced levels of TET2 (Score 1) also exhibited high miR-22 levels (Score 3) (Figure 7D and Figure S6G and see also Figure 1A).

Furthermore, because our *in vivo* analysis demonstrated that miR-22 triggers overt leukemia (Figure 4G), we investigated the miR-22/TET2 interactive relationship in AML with multiple lineage dysplasia (MLD) ($n = 18$). Our combined analyses of *in situ* hybridization and immunohistochemistry revealed a clear anticorrelation between miR-22 expression and the levels of TET2 (Pearson's $r = -0.75$, $p = 2.9 \times 10^{-4}$); in addition, 22.2% of AML with MLD patients showing the reduced levels of TET2 (Score 1) displayed a high expression of miR-22 (Score 3) (Figure 7E). Western blot and real-time qPCR analyses also corroborated a relationship between miR-22 and TET2 in primary AML patients (Figure S6H). Finally, we confirmed these findings by analyzing the expression levels of miR-22 in a very large cohort of AML patients from a previously reported real-time qPCR database (Jongen-Lavrencic et al., 2008). Here miR-22 was found highly expressed in 58.1% of 214 AML patient samples compared to normal bone marrow from healthy donors ($p = 0.028$, Figure 7F). Taken together, these data strongly suggest that aberration of the miR-22-TET2 regulatory network is one of the common events in hematopoietic malignancies.

DISCUSSION

The findings presented in this study point to a number of important conclusions.

The first of these is that miR-22 is a powerful regulator of HSPC maintenance and self-renewal. These effects are so pronounced that miR-22-expressing HSPCs out-competed their wild-type counterparts in all the assays that we performed *in vivo* as well as *in vitro*. Importantly, in so doing miR-22 also skews hematopoietic differentiation, mainly toward myeloid compartment.

We have also identified miR-22 as a key determinant of the pathogenesis of both MDS and hematological malignancies. Remarkably, miR-22 plays a role in the initiation of MDS as well as full-blown leukemia; hence it exerts a bona fide proto-oncogenic activity. MDS is a genetically heterogeneous disease and we are only now beginning to fully appreciate its diversity.

(D) miR-22 expression leads to a significant reduction in global 5-hmC expression levels in the genome in BM cells. 5-hmC levels were analyzed by quantitative dot blot assay with genomic DNA purified from BM cells of *miR-22^{F/+};Mx1-Cre* or littermate control mice.

(E) Plots depict relative expression levels of putative *Tet2* target genes (*Aim2*, *Sp140*, *Igfbt2*, and *Hal*) in CD150^{pos}CD48^{neg}Flt-3^{neg}CD34^{neg}KSL cells of *miR-22^{F/+};Mx1-Cre* mice and littermate controls ~2–3 weeks after plpC administration ($n = 4$). *Ly6c1* and *Actb* were used as a *Tet2* unrelated gene control and an internal control, respectively.

(F and G) Repression of putative downstream genes of *Tet2* in miR-22 transgenic mice. Expression levels of *Tet2*, *Sp140*, and *Aim2* mRNAs (F) or proteins (G) in BM of *miR-22^{F/+};Mx1-Cre* mice or littermate controls 2–3 weeks after plpC administration were evaluated ($n = 3$).

All error bars indicate \pm SD. See also Figures S4A–S4F.




Figure 6. miR-22/TET2 Regulatory Network Affects HSC Function and Hematological Transformation

(A and B) Ectopic expression of TET2 reduces the colony-forming capacity of miR-22-expressing HSPCs. The sorted KSL cells from miR-22^{F/+} mice were infected with Cre-GFP or GFP control vector. GFP⁺KSL cells were resorted and infected with empty vector or TET2-expressing lentiviral particles. Cells were then incubated in semisolid medium. Colony replating assay was performed and the resulting colonies were counted at the indicated replatings (n = 3) (A). Some of TET2 infected cells were also subjected to LTC-IC assay. Representative flow cytometry data of the positivity of c-Kit/Sca-1 in Lin^{neg}CD45^{pos}-gated cells 2 weeks after coculture with stromal cells and mean percentages ± SD of KSL cells in Lin^{neg}CD45^{pos} (in brackets) are shown (n = 3) (B, left). Colony-forming capacities were also determined at the indicated weeks after coculture with stromal cells (n = 3) (B, right).

(legend continued on next page)

This study will therefore help in the reclassification of subtypes of MDS, and possibly AML, on the basis of miR-22 expression.

Furthermore, our comprehensive analysis of MDS patients has identified TET2 protein downregulation as a common event in hemopoietic malignancy (35.5% of total MDS patient samples, and 22.2% of AML samples with MLD, with most of those patients exhibiting high expression of miR-22) and a factor in poor prognosis, findings which directly impact the management and therapeutic decisions for patients with myeloid malignancies. Our findings thus imply that the contribution of TET2 to the pathogenesis of myeloid malignancies could be related to more than its mutations and identify the deregulation of the miR-22-TET2 pathway as one of the most frequent events in hematological malignancies. Importantly, they also provide a compelling rationale for targeted therapies that impact the mechanisms of regulation responsible for the control of TET2 levels (e.g., through miR-22 decoying).

We show that miR-22 negatively regulates the expression of TET2 and that its overexpression closely phenocopies, both in vitro and in vivo, many of the characteristics observed upon inactivation of TET2 (Figure 7G). Extensive mouse modeling efforts support these similarities both in our current study and in other Tet2-related studies (Li et al., 2011; Moran-Crusio et al., 2011; Quivoron et al., 2011). Furthermore, our comprehensive analyses of a large cohort of MDS and AML patients likewise have demonstrated this antagonistic crosstalk.

miR-22 also appears to exert a tumor-promoting role in other tissues, including mammary glands, through its ability to suppress the expression of other TET family members such as TET1 and TET3 (Figure S4D and Song et al., 2013). Thus, miR-22 contributes to inactivation of TET2 and other TET family members in tumorigenesis, representing a tumorigenic pathway in addition to the more familiar ones of TET2 and TET family mutations and deletions.

We previously demonstrated that miR-22 also targets the PTEN tumor suppressor (Poliseno et al., 2010). *Pten* loss has also been implicated in the pathogenesis of hematological malignancies (Yilmaz et al., 2006; Zhang et al., 2006). However, miR-22 through TET2 regulation can bestow to hemopoietic cells

further competitive advantage as also supported by the fact that miR-22 decoying suppresses proliferation even in a *PTEN* mutant setting (Figure S5I). This may in turn explain the potent proto-oncogenic activity of miR-22. Indeed, there are several intriguing phenotypical differences between *Pten*^{F/F}; *Mx1-Cre* and *miR-22*^{F/+}; *Mx1-Cre* mouse models that could illuminate a unique and powerful role exerted by miR-22 in leukemogenesis. First, miR-22 increased the ability of HSCs to repopulate, which was then followed by leukemia development, whereas loss of *Pten* resulted in the exhaustion of normal HSCs. In addition, while *Pten* loss induces short-term expansion in HSCs through excessive cell cycle entry followed by exhaustion, no significant alteration in the stem cell compartment was observed at 2–3 weeks after the introduction of miR-22 (Figure S7). Additionally, while loss of *Pten* in HSCs is known to lead to MPN-like disease followed by leukemia development, hematological dysplasia with ineffective hematopoiesis is one of the main phenotypes caused by miR-22 expression, with bicytopenia or pancytopenia observed in miR-22 transgenic mice. Furthermore, although HSCs from *Pten*^{F/+}; *Mx1-Cre* mice are known to be transplantable, the recipients do not develop hematopoietic syndromes (Yilmaz et al., 2006). In contrast, recipient mice transplanted with HSCs purified from *miR-22*^{F/+}; *Mx1-Cre* mice developed aberrant hematopoiesis followed by MDS and leukemia. Hence, the ability of miR-22 to concomitantly downregulate TET2 and PTEN provides it a unique mode of action and confers on it the ability to overcome *PTEN*-loss-driven stem cell exhaustion while favoring MDS development and leukemogenesis. This in turn raises the possibility that concomitant silencing of TET2 and PTEN by miR-22 could represent a powerful event underlying the pathogenesis of human cancer, as it may overcome the fail-safe mechanisms elicited by loss of *PTEN* (Berger et al., 2011).

Finally, since the technology to inhibit miRNAs for therapy is rapidly evolving (Elmén et al., 2008; Stenvang et al., 2008), this will impact treatment options for a range of diseases and disorders in the years to come. Indeed, LNA-based miRNA decoying has already come of age for the treatment of hepatitis C (Lanford et al., 2010) and may prove very effective for the treatment of

(C) TET2 attenuates the hematopoietic malignancies induced by miR-22 overexpression. KSL cells sorted from *miR-22*^{F/+} mice were infected with Cre-GFP or GFP control vector before TET2 infection, and 1,500 GFP⁺KSL cells were transplanted into recipient mice with 2.0×10^5 Ly45.1 competitor bone marrow mononuclear cells (BM MNCs). Disease-free survival of recipient mice was examined by Kaplan-Meier survival curves. Log-rank test was used to generate p values.

(D) Overview of the experimental design for introduction of TET2 into KSL cells from miR-22 transgenic mice. KSL cells purified from miR-22 transgenic mice were subjected to infection with TET2-expressing vector, and their characteristics were evaluated in vivo and in vitro.

(E) Colony-forming capacity of miR-22 transgenic progenitor cells is attenuated by ectopic expression of TET2. Two weeks after plpC administration, sorted KSL cells from *miR-22*^{F/+}; *Mx1-Cre* mice were infected with TET2-expressing vector and resorted KSL cells were then subjected to incubation in semisolid medium. The resulting colony numbers were scored in three independent littermate pairs in the indicated platings (n = 3). N.D., nondetectable.

(F and G) Ectopic expression of TET2 causes a reduction in LTC-IC capacity of miR-22 transgenic progenitor cells. TET2-infected KSL cells from *miR-22*^{F/+}; *Mx1-Cre* mice or littermate controls, as shown in Figure 6D, were cocultured with stromal cells for the indicated weeks. Maintained KSL cells were evaluated 2 weeks after coculture (n = 3) (F). At the indicated times after coculture, the capacity of colony formation was also determined in semisolid medium (n = 3) (G).

(H) Increased reconstitution capacity of KSL cells of miR-22 transgenic mice is reduced by ectopic expression of TET2. The sorted KSL cells from *miR-22*^{F/+}; *Mx1-Cre* mice or littermate controls were infected with TET2-expressing vector. 1×10^4 KSL cells were resorted and subjected to bone marrow transplantation with 5×10^5 competitor BM MNCs. Representative flow cytometry data of the CD45.1/CD45.2 positivity at 6 weeks after transplantation are shown.

(I) Inhibition of miR-22 increases the expression of TET2 and its putative target genes in human leukemic cells. K562 cells were infected with the vector encoding miR-22 sponge or control sponge, and 48 hr after the infection, cell lysates were subjected to immunoblot analysis for the indicated proteins.

(J) Inhibition of miR-22 suppresses the proliferation of human leukemic cells. K562 cells were infected with the vector encoding miR-22 sponge or control sponge. Forty-eight hours after puromycin selection, 5×10^4 cells were incubated for seven days. Optical densities of the cells were determined at the indicated times (n = 3).

All error bars indicate \pm SD. See also Figures S4G–S4K and S5.




Figure 7. miR-22 Overexpression Directly Correlates with the Silencing of TET2 in Human MDS and AML Patients

(A) TET2 expression is downregulated in human MDS patients. Expression levels of TET2 protein in CD34^{pos} blasts in subtypes of MDS were evaluated by immunohistochemical analysis. Representative images of immunohistochemical analysis for TET2 protein in patients with subtypes of MDS are shown (left). Insets represent TET2 staining in blasts (arrow with B) with a high magnification. Scoring the expression levels of TET2 in blasts was performed as described in Supplemental Experimental Procedures (right). The data were analyzed by Chi-square test. *r*, Pearson's *r*. Scale bars, 60 μ m.

(B) Downregulation of TET2 levels correlates with poor survival rates of human MDS patients. Overall survival of patients that express low TET2 (TET2 Score 1, n = 38, red) or high TET2 (TET2 Score 2 or 3, n = 69, blue) is shown. *p* value was generated by a log-rank test.

(C) Survival rates of low TET2 (Score 1) and high TET2 (Score 2 or 3) patients with RA (left), RAEB-1 (middle), and RAEB-2 (right). Survival rate of the *p* value was generated by a log-rank test.

(legend continued on next page)

blood-borne diseases (Zhang et al., 2012). The data presented in this study thus support the notion that an LNA-based therapeutic targeting of miR-22 may represent an effective strategy for TET2 reactivation as a treatment modality for incurable diseases such as MDS and leukemia.

EXPERIMENTAL PROCEDURES

Mice

The details of generation of miR-22 transgenic mice are described in the [Supplemental Experimental Procedures](#). C57BL/6 mice (B6-CD45.2) and C57BL/6 mice congenic for the CD45 locus (B6-CD45.1) were purchased from The Jackson Laboratory and used as recipients in the transplantation assay.

Real-Time qPCR

The details of RNA isolation and real-time qPCR analysis are described in the [Supplemental Experimental Procedures](#).

Immunofluorescence

Cells were fixed and stained as described (Ko et al., 2010). The details of staining of cells and the sources of antibodies used are also described in the [Supplemental Experimental Procedures](#).

Flow Cytometry

The sources of antibodies used for flow cytometric analysis are described in the [Supplemental Experimental Procedures](#).

Long-Term Cultures and In Vitro Colony-Forming Assays

The details of long-term culture assays are described in the [Supplemental Experimental Procedures](#).

Competitive Reconstitution Assay

The details of competitive reconstitution assay are described in the [Supplemental Experimental Procedures](#).

Human Sample Analysis

Human patient samples were analyzed as described in detail in the [Supplemental Experimental Procedures](#). This study was approved by the Institutional Review Board waiver and Human Tissue Utilization Committee, and clinical and hematological data were recorded after patients provided their informed consent in accordance with the Declaration of Helsinki.

miRNA In Situ Hybridization

The details of in situ hybridization analysis for miR-22 are described in the [Supplemental Experimental Procedures](#).

Immunohistochemistry

The details of immunohistochemical analysis are described in the [Supplemental Experimental Procedures](#).

Dynamic Transcriptome Analysis

The details of dynamic transcriptome analysis are described in the [Supplemental Experimental Procedures](#).

Methylation-Specific PCR

DNA methylation status of *Sp140* in purified KSL cells (or Lin^{neg} cells) was examined by the methylation-specific PCR with genomic DNA treated with sodium bisulfite using the EZ DNA Methylation-Direct kit (Zymo Research). Primer sequences were previously described (Yamazaki et al., 2012). β -actin was used as an internal control of genomic DNA.

Luciferase Reporter Assay

The details of the luciferase reporter assay are described in the [Supplemental Experimental Procedures](#).

Statistical Analysis

Student's *t* test was utilized to determine statistical significance unless otherwise specified. $p < 0.05$ was considered statistically significant.

SUPPLEMENTAL INFORMATION

Supplemental Information for this article includes Supplemental Experimental Procedures and seven figures and can be found with this article online at <http://dx.doi.org/10.1016/j.stem.2013.06.003>.

ACKNOWLEDGMENTS

We thank all the members of the Pandolfi laboratory for their comments and discussion. We are grateful to Thomas Garvey and Min Sup Song for critical editing of the manuscript. We are also especially thankful to Akinobu Matsumoto, Dina Stroopinsky, and Mesruh Turkekel for technical support and help on in situ hybridization; to Lourdes Mendez for comments and help on hematological analysis; to Aviv Bergman for comments and support on statistical analysis of human patients' data; and to Miriam Fayad for support on cohort study of patients with hematological malignancies. K.I. was supported by NIH grants R00CA139009 and R01DK10068901. L.K. was supported by an NHMRC Overseas Biomedical Fellowship. This work was supported by CA141457-03 NIH grants to P.P.P.

Received: May 4, 2012

Revised: March 7, 2013

Accepted: June 11, 2013

Published: July 3, 2013

REFERENCES

- Abdel-Wahab, O., Mullally, A., Hedvat, C., Garcia-Manero, G., Patel, J., Wadleigh, M., Malinge, S., Yao, J., Kilpivaara, O., Bhat, R., et al. (2009). Genetic characterization of TET1, TET2, and TET3 alterations in myeloid malignancies. *Blood* 114, 144–147.
- Abdel-Wahab, O., Manshouri, T., Patel, J., Harris, K., Yao, J., Hedvat, C., Heguy, A., Bueso-Ramos, C., Kantarjian, H., Levine, R.L., and Verstovsek, S. (2010). Genetic analysis of transforming events that convert chronic myeloproliferative neoplasms to leukemias. *Cancer Res.* 70, 447–452.
- Arnold, C.P., Tan, R., Zhou, B., Yue, S.B., Schaffert, S., Biggs, J.R., Doyonnas, R., Lo, M.C., Perry, J.M., Renault, V.M., et al. (2011). MicroRNA programs in normal and aberrant stem and progenitor cells. *Genome Res.* 21, 798–810.

(D) miR-22 overexpression correlates with downregulation of TET2 in MDS patients. One-hundred and seven MDS patient specimens were subjected to in situ hybridization and immunohistochemical analyses, and the expression levels of miR-22 and TET2 were evaluated. The data were analyzed by Chi-square test. *r*, Pearson's *r*.

(E) The expression levels of miR-22 and TET2 in 18 AML with MLD patient samples were evaluated. The data were analyzed by Chi-square test. *r*, Pearson's *r*. AML, acute myeloid leukemia; MLD, multiple lineage dysplasia.

(F) miR-22 is highly expressed in human AML patients. Expression levels of miR-22 in CD34^{pos} bone marrow cells from healthy donors ($n = 9$) and AML patients ($n = 215$) are shown.

(G) Proposed model of oncogenic role of miR-22 in hematopoiesis. miR-22 negatively regulates TET2 tumor suppressor, leading to a reduction of 5-hmC and a potentiation of global gene methylation. This genetic remodeling in turn enhances HSC function and promotes hematological transformation. HSCs, hematopoietic stem cells; LIC, leukemia-initiating cells.

See also [Figure S6](#).

- Bartel, D.P. (2004). MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* 116, 281–297.
- Berger, A.H., Knudson, A.G., and Pandolfi, P.P. (2011). A continuum model for tumour suppression. *Nature* 476, 163–169.
- Betel, D., Wilson, M., Gabow, A., Marks, D.S., and Sander, C. (2008). The microRNA.org resource: targets and expression. *Nucleic Acids Res.* 36(Database issue), D149–D153.
- Branco, M.R., Ficiz, G., and Reik, W. (2012). Uncovering the role of 5-hydroxymethylcytosine in the epigenome. *Nat. Rev. Genet.* 13, 7–13.
- Calin, G.A., and Croce, C.M. (2006). MicroRNA signatures in human cancers. *Nat. Rev. Cancer* 6, 857–866.
- Chen, C.Z., Li, L., Lodish, H.F., and Bartel, D.P. (2004). MicroRNAs modulate hematopoietic lineage differentiation. *Science* 303, 83–86.
- Delhommeau, F., Dupont, S., Della Valle, V., James, C., Trannoy, S., Massé, A., Kosmider, O., Le Couedic, J.P., Robert, F., Alberdi, A., et al. (2009). Mutation in TET2 in myeloid cancers. *N. Engl. J. Med.* 360, 2289–2301.
- Di Bernardo, M.C., Crowther-Swanepoel, D., Broderick, P., Webb, E., Sellick, G., Wild, R., Sullivan, K., Vijayakrishnan, J., Wang, Y., Pittman, A.M., et al. (2008). A genome-wide association study identifies six susceptibility loci for chronic lymphocytic leukemia. *Nat. Genet.* 40, 1204–1210.
- Dostalova Merkerova, M., Krejcik, Z., Votavova, H., Belickova, M., Vasikova, A., and Cermak, J. (2011). Distinctive microRNA expression profiles in CD34+ bone marrow cells from patients with myelodysplastic syndrome. *Eur. J. Hum. Genet.* 19, 313–319.
- Elmén, J., Lindow, M., Schütz, S., Lawrence, M., Petri, A., Obad, S., Lindholm, M., Hedtjärn, M., Hansen, H.F., Berger, U., et al. (2008). LNA-mediated microRNA silencing in non-human primates. *Nature* 452, 896–899.
- Esteller, M. (2008). Epigenetics in cancer. *N. Engl. J. Med.* 358, 1148–1159.
- Ficiz, G., Branco, M.R., Seisenberger, S., Santos, F., Krueger, F., Hore, T.A., Marques, C.J., Andrews, S., and Reik, W. (2011). Dynamic regulation of 5-hydroxymethylcytosine in mouse ES cells and during differentiation. *Nature* 473, 398–402.
- Figuerola, M.E., Abdel-Wahab, O., Lu, C., Ward, P.S., Patel, J., Shih, A., Li, Y., Bhagwat, N., Vasanthakumar, A., Fernandez, H.F., et al. (2010). Leukemic IDH1 and IDH2 mutations result in a hypermethylation phenotype, disrupt TET2 function, and impair hematopoietic differentiation. *Cancer Cell* 18, 553–567.
- Gangaraju, V.K., and Lin, H. (2009). MicroRNAs: key regulators of stem cells. *Nat. Rev. Mol. Cell Biol.* 10, 116–125.
- Griffiths-Jones, S., Grocock, R.J., van Dongen, S., Bateman, A., and Enright, A.J. (2006). miRBase: microRNA sequences, targets and gene nomenclature. *Nucleic Acids Res.* 34 (Database issue), D140–D144.
- Ito, S., D'Alessio, A.C., Taranova, O.V., Hong, K., Sowers, L.C., and Zhang, Y. (2010). Role of Tet proteins in 5mC to 5hmC conversion, ES-cell self-renewal and inner cell mass specification. *Nature* 466, 1129–1133.
- Ito, S., Shen, L., Dai, Q., Wu, S.C., Collins, L.B., Swenberg, J.A., He, C., and Zhang, Y. (2011). Tet proteins can convert 5-methylcytosine to 5-formylcytosine and 5-carboxylcytosine. *Science* 333, 1300–1303.
- Jankowska, A.M., Szpurka, H., Tiu, R.V., Makishima, H., Afable, M., Huh, J., O'Keefe, C.L., Ganetzky, R., McDevitt, M.A., and Maciejewski, J.P. (2009). Loss of heterozygosity 4q24 and TET2 mutations associated with myelodysplastic/myeloproliferative neoplasms. *Blood* 113, 6403–6410.
- Jongen-Lavrencic, M., Sun, S.M., Dijkstra, M.K., Valk, P.J., and Löwenberg, B. (2008). MicroRNA expression profiling in relation to the genetic heterogeneity of acute myeloid leukemia. *Blood* 111, 5078–5085.
- Kallin, E.M., Rodríguez-Ubrea, J., Christensen, J., Cimmino, L., Aifantis, I., Helin, K., Ballestar, E., and Graf, T. (2012). Tet2 facilitates the derepression of myeloid target genes during CEPB α -induced transdifferentiation of pre-B cells. *Mol. Cell* 48, 266–276.
- Ko, M., Huang, Y., Jankowska, A.M., Pape, U.J., Tahiliani, M., Bandukwala, H.S., An, J., Lamperti, E.D., Koh, K.P., Ganetzky, R., et al. (2010). Impaired hydroxylation of 5-methylcytosine in myeloid cancers with mutant TET2. *Nature* 468, 839–843.
- Kosmider, O., Gelsi-Boyer, V., Cheok, M., Grabar, S., Della-Valle, V., Picard, F., Vigué, F., Quesnel, B., Beyne-Rauzy, O., Solary, E., et al.; Groupe Francophone des Myélobdysplasies. (2009). TET2 mutation is an independent favorable prognostic factor in myelodysplastic syndromes (MDSs). *Blood* 114, 3285–3291.
- Lanford, R.E., Hildebrandt-Eriksen, E.S., Petri, A., Persson, R., Lindow, M., Munk, M.E., Kauppinen, S., and Ørum, H. (2010). Therapeutic silencing of microRNA-122 in primates with chronic hepatitis C virus infection. *Science* 327, 198–201.
- Lewis, B.P., Burge, C.B., and Bartel, D.P. (2005). Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *Cell* 120, 15–20.
- Li, Z., Cai, X., Cai, C.L., Wang, J., Zhang, W., Petersen, B.E., Yang, F.C., and Xu, M. (2011). Deletion of Tet2 in mice leads to dysregulated hematopoietic stem cells and subsequent development of myeloid malignancies. *Blood* 118, 4509–4518.
- Liu, T.C., Lin, P.M., Chang, J.G., Lee, J.P., Chen, T.P., and Lin, S.F. (2000). Mutation analysis of PTEN/MMAC1 in acute myeloid leukemia. *Am. J. Hematol.* 63, 170–175.
- Moran-Crusio, K., Reavie, L., Shih, A., Abdel-Wahab, O., Ndiaye-Lobry, D., Lobry, C., Figueroa, M.E., Vasanthakumar, A., Patel, J., Zhao, X., et al. (2011). Tet2 loss leads to increased hematopoietic stem cell self-renewal and myeloid transformation. *Cancer Cell* 20, 11–24.
- Patsos, G., Germann, A., Gebert, J., and Dihlmann, S. (2010). Restoration of absent in melanoma 2 (AIM2) induces G2/M cell cycle arrest and promotes invasion of colorectal cancer cells. *Int. J. Cancer* 126, 1838–1849.
- Petersen, M., and Wengel, J. (2003). LNA: a versatile tool for therapeutics and genomics. *Trends Biotechnol.* 21, 74–81.
- Poliseno, L., Salmena, L., Riccardi, L., Fornari, A., Song, M.S., Hobbs, R.M., Sportoletti, P., Varmeh, S., Egia, A., Fedele, G., et al. (2010). Identification of the miR-106b~25 microRNA cluster as a proto-oncogenic PTEN-targeting intron that cooperates with its host gene MCM7 in transformation. *Sci. Signal.* 3, ra29.
- Quivoron, C., Couronné, L., Della Valle, V., Lopez, C.K., Plo, I., Wagner-Ballon, O., Do Cruzeiro, M., Delhommeau, F., Arnulf, B., Stern, M.H., et al. (2011). TET2 inactivation results in pleiotropic hematopoietic abnormalities in mouse and is a recurrent event during human lymphomagenesis. *Cancer Cell* 20, 25–38.
- Rodríguez-Paredes, M., and Esteller, M. (2011). Cancer epigenetics reaches mainstream oncology. *Nat. Med.* 17, 330–339.
- Sasaki, M., Knobbe, C.B., Munger, J.C., Lind, E.F., Brenner, D., Brüstle, A., Harris, I.S., Holmes, R., Wakeham, A., Haight, J., et al. (2012). IDH1(R132H) mutation increases murine haematopoietic progenitors and alters epigenetics. *Nature* 488, 656–659.
- Shen, L., Kantarjian, H., Guo, Y., Lin, E., Shan, J., Huang, X., Berry, D., Ahmed, S., Zhu, W., Pierce, S., et al. (2010). DNA methylation predicts survival and response to therapy in patients with myelodysplastic syndromes. *J. Clin. Oncol.* 28, 605–613.
- Shih, A.H., Abdel-Wahab, O., Patel, J.P., and Levine, R.L. (2012). The role of mutations in epigenetic regulators in myeloid malignancies. *Nat. Rev. Cancer* 12, 599–612.
- Smith, A.E., Mohamedali, A.M., Kulasekararaj, A., Lim, Z., Gäken, J., Lea, N.C., Przychodzen, B., Mian, S.A., Nasser, E.E., Shooter, C., et al. (2010). Next-generation sequencing of the TET2 gene in 355 MDS and CMML patients reveals low-abundance mutant clones with early origins, but indicates no definite prognostic value. *Blood* 116, 3923–3932.
- Song, S.J., Poliseno, L., Song, M.S., Ala, U., Webster, K., Ng, C., Beringer, G., Brikbak, N.J., Yuan, X., Cantley, L.C., et al. (2013). microRNA-antagonism regulates breast cancer stemness and metastasis via TET family dependent chromatin remodelling. *Cell* 154, in press. <http://dx.doi.org/10.1016/j.cell.2013.06.026>.
- Stenvang, J., Silaharoglu, A.N., Lindow, M., Elmen, J., and Kauppinen, S. (2008). The utility of LNA in microRNA-based cancer diagnostics and therapeutics. *Semin. Cancer Biol.* 18, 89–102.

Tefferi, A., Lim, K.H., Abdel-Wahab, O., Lasho, T.L., Patel, J., Patnaik, M.M., Hanson, C.A., Pardanani, A., Gilliland, D.G., and Levine, R.L. (2009). Detection of mutant TET2 in myeloid malignancies other than myeloproliferative neoplasms: CMML, MDS, MDS/MPN and AML. *Leukemia* 23, 1343–1345.

Wang, J., Xiang, G., Zhang, K., and Zhou, Y. (2012). Expression signatures of intragenic miRNAs and their corresponding host genes in myeloid leukemia cells. *Biotechnol. Lett.* 34, 2007–2015.

Yamazaki, J., Taby, R., Vasanthakumar, A., Macrae, T., Ostler, K.R., Shen, L., Kantarjian, H.M., Estecio, M.R., Jelinek, J., Godley, L.A., and Issa, J.P. (2012). Effects of TET2 mutations on DNA methylation in chronic myelomonocytic leukemia. *Epigenetics* 7, 201–207.

Yilmaz, O.H., Valdez, R., Theisen, B.K., Guo, W., Ferguson, D.O., Wu, H., and Morrison, S.J. (2006). Pten dependence distinguishes haematopoietic stem cells from leukaemia-initiating cells. *Nature* 441, 475–482.

Zhang, J., Grindley, J.C., Yin, T., Jayasinghe, S., He, X.C., Ross, J.T., Haug, J.S., Rupp, D., Porter-Westpfahl, K.S., Wiedemann, L.M., et al. (2006). PTEN maintains haematopoietic stem cells and acts in lineage choice and leukaemia prevention. *Nature* 441, 518–522.

Zhang, Y., Roccaro, A.M., Rombaoa, C., Flores, L., Obad, S., Fernandes, S.M., Sacco, A., Liu, Y., Ngo, H., Quang, P., et al. (2012). LNA-mediated anti-miR-155 silencing in low-grade B-cell lymphomas. *Blood* 120, 1678–1686.