

Available online at www.sciencedirect.com**ScienceDirect**

Procedia Computer Science 96 (2016) 879 – 886

Procedia
Computer Science

20th International Conference on Knowledge Based and Intelligent Information and Engineering Systems

The cycle inconsistency index in pairwise comparisons matrices

Jacek Szybowski

*AGH University of Science and Technology, Address, al. Mickiewicza 30, 30-059 Krakow, Poland**State Higher Vocational School in Nowy Sacz, Staszica 1, 33-300, Nowy Sacz, Poland*

Abstract

The method of pairwise comparisons is widely applied in the decision making process. The inconsistency of data may significantly affect the final result. Since the notion of consistency is based on triads or cycles, there is a great need for defining the measure of a triad or cycle inconsistency.

In the paper a set of properties of a good cycle inconsistency index is proposed. Two construction methods of a cycle-based inconsistency index for a pairwise comparisons matrix are introduced. All is supported by the examples.

© 2016 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Peer-review under responsibility of KES International

Keywords: pairwise comparisons, cycle, inconsistency;

1. Introduction

The first use of the method of Pairwise Comparisons (PC method) is attributed to Ramon Llull. In his recently discovered manuscript *Ars electionis*¹³ this Catalan philosopher, logician and a pioneer of computation theory, described the Borda count (an election system) and the Condorcet criterion (a criterion for a voting system), which Jean-Charles de Borda¹ and Nicolas de Condorcet⁵ independently discovered centuries later. In the 20th century the PC method was used by Thurstone¹⁶, Kendall and Babington-Smith⁶, as well as many others.

The method is used when we are supposed to order a set of alternatives. We compare them pairwise and write the numerical result of these comparisons to a square matrix called a pairwise comparisons matrix (a PC matrix). The problem occurs when it appears that an alternative A is considered to be better than B, B - better than C, and C - better than A. This situation was described as a Condorcet's paradox⁵ and it results in the so-called triad inconsistency.

Obviously, this paradox may involve more alternatives. For instance, A wins with B and C, B - with C and D, C - with D, but A loses with D. This results in the so-called cycle inconsistency.

* Jacek Szybowski. Tel.: +48-12-617-3587.

E-mail address: szybowski@agh.edu.pl

Many researchers describe different types of measures of inconsistency. Recently, some attempts to give a set of axioms for a good inconsistency index have been undertaken by Cavallo and d’Apuzzo⁴ or Koczkodaj, Szybowski and Wajch^{11,10} for the case of triads, and by Koczkodaj and Szwarz⁸ or Brunelli and Fedrizzi³ for the case of whole PC matrices. However, there has been no axiomatization for a cycle inconsistency index, so this paper tries to fill this gap.

2. Preliminaries

An $n \times n$ matrix $\mathbf{A} = [a_{ij}]$ with all positive elements is called a *pairwise comparisons matrix* (a PC matrix) if $n \geq 3$. A PC matrix \mathbf{A} is called *reciprocal* if $a_{ij} = \frac{1}{a_{ji}}$ for every $i, j = 1, \dots, n$ (then obviously $a_{ii} = 1$ for every $i = 1, \dots, n$). When we compare two entities X and Y and we judge, that X is twice better (longer, more expensive etc.), it is quite natural to accept that Y is half as good (long, expensive) as X. That is why we will assume that each PC matrix is reciprocal.

Fix a PC matrix \mathbf{A} and $s \in \mathbb{N}$, such that $3 \leq s \leq n$.

Definition 2.1. An ordered sequence (a_{ij}, a_{ik}, a_{jk}) of 3 elements of \mathbf{A} is called a *triad* if i, j, k are pairwise different elements of the set $\{1, \dots, n\}$.

We will denote the set of all triads in \mathbf{A} by $\mathcal{T}_{\mathbf{A}}$.

Definition 2.2. A triad (a_{ij}, a_{ik}, a_{jk}) is said to be *consistent* if $a_{ij} \cdot a_{jk} = a_{ik}$.

Remark 2.3. A triad (a_{ij}, a_{ik}, a_{jk}) is consistent $\Leftrightarrow a_{ij} \cdot a_{jk} \cdot a_{ki} = 1$.

Definition 2.4. An ordered sequence $(a_{i_1 i_2}, a_{i_2 i_3}, \dots, a_{i_{s-1} i_s}, a_{i_1 i_s})$ of s elements of \mathbf{A} is called an *s-cycle* if i_1, \dots, i_s are pairwise different elements of the set $\{1, \dots, n\}$.

We will denote the set of all s -cycles in \mathbf{A} by $C_{\mathbf{A}}^s$.

Definition 2.5. An s -cycle $(a_{i_1 i_2}, a_{i_2 i_3}, \dots, a_{i_{s-1} i_s}, a_{i_1 i_s})$ is said to be *consistent* if $a_{i_1 i_2} \cdot a_{i_2 i_3} \cdot \dots \cdot a_{i_{s-1} i_s} = a_{i_1 i_s}$.

Remark 2.6. An s -cycle $(a_{i_1 i_2}, a_{i_2 i_3}, \dots, a_{i_{s-1} i_s}, a_{i_1 i_s})$ is consistent $\Leftrightarrow a_{i_1 i_2} \cdot a_{i_2 i_3} \cdot \dots \cdot a_{i_{s-1} i_s} \cdot a_{i_s i_1} = 1$.

Definition 2.7. A PC matrix is *consistent* if all its triads are consistent.

Proposition 2.8. A PC matrix \mathbf{A} is consistent \Leftrightarrow all its s -cycles are consistent.

Proof. The proof is inductive. It is enough to show that $\forall p \in \{3, \dots, s - 1\}$

$$\text{All } p\text{-cycles of } \mathbf{A} \text{ are consistent} \Leftrightarrow \text{all } (p + 1)\text{-cycles of } \mathbf{A} \text{ are consistent.}$$

To prove the \Rightarrow notice that

$$a_{i_1 i_2} \cdot a_{i_2 i_3} \cdot \dots \cdot a_{i_{p-1} i_p} \cdot a_{i_p i_{p+1}} \cdot a_{i_{p+1} i_1} = a_{i_1 i_2} \cdot a_{i_2 i_3} \cdot \dots \cdot a_{i_{p-1} i_p} \cdot a_{i_p i_1} = 1$$

For the reverse implication take any $j \notin \{i_1, \dots, i_p\}$ and multiply the below equations by themselves:

$$\begin{aligned} a_{i_1 j} \cdot a_{j i_2} \cdot a_{i_2 i_3} \cdot \dots \cdot a_{i_{p-1} i_p} \cdot a_{i_p i_1} &= 1 \\ a_{i_1 i_2} \cdot a_{i_2 j} \cdot a_{j i_3} \cdot \dots \cdot a_{i_{p-1} i_p} \cdot a_{i_p i_1} &= 1 \\ &\dots \\ a_{i_1 i_2} \cdot a_{i_2 i_3} \cdot a_{i_3 i_4} \cdot \dots \cdot a_{i_{p-1} j} \cdot a_{j i_1} &= 1. \end{aligned}$$

It follows that

$$a_{i_1 j} \cdot a_{j i_1} \cdot \dots \cdot a_{i_{p-1} j} \cdot a_{j i_{p-1}} \cdot a_{i_1 i_2}^{p-2} \cdot a_{i_2 i_3}^{p-2} \cdot \dots \cdot a_{i_{p-1} i_p}^{p-2} \cdot a_{i_p i_1}^{p-2} = 1,$$

so

$$(a_{i_1 i_2} \cdot a_{i_2 i_3} \cdot \dots \cdot a_{i_{p-1} i_p} \cdot a_{i_p i_1})^{p-2} = 1,$$

and finally,

$$a_{i_1 i_2} \cdot a_{i_2 i_3} \cdot \dots \cdot a_{i_{p-1} i_p} \cdot a_{i_p i_1} = 1.$$



3. The triad inconsistency index

Recall some definitions and results from¹⁰.

Definition 3.1. A function $ii : \mathbb{R}_+^3 \rightarrow [0, +\infty)$ is called a *triad inconsistency index*, if there exists a metric

$$d : \mathbb{R}_+^2 \rightarrow [0, +\infty)$$

such that $\forall x, y, z$ it holds

$$ii(x, y, z) = d(xz, y). \tag{1}$$

Definition 3.2. We say that a triad inconsistency index ii satisfying (1) is *induced by a metric d* .

Straight from the definition of a triad inconsistency index we get

Proposition 3.3. For all numbers a, b, c, d and e a triad inconsistency index ii satisfies the conditions

$$ii(a, b, c) = 0 \Leftrightarrow ac = b \tag{2}$$

$$ii(a, b, c) = ii(b, ac, 1) \tag{3}$$

$$ii(a, de, c) \leq ii(a, b, c) + ii(d, b, e) \tag{4}$$

Example 3.4. Since function $d : \mathbb{R}_+^2 \rightarrow [0, 1)$ given by formula

$$d(x, y) = 1 - \min\left(\frac{x}{y}, \frac{y}{x}\right) \tag{5}$$

is a metric, the Koczkodaj triad inconsistency index

$$KI(a, b, c) = 1 - \min\left(\frac{b}{ac}, \frac{ac}{b}\right),$$

is a triad inconsistency index induced by d .

4. The cycle inconsistency index

Now we may define the inconsistency index for cycles by analogy to the case of triads.

Definition 4.1. A function $ii : \mathbb{R}_+^s \rightarrow [0, +\infty)$ is called a *cycle inconsistency index*, if there exists a metric

$$d : \mathbb{R}_+^2 \rightarrow [0, +\infty)$$

such that $\forall x_1, \dots, x_s$ it holds

$$ii(x_1, \dots, x_s) = d(x_1 \cdot \dots \cdot x_{s-1}, x_s). \tag{6}$$

Definition 4.2. We say that a cycle inconsistency index ii satisfying (6) is *induced by a metric d* .

Straight from the definition of a cycle inconsistency index we get

Proposition 4.3. For all numbers x_1, \dots, x_{2s-1} a cycle inconsistency index ii satisfies the conditions

$$ii(x_1, \dots, x_s) = 0 \Leftrightarrow x_1 \cdot \dots \cdot x_{s-1} = x_s \tag{7}$$

$$ii(x_1, \dots, x_s) = ii(x_s, 1, \dots, 1, x_1 \cdot \dots \cdot x_{s-1}) \tag{8}$$

$$ii(x_1, \dots, x_{s-1}, x_{s+1} \cdot \dots \cdot x_{2s-1}) \leq ii(x_1, \dots, x_{s-1}, x_s) + ii(x_{s+1}, \dots, x_{2s-1}, x_s) \tag{9}$$

Proof. The first statement is obvious. For the proof of the second one notice that

$$ii(x_1, \dots, x_s) = d(x_1 \cdot \dots \cdot x_{s-1}, x_s) = d(x_s, x_1 \cdot \dots \cdot x_{s-1}) = d(x_s \cdot 1 \cdot \dots \cdot 1, x_1 \cdot \dots \cdot x_{s-1}) = ii(x_s, 1, \dots, 1, x_1 \cdot \dots \cdot x_{s-1}).$$

Finally,

$$\begin{aligned} ii(x_1, \dots, x_{s-1}, x_{s+1} \cdot \dots \cdot x_{2s-1}) &= d(x_1 \cdot \dots \cdot x_{s-1}, x_{s+1} \cdot \dots \cdot x_{2s-1}) \leq \\ &\leq d(x_1 \cdot \dots \cdot x_{s-1}, x_s) + d(x_{s+1} \cdot \dots \cdot x_{2s-1}, x_s) = \\ &= ii(x_1, \dots, x_{s-1}, x_s) + ii(x_{s+1}, \dots, x_{2s-1}, x_s). \end{aligned}$$

□

Remark 4.4. We may treat conditions (7) – (9) as axioms for a cycle inconsistency index.

As a consequence of (8) we obtain

Corollary 4.5. For all numbers x_1, \dots, x_s and any permutation j of a set $\{1, \dots, s-1\}$

$$ii(x_1, \dots, x_{s-1}, x_s) = ii(x_{j(1)}, \dots, x_{j(s-1)}, x_s). \quad (10)$$

Proposition 4.6. For each function $ii : \mathbb{R}_+^s \rightarrow [0, +\infty)$ satisfying (7) – (9) function $d_{ii} : \mathbb{R}_+^2 \rightarrow [0, +\infty)$ given by

$$d_{ii}(x, y) = ii(x, 1, \dots, 1, y) \quad (11)$$

is a metric. Moreover, index ii is induced by the metric d_{ii} .

Definition 4.7. We say that a metric d_{ii} satisfying (11) is induced by a cycle inconsistency index ii .

Definition 4.8. We say that a cycle inconsistency index ii is bounded if $\exists M > 0 \forall x_1, \dots, x_s \in \mathbb{R}_+ ii(x_1, \dots, x_s) \leq M$.

Fix a natural number $s \geq 3$.

Example 4.9. The function $D : \mathbb{R}_+^s \rightarrow \{0, 1\}$ given by

$$D(x_1, \dots, x_{s-1}, y) = \begin{cases} 0, & x_1 \cdot \dots \cdot x_{s-1} = y \\ 1, & \text{otherwise} \end{cases}$$

is a bounded s -cycle inconsistency index induced by a discrete metric.

Example 4.10. The function $E : \mathbb{R}_+^s \rightarrow [0, +\infty)$ given by

$$E(x_1, \dots, x_{s-1}, y) = |x_1 \cdot \dots \cdot x_{s-1} - y|$$

is an unbounded s -cycle inconsistency index induced by a Euclidean metric.

Example 4.11. The function $I_1 : \mathbb{R}_+^s \rightarrow [0, 1)$ given by

$$I_1(x_1, \dots, x_{s-1}, y) = \frac{|x_1 \cdot \dots \cdot x_{s-1} - y|}{1 + |x_1 \cdot \dots \cdot x_{s-1} - y|}$$

is a bounded s -cycle inconsistency index induced by a metric d_1 given by formula $d_1(x, y) = \frac{|x-y|}{1+|x-y|}$.

Example 4.12. The function $K : \mathbb{R}_+^s \rightarrow [0, 1)$ given by

$$K(x_1, \dots, x_{s-1}, y) = 1 - \min\left(\frac{y}{x_1 \cdot \dots \cdot x_{s-1}}, \frac{x_1 \cdot \dots \cdot x_{s-1}}{y}\right)$$

is a bounded s -cycle inconsistency index induced by the metric (5).

Notice, that a given triad (cycle) inconsistency index ii may be applied to the triads (cycles) of a PC matrix \mathbf{A} . Basing on the inconsistency indices for triads or s-cycles, we can introduce the inconsistency s -indices for the whole PC-matrices. Obviously, we can do it in various ways. However, two approaches seem the most natural.

The first one, introduced in¹⁴ or⁴ (for the case of triads) involves the arithmetic or geometric means of triad indices.

Example 4.13. The Peláez-Lamata index was defined as

$$PLI(\mathbf{A}) = \frac{6 \sum_{(a_{ij}, a_{ik}, a_{jk}) \in \mathcal{T}_A} PL(a_{ij}, a_{ik}, a_{jk})}{n(n-1)(n-2)},$$

where

$$PL(a_{ij}, a_{ik}, a_{jk}) = \frac{a_{ik}}{a_{ij}a_{jk}} + \frac{a_{ij}a_{jk}}{a_{ik}} - 2.$$

It is natural to generalize the above definition to a cycle-based Peláez-Lamata inconsistency index:

$$PLI_s(\mathbf{A}) = \frac{\sum_{(a_{i_1 i_2}, \dots, a_{i_{s-1} i_s}, a_{i_1 i_s}) \in C_A^s} PL_s(a_{i_1 i_2}, \dots, a_{i_{s-1} i_s}, a_{i_1 i_s})}{\binom{n}{s}},$$

where

$$PL_s(a_{i_1 i_2}, \dots, a_{i_{s-1} i_s}, a_{i_1 i_s}) = \frac{a_{i_1 i_s}}{a_{i_1 i_2} \cdot \dots \cdot a_{i_{s-1} i_s}} + \frac{a_{i_1 i_2} \cdot \dots \cdot a_{i_{s-1} i_s}}{a_{i_1 i_s}} - 2.$$

Proposition 4.14. PL_s does not satisfy (9).

Proof. Consider $s = 4$.

$$PL_4(1, 4, 1, 1, 1 \cdot 2 \cdot 1) = \frac{1 \cdot 2 \cdot 1}{1 \cdot 4 \cdot 1} + \frac{1 \cdot 4 \cdot 1}{1 \cdot 2 \cdot 1} - 2 = \frac{1}{2} > \frac{1}{4} = \frac{3}{4} + \frac{4}{3} - 2 + \frac{3}{2} + \frac{2}{3} - 2 = PL_4(1, 4, 1, 3) + PL_4(1, 2, 1, 3).$$

□

The second type of definitions uses the maximum function. This approach was used (for triads and cycles) for the Koczkodaj inconsistency indices proposed in⁷ and simplified in⁸.

Example 4.15.

$$KI(\mathbf{A}) = \max_{(a_{ij}, a_{ik}, a_{jk}) \in \mathcal{T}_A} K(a_{ij}, a_{ik}, a_{jk}),$$

where

$$K(a_{ij}, a_{ik}, a_{jk}) = 1 - \min\left(\frac{a_{ik}}{a_{ij}a_{jk}}, \frac{a_{ij}a_{jk}}{a_{ik}}\right),$$

and

$$KI_s(\mathbf{A}) = \max_{(a_{i_1 i_2}, \dots, a_{i_{s-1} i_s}, a_{i_1 i_s}) \in C_A^s} K(a_{i_1 i_2}, \dots, a_{i_{s-1} i_s}, a_{i_1 i_s}),$$

where

$$K(a_{i_1 i_2}, \dots, a_{i_{s-1} i_s}, a_{i_1 i_s}) = 1 - \min\left(\frac{a_{i_1 i_s}}{a_{i_1 i_2} \cdot \dots \cdot a_{i_{s-1} i_s}}, \frac{a_{i_1 i_2} \cdot \dots \cdot a_{i_{s-1} i_s}}{a_{i_1 i_s}}\right).$$

Remark 4.16. As it was shown in¹² indices PLI and KI are not equivalent, which means that there are no positive constants α and β such that

$$\alpha KI \leq PLI \leq \beta KI.$$

Example 4.17. Similarly, using the inconsistency index from Ex. 4.9 we can define

$$DI(\mathbf{A}) = \max_{(a_{ij}, a_{ik}, a_{jk}) \in \mathcal{T}_{\mathbf{A}}} D(a_{ij}, a_{jk}, a_{ik})$$

and

$$DI_s(\mathbf{A}) = \max_{(a_{i_1 i_2}, \dots, a_{i_{s-1} i_s}, a_{i_1 i_s}) \in C_{\mathbf{A}}^s} D(a_{i_1 i_2}, \dots, a_{i_{s-1} i_s}, a_{i_1 i_s})$$

However, from Proposition 2.8, we get

Corollary 4.18. $\forall s \in \{4, \dots, n\}$

$$DI(\mathbf{A}) = DI_s(\mathbf{A}).$$

This means that defining discrete cycle inconsistency index DI_s is useless, since it carries the same information as the simpler one based on triads.

On the other hand, this is not the case for other indices, for example KI, as the following example shows.

Example 4.19. Consider the PC matrices

$$A = \begin{bmatrix} 1 & \mathbf{2} & 1 & \mathbf{1} \\ \frac{1}{2} & 1 & \mathbf{3} & \mathbf{1} \\ 1 & \frac{1}{3} & 1 & \mathbf{5} \\ 1 & 1 & \frac{1}{5} & 1 \end{bmatrix}$$

and

$$B = \begin{bmatrix} 1 & \mathbf{9} & \mathbf{1} & \mathbf{2} \\ \frac{1}{9} & 1 & \mathbf{9} & \mathbf{2} \\ 1 & \frac{1}{9} & 1 & \mathbf{1} \\ \frac{1}{2} & \frac{1}{2} & 1 & 1. \end{bmatrix}$$

A simple calculation shows that

$$K(a_{23}, a_{24}, a_{34}) = 1 - \min\left(\frac{1}{3 \cdot 5}, \frac{3 \cdot 5}{1}\right) = \frac{14}{15}$$

(a triad in bold), and

$$K(a_{12}, a_{23}, a_{34}, a_{14}) = 1 - \min\left(\frac{1}{2 \cdot 3 \cdot 5}, \frac{2 \cdot 3 \cdot 5}{1}\right) = \frac{29}{30}$$

(a cycle marked in frames).

It is easy to notice that

$$KI(\mathbf{A}) = K(a_{23}, a_{24}, a_{34}) = \frac{14}{15} < \frac{29}{30} = K(a_{12}, a_{23}, a_{34}, a_{14}) = KI_4(\mathbf{A})$$

On the other hand,

$$AV(\mathbf{A}) := \frac{1}{|\mathcal{T}_{\mathbf{A}}|} \sum_{(a_{ij}, a_{ik}, a_{jk}) \in \mathcal{T}_{\mathbf{A}}} K(a_{ij}, a_{ik}, a_{jk}) = \frac{1}{24} \cdot 6 \cdot \left(\frac{5}{6} + \frac{14}{15} + \frac{1}{2} + \frac{4}{5}\right) = \frac{23}{30} \approx 0.7667$$

and

$$AV_4(\mathbf{A}) := \frac{1}{|C_{\mathbf{A}}^4|} \sum_{(a_{ij}, a_{jk}, a_{kl}, a_{il}) \in C_{\mathbf{A}}^4} K(a_{ij}, a_{jk}, a_{kl}, a_{il}) = \frac{1}{24} \cdot 8 \cdot \left(\frac{29}{30} + \frac{3}{5} + \frac{2}{3}\right) = \frac{67}{90} \approx 0.7444$$

so

$$AV(\mathbf{A}) > AV_4(\mathbf{A}).$$

Similarly,

$$K(b_{12}, b_{13}, b_{23}) = 1 - \min\left(\frac{1}{9 \cdot 9}, \frac{9 \cdot 9}{1}\right) = \frac{80}{81}$$

(a triad in bold), and

$$K(b_{34}, b_{41}, b_{12}, b_{32}) = 1 - \min\left(\frac{\frac{1}{9}}{1 \cdot \frac{1}{2} \cdot 9}, \frac{1 \cdot \frac{1}{2} \cdot 9}{\frac{1}{9}}\right) = \frac{79}{81}$$

(a cycle marked in frames).

It is also easy to notice that

$$KI(\mathbf{B}) = K(b_{12}, b_{13}, b_{23}) = \frac{80}{81} > \frac{79}{81} = K(b_{34}, b_{41}, b_{12}, b_{32}) = KI_4(\mathbf{B}).$$

On the other hand,

$$AV(\mathbf{B}) = \frac{1}{24} \cdot 6 \cdot \left(\frac{80}{81} + \frac{7}{9} + \frac{8}{9} + \frac{1}{2}\right) = \frac{511}{648} \approx 0.7886$$

and

$$AV_4(\mathbf{B}) = \frac{1}{24} \cdot 8 \cdot \left(\frac{79}{81} + \frac{17}{18} + \frac{8}{9}\right) = \frac{455}{486} \approx 0.9362$$

so

$$AV(\mathbf{B}) < AV_4(\mathbf{B}).$$

Note that definitions of some inconsistency indices are based neither on triads nor on cycles, like the Saaty’s consistency index defined in¹⁵ as

$$CI(\mathbf{A}) = \frac{\lambda_{max} - n}{n - 1},$$

where λ_{max} is the principle eigenvalue of \mathbf{A} .

However, as indicated in¹⁰, one may easily identify a triad with a 3×3 PC matrix, so we can introduce a triad inconsistency index on the base of a matrix inconsistency index. A detailed comparison of indices CI and KI for 3×3 matrices was done in².

5. Conclusion

It seems very natural to measure the level of a cycle $(a_{i_1 i_2}, \dots, a_{i_{s-1} i_s}, a_{i_1 i_s})$ inconsistency by means of a distance between $a_{i_1 i_2} \cdot \dots \cdot a_{i_{s-1} i_s}$ and $a_{i_1 i_s}$. The formulas (7) – (9) allow to easily check if a given index is distance-based. Applying different metrics gives an opportunity to define various triad, cycle or PC matrix inconsistency indices. Although the definition of a matrix consistency bases on triads, Example 4.19 shows that the level of inconsistency may increase or decrease if we consider longer cycles instead.

The high level of inconsistency indicates errors in comparison of alternatives. This is why the inconsistency reduction is desirable before prioritization. An algorithm of the reconstruction of a consistent PC matrix from an $(n - 1)$ - subset of its elements (named a base) was proposed in⁹. A cycle inconsistency index might be applied to choose the least inconsistent n -cycle as a base for the construction of the most reliable consistent PC matrix.

Acknowledgements

The research is supported by the Polish Ministry of Science and Higher Education.

References

1. Borda M de. Mémoire sur les élections au scrutin. *Histoire de l'Académie Royale des Sciences*. Paris 1781.
2. Bozóki S, Rapsák T. On Saaty's and Koczkodaj's inconsistencies of pairwise comparisons matrices. *J. Global Optim.* 2008; **42(2)**: 157–175.
3. Brunelli M, Fedrizzi M. Axiomatic properties of inconsistency indices for pairwise comparisons. *Journal of the Operational Research Society* 2015; **66(1)**: 1–15.
4. Cavallo B, D'Apuzzo L. A general unified framework for pairwise comparisons matrices in multicriterial methods. *Int. J. Intell. Syst.* 2009; **24**: 377–398.
5. Condorcet MJA de. Essai sur l'application de l'analyse à la probabilité des décisions rendues à la pluralité des voix. Paris: l'Imprimerie Royale 1785; [fascimile edition New York: Chelsea, 1972].
6. Kendall MG, Babington-Smith B. On the method of paired comparisons. *Biometrika* 1939; **31(3/4)**: 324–345.
7. Koczkodaj WW. A New Definition of Consistency of Pairwise Comparisons. *Mathematical and Computer Modelling* 1993; **18(7)**: 79–84.
8. Koczkodaj WW, Szwarz R. On Axiomatization of Inconsistency Indicators for Pairwise Comparisons. *Fundamenta Informaticae* 2014; **132(4)**: 485–500.
9. Koczkodaj WW, Szybowski J. Pairwise comparisons simplified. *Applied Mathematics and Computation* 2015; **253**: 387–394.
10. Koczkodaj WW, Szybowski J. The key properties of inconsistency for a triad in pairwise comparisons matrices. Submitted. Available at <http://arxiv.org/abs/1505.05220>.
11. Koczkodaj WW, Szybowski J, Wajch E. Inconsistency indicator maps on groups for pairwise comparisons. *International Journal of Approximate Reasoning* 2016; **69(2)**: 81–90.
12. Kulakowski K, Szybowski J. The new triad based inconsistency indices for pairwise comparisons. *Knowledge-Based and Intelligent Information and Engineering Systems 18th Annual Conference, KES-2014, Procedia Computer Science* 2014; **35**: 1132–1137.
13. Llull R. *Ars Electionis* (manuscript) 1299.
14. Peláez JI, Lamata MT. A new measure of consistency for positive reciprocal matrices. *Computers and Mathematics with Applications* 2003; **46(12)**: 1839–1845.
15. Saaty TL. A scaling method for priorities in hierarchical structures. *Journal of Mathematical Psychology* 1977; **15(3)**: 234–281.
16. Thurstone LL. A law of comparative judgments. *Psychol. Rev.* 1927; **34**: 273–286.