



ELSEVIER

Contents lists available at ScienceDirect

Journal of Algebra

[www.elsevier.com/locate/jalgebra](http://www.elsevier.com/locate/jalgebra)

# Integral closure and generic elements

Cătălin Ciupercă

Department of Mathematics 2750, North Dakota State University, PO Box 6050, Fargo, ND 58108-6050, USA

## ARTICLE INFO

### Article history:

Received 26 March 2010

Available online 7 August 2010

Communicated by Luchezar L. Avramov

### MSC:

13B22

13A30

13H05

### Keywords:

Integral closure of ideals

Generic elements

Formally equidimensional rings

Complete intersection ideals

Almost complete intersection ideals

General grade reductions

## ABSTRACT

Let  $(R, \mathfrak{m})$  be a formally equidimensional local ring with depth  $R \geq 2$  and  $I = (a_1, \dots, a_n)$  an  $\mathfrak{m}$ -primary ideal in  $R$ . The main result of this paper shows that if  $I$  is integrally closed, then so is its image modulo a generic element, that is, if  $T = R[X_1, \dots, X_n]/(a_1X_1 + \dots + a_nX_n)$ , then  $\overline{IT} = I\overline{T}$ .

© 2010 Elsevier Inc. All rights reserved.

## 1. Introduction

Let  $R$  be a commutative noetherian ring and  $I$  an ideal in  $R$ . An element  $x \in R$  is said to be integral over  $I$  if it satisfies an equation  $x^n + b_1x^{n-1} + \dots + b_{n-1}x + b_n = 0$  with  $b_i \in I^i$  for all  $i$ . The set of all the elements that are integral over  $I$  is an ideal  $\overline{I}$ , the integral closure of  $I$ . If  $\varphi: R \rightarrow S$  is a ring homomorphism, then  $\overline{IS} \subseteq \overline{I\overline{S}}$ , a property referred to as *persistence* (see [7, 1.1.3]). The equality does not necessarily hold; however, note that  $\overline{IS} = \overline{I\overline{S}}$  if and only if  $\overline{IS}$  is integrally closed.

For a formally equidimensional local ring  $(R, \mathfrak{m})$  with depth  $R \geq 2$  and an  $\mathfrak{m}$ -primary ideal  $I$ , we prove that the integral closure of  $I$  is preserved under specialization modulo generic elements. In the language of first general grade reductions (2.2) introduced by Hochster [4], this means that the extension of an integrally closed ideal  $I$  to a first general grade reduction of  $(R, I)$  is integrally closed, too. More precisely, we prove the following theorem.

---

E-mail address: [catalin.ciuperca@ndsu.edu](mailto:catalin.ciuperca@ndsu.edu).

**Theorem 1.** Let  $(R, \mathfrak{m})$  be a local ring with depth  $R \geq 2$  and  $I = (a_1, a_2, \dots, a_n)$  an  $\mathfrak{m}$ -primary ideal in  $R$ . Let  $S = R[X_1, X_2, \dots, X_n]$ ,  $\alpha = a_1X_1 + a_2X_2 + \dots + a_nX_n \in S$  and  $T = S/\alpha S$ . The following hold:

- (a) If  $R$  is formally equidimensional, then  $\overline{IT} = \widehat{IT}$ .
- (b) If  $R$  is analytically unramified and Cohen–Macaulay, then  $\overline{IT} = \widehat{IT}$  for  $\mathfrak{m} \gg 0$ .

This implies the following local version.

**Corollary 2.** Let  $(R, \mathfrak{m})$  be a local ring with depth  $R \geq 2$  and  $I = (a_1, a_2, \dots, a_n)$  an  $\mathfrak{m}$ -primary ideal in  $R$ . Let  $U = R[X_1, X_2, \dots, X_n]_{\mathfrak{m}[X_1, \dots, X_n]}$ ,  $\alpha = a_1X_1 + a_2X_2 + \dots + a_nX_n \in U$  and  $V = U/\alpha U$ . The following hold:

- (a) If  $R$  is formally equidimensional, then  $\overline{IV} = \widehat{IV}$ .
- (b) If  $R$  is analytically unramified and Cohen–Macaulay, then  $\overline{IV} = \widehat{IV}$  for  $\mathfrak{m} \gg 0$ .

Under the assumption that  $R$  is an equidimensional, universally catenary ring such that  $R/\sqrt{0}$  is analytically unramified, part (a) of Theorem 1 also appears in a 2006 preprint of Hong and Ulrich [5, Theorem 2.1]. Both parts of Corollary 2 were also proved by Itoh [10, Theorem 1] for ideals generated by a system of parameters in analytically unramified Cohen–Macaulay local rings of dimension at least two.

One application of these theorems is their use in proofs based on induction. We exemplify this in the final section of this paper by extending some results regarding integrally closed almost complete intersection ideals in regular local rings obtained by the author in [2].

## 2. Preliminary results

All the rings considered in this paper are commutative with identity. If the ring  $R$  is noetherian and  $I$  is an ideal in  $R$ , we denote by  $\text{grade } I$  the common length of all the maximal regular sequences contained in  $I$ . If the ring  $R$  is local with maximal ideal  $\mathfrak{m}$ , then  $\text{grade } \mathfrak{m}$  will be denoted  $\text{depth } R$ . We also say that the local ring  $R$  is formally equidimensional if its completion is an equidimensional ring, that is,  $\dim \widehat{R}/P = \dim R$  for all the minimal prime ideals  $P \in \text{Spec } \widehat{R}$ . In the literature, the local rings with this property are also called quasi-unmixed.

In this section we prove several lemmas which will be used in the proof of the main result.

**Remark 2.1.** With the notation used in Theorem 1 and Corollary 2, both  $S$  and  $U$  are faithfully flat extensions of  $R$  and  $\dim U = \dim R$ . In particular, if  $I$  is an ideal in  $R$ , then  $\overline{IS} = \widehat{IS}$  and  $\overline{IU} = \widehat{IU}$  [7, 8.4.2(9)].

The next discussion shows that if Theorem 1 is true for *some* set of generators of  $I$ , then the theorem holds for *every* set of generators of  $I$ .

### 2.2. General grade reductions

Let  $I = (a_1, \dots, a_n)$  be an ideal in a noetherian ring  $R$  with  $\text{grade } I > 0$ . The element  $\alpha = a_1X_1 + a_2X_2 + \dots + a_nX_n$  is a non-zero-divisor on  $S = R[X_1, \dots, X_n]$  (it follows inductively from [12, (6.13)]), which implies that the grade of  $IS/\alpha S$  in  $S/\alpha S$  is one less than the grade of  $I$  in  $R$ . Introduced by Hochster in [4],  $(S/\alpha S, IS/\alpha S)$  is called a first general grade reduction of  $(R, I)$ . While a first general grade reduction does depend on the choice of generators of  $I$ , it can be shown that any two first general grade reductions  $(T_1, IT_1)$  and  $(T_2, IT_2)$  of  $(R, I)$  are equivalent in the following sense: there exist indeterminates  $Y_1, \dots, Y_r$  over  $T_1$  and  $Z_1, \dots, Z_s$  over  $T_2$ , and an  $R$ -algebra isomorphism  $\phi : T_1[Y_1, \dots, Y_r] \xrightarrow{\cong} T_2[Z_1, \dots, Z_s]$ . Note that this implies that  $\phi(IT_1[Y_1, \dots, Y_r]) = IT_2[Z_1, \dots, Z_s]$ , since  $\phi$  is an  $R$ -algebra isomorphism. The existence of this  $R$ -algebra isomorphism follows from the proof of [4, Proposition 1]; for the convenience of the reader, we repeat the argument here. Assume that  $(T_1, IT_1)$  is obtained with respect to the sequence of generators  $b_1, \dots, b_m$  of  $I$  and  $(T_2, IT_2)$

with respect to  $c_1, \dots, c_p$ . Let  $(T_3, IT_3)$  be the first general grade reduction of  $(R, I)$  obtained by using  $b_1, \dots, b_m, c_1, \dots, c_p$  as generators of  $I$ . Since it is enough to show that  $(T_1, IT_1)$  is equivalent to  $(T_3, IT_3)$  and  $(T_2, IT_2)$  is equivalent to  $(T_3, IT_3)$ , we can assume from the beginning that  $m < p$  and  $b_k = c_k$  for  $k = 1, \dots, m$ . By induction, we can also assume that  $p = m + 1$ , in which case we can take  $T_1 = R[X_1, \dots, X_m]/(b_1X_1 + \dots + b_mX_m)$  and  $T_2 = R[X_1, \dots, X_m, Z]/(b_1X_1 + \dots + b_mX_m + c_{m+1}Z)$ , where  $X_1, \dots, X_m, Z$  are indeterminates over  $R$ . Write  $c_{m+1} = r_1b_1 + \dots + r_mb_m$  ( $r_i \in R$ ) and let  $X'_k = X_k + r_kZ$  for  $k = 1, \dots, m$ . Then  $X'_1, \dots, X'_m, Z$  are algebraically independent over  $R$  and we have the following  $R$ -algebra isomorphisms

$$T_2 \cong (R[X'_1, \dots, X'_m]/(b_1X'_1 + \dots + b_mX'_m))[Z] \cong T_1[Z].$$

Now let us observe that if  $(T_1, IT_1)$  and  $(T_2, IT_2)$  are two first general grade reductions of  $(R, I)$  and  $m \geq 1$ , then  $\overline{I^m T_1} = \overline{I^m T_2}$  if and only if  $\overline{I^m T_2} = \overline{I^m T_1}$ , or equivalently,  $\overline{I^m T_1}$  is integrally closed if and only if  $\overline{I^m T_2}$  is integrally closed. Indeed, if  $\phi : T_1[Y_1, \dots, Y_r] \xrightarrow{\cong} T_2[Z_1, \dots, Z_s]$  is an  $R$ -algebra isomorphism, then  $\phi(\overline{I^m T_1}[Y_1, \dots, Y_r]) = \overline{I^m T_2}[Z_1, \dots, Z_s]$  and, by Remark 2.1, it follows that  $\overline{I^m T_1}$  is integrally closed if and only if  $\overline{I^m T_2}$  is integrally closed.

This shows that if Theorem 1 holds for some set of generators of  $I$ , then the theorem holds for every set of generators of  $I$ . When proving the main result, this observation allows us to choose the set of generators of  $I$  with some extra properties enabled by the assumption that  $I$  is an  $m$ -primary ideal in a local ring with depth  $R \geq 2$ .

**Lemma 2.3.** *Let  $R$  be a noetherian ring and  $I = (a_1, a_2, \dots, a_n)$  an ideal in  $R$  with  $\text{grade } I > 0$ . Let  $a \in R$  be a non-zero-divisor such that  $\text{grade}(I + aR) \geq 2$  and let  $\alpha = a_1X_1 + \dots + a_nX_n \in S = R[X_1, \dots, X_n]$ . Then  $a, \alpha$  is a permutable regular sequence on  $S$ . In particular, if the elements  $a_1, \dots, a_n$  are non-zero-divisors and  $\text{grade } I \geq 2$ , then  $a_j, \alpha$  is a permutable regular sequence on  $S$  for all  $j$ .*

**Proof.** Both  $a$  and  $\alpha$  are non-zero-divisors on  $S$  (2.2), so it is enough to prove that  $\alpha$  is a non-zero-divisor on  $S/aS$ . Since  $\text{grade}(I + aR) \geq 2$ , there exists  $c = r_1a_1 + \dots + r_na_n \in I$  ( $r_i \in R$ ) such that  $a, c$  is a regular sequence on  $R$ . By applying the  $R$ -algebra automorphism of  $S$  that maps  $X_i$  to  $X_i + r_i$  ( $i = 1, \dots, n$ ), it follows that it is enough to prove that  $a, \alpha + c$  is a regular sequence on  $S$ .

Let  $f \in S$  such that

$$(a_1X_1 + \dots + a_nX_n + c)f \in aS. \tag{2.3.1}$$

We want to prove that  $f \in aS$ . Considering a monomial order on  $S$ , let  $bX_1^{\alpha_1} \dots X_n^{\alpha_n}$  be the smallest term of  $f$ . The coefficient of the smallest term in the left-hand side of (2.3.1) is  $bc$ , so  $bc \in aR$ , and since  $a, c$  is a regular sequence on  $R$ , we obtain  $b \in aR$ . Now

$$(f - bX_1^{\alpha_1} \dots X_n^{\alpha_n})(a_1X_1 + \dots + a_nX_n + c) \in aS,$$

and repeating the argument with  $f - bX_1^{\alpha_1} \dots X_n^{\alpha_n}$  instead of  $f$  we eventually get that all the coefficients of  $f$  belong to  $aR$ , and hence  $f \in aS$ . This proves that  $\alpha + c$  is a non-zero-divisor on  $S/aS$ .  $\square$

Keeping the notation introduced in 2.2 we have the following lemma.

**Lemma 2.4.** *Let  $R$  be a noetherian ring and  $I = (a_1, a_2, \dots, a_n)$  an ideal in  $R$  with  $a_1, a_2, \dots, a_n$  non-zero-divisors ( $n \geq 2$ ) and  $\text{grade } I \geq 2$ . Let  $\alpha = a_1X_1 + \dots + a_nX_n \in S = R[X_1, \dots, X_n]$ . Then for each  $i$  we have the following isomorphisms of  $R$ -algebras*

$$T = S/\alpha S \cong R[X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_n] \left[ \frac{a_1}{a_i}X_1 + \dots + \frac{a_{i-1}}{a_i}X_{i-1} + \frac{a_{i+1}}{a_i}X_{i+1} + \dots + \frac{a_n}{a_i}X_n \right]$$

and

$$T_i := T \left[ \frac{a_1}{a_i}, \dots, \frac{a_n}{a_i} \right] \cong R[X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_n] \left[ \frac{a_1}{a_i}, \dots, \frac{a_n}{a_i} \right].$$

**Proof.** We begin by recalling a well-known result: if  $R$  is a commutative ring and  $b, c$  is a regular sequence on  $R$ , then the  $R$ -algebra homomorphism  $\tau : R[X] \rightarrow R[c/b]$  with  $\tau(X) = -c/b$  induces an  $R$ -algebra isomorphism  $R[X]/(bX + c) \cong R[c/b]$  (see, for example, [14, (7.1)]).

Since  $\alpha S \cap R = (0)$ , we may regard  $R$  as embedded in  $S/\alpha S$ . We may also assume that  $i = n$ . By Lemma 2.3,  $a_1 X_1 + \dots + a_{n-1} X_{n-1}, a_n$  is a permutable regular sequence on  $R[X_1, \dots, X_{n-1}]$  and hence we obtain the  $R$ -algebra isomorphism

$$\phi : S/\alpha S \xrightarrow{\cong} R[X_1, \dots, X_{n-1}] \left[ \frac{a_1}{a_n} X_1 + \dots + \frac{a_{n-1}}{a_n} X_{n-1} \right]$$

which maps  $X_i + \alpha S$  to  $X_i$ , for  $1 \leq i \leq n - 1$ , and  $X_n + \alpha S$  to  $-\frac{a_1}{a_n} X_1 - \dots - \frac{a_{n-1}}{a_n} X_{n-1}$ . Since  $\phi(a_i) = a_i$  for all  $i$ , this induces the isomorphism

$$(S/\alpha S) \left[ \frac{a_1}{a_n}, \dots, \frac{a_{n-1}}{a_n} \right] \cong R[X_1, \dots, X_{n-1}] \left[ \frac{a_1}{a_n} X_1 + \dots + \frac{a_{n-1}}{a_n} X_{n-1} \right] \left[ \frac{a_1}{a_n}, \dots, \frac{a_{n-1}}{a_n} \right],$$

that is,

$$(S/\alpha S) \left[ \frac{a_1}{a_n}, \dots, \frac{a_{n-1}}{a_n} \right] \cong R[X_1, \dots, X_{n-1}] \left[ \frac{a_1}{a_n}, \dots, \frac{a_{n-1}}{a_n} \right]. \quad \square$$

### 2.5. Superficial elements

Let  $R$  be a noetherian ring and  $I$  an ideal in  $R$ . An element  $x \in I$  is said to be a superficial element of  $I$  if there exists  $c \in \mathbb{N}$  such that  $(I^{n+1} : x) \cap I^c = I^n$  for all  $n \geq c$ . Such elements exist, for instance, when the ring  $R$  has infinite residue fields [7, 8.5.7]. Furthermore, superficial elements of  $I$  exist even when we require some extra properties: if  $R$  is a local ring with infinite residue field and  $K_1, \dots, K_m$  are ideals in  $R$  not containing  $I$ , then there exists a superficial element of  $I$  that is not contained in  $K_1 \cup \dots \cup K_m$  [7, 8.5.9].

In particular, if  $R$  is a local ring with infinite residue field and  $I$  is an ideal in  $R$  with grade  $I > 0$ , then there exists  $x \in I$  a non-zero-divisor superficial element of  $I$ . Moreover, there exist elements  $a_1, \dots, a_n$  generating  $I$  that are non-zero-divisors and superficial elements. Indeed, we can choose  $a_1 \in I$  a non-zero-divisor superficial element of  $I$  and then successively take  $a_i \in I$  a superficial element outside of  $(a_1, \dots, a_{i-1})$  that avoids all the associated primes of  $R$ . Since the ring is noetherian, we eventually have  $I = (a_1, \dots, a_n)$  for some  $n$ .

In the same vein, if  $R$  is a local ring with infinite residue field and  $I$  is an ideal in  $R$  with grade  $I \geq 2$ , then we can choose  $a_1, \dots, a_n$  generators for  $I$  such that  $a_i, a_j$  is a regular sequence for all  $i \neq j$  and all the elements  $a_i$  are superficial. As shown above, start by choosing  $a_1 \in I$  a non-zero-divisor superficial element of  $I$  and then take  $a_i \in I$  a superficial element outside of  $(a_1, \dots, a_{i-1})$  that avoids all the associated prime ideals of the ideals  $(a_1), \dots, (a_{i-1})$ . (Since grade  $I \geq 2$ , the ideal  $I$  is not contained in any of those prime ideals.) Eventually we obtain  $I = (a_1, \dots, a_n)$  for some  $n$ . Note that since  $R$  is local, a permutation of a regular sequence is a regular sequence, too.

We mention here that a non-zero-divisor  $x \in I$  is a superficial element of  $I$  if and only if, for all  $n$  sufficiently large,  $(I^{n+1} : x) = I^n$  [7, 8.5.3]. This also implies that for a non-zero-divisor superficial element  $x$  of  $I$  we have  $(I^{n+1} : x) = I^n$  for all  $n$ . Indeed, let  $N$  be such that  $(I^{k+1} : x) = I^k$  for  $k \geq N$  and let  $y \in (I^{N+1} : x)$ . By [7, 6.8.12], there exists  $c$  that avoids all the minimal prime ideals of  $R$  and a

positive  $M$  such that  $cy^t x^t \in I^{(n+1)t}$  for all  $t \geq M$ . If  $t \geq \max\{M, N/n\}$ , we have  $\overline{cy^t} \in (I^{(n+1)t} : x^t) = I^{nt}$ , and by using again the characterization of the integral closure from [7, 6.8.12] we obtain  $y \in \overline{I^m}$ .

If  $I = (a_1, \dots, a_n)$ , by passing to  $S = R[X_1, \dots, X_n]$ , the element  $\alpha = a_1 X_1 + \dots + a_n X_n \in IS$  is sufficiently general in the above sense. More exactly we have the following proposition.

**Proposition 2.6.** *Let  $R$  be a noetherian ring,  $I = (a_1, \dots, a_n)$  an ideal with grade  $I > 0$  and  $\alpha = a_1 X_1 + \dots + a_n X_n \in S = R[X_1, \dots, X_n]$ . Then  $(\overline{I^m} S : \alpha) = \overline{I^{m-1}} S$  for all  $m$  and  $(I^m S : \alpha) = I^{m-1} S$  for  $m \gg 0$ .*

**Proof.** By localizing at the prime ideals that contain  $I$  we may assume that  $R$  is a local ring with maximal ideal  $\mathfrak{m}$ . Moreover, by replacing  $R$  with the faithfully flat extension  $R(Z) := R[Z]_{\mathfrak{m}R[Z]}$ , we may also assume that  $R$  has infinite residue field. Now let us observe that we may assume that  $a_1$  is a non-zero-divisor and a superficial element of  $I$ . Indeed, since grade  $I > 0$ , by a refinement of a prime avoidance argument, it follows that there exist  $r_2, \dots, r_n \in R$  such that  $a'_1 := a_1 + r_2 a_2 + \dots + r_n a_n$  is a non-zero-divisor and a superficial element of  $I$ . Note that  $I = (a'_1, a_2, \dots, a_n)$ . If we consider the  $R$ -algebra automorphism  $\phi : S \rightarrow S$  that maps  $X_1$  to  $X_1$  and  $X_i$  to  $X_i + r_i X_1$  for  $i \geq 2$ , we have  $\phi(\alpha) = a'_1 X_1 + a_2 X_2 + \dots + a_n X_n$ , and therefore we may assume that the first generator  $a_1$  of  $I$  is a non-zero-divisor superficial element of  $I$ . (By repeating the process we can actually make all the generators of  $I$  satisfy this property.)

Let  $f \in (\overline{I^m} S : \alpha)$ . If we consider a monomial order on  $S$  with  $X_1 < \dots < X_n$  and  $bX_1^{\alpha_1} \dots X_n^{\alpha_n}$  is the smallest term that appears in  $f$ , from  $f\alpha \in \overline{I^m} S$  we obtain  $ba_1 \in \overline{I^m}$ , and hence  $b \in (\overline{I^m} : a_1) = \overline{I^{m-1}}$ , where the last equality holds because  $a_1$  is a non-zero-divisor superficial element of  $I$  (2.5). Replacing  $f$  by  $f - bX_1^{\alpha_1} \dots X_n^{\alpha_n}$  and repeating the argument we eventually obtain that all the coefficients of  $f$  are in  $\overline{I^{m-1}}$ . Similarly, if  $f \in (I^m S : \alpha)$ , all the coefficients of  $f$  are in  $(I^m : a_1) = I^{m-1}$  for  $m \gg 0$ .  $\square$

**Remark 2.7.** If  $I$  is an ideal generated by a regular sequence  $a_1, \dots, a_n$ , then  $(I^k : a_i^l) = I^{k-l}$  for all  $i$  and all  $k \geq l$ . This follows, for instance, from the  $R$ -algebra isomorphism between the polynomial ring  $(R/I)[X_1, \dots, X_n]$  and the associated graded ring  $\mathcal{G} = \bigoplus_{n \geq 0} I^n / I^{n+1}$  that sends  $X_j$  to  $a_j + I^2 \in I/I^2$  (see [1, (1.1.8, 1.1.15)]). Moreover, under the same assumptions,  $(\overline{I^k} : a_i^l) = \overline{I^{k-l}}$ . Indeed, if  $ya_i^l \in \overline{I^k}$ , then there exists  $c \in R$  that avoids all the minimal prime ideals such that  $cy^m a_i^{lm} \in I^{km}$  for  $m \gg 0$  (cf. [7, 6.8.12]). Then  $cy^m \in (I^{km} : a_i^{lm}) = I^{(k-l)m}$  for  $m \gg 0$ , thus  $y \in \overline{I^{k-l}}$ .

In the next lemma we use the notation established in Lemma 2.4.

**Lemma 2.8.** *Let  $R$  be a noetherian ring and  $I = (a_1, a_2, \dots, a_n)$  an ideal in  $R$  with  $a_1, \dots, a_n$  non-zero-divisors and superficial elements of  $I$  and grade  $I \geq 2$ . Let  $\alpha = a_1 X_1 + \dots + a_n X_n \in S = R[X_1, \dots, X_n]$ ,  $T = S/\alpha S$  and  $T_i := T[a_1/a_i, \dots, a_n/a_i]$ . Then  $X_j, a_i$  is a permutable regular sequence on  $T_i$  for all  $i, j$ .*

**Proof.** Clearly  $a_i$  is a non-zero-divisor on  $T_i$ , so it is enough to prove that  $X_j, a_i$  is a regular sequence on  $T_i$ . If  $j \neq i$ , by the second isomorphism from Lemma 2.4, it follows that  $X_j$  is a non-zero-divisor on  $T_i$ , and clearly  $a_i$  is a non-zero-divisor on

$$T_i / X_j T_i \cong R[X_1, \dots, X_{j-1}, X_{j+1}, \dots, X_{i-1}, X_{i+1}, \dots, X_n] \left[ \frac{a_1}{a_i}, \dots, \frac{a_n}{a_i} \right].$$

In the case  $i = j$ , without loss of generality we may assume that  $i = j = n$ . The second isomorphism from Lemma 2.4 maps  $X_n$  to  $-\frac{a_1}{a_n} X_1 - \dots - \frac{a_{n-1}}{a_n} X_{n-1}$ , which, as shown in 2.2, is a non-zero-divisor on  $R[\frac{a_1}{a_n}, \dots, \frac{a_{n-1}}{a_n}][X_1, \dots, X_{n-1}]$ . To prove that  $\frac{a_1}{a_n} X_1 + \dots + \frac{a_{n-1}}{a_n} X_{n-1}, a_n$  is a regular sequence on  $R[\frac{a_1}{a_n}, \dots, \frac{a_{n-1}}{a_n}][X_1, \dots, X_{n-1}]$  we apply Lemma 2.3 in the ring  $R[\frac{a_1}{a_n}, \dots, \frac{a_{n-1}}{a_n}]$  with  $a = a_n$  and the ideal  $J = (\frac{a_1}{a_n}, \dots, \frac{a_{n-1}}{a_n})$ . To be able to do this we will show that  $a_n, a_1/a_n$  is a regular sequence on  $R[\frac{a_1}{a_n}, \dots, \frac{a_{n-1}}{a_n}]$  and hence grade  $(J, a_n) \geq 2$ . Since  $a_1$  is superficial element of  $I$  and a non-zero-divisor

on  $R$ , there exists  $n_0$  such that  $I^{n_0+k} : a_1 = I^{n_0+k-1}$  for all  $k \geq 0$ . Assume that  $(a_1/a_n)f = a_n g$  with  $f, g \in R[\frac{a_1}{a_n}, \dots, \frac{a_{n-1}}{a_n}]$ . There exists  $N_0$  such that for  $N \geq N_0$  we have  $f' = f a_n^N \in I^N \subseteq R$  and  $g' = g a_n^N \in I^N \subseteq R$ . Then  $a_1 f' = a_n^2 g'$  and hence, for  $N \geq \max\{n_0 - 2, N_0\}$ , we have  $f' \in (I^{N+2} : a_1) = I^{N+1}$ . Finally,  $f = f'/a_1^N \in IR[\frac{a_1}{a_n}, \dots, \frac{a_{n-1}}{a_n}] = a_n R[\frac{a_1}{a_n}, \dots, \frac{a_{n-1}}{a_n}]$ .  $\square$

The following lemma will play a crucial role in the proof of the main result.

**Lemma 2.9.** *Let  $R$  be a noetherian ring and  $I = (a_1, a_2, \dots, a_n)$  an ideal in  $R$  with  $a_1, \dots, a_n$  non-zero-divisors and superficial elements of  $I$ . Let  $S = R[X_1, X_2, \dots, X_n]$ ,  $\alpha = a_1 X_1 + a_2 X_2 + \dots + a_n X_n \in S$  and  $T = S/\alpha S$ . If  $\text{grade } I \geq 2$ , then  $X_j$  is a non-zero-divisor on  $T/\overline{I^m T}$  for all  $j$  and all  $m \geq 1$ .*

**Proof.** We may assume  $j = 1$  and let  $f \in T$  such that  $f X_1 \in \overline{I^m T}$ . Then there exists  $c \in T$  that avoids all the minimal prime ideals of  $T$  such that  $c f^k X_1^k \in I^{mk} T$  for  $k \gg 0$ . Let  $T_i = T[\frac{a_1}{a_i}, \dots, \frac{a_n}{a_i}]$  ( $i = 1, \dots, n$ ). Then  $c f^k X_1^k \in I^{mk} T_i = a_i^{mk} T_i$  for all  $i$  and  $k \gg 0$ . By Lemma 2.8,  $X_1, a_i$  is a permutable regular sequence on  $T_i$ , and hence  $c f^k \in a_i^{mk} T_i = I^{mk} T_i$  for  $k \gg 0$ . We note here that for an arbitrary ideal  $J$  in  $T$  we have  $\bigcap_{i=1}^n J T_i \cap T = \bigcup_{r=1}^\infty (J I^r T : I^r T)$ . Therefore, for  $k \gg 0$ , we obtain

$$c f^k \in \bigcap_{i=1}^n I^{mk} T_i \cap T = \bigcup_{r=1}^\infty (I^{mk+r} T : I^r T) = \widetilde{I^{mk} T},$$

the Ratliff–Rush closure of  $I^{mk} T$  [15, 2.3.1]. Since  $\widetilde{I^{mk} T} = I^{mk} T$  for  $k \gg 0$  [15, 2.3.2], we have  $c f^k \in I^{mk} T$  for  $k \gg 0$ , or equivalently,  $f \in \overline{I^m T}$ . This finishes the proof that  $X_1$  is a non-zero-divisor on  $T/\overline{I^m T}$ .  $\square$

### 3. The proof of the main result

With the preliminaries in place we are now prepared to prove the main result. The next proposition, a particular case of Theorem 1(a), is essentially due to Itoh. It is proved by Itoh in [10] only for ideals generated by regular sequences. However, by using a result also due to Itoh [9, Lemma 3], it can be proved in the generality stated below.

**Proposition 3.1.** *Let  $R$  be a locally formally equidimensional ring and  $I = (a_1, a_2, \dots, a_n)$  a parameter ideal in  $R$  of height  $n$  with  $\text{grade } I \geq 2$ . Let  $S = R[X_1, \dots, X_n]$ ,  $\alpha = a_1 X_1 + \dots + a_n X_n \in S$  and  $T = S/\alpha S$ . Then  $\overline{IT} = IT$ .*

**Proof.** By localizing at the prime ideals that contain  $I$ , we may assume that the ring  $R$  is local. Let  $A = R[X_1, \dots, X_{n-1}]_{\mathfrak{m}[X_1, \dots, X_{n-1}]}$ , where  $\mathfrak{m}$  is the maximal ideal of  $R$ ,  $J = IA$ ,  $\beta = a_1 X_1 + \dots + a_{n-1} X_{n-1}$  and  $B = A[\beta/a_n]$ . Note that by the discussion 2.2 it is enough to prove the proposition for a special set of generators for  $I$ . Since  $\text{grade } I > 0$ , by a refinement of a prime avoidance argument, there exist  $r_1, \dots, r_{n-1} \in R$  such that  $a'_n := a_n + r_1 a_1 + \dots + r_{n-1} a_{n-1}$  is a non-zero-divisor. Since  $I = (a_1, \dots, a_{n-1}, a'_n)$ , by replacing  $a_n$  with  $a'_n$  we may assume that  $a_n$  is a non-zero-divisor. Moreover, by Lemma 2.3,  $\beta, a_n$  is a permutable regular sequence on  $A$ , and hence, by the result mentioned at the beginning of the proof of Lemma 2.4, we have  $B \cong A[X_n]/\alpha A[X_n]$ . First we claim that  $\overline{J^k B} \cap A = \overline{J^k}$  for every  $k$ . Note that  $J$  is a parameter ideal in  $A$  and  $a_n, \beta$  are part of a minimal set of generators for  $J$ . For  $w \in \overline{J^k B} \cap A$  we have  $w a_n^N \in \overline{J^{k+N}}$  for  $N \gg 0$ , hence  $w \in (\overline{J^{k+N}} : a_n^N) = \overline{J^k}$  (cf. [7, Corollary 6.8.13]).

Next we claim that  $\overline{JB} = \overline{JB}$ . For  $y \in \overline{JB}$ , write  $y = z/a_n^t$  with  $z \in (\beta, a_n)^t \subseteq A$ , so that  $z \in (\beta, a_n)^t \cap (\overline{J^{t+1} B} \cap A) = (\beta, a_n)^t \cap \overline{J^{t+1}}$ . Since  $R$  is locally formally equidimensional, by [9, Lemma 3] it follows that  $(\beta, a_n)^t \cap \overline{J^{t+1}} = (\beta, a_n)^t \overline{J}$ , and therefore  $y \in \overline{JB}$ .

Now let  $f \in S$  with  $\bar{f} = f + \alpha S \in \overline{IS/\alpha S}$ . If  $\phi$  denotes the isomorphism between  $S/\alpha S$  and  $R[X_1, \dots, X_{n-1}][\beta/a_n] \subseteq A[\beta/a_n] = B$  we have  $\phi(\bar{f}) \in \overline{JB} = \bar{J}B = \bar{I}B$ . Then  $f \in \bar{I}A[X_n]$  and hence  $f \in \bar{I}R[X_1, \dots, X_n]_{\mathfrak{m}[X_1, \dots, X_n]} \cap R[X_1, \dots, X_n] = \bar{I}R[X_1, \dots, X_n]$ .  $\square$

**Remark 3.2.** If the ideal  $I$  is generated by a regular sequence, then so is  $J$ , and from [8, Proposition 6] it follows that the equality  $(\beta, a_n)^t \cap \overline{J^{t+1}} = (\beta, a_n)^t \bar{J}$  holds in an arbitrary ring. Therefore in this case the conclusion of Proposition 3.1 is true without assuming that  $R$  is locally formally equidimensional.

We now prove the main result of the paper.

**Theorem 3.3.** *Let  $(R, \mathfrak{m})$  be a formally equidimensional local ring with depth  $R \geq 2$  and  $I = (a_1, a_2, \dots, a_n)$  an  $\mathfrak{m}$ -primary ideal in  $R$ . Let  $S = R[X_1, X_2, \dots, X_n]$ ,  $\alpha = a_1X_1 + a_2X_2 + \dots + a_nX_n \in S$  and  $T = S/\alpha S$ . Then  $\overline{IT} = \bar{I}T$ .*

**Proof.** By replacing  $R$  with the faithfully flat extension  $R(Z) := R[Z]_{\mathfrak{m}R[Z]}$  we may assume that  $R$  is a local ring with infinite residue field.

We have noted in 2.2 that it is enough to prove the theorem for a special set of generators for  $I$ . For a local ring with infinite residue field, one can successively find sufficiently general elements  $x_1, \dots, x_d \in I$  that form a minimal reduction of  $I$  ( $d = \dim R$ ). Therefore we can choose generators  $a_1, \dots, a_n$  for  $I$  such that  $a_1, \dots, a_d$  is a minimal reduction of  $I$  and all the elements  $a_1, \dots, a_n$  are non-zero-divisors and superficial elements of  $I$  (see discussion in 2.5).

Let  $f \in S$  with  $f + \alpha S \in \overline{IT}$ . To prove that  $f \in \overline{IS} = \bar{I}S$ , without loss of generality we may assume that  $f$  is a homogeneous polynomial. Since  $f + \alpha S \in \overline{IT}$ , there exist a positive integer  $k$  and polynomials  $g_i = g_i(X_1, \dots, X_n) \in I^k S$ ,  $h = h(X_1, \dots, X_n) \in S$  such that

$$f^s + g_1 f^{s-1} + \dots + g_s = (a_1 X_1 + \dots + a_n X_n) h. \tag{3.3.1}$$

We will proceed by induction on the degree of  $f$ . If the degree of  $f$  is zero, i.e.,  $f \in R$ , by evaluating (3.3.1) at  $X_1 = \dots = X_n = 0$  we obtain an equation of integral dependence of  $f$  over  $I$ , so  $f \in \bar{I} \subseteq \overline{IS}$ . Now assume that (3.3.1) implies  $f \in \overline{IS}$  for any homogeneous polynomial  $f \in S$  of degree at most  $k - 1$  ( $k \geq 1$ ) and let  $f$  be a homogeneous polynomial of degree  $k$  that satisfies (3.3.1). We will show that  $f \in \overline{IS}$  using again induction, but on  $e := n - d$ . If  $n = d$ , then  $I$  is a parameter ideal and the conclusion follows from Proposition 3.1. Now assume that  $e \geq 1$ . By evaluating (3.3.1) at  $X_n = 0$  we obtain an equality that shows that the coset of  $f(X_1, \dots, X_{n-1}, 0)$  belongs to  $\overline{IT'}$  where  $T' = R[X_1, \dots, X_{n-1}]/(a_1 X_1 + \dots + a_{n-1} X_{n-1})$ . Since  $\overline{IT'} = \overline{(a_1, \dots, a_{n-1})T'}$  and  $f(X_1, \dots, X_{n-1}, 0)$  is either zero or a homogeneous polynomial of degree  $k$ , by the induction hypothesis we have  $f(X_1, \dots, X_{n-1}, 0) \in \overline{IR[X_1, \dots, X_{n-1}]} \subseteq \overline{IS}$ . On the other hand,

$$f(X_1, \dots, X_n) = f(X_1, X_2, \dots, X_{n-1}, 0) + X_n f_1(X_1, \dots, X_n) \tag{3.3.2}$$

for some homogeneous polynomial  $f_1 = f_1(X_1, \dots, X_n) \in S$  of degree  $k - 1$ . Since both  $f + \alpha S$  and  $f(X_1, X_2, \dots, X_{n-1}, 0) + \alpha S$  belong to  $\overline{IT}$ , we have  $X_n f_1 + \alpha S \in \overline{IT}$ , and by Lemma 2.9 we obtain  $f_1 + \alpha S \in \overline{IT}$ . Since the degree of  $f_1$  is  $k - 1$ , by the induction hypothesis we have  $f_1 \in \overline{IS}$ , and (3.3.2) implies that  $f \in \overline{IS}$ .  $\square$

By localizing at the maximal ideal  $\mathfrak{m}[X_1, \dots, X_n]$  of  $S$  we also obtain the following local version.

**Corollary 3.4.** *Let  $(R, \mathfrak{m})$  be a formally equidimensional local ring with depth  $R \geq 2$  and  $I = (a_1, a_2, \dots, a_n)$  an  $\mathfrak{m}$ -primary ideal in  $R$ . Let  $U = R[X_1, X_2, \dots, X_n]_{\mathfrak{m}[X_1, \dots, X_n]}$ ,  $\alpha = a_1 X_1 + a_2 X_2 + \dots + a_n X_n \in U$  and  $V = U/\alpha U$ . Then  $\overline{IV} = \bar{I}V$ .*

**Remark 3.5.** Assume that  $I$  is an ideal generated by a regular sequence  $a_1, \dots, a_n$ . Itoh [8] and Huneke [6] (in the case of rings containing a field) proved that  $\overline{I^{m+1}} \cap I^m = \overline{I}I^m$  for all  $m$ . Based on this, in a subsequent paper [10], Itoh proved that  $\overline{IT} = \overline{I}T$ . Let us observe the statement  $\overline{IT} = \overline{I}T$  can be used to recover the equality  $\overline{I^{m+1}} \cap I^m = \overline{I}I^m$  for all  $m$ .

We will use induction on  $m + n$ . For  $m = 0$  there is nothing to prove. For  $n = 1$ , let  $I = (a)$ . If  $ra^m \in (\overline{a^{m+1}}) \cap (a^m)$ , then there exists  $c \in R$  avoiding all the minimal prime ideals of  $R$  such that  $cr^s a^{ms} \in (a^{ms+s})$  for  $s \gg 0$ . Then  $cr^s \in (a^s)$  for  $s \gg 0$ , and hence  $r \in (\overline{a})$ . For  $n \geq 2$ , as in Corollary 3.4, let  $V = U/\alpha U$  where  $U = R[X_1, X_2, \dots, X_n]_{\mathfrak{m}[X_1, \dots, X_n]}$  and  $\alpha = a_1 X_1 + a_2 X_2 + \dots + a_n X_n \in U$ . Since  $\alpha$  is a non-zero-divisor on  $U$  (2.2),  $\text{grade}(IV) = \text{grade } I - 1 = n - 1$ . Also,  $IV$  can be generated by  $n - 1$  elements  $(\alpha \notin \text{Im } U)$ , and therefore  $IV$  is a complete intersection ideal. Let  $y \in \overline{I^{m+1}} \cap I^m$ . Then  $y + \alpha U \in \overline{I^{m+1}}V \cap I^m V$  and by the induction hypothesis we obtain  $y + \alpha U \in I^m \overline{I}V$ . Applying Corollary 3.4 we obtain  $y \in \overline{I^m I}U + \alpha U$ , so that we can write  $y = z + u\alpha$  where  $z \in I^m \overline{I}U$  and  $u \in U$ . Then  $u\alpha = y - z \in I^m U \cap \overline{I^{m+1}}U$  and using Remark 2.7 we obtain  $u \in (I^m U : \alpha) \cap (\overline{I^{m+1}}U : \alpha) = I^{m-1}U \cap \overline{I^m}U = (I^{m-1} \cap \overline{I^m})U$ . By the induction hypothesis we then obtain  $u \in I^{m-1} \overline{I}U$ , so  $y \in I^m \overline{I}U \cap R = I^m \overline{I}$ .

For an analytically unramified Cohen–Macaulay local ring  $(R, \mathfrak{m})$  of dimension  $d \geq 2$  and an ideal  $I$  generated by a maximal regular sequence, Itoh [10, Theorem 1(3)] also proved that  $\overline{I^m}V = \overline{I^m}V$  for  $m \gg 0$ . Note that this implies that  $\overline{I^m T} = \overline{I^m}T$  for  $m \gg 0$ . Indeed, all the zero-divisors of the  $S$ -module  $S/(\overline{I^m}, \alpha)$  are contained in  $\mathfrak{m}[X_1, \dots, X_n]$ , and hence  $\overline{I^m}V \cap T = \overline{I^m}T$ , which implies that  $\overline{I^m T} \subseteq \overline{I^m}V \cap T = \overline{I^m}T$ .

We will extend this result to arbitrary  $\mathfrak{m}$ -primary ideals.

**Theorem 3.6.** Let  $(R, \mathfrak{m})$  be an analytically unramified Cohen–Macaulay local ring of dimension  $d \geq 2$  and  $I = (a_1, a_2, \dots, a_n)$  an  $\mathfrak{m}$ -primary ideal in  $R$ . Let  $S = R[X_1, X_2, \dots, X_n]$ ,  $\alpha = a_1 X_1 + a_2 X_2 + \dots + a_n X_n \in S$  and  $T = S/\alpha S$ . Then  $\overline{I^m T} = \overline{I^m}T$  for  $m \gg 0$ .

**Proof.** We begin the proof as in Theorem 3.3. We may assume that  $R$  has infinite residue field and since it is enough to prove the theorem for some set of generators for  $I$  (2.2), choose  $a_1, \dots, a_n$  generators for  $I$  such that  $(a_1, \dots, a_d)$  is a minimal reduction of  $I$  and all the elements  $a_i$  are non-zero-divisors and superficial elements of  $I$  (see discussion in 2.5).

Given an element  $f \in S$  with  $f + \alpha S \in \overline{I^m T}$ , we obtain an equation

$$f^s + g_1 f^{s-1} + \dots + g_s = (a_1 X_1 + \dots + a_n X_n)h \tag{3.6.1}$$

where  $g_i = g_i(X_1, \dots, X_n) \in I^{mi}S$  and  $h = h(X_1, \dots, X_n) \in S$ . We want to prove that for  $m \gg 0$  this implies that  $f \in \overline{I^m}S + \alpha S$ . Note that we may assume that  $f$  is homogeneous. We will use induction on the degree of  $f$ . If  $\text{deg } f = 0$ , by setting  $X_1 = \dots = X_n = 0$  in (3.6.1) we obtain  $f \in \overline{I^m}$ . Now assume that the claim is true for every homogeneous polynomial of degree at most  $k - 1$  and let  $f$  be a homogeneous polynomial of degree  $k$  that satisfies (3.6.1). We use again induction on  $e := n - d$ . If  $n = d$ , the conclusion follows from the result of Itoh that we mentioned in the discussion preceding this theorem. Assume that  $e \geq 1$ . By setting  $X_n = 0$  in (3.6.1), the coset of  $f(X_1, \dots, X_{n-1}, 0)$  is an element of  $\overline{I^m T'}$  where  $T' = R[X_1, \dots, X_{n-1}]/(a_1 X_1 + \dots + a_{n-1} X_{n-1})$ . Since  $\overline{I^m T'} = (\overline{a_1}, \dots, \overline{a_{n-1}})^{\overline{I^m T'}}$  and  $f(X_1, \dots, X_{n-1}, 0)$  is either zero or a homogeneous polynomial of degree  $k$  in  $n - 1$  variables, by the induction hypothesis we have  $f(X_1, \dots, X_{n-1}, 0) \in \overline{I^m}R[X_1, \dots, X_{n-1}] + (a_1 X_1 + \dots + a_{n-1} X_{n-1})R[X_1, \dots, X_{n-1}]$  so that we can write

$$f(X_1, \dots, X_{n-1}, 0) = p + (a_1 X_1 + \dots + a_{n-1} X_{n-1})q$$

where  $p \in \overline{I^m}S$  is a homogeneous polynomial of degree  $k$  and  $q \in S$  is a homogeneous polynomial of degree  $k - 1$ . On the other hand,



$$f(X_1, \dots, X_n) = f(X_1, \dots, X_{n-1}, 0) + X_n f_1(X_1, \dots, X_n)$$

where  $f_1 \in S$  is a homogeneous polynomial of degree  $k - 1$ , and therefore we have

$$f(X_1, \dots, X_n) = p + (a_1 X_1 + \dots + a_{n-1} X_{n-1} + a_n X_n)q + X_n(f_1 - a_n q). \tag{3.6.2}$$

Since  $f + \alpha S \in \overline{I^m T}$  and  $p \in \overline{I^m S}$  we have  $X_n(f_1 - a_n q) + \alpha S \in \overline{I^m T}$ , and by Lemma 2.9 we obtain  $(f_1 - a_n q) + \alpha S \in \overline{I^m T}$ . Since the polynomial  $(f_1 - a_n q)$  is homogeneous of degree at most  $k - 1$ , by the induction hypothesis we have  $(f_1 - a_n q) \in \overline{I^m S} + \alpha S$ , and by (3.6.2) we obtain  $f \in \overline{I^m S} + \alpha S$ , which finishes the proof.  $\square$

**Corollary 3.7.** *Let  $(R, \mathfrak{m})$  be an analytically unramified Cohen–Macaulay local ring of dimension  $d \geq 2$  and let  $I = (a_1, a_2, \dots, a_n)$  be an  $\mathfrak{m}$ -primary ideal in  $R$ . Denote  $U = R[X_1, X_2, \dots, X_n]_{\mathfrak{m}[X_1, \dots, X_n]}$ ,  $\alpha = a_1 X_1 + a_2 X_2 + \dots + a_n X_n \in U$  and  $V = U/\alpha U$ . Then  $\overline{I^m V} = \overline{I^m} V$  for  $m \gg 0$ .*

**Remark 3.8.** Theorems 3.3 and 3.6 and their corollaries can be extended to equimultiple ideals. (In general, the analytic spread  $\ell(I)$  of the ideal  $I$  is at most the height of  $I$ ; when the equality holds, we say that  $I$  is equimultiple.) More precisely, if  $R$  is a locally formally equidimensional ring and  $I$  is an equimultiple ideal with grade  $I \geq 2$ , then  $\overline{I T} = \overline{I} T$  and  $\overline{I^m T} = \overline{I^m} T$  for  $m \gg 0$ . Indeed, if  $I$  is equimultiple, then the ideals  $\overline{I^m}$  ( $m \geq 1$ ) have no embedded components [11] and, by localizing at the minimal prime ideals of  $I$ , the conclusions follow.

### 4. Applications

As mentioned in the introduction, the results that show that the integral closure is preserved modulo a generic element are useful for proving integral closure statements by induction. As an example of such an application, we extend some results obtained by the author in [2].

#### 4.1. Almost complete intersection ideals in regular local rings

Let  $(A, \mathfrak{m})$  be a regular local ring of dimension  $d$  and  $I$  an  $\mathfrak{m}$ -primary almost complete intersection ideal, that is, minimally generated by  $d + 1$  elements. Under the assumption that  $A$  contains a field, the author [2, 3.3, 3.5, 3.7] proved that if  $I$  is integrally closed, then  $I$  contains  $d - 2$  regular parameters  $x_1, \dots, x_{d-2}$  such that in the 2-dimensional regular local ring  $A' = A/(x_1, \dots, x_{d-2})$  the ideal  $IA'$  is integrally closed and generated by three elements. The structure of such ideals in 2-dimensional regular local rings is then completely characterized by results of Noh [13]. In addition, we proved that the Rees algebra  $\mathcal{R} = \bigoplus_{n \geq 0} I^n t^n$  is a Cohen–Macaulay normal domain and the associated graded ring  $\mathcal{G} = \bigoplus_{n \geq 0} I^n / I^{n+1}$  is a Cohen–Macaulay ring with  $a(\mathcal{G}) = 1 - d$ , where  $a(\mathcal{G})$ , the  $a$ -invariant of  $\mathcal{G}$ , is defined by  $a(\mathcal{G}) = \sup\{i \mid H_d^M(\mathcal{G})_i \neq 0\}$  with  $M$  being the maximal homogeneous ideal of  $\mathcal{G}$ .

The assumption that  $A$  contains a field was needed because the proofs relied upon the following lemma: if  $A$  is a regular local ring containing a field,  $I$  is an ideal in  $A$  with  $I \not\subseteq \mathfrak{m}^2$  and  $x \in I \setminus \mathfrak{m}^2$ , then  $\overline{(I/xA)} = \overline{I}/xA$  [2, 3.4]. Using Corollary 3.4, we will prove this lemma without assuming that  $A$  contains a field, and hence all the results about integrally closed  $\mathfrak{m}$ -primary almost complete intersection ideals that we mentioned above hold in an arbitrary regular local ring.

**Lemma 4.2.** *Let  $A$  be a regular local ring,  $I$  an ideal in  $A$  with  $I \not\subseteq \mathfrak{m}^2$  and let  $x \in I \setminus \mathfrak{m}^2$ . Then  $\overline{(I/xA)} = \overline{I}/xA$ .*

**Proof.** It is enough to prove this result for  $\mathfrak{m}$ -primary ideals  $I$ . Indeed, for  $f + xA \in \overline{(I/xA)}$  we have  $f + xA \in \overline{(I + \mathfrak{m}^k)/xA}$  for all  $k$ , and hence  $f \in \overline{I + \mathfrak{m}^k}$  for all  $k$ . Since  $\bigcap_k \overline{I + \mathfrak{m}^k} = \overline{I}$  [7, 6.8.5], we obtain  $f \in \overline{I}$ .

Let  $I = (a_1, \dots, a_n)$  and  $\alpha = a_1 X_1 + \dots + a_n X_n \in U = A[X_1, \dots, X_n]_{\mathfrak{m}[X_1, \dots, X_n]}$ . Set  $V = U/\alpha U$ , denote by  $M$  the maximal ideal of  $V$ , and let  $\phi: A/xA \rightarrow V/xV$  be the homomorphism induced by the embedding  $A \rightarrow V$ .

We will use induction on the dimension of  $A$ . If  $\dim A = 1$ , the statement is clear. Assume that  $\dim A \geq 2$ . Since  $\alpha \notin \mathfrak{m}^2 U$ ,  $V$  is a regular local ring of dimension one less than the dimension of  $A$ . Also, note that  $x \in IV \setminus M^2$ . By the induction hypothesis, if  $y \in A$  with  $\bar{y} \in \overline{IV/xA}$ , then  $\phi(\bar{y}) \in \overline{IV/xV} = \overline{IV}/xV$ . Furthermore, by Corollary 3.4, we have  $\phi(\bar{y}) \in \overline{IV}/xV \cong \overline{IU}/(xU + \alpha U)$ , which implies that  $y \in \overline{IU}$ , and hence  $y \in \overline{IU} \cap A = \overline{I}$ .  $\square$

The following consequence was also noted in [2, 3.5], but now we can drop the assumption that  $A$  contains a field. The same proof from [2] will work.

**Corollary 4.3.** *Let  $(A, \mathfrak{m})$  be a  $d$ -dimensional regular local ring, and let  $I$  be an ideal of  $A$  such that the embedding dimension of  $A/I$  is at most two. Then  $\overline{I}^n = I^{n-1}\overline{I}$  for all  $n \geq 1$ .*

**Remark 4.4.** The above corollary also shows that an integrally closed  $\mathfrak{m}$ -primary complete intersection ideal in a regular local ring is normal. This is a well-known result proved by Goto [3, Theorem 3.1] by using different methods.

## References

- [1] W. Bruns, J. Herzog, *Cohen–Macaulay Rings*, Cambridge Stud. Adv. Math., vol. 39, Cambridge University Press, Cambridge, 1993.
- [2] C. Ciupercă, Integrally closed almost complete intersection ideals, *J. Algebra* 302 (2006) 720–728.
- [3] S. Goto, Integral closedness of complete intersection ideals, *J. Algebra* 108 (1987) 151–160.
- [4] M. Hochster, Properties of Noetherian rings stable under general grade reduction, *Arch. Math. (Basel)* 24 (1973) 393–396.
- [5] J. Hong, B. Ulrich, Specialization and integral closure, arXiv:math/0611773v1 [math.AC].
- [6] C. Huneke, Hilbert functions and symbolic powers, *Michigan Math. J.* 34 (1987) 293–318.
- [7] C. Huneke, I. Swanson, *Integral Closure of Ideals, Rings, and Modules*, London Math. Soc. Lecture Note Ser., vol. 336, Cambridge University Press, Cambridge, 2006.
- [8] S. Itoh, Integral closures of ideals generated by regular sequences, *J. Algebra* 117 (1988) 390–401.
- [9] S. Itoh, Integral closures of ideals of the principal class, *Hiroshima Math. J.* 17 (1987) 373–375.
- [10] S. Itoh, Coefficients of normal Hilbert polynomials, *J. Algebra* 150 (1992) 101–117.
- [11] S. McAdam, Asymptotic prime divisors and analytic spreads, *Proc. Amer. Math. Soc.* 80 (1980) 555–559.
- [12] M. Nagata, *Local Rings*, Intersci. Tracts in Pure Appl. Math., vol. 13, Interscience Publishers, 1962.
- [13] S. Noh, Valuation ideals of order two in 2-dimensional regular local rings, *Math. Nachr.* 261/262 (2003) 123–140.
- [14] L.J. Ratliff Jr., Conditions for  $\text{Ker}(R[X] \rightarrow R[c/b])$  to have a linear base, *Proc. Amer. Math. Soc.* 39 (1973) 509–514.
- [15] L.J. Ratliff Jr., D.E. Rush, Two notes on reductions of ideals, *Indiana Univ. Math. J.* 27 (1978) 929–934.