

ORIGINAL RESEARCH

Impact of Degree of Commissural Opening After Percutaneous Mitral Commissurotomy on Long-Term Outcome

David Messika-Zeitoun, MD,*† Julie Blanc, MD,* Bernard Iung, MD,* Eric Brochet, MD,* Bertrand Cormier, MD,* Dominique Himbert, MD,* Alec Vahanian, MD*

Paris, France

OBJECTIVES We sought to evaluate the prognostic value of the degree of commissural opening (CO) on outcome.

BACKGROUND Commissural opening is the main mechanism by which the mitral valve area (MVA) increases after percutaneous mitral commissurotomy (PMC) but its impact on long-term outcome has never been evaluated.

METHODS Of 1,024 consecutive patients with severe MS who underwent PMC, degree of CO was prospectively evaluated in 875 patients (age 48 ± 13 years, female 83%, New York Heart Association (NYHA) functional class III/IV 75%) with good immediate PMC results (MVA ≥ 1.5 cm² and no mitral regurgitation $>2/4$). These 875 patients were divided into 3 groups: both commissures only partially opened or not split (Group 1, n = 189), 1 commissure completely split (Group 2; n = 459), and both commissures completely split (Group 3; n = 227). During a follow-up of 55 ± 28 months, following clinical end points were collected: death, cardiovascular death, need for mitral valve surgery or repeat dilation, and NYHA functional class.

RESULTS Before PMC, patients in Group 1 were older, more often in NYHA functional class III/IV, but MVA and mean gradient were not different ($p \geq 0.50$). Immediately after PMC, there were significant differences between groups with regard to mean gradient (Group 1, 5.1 ± 2.1 mm Hg; Group 2, 4.5 ± 1.7 mm Hg; Group 3, 4.0 ± 1.6 mm Hg; $p < 0.0001$) and MVA (Group 1, 1.8 ± 0.2 cm²; Group 2, 1.9 ± 0.2 cm²; Group 3, 2.1 ± 0.3 cm²; $p < 0.0001$). The 10-year rate of good functional results (survival without need for mitral surgery or repeat dilation and NYHA functional class I or II at last follow-up) was significantly higher in Group 3 ($76 \pm 5\%$) than in Groups 1 and 2 ($39 \pm 8\%$ and $57 \pm 11\%$, respectively; $p < 0.0001$). In multivariable analysis, either the degree of CO or the MVA was an independent predictor of good late functional results ($p < 0.05$).

CONCLUSIONS Complete CO is associated with larger MVA, smaller gradients, and functional improvement. The degree of CO provides important prognostic information and thus should be systematically evaluated during and after PMC and considered as a complementary measure of the procedural success in addition to the MVA, not always easy to assess. (J Am Coll Cardiol Img 2009;2:1-7) © 2009 by the American College of Cardiology Foundation

From the *AP-HP, Cardiology Department, Bichat Hospital and †INSERM U698, University Paris 7, Paris, France. Dr. Messika-Zeitoun was supported by a contrat d'interface INSERM U698.

Manuscript received June 6, 2008; revised manuscript received October 14, 2008, accepted October 16, 2008.

Since its development by Inoue in 1989, percutaneous mitral commissurotomy (PMC) has become an efficient and frequently used procedure to treat mitral stenosis (MS). Its safety and immediate efficacy have been widely demonstrated (1), and PMC is considered as the first-line treatment for symptomatic patients with a favorable anatomy (2,3).

See page 8

Commissural splitting is the main mechanism by which the mitral valve area (MVA) increases after PMC. Due to the heterogeneity of mitral valve alterations (degree of commissural fusion, valve fibrosis, and leaflet calcification), the anatomy of the mitral orifice after PMC is not uniform and widely varies from 1 patient to another with various degrees of commissural opening (CO): 1 or both commissures; complete, partial, or no commissural opening (4). In small pioneering series, CO has been associated with larger MVA (5-7), better leaflet mobility, and a better outcome (4). However, the impact of the degree of CO on long-term outcome has never been evaluated in a large population. We hypothesized that it may provide important prognostic information.

After a successful PMC, defined as an MVA ≥ 1.5 cm² or ≥ 1 cm²/m² with no regurgitation $>2/4$, we identified 7 predictive factors of poor late functional results in 912 patients. Four are pre-procedural characteristics (advanced age, high New York Heart Association [NYHA] functional class, less favorable mitral valve anatomy, and atrial fibrillation [AF]), and 3 are related to the immediate results (low post-procedure MVA, high post-procedure mean transmitral gradient, and grade 2 post-procedure mitral regurgitation [MR]) (8). The aim of the present study was to evaluate, in this population, the prognostic value of the degree of CO.

METHODS

Study population. From March 1986 to March 1995, 1,024 consecutive patients with severe MS (valve area <1.5 cm²) residing in France underwent a PMC in our department. Contraindications to PMC were severe calcification of both commissures, left atrial thrombus on transesophageal echo-

cardiography, and mitral regurgitation of grade >2 . A good immediate result (MVA ≥ 1.5 cm² or ≥ 1 cm²/m² with no regurgitation $>2/4$) was obtained in 912 patients. The degree of CO was prospectively recorded in 875 (96%) patients. These 875 patients constituted our study population.

Procedure. All procedures were performed by the anterograde trans-septal approach. A double balloon was used in 541 cases, and the Inoue balloon in the 334 cases (after October 1990), according to the stepwise technique, under echocardiographic guidance.

Echocardiographic measurements. Echocardiography was performed the day before and 24 to 48 hours after the procedure by experienced operators. MVA was measured by 2-dimensional echocardiography in parasternal short-axis view (planimetry). Mitral valve anatomy was classified into 3 classes according to transthoracic echocardiography and fluoroscopy, as previously described (1): flexible valves and mild subvalvular disease (chordae ≥ 10 mm long), flexible valves and extensive subvalvular disease (chordae <10 mm long), and calcified valves confirmed by fluoroscopy. Of note, in a subset of 40 patients, the mean \pm SD (range) the Wilkins (9) score was 8.0 ± 0.8 (7 to 9) for echocardiographic Class 1, 9.9 ± 1.3 (8 to 12) for Class 2, and 12.5 ± 1.3 (10 to 15) for Class 3. Mean transmitral gradient was assessed by continuous-wave Doppler. Measurement of the systolic pulmonary artery pressure was based on the maximal velocity of the tricuspid regurgitation. Mitral regurgitation was semiquantitatively graded from 0 to 4 (10). The degree of CO was prospectively and semiquantitatively evaluated as none, partial (up from only several millimeters from the valve orifice), or complete (up to the level of the mitral annulus) in a parasternal short-axis view from a detailed echocardiographic examination with multiple scanning of the entire mitral valve apparatus from the left ventricle to the left atrium. A complete CO was defined as a complete disruption of the mitral valve co-optation line at the level of the annulus in diastole; a partial CO was defined as a partial disruption with a clear co-optation point between the anterior and posterior leaflets (Fig. 1). Patients were divided into 3 groups: Group 1 if both commissures were either not split or only partially open, Group 2 if 1 commissure was completely split, and Group 3 if both commissures were completely split (Fig. 1).

Follow-up. Follow-up was based on visits to the department or on a standardized questionnaire sent to

ABBREVIATIONS AND ACRONYMS

| | |
|-------------|--------------------------------------|
| AF | = atrial fibrillation |
| CO | = commissure opening |
| MR | = mitral regurgitation |
| MS | = mitral stenosis |
| MVA | = mitral valve area |
| NYHA | = New York Heart Association |
| PMC | = percutaneous mitral commissurotomy |




Figure 1. Mitral Orifice Before Commissurotomy and Commissural Openings

(A) Examples of mitral orifice before commissurotomy and commissural openings in the 3 groups. (B) In Group 1, both commissures were either not split or only partially open. (C) In Group 2, 1 commissure is completely open (arrow). (D) In Group 3, both commissures are completely open (arrows).

the patient's cardiologist. The following clinical end-points were collected: death, cardiovascular death, need for mitral surgery or repeat dilation, and NYHA functional class. A good functional result (composite end point) was defined as survival considering only cardiovascular death with no need for mitral surgery or repeat dilation and a patient in NYHA functional class I or II at most recent follow-up.

Statistical analysis. Results were expressed as mean \pm SD or percentage. Group comparisons were performed with analysis of variance, and the chi-square or *t* test as appropriate. Cumulative survival curves were determined for the clinical events and the composite end point of good functional results according to the Kaplan-Meier method. Survival status was censored at the time of surgery or repeat dilation. The log-rank test was used for comparison of rates of long-term functional results among the 3 groups of patients. A Cox proportional hazards analysis was performed to evaluate the predictive value of the degree of CO after adjustment for the 7 previously identified predictive factors of long-term functional results (age, NYHA functional class, mitral valve anatomy class, rhythm, post-procedure MVA, post-procedure MR grade, and post-procedure mean gradient). The multivariable analysis was also performed including all pre- and post-procedural variables. When included in the multivariable analysis, continuous variables were divided into subgroups with the same clinically

chosen cutoff points that have been used in a previous analysis of this series (8).

Agreement between operators and for the same operator (interobserver and intraobserver variability) of the degree of CO assessment was evaluated by the kappa value. Analyses were performed using SAS statistical software (SAS Institute, Cary, North Carolina). A *p* value <0.05 was considered significant.

RESULTS

Baseline characteristics. The mean age of the 875 patients was 48 ± 13 years (range 16 to 86 years) and 726 were females (83%). Approximately one-third were in AF. Most of the patients were in NYHA functional class III/IV. A previous commissurotomy had been performed in 130 (15%) patients, a mean of 15 ± 8 years earlier (surgical commissurotomy in 120 patients and PMC in 10 patients). In all, 130 (15%) patients had flexible valve and mild subvalvular disease (chordae ≥ 10 mm long), 513 (59%) patients had flexible valve and extensive subvalvular disease (chordae <10 mm long), and 232 (26%) patients had calcified mitral valve confirmed by fluoroscopy. Baseline characteristics of these 875 patients are summarized in Table 1.

| Table 1. Clinical and Echocardiographic Characteristics of the Study Population Before and After PMC | | | | | |
|---|------------------------------|------------------------------|------------------------------|------------------------------|-----------------------------------|
| | Overall (n = 875) | Group 1 (n = 189) | Group 2 (n = 459) | Group 3 (n = 227) | p Value Between Groups |
| Clinical variables | | | | | |
| Age, yrs | 48 ± 13 | 53 ± 13 | 47 ± 13 | 47 ± 13 | <0.0001 |
| Female | 726 (83%) | 161 (85%) | 382 (83%) | 183 (81%) | 0.46 |
| NYHA functional class III/IV | 657 (75%) | 155 (82%) | 333 (72%) | 169 (75%) | 0.04 |
| Sinus rhythm | 543 (62%) | 105 (56%) | 291 (63%) | 147 (65%) | 0.11 |
| Previous commissurotomy | 130 (15%) | 30 (16%) | 64 (14%) | 36 (16%) | 0.73 |
| Cardiothoracic index | 51 ± 5 | 52 ± 5 | 51 ± 5 | 50 ± 5 | 0.0009 |
| Echocardiographic variables | | | | | |
| Before PMC | | | | | |
| Mitral valve area, cm ² | 1.08 ± 0.22 | 1.09 ± 0.22 | 1.07 ± 0.22 | 1.08 ± 0.23 | 0.55 |
| Mean gradient, mm Hg | 10.1 ± 4.5 | 9.9 ± 4.2 | 10.2 ± 4.7 | 10.1 ± 4.1 | 0.68 |
| Mitral regurgitation grade 1 or 2 | 311 (36%) | 78 (41%) | 162 (35%) | 71 (31%) | 0.10 |
| After PMC | | | | | |
| Mitral valve area, cm ² | 1.94 ± 0.27 | 1.77 ± 0.19 | 1.93 ± 0.25 | 2.10 ± 0.27 | <0.0001 |
| Mean gradient, mm Hg | 4.5 ± 1.8 | 5.1 ± 2.1 | 4.5 ± 1.7 | 4.0 ± 1.6 | <0.0001 |
| Mitral regurgitation grade 2 | 184 (21%) | 42 (22%) | 111 (24%) | 31 (14%) | 0.0006 |

Data presented are number of patients (%) or mean ± SD.
NYHA = New York Heart Association; PMC = percutaneous mitral commissurotomy.

Immediate results. After PMC, the mean MVA increased from 1.08 ± 0.22 cm² to 1.94 ± 0.27 cm², and the mean transmitral gradient decreased from 10.1 ± 4.5 mm Hg to 4.5 ± 1.8 mm Hg (both $p < 0.0001$). Both commissures were either only partially open or not split in 189 (22%) patients (Group 1), 1 commissure was completely split in 459 (52%) patients (Group 2), and both commissures were completely split in 227 (26%) patients (Group 3). Intraobserver and interobserver variability was assessed in 20 patients. Agreement was good, with a kappa value of 0.74 and 0.78, respectively. Of note, most of the discrepancies observed were between partial and not opened commissures.

Patients in Group 1 were older and more often in NYHA functional class III/IV, but pre-procedure mean gradient and MVA were not different between groups. In contrast, there were significant differences with regard to post-procedure mean transmitral gradient (5.1 ± 2.1 mm Hg in Group 1, 4.5 ± 1.7 mm Hg in Group 2, and 4.0 ± 1.6 mm Hg in Group 3; $p < 0.0001$) and MVA (1.77 ± 0.19 cm² in Group 1, 1.93 ± 0.25 cm² in Group 2, and 2.10 ± 0.27 cm² in Group 3; $p < 0.0001$) (Table 1, Fig. 2). Thus, mean transmitral gradient decreased and MVA increased with the degree of CO.

Impact of commissural splitting on late functional results. Follow-up was complete in all patients. Mean follow-up duration was 55 ± 28 months




Figure 2. Differences in Post-Procedure Mean Mitral Gradient and Mitral Valve Area Among the 3 Groups

Post-procedure mean mitral gradient and mitral valve area according to groups as defined in Figure 1. Mitral valve area progressively increased and mean gradient progressively decreased with the number and the extent of commissural openings.

(range 1 to 132 months). During follow-up, 40 deaths (20 cardiovascular deaths) were observed, 102 patients underwent repeat PMC or mitral valve surgery, and 142 patients had worsening symptoms. In the entire population, 5- and 10-year event rates were, respectively, $95 \pm 1\%$ and $88\% \pm 4\%$ for global survival (40 deaths), $97 \pm 1\%$ and $95 \pm 2\%$ for survival considering only cardiovascular-related death (20 deaths), $86 \pm 1\%$ and $68 \pm 5\%$ for survival with no need for surgery or repeat dilation (122 events), and $81 \pm 2\%$ and $62 \pm 5\%$ for good functional results (survival considering only cardiovascular death with no need for mitral surgery or repeat dilation and the patient in NYHA functional class I or II at most recent follow-up; 162 events). Rates for good functional results were significantly different among the 3 groups at 10 years (5 years): $39 \pm 8\%$ ($73 \pm 4\%$) in Group 1, $57 \pm 11\%$ ($83 \pm 2\%$) in Group 2, and $76 \pm 5\%$ ($87 \pm 3\%$) in Group 3 ($p < 0.0001$) (Fig. 3).

In multivariable analysis using a Cox model and the 7 previously identified predictive factors of poor late functional results (age, NYHA functional class, mitral valve anatomy, AF, post-procedure MVA, post-procedure mean transmitral gradient, and post-procedure MR grade), either the degree of CO or the MVA was an independent predictor of good late functional results ($p < 0.05$). Of note, similar results were observed when the multivariable analysis was performed including all pre- and post-procedure variables in the model.

DISCUSSION

In this large series of patients with MS, we show that complete CO is associated with larger MVA, smaller mean transmitral gradient, and functional improvement, and that the degree of CO provides important prognostic information.

Since its introduction by Inoue *et al.* (11) in 1984, PMC has been successfully and safely performed in large series of patients at numerous centers (8,12-15). PMC is now the treatment of choice for patients with MS and favorable anatomy (2,3). It provides results similar to those of surgical commissurotomy, even if the comparison of percutaneous and surgical series may be biased owing to differences in patients' clinical and mitral valve characteristics (16-18). Pioneering clinical and pathology studies have shown that CO is the main mechanism by which MVA increases after balloon valvuloplasty (percutaneous commissurotomy) (4). Commissural calcification, an important predictor




Figure 3. 10-Year Rates of Good Functional Results in the 3 Groups

Impact of commissural opening on late functional results. During a mean follow-up duration of 55 ± 28 months, the 10-year rates of good functional results (survival considering only cardiovascular death with no need for mitral surgery or repeat dilation and patient in New York Heart Association functional class I or II at most recent follow-up) was significantly different among the 3 groups ($p < 0.0001$). Groups are as defined in Figure 1.

of unsuccessful commissurotomy and failure of CO (19,20), is associated with a poor midterm event-free survival and a high incidence of mitral valve replacement. However, the relationship between the degree of CO and post-procedural MVA has never been extensively evaluated. In this large-scale study, we demonstrate that MVA progressively increases with the number and the extent of CO. It is worth noting that approximately two-thirds of patients referred for PMC had at least 1 completely open commissure.

Several important prognostic factors have been identified after successful PMC. Older age, less favorable anatomy (valve calcifications, severe subvalvular disease, echocardiography score), long-lasting disease (previous commissurotomy, atrial fibrillation, NYHA functional class IV, high systolic pulmonary artery pressure, severe tricuspid regurgitation), and less satisfactory immediate results (low post-procedure MVA and high post-procedure mean transmitral gradient) have been uniformly associated with a worst outcome (2,3,8,13,15). Few studies have aimed at evaluating the prognostic impact of the degree of CO on outcome. In a small pioneering study of 30 patients, Fatkin *et al.* (4) observed a better outcome for patients with than without CO. In a large and more recent study, commissural MR, as a surrogate for complete CO, was an independent predictor of restenosis and survival (21). In the present study, complete CO was associated with larger MVA, smaller mean gradient, and better functional out-

come. In multivariable analysis, either the degree of CO or the MVA was an independent predictor of good late functional results ($p < 0.05$).

Our study, showing for the first time the prognostic value of the degree of CO, has important clinical implications. Pre- and post-procedural evaluations of PMC results are based on echocardiographic assessment of the MVA. PMC is usually guided by echocardiography, and a decision to stop the procedure, in the absence of complications, is often based on MVA measurement. Planimetry is considered as the reference method but must be precisely performed at the tip of the leaflets, and therefore requires experienced operators (2). Moreover, the planimetry is not always feasible. Other methods also have their own limitations. The pressure half-time method (22) is commonly used, but it has been demonstrated that this method should be used cautiously especially for patients older than 60 years, in AF, or immediately after percutaneous balloon commissurotomy (23,24). The continuity equation is invalidated by the commonly associated aortic or mitral regurgitation (25). The proximal isovelocity surface area method is accurate and reproducible (26) but reputedly technically demanding and time-consuming; consequently, it is rarely used in practice.

All these limitations are even more important during the procedure in the catheterization laboratory where echocardiographic conditions are suboptimal. In contrast, degree of CO is relatively easy to assess and can be considered as a complementary measure of procedural success. These results also strengthen the importance of echocardiographic guidance of the procedure because hemodynamic variables are still used by some operators as the only criteria to determine when the procedure should be stopped. We recommend that further inflations with larger balloon volumes (30-mm balloon fully inflated with 30 ml) should be performed, and the procedure (in the absence of complication) not terminated before a complete opening of at least 1 commissure is achieved. It is worth noting that even with this aggressive attitude, we could not obtain a complete CO in 22% of our patients. Thus, the achievement of a complete and bicommissural opening must remain a major goal during PMC, and the degree of CO is an easy and simple predictor of long-term functional result.

The present study deserves several comments. First, degree of CO is the cornerstone of our results. In contrast to the MVA, which is quantitative and

continuous, the degree of CO was assessed semi-quantitatively, as described in Methods, but the intraobserver and interobserver variability was good. In our opinion, the degree of CO is difficult to measure quantitatively because it is hard to define exactly where the commissure starts. Second, the degree of CO was evaluated using 2-dimensional echocardiography. We have previously shown that 3-dimensional echocardiography provides a better assessment of the degree of CO than 2-dimensional echocardiography (27). The impact of this new modality of commissural assessment needs further evaluation. Third, when both the MVA and degree of CO were entered in the model, the degree of CO was not an independent predictor of outcome, meaning that the degree of CO and MVA is closely related and that its prognostic value is superseded by that of the MVA. The same results were observed overall and for patients above or below the median MVA. Furthermore, when the multivariable analysis was performed in the total population (whether or not PMC was successful, excluding patients with severe mitral regurgitation), the degree of CO provided additional prognostic value. A possible mechanistic explanation is that in patients with intermediate PMC results, a complete CO may provide a better adaptation to exercise. These results further emphasize the importance of achieving at least 1 complete CO during the procedure. Finally, we did not assess the morphology of the commissure before the procedure and thus examine the relationship between pre-procedural commissural anatomy and the degree of CO after PMC.

CONCLUSIONS

The present study demonstrates, in a large series of patients, that CO degree and MVA are closely related. After PMC, MVA increases and mean transmitral gradient decreases with the degree of CO. The degree of CO provides important prognostic information, and complete CO is associated with better late functional results. Thus, degree of CO should be systematically evaluated during and after PMC and can be considered as a complementary measure of the procedural success in addition to the MVA, which is not always easy to assess.

Reprint requests and correspondence: Dr. David Messika-Zeitoun, Cardiovascular Division, Bichat Hospital, 46 rue Henri Huchard, Paris 75018, France. *E-mail:* david.messika-zeitoun@bch.aphp.fr.

REFERENCES

1. Lung B, Cormier B, Ducimetiere P, et al. Immediate results of percutaneous mitral commissurotomy. A predictive model on a series of 1514 patients. *Circulation* 1996;94:2124-30.
2. Bonow RO, Carabello BA, Chatterjee K, et al. ACC/AHA 2006 guidelines for the management of patients with valvular heart disease: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Revise the 1998 Guidelines for the Management of Patients With Valvular Heart Disease): developed in collaboration with the Society of Cardiovascular Anesthesiologists; endorsed by the Society for Cardiovascular Angiography and Interventions and The Society of Thoracic Surgeons. *J Am Coll Cardiol* 2006;48:e1-148.
3. Vahanian A, Baumgartner H, Bax J, et al. Guidelines on the management of valvular heart disease: the Task Force on the Management of Valvular Heart Disease of the European Society of Cardiology. *Eur Heart J* 2007;28:230-68.
4. Fatkin D, Roy P, Morgan JJ, Feneley MP. Percutaneous balloon mitral valvotomy with the Inoue single-balloon catheter: commissural morphology as a determinant of outcome. *J Am Coll Cardiol* 1993;21:390-7.
5. McKay CR, Kawanishi DT, Rahimtoola SH. Catheter balloon valvuloplasty of the mitral valve in adults using a double-balloon technique. Early hemodynamic results. *JAMA* 1987;257:1753-61.
6. Kaplan JD, Isner JM, Karas RH, et al. In vitro analysis of mechanisms of balloon valvuloplasty of stenotic mitral valves. *Am J Cardiol* 1987;59:318-23.
7. Block PC, Palacios IF, Jacobs ML, Fallon JT. Mechanism of percutaneous mitral valvotomy. *Am J Cardiol* 1987;59:178-9.
8. Lung B, Garbarz E, Michaud P, et al. Late results of percutaneous mitral commissurotomy in a series of 1024 patients. Analysis of late clinical deterioration: frequency, anatomic findings, and predictive factors. *Circulation* 1999;99:3272-8.
9. Abascal VM, Wilkins GT, Choong CY, et al. Echocardiographic evaluation of mitral valve structure and function in patients followed for at least 6 months after percutaneous balloon mitral valvuloplasty. *J Am Coll Cardiol* 1988;12:606-15.
10. Zoghbi WA, Enriquez-Sarano M, Foster E, et al. Recommendations for evaluation of the severity of native valvular regurgitation with two-dimensional and Doppler echocardiography. *J Am Soc Echocardiogr* 2003;16:777-802.
11. Inoue K, Owaki T, Nakamura T, Kitamura F, Miyamoto N. Clinical application of transvenous mitral commissurotomy by a new balloon catheter. *J Thorac Cardiovasc Surg* 1984;87:394-402.
12. Nobuyoshi M, Hamasaki N, Kimura T, et al. Indications, complications, and short-term clinical outcome of percutaneous transvenous mitral commissurotomy. *Circulation* 1989;80:782-92.
13. Palacios IF, Sanchez PL, Harrell LC, Weyman AE, Block PC. Which patients benefit from percutaneous mitral balloon valvuloplasty? Prevalvuloplasty and postvalvuloplasty variables that predict long-term outcome. *Circulation* 2002;105:1465-71.
14. Bassand JP, Schiele F, Bernard Y, et al. The double-balloon and Inoue techniques in percutaneous mitral valvuloplasty: comparative results in a series of 232 cases. *J Am Coll Cardiol* 1991;18:982-9.
15. Hernandez R, Banuelos C, Alfonso F, et al. Long-term clinical and echocardiographic follow-up after percutaneous mitral valvuloplasty with the Inoue balloon. *Circulation* 1999;99:1580-6.
16. Reyes VP, Raju BS, Wynne J, et al. Percutaneous balloon valvuloplasty compared with open surgical commissurotomy for mitral stenosis. *N Engl J Med* 1994;331:961-7.
17. Patel JJ, Shama D, Mitha AS, et al. Balloon valvuloplasty versus closed commissurotomy for pliable mitral stenosis: a prospective hemodynamic study. *J Am Coll Cardiol* 1991;18:1318-22.
18. Ben Farhat M, Ayari M, Maatouk F, et al. Percutaneous balloon versus surgical closed and open mitral commissurotomy: 7-year follow-up results of a randomized trial. *Circulation* 1998;97:245-50.
19. Cannan CR, Nishimura RA, Reeder GS, et al. Echocardiographic assessment of commissural calcium: a simple predictor of outcome after percutaneous mitral balloon valvotomy. *J Am Coll Cardiol* 1997;29:175-80.
20. Sutaria N, Northridge DB, Shaw TR. Significance of commissural calcification on outcome of mitral balloon valvotomy. *Heart* 2000;84:398-402.
21. Kang DH, Park SW, Song JK, et al. Long-term clinical and echocardiographic outcome of percutaneous mitral valvuloplasty: randomized comparison of Inoue and double-balloon techniques. *J Am Coll Cardiol* 2000;35:169-75.
22. Hatle L, Angelsen B, Tromsdal A. Noninvasive assessment of atrioventricular pressure half-time by Doppler ultrasound. *Circulation* 1979;60:1096-104.
23. Thomas J, Wilkins G, Choong C, et al. Inaccuracy of mitral pressure half-time immediately after percutaneous mitral valvotomy: dependence on transmitral gradient and left atrial and ventricular compliance. *Circulation* 1988;78:980-93.
24. Messika-Zeitoun D, Meizels A, Cachier A, et al. Echocardiographic evaluation of the mitral valve area before and after percutaneous mitral commissurotomy—the pressure half-time method revisited. *J Am Soc Echocardiogr* 2005;18:1409-14.
25. Palacios IF. What is the gold standard to measure mitral valve area post mitral balloon valvuloplasty? *Cathet Cardiovasc Diagn* 1994;33:315-6.
26. Messika-Zeitoun D, Fung Yiu S, Cormier B, et al. Sequential assessment of mitral valve area during diastole using colour M-mode flow convergence analysis: new insights into mitral stenosis physiology. *Eur Heart J* 2003;24:1244-53.
27. Messika-Zeitoun D, Brochet E, Holmin C, et al. Three-dimensional evaluation of the mitral valve area and commissural opening before and after percutaneous mitral commissurotomy in patients with mitral stenosis. *Eur Heart J* 2007;28:72-9.

Key Words: mitral stenosis ■ percutaneous mitral commissurotomy ■ functional improvement ■ echocardiography ■ mitral valve area.