

Impact of Regulated Secretion on Antiparasitic CD8 T Cell Responses

Harshita Satija Grover,^{1,4} H. Hamlet Chu,^{1,4} Felice D. Kelly,² Soo Jung Yang,¹ Michael L. Reese,^{2,5} Nicolas Blanchard,³ Federico Gonzalez,¹ Shiao Wei Chan,¹ John C. Boothroyd,² Nilabh Shastri,^{1,*} and Ellen A. Robey^{1,*}

¹Division of Immunology and Pathogenesis, Department of Molecular and Cellular Biology, University of California, Berkeley, Berkeley, CA 94720-3200, USA

²Department of Microbiology and Immunology, Stanford University School of Medicine, Stanford, CA 94305-5124, USA

³Center of Pathophysiology of Toulouse-Purpan, INSERM UMR1043-CNRS UMR5282, University of Toulouse, 31024 Toulouse Cedex 3, France

⁴Co-first author

⁵Present address: Department of Pharmacology, University of Texas, Southwestern Medical Center, Dallas, TX 75390-9003, USA

*Correspondence: nshastri@berkeley.edu (N.S.), erobey@berkeley.edu (E.A.R.)

<http://dx.doi.org/10.1016/j.celrep.2014.04.031>

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/3.0/>).

SUMMARY

CD8 T cells play a key role in defense against the intracellular parasite *Toxoplasma*, but why certain CD8 responses are more potent than others is not well understood. Here, we describe a parasite antigen, ROP5, that elicits a CD8 T cell response in genetically susceptible mice. ROP5 is secreted via parasite organelles termed rhoptries that are injected directly into host cells during invasion, whereas the protective, dense-granule antigen GRA6 is constitutively secreted into the parasitophorous vacuole. Transgenic parasites in which the ROP5 antigenic epitope was targeted for secretion through dense granules led to enhanced CD8 T cell responses, whereas targeting the GRA6 epitope to rhoptries led to reduced CD8 responses. CD8 T cell responses to the dense-granule-targeted ROP5 epitope resulted in reduced parasite load in the brain. These data suggest that the mode of secretion affects the efficacy of parasite-specific CD8 T cell responses.

INTRODUCTION

CD8 T cells are key for the control of intracellular pathogens, including the protozoan parasite *Toxoplasma gondii*. *T. gondii* infects a wide array of warm-blooded hosts, including one-third of humans worldwide (Carruthers, 2002; Montoya and Liesenfeld, 2004), but typically causes little pathology, due in part to a robust T cell response (Brown and McLeod, 1990; Denkers and Gazzinelli, 1998; Hakim et al., 1991; Lindberg and Frenkel, 1977). However, not all CD8 T cell responses are equally effective in controlling the parasite, as dramatically illustrated by the differential sensitivity to infection in two inbred mouse strains, BALB/c and C57BL/6 (B6). BALB/c mice show strong resistance to infection due to the presence of the protective MHC class I allele H-2L^d, whereas B6 mice, which lack this particular allele,

are highly sensitive (Brown et al., 1995; Brown and McLeod, 1990; Suzuki et al., 1991, 1994). We recently showed that the protective effect of MHC class I H-2L^d is due to a potent CD8 T cell response directed against a single parasite protein, GRA6 (Blanchard et al., 2008). H-2L^d-GRA6-specific T cells account for the majority of CD8 T cells in the brains of infected H-2^d mice and effectively control the parasite load. In contrast, B6 (H-2^b) mice exhibit much higher parasite loads in the brain and eventually succumb to infection despite the presence of parasite-specific CD8 T cells (Schaeffer et al., 2009). Understanding why particular CD8 T cell responses predominate over others, and why some responses provide more effective protection is critical for designing improved vaccines and other therapies against intracellular pathogens.

One factor that may influence the immunogenicity and immunoprotection of potential CD8 antigens is the intracellular pathway by which pathogen-derived antigens are processed and presented in infected host cells. For cytosolic antigens, such as many viral antigens, presentation occurs via the classical class I MHC presentation pathway. In this pathway, proteins are degraded in the host cytosol by proteasomes and the resulting peptides are transported into the endoplasmic reticulum (ER) via the TAP transporter, receive a final trimming by the ERAP, are loaded onto MHC class I, and finally are transported to the surface as peptide-MHC complexes for recognition by a CD8 T cell. In contrast, for pathogen proteins that enter the cell via phagocytosis, antigen presentation occurs by an alternative “cross-presentation” pathway that requires a phagosome in addition to the ER vesicular transport step (Joffre et al., 2012). The importance of antigen compartmentalization for the CD8 T cell response is illustrated by the protective response to intracellular bacteria that occurs when the antigen is secreted into the cytosol, but not when the antigen is retained inside the bacteria (Shen et al., 1998).

For intracellular parasites, the pathways by which potential antigens traffic from the pathogen into the host cell may also impact CD8 T cell responses. For example, *T. gondii* resides within a specialized nonfusogenic compartment, the parasitophorous vacuole, which restricts the movement of material




Figure 1. *T. gondii*-Specific CD8 T Cell Responses in B6 Mice Ex Vivo and after In Vitro Expansion

C57BL/6 mice were immunized with irradiated *T. gondii* tachyzoites. Splenocytes were harvested from mice at 2 weeks postimmunization, and *T. gondii*-specific CD8 and CD4 T cell responses were measured by intracellular cytokine staining for IFN- γ . APCs \pm *T. gondii* were used as stimulators for ex vivo and in vitro restimulations. (A–C) Ex vivo and in vitro restimulation results are shown as (A) representative flow-cytometry plots and (B and C) plots depicting the expansion of CD8 T cells (B) and CD4 T cells (C) over the course of ten in vitro restimulations. Data are representative of three experiments.

(D) LacZ response of subcloned CD8 hybridoma (BTg45Z) after an overnight culture with wild-type APCs \pm *T. gondii* or APCs blocked with antibodies against H-2K^b or H-2D^b MHC I molecules + *T. gondii*. Data are representative of at least two independent experiments.

into the host cytosol and thus poses a barrier to antigen presentation. Nevertheless, studies with model antigens have shown that proteins that are constitutively secreted into the parasitophorous vacuole lumen via parasite organelles termed dense granules can elicit strong CD8 T cell responses (Gregg et al., 2011; Gubbels et al., 2005). Moreover, the potent endogenous CD8 antigen GRA6 is also constitutively secreted via dense granules (Blanchard et al., 2008). *Toxoplasma* also possesses distinct secretory organelles termed rhoptries that are injected directly into the host cell during productive and abortive invasion events (Blader and Saeij, 2009; Boothroyd and Dubremetz, 2008; Koshy et al., 2012), and this distinct spatial and temporal pattern of secretion could affect the ability of a parasite protein to be presented by MHC class I. All endogenous *T. gondii* CD8 antigens identified to date possess secretory signals, and include both dense-granule and rhoptry proteins (Frickel et al., 2008; Wilson et al., 2010; Blanchard et al., 2008). However, how the mode of secretion of potential *T. gondii* antigens affects the nature of the CD8 T cell response has not been investigated.

A comparison between the immunodominant, protective H-2L^d-GRA6 response and the endogenous T cell response from susceptible H-2^b mice should reveal clues about what constitutes an optimal CD8 T cell response. Aside from GRA6, all other defined endogenous MHC-I-restricted *T. gondii* antigens have been identified with the use of prediction strategies (Frickel et al., 2008; Khan et al., 1988, 1994; Wilson et al., 2010). These approaches, while useful, may miss many endogenous responses that do not fit into a given set of assumptions. In contrast, the alternative approach of using T cells from infected mice to probe expression libraries provides a relatively unbiased way to characterize endogenous T cell responses (Blanchard et al., 2008; Malarkannan et al., 2001).

Here, we used an expression-cloning approach to characterize the CD8 T cell response in H-2^b mice, which led to the identification of a nonamer peptide, YAL9, derived from the

rhoptry-targeted parasite protein ROP5. In comparison with the H-2L^d-GRA6 response, the H-2D^b-ROP5 (YAL9) response accounts for a relatively small proportion of the endogenous CD8 T cell response to the parasite and does not confer strong protection. To test the impact of regulated secretion on the T cell response, we engineered parasites in which the YAL9 epitope from ROP5 was altered from its normal rhoptry targeting to secretion via dense granules, and we examined the impact of this alteration on the T cell response. Redirecting the ROP5 epitope to the dense granules led to a dramatic increase in the number and protective capacity of H-2D^b-ROP5-specific T cells, and to a reduced cyst burden in chronically infected mice. Conversely, retargeting the T cell-stimulatory epitope from GRA6 (HF10) for secretion via rhoptries led to a substantially reduced T cell response. Our data suggest that the mode of protein secretion can have a dramatic effect on the magnitude and efficacy of the CD8 T cell response, with important implications for immunity to *T. gondii* and related malaria-causing intracellular parasites.

RESULTS

T. gondii-Immunized C57BL/6 Mice Exhibit Potent CD4 T Cell Responses but Weaker CD8 T Cell Responses

To characterize the T cell response in C57BL/6 (B6, H-2^b) mice, we immunized animals with irradiated tachyzoites of the type II *T. gondii* strain *Prugniaud* (Pru) and examined *T. gondii*-specific CD4 and CD8 T cell responses ex vivo at 2 weeks. In contrast to H-2^d mice, in which the CD8 T cell response dominates over the CD4 T cell response (Blanchard et al., 2008), B6 mice generated a robust CD4 T cell response but a weaker CD8 response toward *T. gondii* (Figure 1A, top panels; Grover et al., 2012). To selectively expand the CD8 T cells, we used MHC class II-deficient dendritic cells (DCs) as antigen-presenting cells (APCs) for in vitro restimulations. The *T. gondii*-specific CD8 T cells progressively expanded




Figure 2. BTg45Z Hybridoma Recognizes the ROP5 Rhoptyr Protein

(A) *T. gondii* cDNA library was screened to identify the antigen recognized by BTg45Z. Two cDNA pools (pTg8E6 and pTg13B7) stimulated the lacZ response of the hybrid.

(B) pTg8E6 was further fractionated to identify a positive cDNA clone pTg8E6.1. The BTg45Z response to pTg8E6.1 was antigen specific and MHC restricted.

(C) *T. gondii* database (<http://toxodb.org/toxo/>) BLAST search results revealed that the antigenic cDNA clone pTg8E6.1 encoded a truncated version of the *T. gondii* protein, TGMe49_018080 (Rhoptyr kinase family protein 5). Alignment of the amino acid sequence encoded by the truncated cDNA clone pTg8E6.1 as compared with the full-length polymorphic clones ROP5IIA and ROP5IIC (Reese et al., 2011) and pTg13B7 is depicted here. (–) represents missing amino acids in the sequence, (*) represents alignment, and (/ or gap) represents alignment between ROP5IIC and pTg8E6.1 and pTg13B7.

See also Figure S1.

was *T. gondii* specific and restricted by the H-2D^b MHC class I molecule.

ROP5 Is the Antigenic Protein Recognized by CD8 T Cell Hybridoma

We identified the *T. gondii* antigenic protein by expression cloning. A *T. gondii*-derived cDNA expression library was screened for BTg45Z-stimulating genes (Figure S1; Blanchard et al., 2008). APCs transfected with two cDNA pools (pTg8E6 and pTg13B7) stimulated the BTg45Z T cell hybrid (Figure 2A). Fractionation of the cDNA pools yielded an antigenic active clone, pTg8E6.1 (Figure 2B), that encoded a member of the rhoptyr protein 5 (ROP5) family of proteins. The ROP5 locus is located on chromosome XII and encodes a polymorphic family of four to ten pseudokinases that differ both between *Toxoplasma* strains

and between individual genes within a single cluster (Reese et al., 2011). Sequence comparison with different polymorphic versions of ROP5 in the type II strain of *T. gondii* (Reese et al., 2011) revealed that pTg8E6.1 matched the ROP5 version IIC, a protein of 549 amino acids (Figure 2C).

Identification of the Minimal Antigenic Peptide

To define the minimal antigenic peptide recognized by the BTg45Z hybrid, we tested candidate peptides predicted to bind D^b MHC (Immune Epitope Database and Analysis Resource, <http://www.iedb.org/>; Figure 3A). Analysis of cDNA deletion constructs (Table 1) suggested that the antigenic

over successive in vitro restimulation cultures (Figures 1A, lower panels, 1B, and 1C).

Generation of a *T. gondii*-Specific CD8 T Cell Hybridoma

To further characterize the *T. gondii*-specific CD8 T cell response at a clonal level, we generated *T. gondii*-specific, lacZ-inducible CD8 T cell hybridomas as described previously (Karttunen et al., 1992; Sanderson and Shastri, 1994). One such hybridoma, referred to as BTg45Z, produced lacZ when cocultured with bone-marrow-derived DCs (BMDCs) infected with *T. gondii* (Figure 1D) and this response was blocked by anti-H-2D^b, but not anti-H-2K^b, monoclonal antibody. Thus, the BTg45Z hybridoma




Figure 3. Identification of the Antigenic Epitope within ROP5IIC that Stimulates BTg45Z Hybridoma

(A) Schematic representation of the N-terminal and C-terminal deletion constructs generated to identify the antigenic epitope within clone pTg8E6.1. The clone was truncated and missing the 5' UTR and nucleotides encoding the first 222 amino acids. The specific amino acids tested in each construct are shown in parentheses and the antigenic activity region is indicated by the filled gray circle. Open circles represent other putative H-2D^b binding epitopes as predicted by the Immune Epitope Database (<http://www.iedb.org/>).

(B) BTg45Z lacZ response after an overnight stimulation to different deletion constructs that were transfected in H-2D^b expressing APCs.

(C) Sequences of antigenic peptide and its N-terminal extended versions. BTg45Z lacZ response to BMDCs alone or BMDCs that were pulsed with varying concentrations of the indicated peptides is shown.

See also [Figures S2](#) and [S3](#).

activity mapped to the YAVANYFFL (YAL9) peptide ([Figure 3B](#)). Moreover, the YAL9 peptide had higher antigenic activity relative to N-terminally extended versions of the peptide ([Figure 3C](#)), and was found in high-performance liquid chromatography (HPLC)-

fractionated extracts of cells transfected with pTg8E6.1 ([Figure S1](#)). We conclude that ROP5IIC was processed by APCs to the YAL9 peptide that was presented by MHC I H-2 D^b and recognized by BTg45Z T cells.

Presentation of YAL9 Requires Proteasome and TAP and Is Partially Dependent on ERAAP

In the classical MHC class I presentation pathway, cytosolic proteins are broken down into peptides by the proteasome and are transported into the ER via TAP. In the ER, some peptides are further processed by the aminopeptidase ERAAP and loaded onto MHC class I molecules ([Serwold et al., 2001](#)). To assess whether presentation of YAL9 peptide required the components of the classic MHC-I presentation pathway, we infected BMDCs from wild-type B6, TAP^{-/-}, or ERAAP^{-/-} mice and used them as APCs to stimulate the BTg45Z hybridoma. The BTg45Z cell response was completely abolished in TAP-deficient cells and was strongly inhibited by the proteasome inhibitor lactacystin ([Figure S2](#)). In contrast, loss of ERAAP only partially inhibited BTg45Z responses. Thus, presentation of YAL9 requires proteasome processing in the cytoplasm and transport by TAP into the ER, but is only partially dependent on further trimming by ERAAP.

Detection of YAL9-Specific CD8 T Cells in *T. gondii*-Infected Mice

We next assessed the contribution of YAL9-specific CD8 T cells to the overall T cell response to the parasite. We orally infected H-2^b (C57BL/6) mice and determined the abundance of specific T cells in the CD8 T cell repertoire at 4 weeks postinfection by H-2D^b-YAL9 MHC tetramer staining and intracellular cytokine staining (ICCS) after ex vivo restimulation with the YAL9 peptide ([Figures 4A–4E](#)). The YAL9-specific T cells, as measured by ICCS, accounted for 0.3% of splenic CD8 T cells ([Figure 4B](#)). Based on the ~3.5% parasite-specific cells among the total CD8 T cells, we estimate that ~10% of parasite-specific CD8 splenocytes are specific for YAL9. Tetramer staining gave a slightly higher value (0.5%) of YAL9-specific T cells among splenic CD8 T cells, perhaps due to higher sensitivity of this assay ([Figure 4C](#)). Among CD8 T cells in the brain, YAL9-specific cells accounted for ~1% of the CD8 response based on both ICCS and tetramer staining, corresponding to ~10% of the total parasite-specific CD8 response as estimated by ICCS ([Figures 4D](#) and [4E](#)). Analyses of mice that were immunized with irradiated parasites gave similar results, with ~0.5% of splenic CD8 T cells showing reactivity to the YAL9 peptide ([Figures 4F–4H](#)). Thus, unlike the response to the GRA6-derived peptide HF10, which dominates the CD8 T cell response to the parasites in H-2^d mice ([Blanchard et al., 2008](#)), the ROP5-derived YAL9 peptide is a detectable but minor component of the CD8 T cell response in H-2^b mice.

Immunization of H-2^d mice with DCs pulsed with the single HF10 peptide derived from the dense-granule protein GRA6 elicited robust protection against challenge with live parasites ([Blanchard et al., 2008](#)). To determine the protective potential of the YAL9-specific CD8 T cells, we immunized B6 mice with DCs pulsed with YAL9 peptide and challenged the mice with a lethal dose of *T. gondii*. We observed no statistically significant

Table 1. Oligonucleotide Primers for ROP5 Deletion Constructs

Construct	Forward (5'–3')	Reverse (5'–3')
ΔNΔC-3	GATCCATGTTACACCTGAAAACCTTTTCATTATGTGAT	CTAGATCACATAATGAAAAGGTTTTTCAGGTGTGAACATG
ΔNΔC-5	GATCCATGAATGCGTGGCAACTGGGTCTTAGCATATGAT	CTAGATCATATGCTAAGACCCAGTTGCCACGCATTTCATG
ΔN-7	CGCGGATCCGCGATGGTTCGGAACCCGAGGACCG	GCTCTAGAGCAGCGACTGAGGGCGCAGC
ΔNΔC-8	CGCGGATCCGCGATGGAGGAGGCACGG	GCTCTAGAGCGCTTGTATGCCGGTCTCTCG
ΔNΔC-9	CGCGGATCCGCGATGCCAGGACAGAGC	GCTCTAGAGCGCTTGTATGCCGGTCTCTCG

difference in survival (Figure S3A), cyst number, or parasite load (Figures S3H and S3I) between mice immunized with the YAL9 versus an irrelevant H-2D^b binding peptide. These results are consistent with evidence that the GRA6 response alone can account for the protective effect of MHC-class I H-2L^d expression on an H-2^d background (Brown et al., 1995; Blanchard et al., 2008; Feliu et al., 2013), implying that none of the parasite antigens presented by H-2^b class I alleles provide as strong protection as H-2L^d-HF10.

Generation of Transgenic Parasites with Altered Intracellular Trafficking of T Cell Epitopes

The ROP5 and GRA6 antigenic precursors differ in their modes of secretion by *T. gondii* (Figure 5A). The ROP5 protein is secreted into host cells during parasite invasion via secretory organelles known as rhoptries, whereas GRA6 is constitutively secreted into the parasitophorous vacuole through dense granules. To assess the influence of differential protein targeting on T cell responses, we generated three hemagglutinin (HA)- or FLAG-tagged constructs encoding type II ROP5 (the so-called IIC allele) and GRA6 genes, along with constructs designed to swap the secretion patterns of the antigenic ROP5 and GRA6 peptides (Figures 5B and S4). These constructs were (1) GRA6-YAL9, which replaces the T cell epitope of GRA6 (HF10) with that of ROP5 (YAL9) to target this normally rhoptry peptide to dense granules; (2) GRA6-ROP5, which has the promoter and signal peptide of the *GRA6* gene (Gendrin et al., 2010) fused upstream of the C-terminal domain of ROP5, including the YAL9 epitope, thus also targeting the bulk of this rhoptry protein to dense granules; and (3) ROP5-HF10, in which the HF10 epitope of GRA6 is fused to the C terminus of ROP5, thus targeting this normally dense-granule peptide for secretion via rhoptries.

We introduced these constructs as transgenes into the type III parasite strain CTG, whose alleles of GRA6 and ROP5 do not encode the YAL9 or HF10 T cell-stimulatory epitopes (Blanchard et al., 2008; Reese et al., 2011; Feliu et al., 2013). To confirm appropriate targeting of the proteins, we performed immunofluorescence analysis of transgenic parasites (Figures S4B and S4C). As expected, parasites expressing GRA6, GRA6-YAL9, and GRA6-ROP5 all displayed transgenic protein colocalization with the dense-granule marker GRA7 (Figures S4B and S4C), whereas parasites expressing ROP5-HF10 displayed accumulation of transgenic protein in structures that colocalized with rhoptry markers ROP2, ROP3, and ROP4 (Figure S4B). On the other hand, and also as expected, CTG harboring the wild-type ROP5 construct showed a typical rhoptry pattern of staining and failed to localize with GRA7 (Figure S4C). Western blot analysis indicated that GRA6-YAL9 was expressed at somewhat lower levels

(~1/3) than what was seen for GRA6-ROP5 (Figure S4D). A comparison of the HA-tagged GRA6-ROP5 and the FLAG-tagged ROP5 indicated that ROP5 was also expressed at a lower level (~1/5–1/10) relative to GRA6-ROP5 (Figure S4E). For the parasite strains expressing the GRA6 epitope HF10, ROP5-HF10 levels were similar to those of transgenic GRA6 (Figure S4F).

The Immunogenicity of Antigenic Peptides Is Enhanced by Targeting the Precursor Protein to Dense Granules

We next assessed the ability of transgenic parasites to stimulate ROP5-specific or GRA6-specific T cells. We expected that the parental CTG strain would not elicit a YAL9-specific T cell response, since this strain harbors ROP5 alleles that lack this antigenic epitope (Reese et al., 2011). To confirm this, we measured the response of the BTg45Z hybridoma to DCs infected with CTG parasites or pulsed with peptides derived from the regions corresponding to the YAL9 peptide in the allelic forms of ROP5 contained within the CTG parasites (Figures S5A–S5C). As expected, we observed no detectable response to CTG parasites or to these alternate ROP5 peptides. Interestingly, a polymorphic peptide from another type II allele within the multi-gene ROP5 locus (ROP5IIA, which is identical to YAL9 except for a L substituting for a F at position 8) stimulated the hybridoma comparably to what was seen with YAL9 from ROP5IIC (Figure S5C). However, and as predicted by their respective genome sequences, neither the CTG nor the RH parasite strains expressed a form of ROP5 that stimulated our ROP5-specific hybridoma (data not shown). We also confirmed that immunization of mice with irradiated CTG parasites did not elicit a detectable response to the YAL9 epitope as measured by both ICCS and tetramer staining (Figures S5D and S5E).

To assess whether the immunogenicity of the YAL9 or HF10 epitopes was affected by altering the secretion of the precursor protein, we infected DCs with the transgenic parasites described above and measured peptide presentation using the corresponding lacZ-inducible hybridoma. The ROP5-specific BTg45Z hybridoma response to transgenic CTG parasites expressing either one of the dense-granule-targeted YAL9 constructs (GRA6-ROP5 and GRA6-YAL9) was substantially higher than those expressing YAL9 in its native form, i.e., as a part of ROP5IIC targeted to its normal location, the rhoptries (Figure 5C, left panel). On the other hand, the GRA6-specific CTgEZ4 hybridoma response to parasites harboring the rhoptry-targeted HF10 peptide (ROP5-HF10) was substantially reduced compared with the response to the normally dense granule-targeted version of the epitope (Figure 5C, right panel).

We compared the different transgenic parasite strains for their ability to elicit CD8 T cell responses in vivo. We infected mice and




Figure 4. YAL9-Specific CD8 T Cell Response Constitutes a Detectable Fraction of the In Vivo *T. gondii*-Specific Response

(A–H) Mice were orally infected with 25–40 *T. gondii* cysts (A–E) or immunized with irradiated 5×10^6 *T. gondii*, ME49 tachyzoites (F–H). Splenocytes and brain leukocytes were harvested from mice 3–4 weeks after oral infection or 2 weeks after immunization. *T. gondii*-specific T cell responses were measured by intracellular cytokine staining for IFN- γ or by staining with peptide-MHC tetramers.

(A) Representative flow-cytometry plots of splenocytes from naive and infected mice. Left panels show intracellular IFN- γ staining after in vitro restimulation with APCs (either +/- *T. gondii* or 1 μ M YAL9 peptide). Right panels show MHC class I H-2D^b-YAL9 tetramer staining.

(B and D) Compiled data showing *T. gondii*-specific and H-2D^b-YAL9-specific responses from (B) spleen and (D) brain of infected mice. Data are corrected for background based on responses by CD8 T cells toward uninfected APCs or APCs pulsed with irrelevant peptide. Each dot represents an individual mouse. (C and E) Compiled data showing flow-cytometry analysis of MHC class I H-2D^b-YAL9 tetramer staining on (C) splenocytes and (E) brain leukocytes. Cells were also costained with CD8 antibody.


(F) Representative flow-cytometry plots of splenocytes from naive and immunized mice. Left panels show intracellular IFN- γ staining after in vitro restimulation with APCs either +/- *T. gondii* or 1 μ M YAL9 peptide. Right panels show MHC class I H-2D^b-YAL9 tetramer staining.

(G) Compiled data showing splenic *T. gondii*-specific and H-2D^b-YAL9-specific responses from immunized mice. Data are corrected for background based on responses by CD8 T cells toward uninfected APCs or APCs pulsed with irrelevant peptide. Each dot represents an individual mouse.

(H) Compiled data showing MHC class I H-2D^b-YAL9 tetramer staining on splenocytes from immunized mice. Cells were also costained with CD8 antibody. Data are representative of at least three independent experiments. Data (mean and SEM) are representative of at least three independent experiments. See also Figure S3.

examined T cell responses in spleen and brain leukocytes at 3–4 weeks postinfection. At day 21 postinfection, ROP5-specific T cells made up ~0.5% of the total CD8 T cell population in the spleen and brains of mice infected with type III CTG parasites expressing the normally targeted ROP5 protein, similar to the results obtained with the type II strains Pru and ME49 (Figure 5D). Strikingly, infection with CTG+GRA6-ROP5 parasites elicited a 10- to 20-fold higher YAL9-specific T cell response compared with CTG+ROP5 parasites, as measured by tetramer staining (Figure 5D). In spite of the increase in the number of specific T cells in the brain, their expression of effector/memory markers KLRG-1 and IL7R remained unchanged (Figure 5E). In addition, a comparison of the YAL9-specific versus parasite-specific CD8 T cells using in vitro restimulation and intracellular IFN- γ staining revealed that the YAL9-specific T cell response accounted for the bulk of the antiparasite CD8 response in mice infected with CTG+GRA6-ROP5 parasites (Figure 5F). This predominance of the YAL9-specific response resulted in a significant decrease in the CD4-to-CD8 ratio among parasite-specific IFN- γ producing T cells in both the spleen and brain (Figure 5G). The YAL9-specific CD8 T cell response to the CTG+GRA6-YAL9 strain was also enhanced compared with CTG+ROP5, although to a lesser extent than with CTG+GRA6-ROP5 parasites (Figures 5D, 5F, and 5G). A similar enhancement of the CD8 T cell response to dense granule-targeted YAL9 was also observed when mice were immunized with irradiated transgenic parasites (Figures 5H and 5I).

We observed a similar correlation between enhanced T cell response and dense-granule targeting with the GRA6 epitope



(legend on next page)

(Figure 5J). GRA6-specific T cells constituted ~10% of the total CD8 T cell population in the spleen and 30% in the brains of mice infected with CTG parasites expressing the normally targeted GRA6 protein, similar to the results obtained with the type II strain Pru (Figure 5J). In contrast, mice infected with parasites harboring the rhoptry-targeted version of the GRA6 epitope (ROP5-HF10) had 3- to 4-fold lower numbers of GRA6-specific T cells. Together, these data indicate that the same T cell epitope evokes a greater T cell response when secreted via dense granules versus rhoptries.

YAL9-Specific T Cells Provide Protection against Parasites in which the Epitope Is Targeted to Dense Granules

In some settings, CD8 T cells may be expanded effectively but still fail to provide strong protection (Shen et al., 1998). To determine whether retargeting of the YAL9 epitope could enhance immune protection as well as CD8 expansion, we examined the ability of peptide-DC immunization to provide protection against transgenic parasites in which the YAL9-containing protein was targeted for secretion via dense granules versus rhoptries. We performed peptide immunization with YAL9 in mice according to the scheme in Figure 6A, and challenged them intraperitoneally (i.p.) with a high dose of CTG+ROP5 or CTG+GRA6-ROP5 transgenic parasites. Strikingly, all of the mice that were immunized with YAL9 and then challenged with CTG+GRA6-ROP5

survived the acute phase of infection. In contrast, the majority of mice that were immunized with YAL9 peptide and then challenged with CTG+ROP5, or were immunized with control peptide (WI9), succumbed to infection within the first week after challenge (Figure 6B). Thus, the combination of peptide immunization to create a preexisting ROP5-specific T cell population and challenge with parasites harboring the epitope targeted for secretion via dense granules provided strong immune protection during the acute phase of infection.

To further explore the mechanism of protection, we measured the T cell responses in the chronic phase of infection in the surviving mice (Figures 6C–6E). Interestingly, there was an enhanced YAL9-specific response in the brains of CTG+GRA6-ROP5-infected animals irrespective of the peptide immunization. This difference was seen as an increased proportion of YAL9-specific CD8 T cells based on tetramer staining (Figure 6C), ex vivo restimulation followed by ICCS for IFN- γ (Figure 6D), and an overall decrease in the CD4-to-CD8 ratio among INF- γ -producing, *T. gondii*-specific T cells (Figure 6E). Similar trends were observed in the splenic T cells of surviving mice (Figures S6A–S6C). We also observed a marked difference in the cyst load of surviving mice chronically infected with CTG+ROP5 relative to CTG+GRA6-ROP5, regardless of the peptide immunization (Figure 6F). This difference is likely due to a more protective CD8 T cell response rather than to reduced overall fitness of the parasite strains, since CTG+GRA6-ROP5, CTG+GRA6, and

Figure 5. Enhanced T Cell Response when the Antigenic Precursor Proteins Are Targeted to Parasite Dense Granules

(A) Schematic depicting distinct modes of secretion for the dense-granule protein GRA6 (depicted in red) (Holtappels et al., 2008) versus rhoptry protein ROP5 (depicted in blue). Also shown are possible routes for antigen processing and presentation via MHC-I. *T. gondii* injects rhoptry proteins directly into the host cell during invasion. ROP5 associates with the cytosolic face of parasitophorous vacuole membrane (PVM) following secretion. Parasites constitutively secrete dense-granule proteins into the PV lumen of invaded host cells. Proteolytic processing of parasite secretory proteins could take place in the host cells' cytosol by the proteasome, and further trimming by ERAAP may occur after TAP-mediated transport of peptides into the ER. Processed peptides are loaded onto MHC class I molecules to be transported to the cell surface for recognition by a CD8 T cell (not depicted).

(B) Left-hand diagrams show wild-type ROP5 and GRA6 genes, indicating the location of their antigenic epitopes (YAL9 and HF10) and placement of HA and FLAG epitope tags. Right-hand diagrams show the transgenic constructs used to retarget expression of the epitopes to different parasite secretory compartments. GRA6-YAL9 has the HF10 epitope of the GRA6 gene replaced by YAL9. An additional ten amino acids from ROP5 flanking the YAL9 epitope were included to allow for efficient processing (not depicted). GRA6-ROP5 contains the GRA6 promoter and signal sequence fused to the C-terminal portion of the ROP5IIIC gene including the YAL9 epitope. ROP5-HF10 contains the HF10 epitope from GRA6 fused to the C terminus of ROP5IIIC gene. All constructs were introduced as transgenes into the parental type III strain, CTG, which harbors allelic forms of ROP5 and GRA6 that lack the T cell stimulatory epitopes.

(C) BMDCs from C57BL/6 (H-2^b) or B10.D2 (H-2^d) mice were infected in vitro with irradiated transgenic *T. gondii* tachyzoites at varying MOIs, and BTg45Z and CTgEZ.4 lacZ response was measured after an overnight stimulation with BMDCs \pm *T. gondii*.

(D) C57BL/6 mice (H-2^b) were infected i.p. with 10⁵ transgenic *T. gondii* tachyzoites. Splenocytes or brain leukocytes were harvested from mice 3–4 weeks postinfection, and *T. gondii*-specific T cell responses were measured by staining with tetramers. Compiled data show MHC class I H-2D^b-YAL9 tetramer staining on gated CD8+ B220– splenocytes (left panel) and brain leukocytes (right panel).

(E) Compiled data showing percent expression of the effector T cell marker KLRG-1 and memory T cell marker IL7R in brain *T. gondii*-specific and H-2D^b-YAL9-specific CD8 T cells from mice infected by 10⁵ transgenic parasites i.p. Brain leukocytes were harvested from mice at 3 weeks postinfection. Each bar represents an average from four to five mice. Data (mean and SEM) are pooled from two independent experiments.

(F) Compiled data showing splenic (left panel) and brain (right panel) *T. gondii*-specific and H-2D^b-YAL9-specific CD8 T cell responses from infected mice as measured by intracellular cytokine staining for IFN- γ . Data are corrected for background based on responses by T cells toward uninfected APCs or APCs pulsed with irrelevant peptide.

(G) Plots depicting the ratio of *T. gondii*-specific CD4-to-CD8 T cell response from spleen (left panel) or brain (right panel) of infected mice.

(H) Mice were immunized with irradiated 5 \times 10⁶ transgenic *T. gondii* tachyzoites. Splenocytes were harvested from mice at 2 weeks postimmunization and *T. gondii*-specific T cell responses were measured using intracellular staining for IFN- γ . Compiled data show splenic *T. gondii* and YAL9-specific CD8 T cell responses from immunized mice. Data are corrected for background based on responses by T cells toward uninfected APCs or APCs pulsed with irrelevant peptide.

(I) Compiled data showing MHC class I H-2D^b-YAL9 tetramer staining on splenocytes from immunized mice. Cells were also costained with CD8 antibody. Each bar represents an average from six mice. Each dot represents an individual mouse. Data (mean and SEM) are pooled from two independent experiments.

(J) B6xB6.C (H-2^{b/d}) mice were infected with the indicated transgenic *T. gondii* tachyzoites. Splenocytes or brain leukocytes were harvested from mice at 21 days postinfection and GRA6-specific T cell responses were measured by staining with H-2L^d-HF10 tetramers. Compiled tetramer staining data on gated CD8+ B220– splenocytes (left panel) and CD8+ brain leukocytes (right panel). *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001.

See also Figures S4 and S5.




Figure 6. Enhanced Protection when the YAL9 Epitope Is Targeted to Parasite Dense Granules

(A) Mice were immunized with lipopolysaccharide-activated BMDCs pulsed with YAL9 or control peptide (W19), boosted after 2 weeks, and after 3 weeks challenged with 5×10^5 live transgenic *T. gondii* tachyzoites i.p. (either CTG+ROP5 or CTG+GRA6-ROP5).

(B) Kaplan-Meier survival curves among the four groups of mice.

(C–F) Mice that survived the acute phase of infection were sacrificed at 7 weeks after challenge.

(C) MHC class I H-2D^b-YAL9 tetramer staining on brain leukocytes from surviving mice. Cells were also costained with CD8 antibody.

(D and E) CD8 T cells (D) and ratio of *T. gondii*-specific CD4-to-CD8 T cells in the brain (E) as measured by ICCS for IFN- γ using flow cytometry after ex vivo restimulation with *T. gondii*-infected APCs or peptide-pulsed APCs. Data are background corrected based on the values from uninfected APCs or APC pulsed with irrelevant peptide.

(F) Number of cysts in the brain as measured by staining a portion of the brain with fluorescent lectin to detect the cysts. Data (mean and SEM) are pooled from two representative experiments with at least five mice per condition in each experiment. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$; NS, not significant.

See also Figure S6.

CTG+ROP5 transgenic parasites all displayed similar growth in TAP^{-/-} mice, which lack a functional CD8 T cell response (data not shown). These data indicate that the YAL9-specific CD8 T cells generated upon challenge with CTG+GRA6-ROP5 parasites can eventually expand and predominate in the brain and control parasites during the chronic phase of infection, provided that the mice can survive the acute phase. Moreover, the generation of preexisting YAL9-specific T cells by peptide immunization can allow for survival of acute infection, but only when the mice are challenged with parasites expressing the dense granule-targeted YAL9. These data imply that both superior T cell expansion and superior protection contribute to the efficacy of the CD8 T cell response to dense granule-targeted YAL9. Moreover, these results imply that constitutively secreted endogenous parasite antigens provide better target antigens to elicit protective CD8 responses.

DISCUSSION

Intracellular protozoan parasites possess distinct secretory pathways by which they introduce their proteins into infected host cells, thus providing a means of immune detection by CD8 T cells. However, the impact of regulated secretion on CD8 T cell responses is not known. Here, we have identified a relatively ineffective CD8 T cell response in the susceptible (H-2^b) strain of mice that is directed toward a peptide (YAL9) derived from the parasite protein ROP5. ROP5 originates in the rhoptries and is injected directly into the cytoplasm of the

host cell during invasion, whereas the highly protective antigen GRA6 is a dense-granule protein that is constitutively secreted into the lumen of the parasitophorous vacuole. To test the hypothesis that the mode of protein secretion contributes to the difference in the CD8 response between ROP5 and GRA6, we engineered parasites in which the epitopes from ROP5 and GRA6 were altered in their secretion pattern. Strikingly, parasites in which the ROP5 epitope was secreted via dense granules elicited a much greater CD8 T cell response, whereas targeting the GRA6 epitope to rhoptries led to decreased responses. Moreover, the combination of immunizing mice with the ROP5 epitope and retargeting the ROP5 epitope to dense granules led to strong protection. These data will help to unravel the differences between effective and weak CD8 T cell responses, and have important implications for host-pathogen interactions and the design of CD8 vaccines against intracellular parasites.

Why does targeting epitopes to dense granules lead to improved CD8 T cell responses? Although we cannot exclude the possibility that this effect is due solely to higher overall levels of the precursor protein, we believe this is unlikely, since endogenous ROP5 is already among the most highly expressed proteins in *Toxoplasma* (Toxoplasma Genomics Resource, <http://toxodb.org/toxo/>) but evokes a relatively weak T cell response. In addition, although increased protein levels may contribute to the enhanced CD8 response elicited by the dense-granule-targeted GRA6-ROP5, this is not the case for another dense-granule-targeted version of the ROP5 epitope, GRA6-YAL9,

which is expressed at similar levels to the rhoptry-targeted ROP5 transgene but also elicits a stronger response. Moreover, the dense-granule-targeted GRA6 protein evokes a substantially higher T cell response compared with the rhoptry-targeted version of the same epitope ROP5-HF10, although both precursor proteins are expressed at similar levels. In addition, since only certain pools of parasite proteins enter host cells and become accessible to the class I MHC processing pathway, there may be little correlation between the amount of precursor protein in the parasite and the number of peptides that are ultimately displayed on the host cell surface. Lastly, although in some settings there is a correlation between antigen expression levels and the magnitude of the CD8 T cell response, this is not always the case. Indeed, some studies have revealed little correlation among the expression levels of the antigenic precursor protein, the amount of peptide that is displayed, and the magnitude of the CD8 responses (Croft et al., 2013; Dolan et al., 2011; Neefjes et al., 2011; Schwab et al., 2003).

A more likely explanation is that enhanced CD8 recognition is related to the altered spatial and/or temporal pattern of entry of the precursor protein into host cells. For example, whereas rhoptry release provides a single bolus of precursor protein during invasion, constitutive secretion via dense granules may eventually lead to a greater accumulation of the antigenic precursor protein in the infected cell, and ultimately to a higher density of peptide-MHC complexes displayed on the host cell surface.

With regard to the spatial pattern of antigen entry into host cells, at first glance it might be predicted that rhoptry targeting would favor CD8 T cell responses, since parasite proteins are directly secreted into the host cytosol and thus should be able to directly enter the classical MHC-I presentation pathway without crossing a membrane. However, reports of a direct interaction between the parasitophorous vacuole and the host ER could provide an efficient route for cross-presentation of dense-granule-targeted antigens (Goldszmid et al., 2009), thus offsetting this potential advantage of rhoptry-targeted antigens.

It is also interesting to consider that rhoptry versus dense-granule targeting may also have an impact on the spatial distribution of host cells presenting parasite antigens *in vivo*. This notion stems from the observation that parasites can secrete rhoptry contents into host cells in the absence of productive invasion (Koshy et al., 2010). Indeed, a recent study that used parasites expressing a rhoptry-targeted CRE reporter and mice with a CRE-inducible fluorescent reporter revealed that a much greater number of host cells had activated the reporter, and thus had received parasite rhoptry proteins, compared with host cells harboring parasites (Koshy et al., 2012). Thus, one could envisage that CD8 T cell antigens that are targeted for secretion via rhoptries would be presented more broadly *in vivo*, including on host cells that do not harbor parasites. Such a mechanism could help to explain the poor protection of CD8 T cells directed against rhoptry-derived antigen, since the resulting CD8 effector T cells could be distracted by interacting with noninvaded host cells rather than delivering protective cytokines to infected cells where they are most needed. Misdirected CD8 effector responses also lead to increased collateral damage by T cells, although the correlation between decreased protection

and increased parasite load argues against this as a major mechanism in this setting.

Although our data indicate that the mode of secretion has an important impact on CD8 responses, this does not imply that secretion is the sole determinant of antigenicity. Indeed, some parasite antigens derived from dense-granule proteins do not trigger strongly immunodominant CD8 T cell responses (Frickel et al., 2008). Other factors, including the route of protein trafficking after secretion, the C-terminal position of the epitope within the source antigen (Feliu et al., 2013), the ability of the peptide to bind MHC, and the affinity and precursor frequency of the responding T cells (Kotturi et al., 2008; Moon et al., 2007; Obar et al., 2008), will undoubtedly have an impact as well. Moreover, although most CD8 T cell responses appear to be targeted to invaded host cells (Goldszmid et al., 2009; Gubbels et al., 2005), the ability of particular antigens to be efficiently cross-presented by noninvaded bystander cells could potentially promote strong CD8 T cell responses independently of the mode of secretion from the parasite (Mashayekhi et al., 2011).

Apicomplexan parasites, including the disease-causing agents *Plasmodium spp.*, *Cryptosporidium*, and *Toxoplasma*, use distinct secretory organelles to deliver parasite effector molecules into the host cells and thereby manipulate host cell physiology. Entry of parasite effector proteins into host cells can potentially alert the host to the presence of an intracellular pathogen, and act as an “Achilles heel” for the parasite. Therefore, knowledge of how the secretion pattern affects CD8 T cell responses is important for the design of effective vaccines against these agents. In particular, there is intense interest in developing vaccines that elicit CD8 T cell responses to the pre-erythrocytic form of the malaria parasite *Plasmodium*. Although proteomic and genomic analyses have yielded a large number of possible candidate targets, screening and prioritizing these candidates remains a major challenge (Duffy et al., 2012; Hill, 2011; Vaughan and Kappe, 2012). Our data indicate that determining the mode of secretion of potential vaccine targets may be a powerful approach to narrow the list of vaccine candidates, and that antigens that reside within parasite dense granules may be particularly promising candidates, whereas rhoptry proteins should be avoided. In addition, our results may aid in the design of more effective vaccine strains of *Toxoplasma gondii*, an approach that is currently being explored to promote antitumor immunity (Baird et al., 2013).

EXPERIMENTAL PROCEDURES

Mice

B6.C, C57BL/6 (B6), B10.D2, DBA, CBA, MHC class II-deficient B6.129S-H2^{dIAb1-Ea}, and TAP-deficient B6.129S2-*Tap1*^{tm1Atp}/J mice were obtained from The Jackson Laboratory. B6xB6.C (H-2^{b/d}), MHC class II-deficient B6.129S-H2^{dIAb1-Ea} and TAP-deficient B6.129S2-*Tap1*^{tm1Atp}/J mice were then bred at the UC Berkeley animal facility. H-2^b ERAAP-deficient mice on the C57BL/6 background have been described previously (Hammer et al., 2006). For all immunization and infection experiments, sex- and age-matched mice were used. Mice were used with the approval of the Animal Care and Use Committee of the University of California.

Generation of Transgenic Parasites

Transgenic parasites were generated using the type III strain CTGΔ*hxgprt* (Donald et al., 1996), engineered to express luciferase and GFP. For the

ROP5 construct, DNA sequences encoding ROP5IIC, including its native promoter, were cloned into pHXGPRT adjacent to the *HXPGR*T cassette and in frame with a C-terminal 3x FLAG. For the ROP5-HF10 construct, the promoter and coding regions of the type III ROP5C allele were fused to a C-terminal HA-tag and GRA6 epitope HF10 (HPGSVNEFD). For the GRA6 construct, DNA sequences encoding GRA6 type II, including its native promoter and entire coding region, were modified by the addition of an HA epitope between amino acids 50 and 51 of GRA6. For the GRA6-YAL9 construct, the last ten amino acids (HF10) of GRA6 were replaced by the amino acids FAQLSPGQSD YAVANYFFL (where the underlined residues represent the epitope YAL9 from ROP5 type IIC). The GRA6 construct was also used as the starting plasmid for generation of GRA6-ROP5, wherein DNA sequences encoding the pseudokinase domain of ROP5 type IIC (Reese et al., 2011) were amplified and cloned downstream of the native GRA6 type II promoter, GRA6 coding sequence up to residue 50, and an in-frame HA tag. The transgenic strains were made by electroporation of linearized plasmids as described previously (Donald et al., 1996).

In Vivo Infection and Immunization

Mice were immunized i.p. with 1–5 × 10⁶ irradiated tachyzoites of the type II Prugnuiad-tomato-OVA strain (Pru) as described previously (Grover et al., 2012). To induce protection, mice were immunized in the footpad with activated B6 BMDCs loaded with synthetic peptide as described previously (Blanchard et al., 2008). After 7–14 days, the mice were given the same dose for boost and then infected 21 days after boost with a lethal dose of live Pru tachyzoites (1 × 10⁴ to 5 × 10⁵) i.p. For chronic infections, the mice were orally fed 25–50 cysts of the Pru strains or by i.p. injection of (5 × 10³ to 5 × 10⁵) live tachyzoites from parental or transgenic CTG strains. Cysts were obtained from CBA mice infected for 3–5 weeks i.p. with live 400 Pru tachyzoites. Parasite load was analyzed by semiquantitative PCR (Kirisits et al., 2000) or by counting individual labeled cysts as described previously (Grover et al., 2012).

Ex Vivo Analysis

Mice were euthanized at 3–7 weeks postinfection or 1–2 weeks postimmunization. Spleen and brain leukocyte populations were prepared as previously described (Blanchard et al., 2008). The proportion of *T. gondii*-specific or antigen-specific cells was monitored by ICCS for IFN-γ on CD4⁺ or CD8α⁺ cells. APCs were either infected the day before or pulsed with antigenic peptide on the same day and used for the ex vivo IFN-γ assay. Antigen-specific CD8α⁺ cells were also detected via staining using MHC class I peptide tetramers (see below).

Generation of *Toxoplasma gondii*-Specific T Cell Hybridomas

C57BL/6 mice were immunized with 1 × 10⁶ irradiated Pru tachyzoites (14,000 rads) i.p. Mice were euthanized at 7 days postinfection and spleens were dissociated into single-cell suspensions. *T. gondii*-specific populations were expanded in vitro by weekly restimulations with irradiated syngeneic splenocytes or irradiated MHC class II-deficient, B6.129S-H2^{dIAb1-Ea} BMDCs (2,000 rads) that were infected with irradiated Pru tachyzoites the day before.

After 8 weeks of in vitro restimulation, T cells were fused to a TCRαβ-negative, lacZ-inducible BWZ.36.CD8α fusion partner, allowing T cell hybridoma activation to be measured by the lacZ response (Malarkannan et al., 2001). The antigen specificity and MHC restriction of the hybridomas were assessed by overnight incubation with infected or uninfected BMDCs from wild-type C57BL/6 mice in the presence of MHC class I blocking antibodies (Anti-H-2K^b antibody clone: 5F1.5; anti-H-2D^b antibody clone: B22.249).

Expression Cloning

Bacteria were transformed with the *T. gondii* cDNA library cloned in pcDNA (Blanchard et al., 2008), and DNA from bacterial transformants (9 cfu/well) was screened to identify the antigen recognized by the T cell hybridoma as described previously (Blanchard et al., 2008). To further identify antigenic peptides, reverse-phase HPLC was used to separate extracts (Blanchard et al., 2008) from COS-7 cells transfected with pTg8E6.1 or pcDNA. Before fractionation, 10 pmol of the antigenic peptide YAL9 was added to the untransfected COS-7 extracts. HPLC fractions were collected in 96-well plates, dried, and

analyzed for antigenicity by addition of 5 × 10⁴ APCs/well and 1 × 10⁵ T cell hybridoma/well.

In Vitro Differentiation and Infection

Bone marrow cells obtained from mouse femurs and tibias were differentiated into DCs as described previously (Grover et al., 2012). After differentiation, the BMDCs were infected overnight with irradiated tachyzoites (14,000 rads) at various multiplicities of infection (MOIs). The next day, cells were washed twice to remove any extracellular parasites and were used in the assays.

Proteasome Inhibition

Proteasome inhibitor lactacystin was titrated (10 μM to 0 μM) onto 96-well plates containing 5 × 10⁴ BMDCs/well in complete RPMI media. Cells were incubated with the inhibitor for 2 hr at 37°C followed by the addition of irradiated tachyzoites. At 8 hr postinfection, the media were removed from treated and infected BMDCs and replaced with fresh complete media, with the addition of 1 × 10⁵ T cell hybridoma/well.

Flow Cytometry

Surface staining with anti-mouse CD4 (RM4-5), CD8α (53-6.7), IFN-γ (XMG1.2; BD Biosciences), and anti-mouse KLRG-1 and IL7R (eBiosciences or BioLegend) antibodies was done according to the manufacturer's instructions. Intracellular cytokine staining for IFN-γ was performed using the Cytotfix/Cytoperm kit (BD Pharmingen). The concentration and time of staining of the fluorescently labeled MHC class I tetramers bound to *T. gondii* antigenic peptide (NIH Tetramer Facility) were optimized and cells were incubated with the tetramer at 4°C for 1 hr. The cells were then washed and followed by surface staining at 4°C for 30 min with anti-mouse CD4 or CD8α, B220 (CD45R), KLRG-1, and IL7R antibodies. All flow-cytometry data were acquired on an XL Analyzer or FC 500 (Coulter) or BD LSRII and analyzed using FlowJo software (Tree Star).

Western Blotting and Immunofluorescence Analysis

Parasites were harvested from infected human foreskin fibroblasts (HFFs) by scraping, passaging through a 27 g needle, and filtering through a 5 μm filter. They were then washed in PBS and resuspended in either Laemmli buffer with dithiothreitol or NP-40 buffer (25 mM Tris, 100 mM NaCl, 1 mM EDTA, 0.1% NP40 + protease inhibitors). Proteins were separated by 10% SDS-PAGE and transferred to PVDF membrane. Membranes were probed with rat anti-HA antibody directly conjugated to horseradish peroxidase (HRP; Roche 3F10) or mouse monoclonal (M2) anti-3xFLAG (Sigma), and then rabbit anti-SAG1 or rabbit anti-GRA7, and probed with HRP-conjugated secondary antibodies. HRP levels were detected by chemiluminescence on film, followed by densitometric analysis (ImageJ, NIH; as described on the How to Western-Blot website, <http://howtowesternblot.net/data-analysis-3/quantification/>).

For protein localization, immunofluorescence of *T. gondii*-infected HFF cells was performed as described previously (Koshy et al., 2012). After membrane permeabilization, HA-tagged proteins were detected by rat anti-HA (Roche) followed by fluorescently conjugated goat anti-rat IgG (Invitrogen). FLAG-tagged proteins were detected with mouse monoclonal (M2) anti-3xFLAG (Sigma) followed by Alexa Fluor 488-conjugated goat anti-mouse IgG (Invitrogen). Rhoptries were marked using mouse anti-ROP2, -ROP3, -ROP4 antibody followed by Alexa Fluor 650-conjugated goat anti-mouse IgG (Invitrogen). Dense granules were marked using rabbit anti-GRA7 followed by Alexa Fluor 650-conjugated goat anti-rabbit IgG (Invitrogen). Images were acquired by standard microscopy at 600× or 1,000× magnification with minimal and equal processing in ImageJ (NIH).

Statistical Analysis

We calculated p values using the two-tailed Student's (nonparametric) t test or one-way ANOVA with Bonferroni post test in Prism (GraphPad).

SUPPLEMENTAL INFORMATION

Supplemental Information includes six figures and can be found with this article online at <http://dx.doi.org/10.1016/j.celrep.2014.04.031>.

ACKNOWLEDGMENTS

We thank Jon Boyle for access to unpublished information prior to publication and Kerry Buchholz for technical assistance and discussion of results. We are grateful to the NIH Tetramer Facility for the YAL9-D^b and the HF10-L^d tetramers, and the Cancer Research Laboratory Flow Facility at UC Berkeley for technical assistance. Imaging for immunofluorescence analysis was performed at the Stanford Neuroscience Microscopy Service, supported by NIH NS069375. This work was supported by NIH grants P01 AI065831 to E.A.R. and N.S., R01 AI065537 to E.A.R., and R01 AI076753 to J.C.B. H.S.G. was supported in part by a Cancer Research Coordinating Committee fellowship. M.L.R. was supported by an American Cancer Society fellowship. H.C. was supported by an American Heart Association postdoctoral fellowship. F.K. was supported by an NIH F32 fellowship.

Received: February 11, 2013

Revised: February 21, 2014

Accepted: April 16, 2014

Published: May 22, 2014

REFERENCES

- Baird, J.R., Fox, B.A., Sanders, K.L., Lizotte, P.H., Cubillos-Ruiz, J.R., Scarlett, U.K., Rutkowski, M.R., Conejo-Garcia, J.R., Fiering, S., and Bzik, D.J. (2013). Avirulent *Toxoplasma gondii* generates therapeutic antitumor immunity by reversing immunosuppression in the ovarian cancer microenvironment. *Cancer Res.* **73**, 3842–3851.
- Blader, I.J., and Saeij, J.P. (2009). Communication between *Toxoplasma gondii* and its host: impact on parasite growth, development, immune evasion, and virulence. *APMIS* **117**, 458–476.
- Blanchard, N., Gonzalez, F., Schaeffer, M., Joncker, N.T., Cheng, T., Shastri, A.J., Robey, E.A., and Shastri, N. (2008). Immunodominant, protective response to the parasite *Toxoplasma gondii* requires antigen processing in the endoplasmic reticulum. *Nat. Immunol.* **9**, 937–944.
- Boothroyd, J.C., and Dubremetz, J.F. (2008). Kiss and spit: the dual roles of *Toxoplasma* rhoptries. *Nat. Rev. Microbiol.* **6**, 79–88.
- Brown, C.R., and McLeod, R. (1990). Class I MHC genes and CD8+ T cells determine cyst number in *Toxoplasma gondii* infection. *J. Immunol.* **145**, 3438–3441.
- Brown, C.R., Hunter, C.A., Estes, R.G., Beckmann, E., Forman, J., David, C., Remington, J.S., and McLeod, R. (1995). Definitive identification of a gene that confers resistance against *Toxoplasma* cyst burden and encephalitis. *Immunology* **85**, 419–428.
- Carruthers, V.B. (2002). Host cell invasion by the opportunistic pathogen *Toxoplasma gondii*. *Acta Trop.* **87**, 111–122.
- Croft, N.P., Smith, S.A., Wong, Y.C., Tan, C.T., Dudek, N.L., Fleisch, I.E., Lin, L.C., Tscharke, D.C., and Purcell, A.W. (2013). Kinetics of antigen expression and epitope presentation during virus infection. *PLoS Pathog.* **9**, e1003129.
- Denkers, E.Y., and Gazzinelli, R.T. (1998). Regulation and function of T-cell-mediated immunity during *Toxoplasma gondii* infection. *Clin. Microbiol. Rev.* **11**, 569–588.
- Dolan, B.P., Bennink, J.R., and Yewdell, J.W. (2011). Translating DRiPs: progress in understanding viral and cellular sources of MHC class I peptide ligands. *Cell. Mol. Life Sci.* **68**, 1481–1489.
- Donald, R.G., Carter, D., Ullman, B., and Roos, D.S. (1996). Insertional tagging, cloning, and expression of the *Toxoplasma gondii* hypoxanthine-xanthine-guanine phosphoribosyltransferase gene. Use as a selectable marker for stable transformation. *J. Biol. Chem.* **271**, 14010–14019.
- Duffy, M.F., Selvarajah, S.A., Josling, G.A., and Petter, M. (2012). The role of chromatin in *Plasmodium* gene expression. *Cell. Microbiol.* **14**, 819–828.
- Feliu, V., Vasseur, V., Grover, H.S., Chu, H.H., Brown, M.J., Wang, J., Boyle, J.P., Robey, E.A., Shastri, N., and Blanchard, N. (2013). Location of the CD8 T cell epitope within the antigenic precursor determines immunogenicity and protection against the *Toxoplasma gondii* parasite. *PLoS Pathog.* **9**, e1003449.
- Frickel, E.M., Sahoo, N., Hopp, J., Gubbels, M.J., Craver, M.P., Knoll, L.J., Ploegh, H.L., and Grotenbreg, G.M. (2008). Parasite stage-specific recognition of endogenous *Toxoplasma gondii*-derived CD8+ T cell epitopes. *J. Infect. Dis.* **198**, 1625–1633.
- Gendrin, C., Bittame, A., Mercier, C., and Cesbron-Delauw, M.F. (2010). Post-translational membrane sorting of the *Toxoplasma gondii* GRA6 protein into the parasite-containing vacuole is driven by its N-terminal domain. *Int. J. Parasitol.* **40**, 1325–1334.
- Goldszmid, R.S., Coppens, I., Lev, A., Caspar, P., Mellman, I., and Sher, A. (2009). Host ER-parasitophorous vacuole interaction provides a route of entry for antigen cross-presentation in *Toxoplasma gondii*-infected dendritic cells. *J. Exp. Med.* **206**, 399–410.
- Gregg, B., Dzierzinski, F., Tait, E., Jordan, K.A., Hunter, C.A., and Roos, D.S. (2011). Subcellular antigen location influences T-cell activation during acute infection with *Toxoplasma gondii*. *PLoS ONE* **6**, e22936.
- Grover, H.S., Blanchard, N., Gonzalez, F., Chan, S., Robey, E.A., and Shastri, N. (2012). The *Toxoplasma gondii* peptide AS15 elicits CD4 T cells that can control parasite burden. *Infect. Immun.* **80**, 3279–3288.
- Gubbels, M.J., Striepen, B., Shastri, N., Turkoz, M., and Robey, E.A. (2005). Class I major histocompatibility complex presentation of antigens that escape from the parasitophorous vacuole of *Toxoplasma gondii*. *Infect. Immun.* **73**, 703–711.
- Hakim, F.T., Gazzinelli, R.T., Denkers, E., Hieny, S., Shearer, G.M., and Sher, A. (1991). CD8+ T cells from mice vaccinated against *Toxoplasma gondii* are cytotoxic for parasite-infected or antigen-pulsed host cells. *J. Immunol.* **147**, 2310–2316.
- Hammer, G.E., Gonzalez, F., Champsaur, M., Cado, D., and Shastri, N. (2006). The aminopeptidase ERAAP shapes the peptide repertoire displayed by major histocompatibility complex class I molecules. *Nat. Immunol.* **7**, 103–112.
- Hill, A.V. (2011). Vaccines against malaria. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **366**, 2806–2814.
- Holtappels, R., Simon, C.O., Munks, M.W., Thomas, D., Deegen, P., Kühnapfel, B., Däubner, T., Emde, S.F., Podlech, J., Grzimek, N.K., et al. (2008). Subdominant CD8 T-cell epitopes account for protection against cytomegalovirus independent of immunodomination. *J. Virol.* **82**, 5781–5796.
- Joffre, O.P., Segura, E., Savina, A., and Amigorena, S. (2012). Cross-presentation by dendritic cells. *Nat. Rev. Immunol.* **12**, 557–569.
- Karttunen, J., Sanderson, S., and Shastri, N. (1992). Detection of rare antigen-presenting cells by the lacZ T-cell activation assay suggests an expression cloning strategy for T-cell antigens. *Proc. Natl. Acad. Sci. USA* **89**, 6020–6024.
- Khan, I.A., Smith, K.A., and Kasper, L.H. (1988). Induction of antigen-specific parasitocidal cytotoxic T cell splenocytes by a major membrane protein (P30) of *Toxoplasma gondii*. *J. Immunol.* **141**, 3600–3605.
- Khan, I.A., Ely, K.H., and Kasper, L.H. (1994). Antigen-specific CD8+ T cell clone protects against acute *Toxoplasma gondii* infection in mice. *J. Immunol.* **152**, 1856–1860.
- Kirisits, M.J., Mui, E., and McLeod, R. (2000). Measurement of the efficacy of vaccines and antimicrobial therapy against infection with *Toxoplasma gondii*. *Int. J. Parasitol.* **30**, 149–155.
- Koshy, A.A., Fouts, A.E., Lodoen, M.B., Alkan, O., Blau, H.M., and Boothroyd, J.C. (2010). *Toxoplasma* secreting Cre recombinase for analysis of host-parasite interactions. *Nat. Methods* **7**, 307–309.
- Koshy, A.A., Dietrich, H.K., Christian, D.A., Melehan, J.H., Shastri, A.J., Hunter, C.A., and Boothroyd, J.C. (2012). *Toxoplasma* co-opts host cells it does not invade. *PLoS Pathog.* **8**, e1002825.
- Kotturi, M.F., Scott, I., Wolfe, T., Peters, B., Sidney, J., Cheroutre, H., von Herrath, M.G., Buchmeier, M.J., Grey, H., and Sette, A. (2008). Naive precursor frequencies and MHC binding rather than the degree of epitope diversity shape CD8+ T cell immunodominance. *J. Immunol.* **181**, 2124–2133.
- Lindberg, R.E., and Frenkel, J.K. (1977). Toxoplasmosis in nude mice. *J. Parasitol.* **63**, 219–221.

- Malarkannan, S., Mendoza, L.M., and Shastri, N. (2001). Generation of antigen-specific, lacZ-inducible T-cell hybrids. *Methods Mol. Biol.* *156*, 265–272.
- Mashayekhi, M., Sandau, M.M., Dunay, I.R., Frickel, E.M., Khan, A., Goldszmid, R.S., Sher, A., Ploegh, H.L., Murphy, T.L., Sibley, L.D., and Murphy, K.M. (2011). CD8 α (+) dendritic cells are the critical source of interleukin-12 that controls acute infection by *Toxoplasma gondii* tachyzoites. *Immunity* *35*, 249–259.
- Montoya, J.G., and Liesenfeld, O. (2004). Toxoplasmosis. *Lancet* *363*, 1965–1976.
- Moon, J.J., Chu, H.H., Pepper, M., McSorley, S.J., Jameson, S.C., Kedl, R.M., and Jenkins, M.K. (2007). Naive CD4(+) T cell frequency varies for different epitopes and predicts repertoire diversity and response magnitude. *Immunity* *27*, 203–213.
- Neeffjes, J., Jongsma, M.L., Paul, P., and Bakke, O. (2011). Towards a systems understanding of MHC class I and MHC class II antigen presentation. *Nat. Rev. Immunol.* *11*, 823–836.
- Obar, J.J., Khanna, K.M., and Lefrançois, L. (2008). Endogenous naive CD8+ T cell precursor frequency regulates primary and memory responses to infection. *Immunity* *28*, 859–869.
- Reese, M.L., Zeiner, G.M., Saeij, J.P., Boothroyd, J.C., and Boyle, J.P. (2011). Polymorphic family of injected pseudokinases is paramount in *Toxoplasma* virulence. *Proc. Natl. Acad. Sci. USA* *108*, 9625–9630.
- Sanderson, S., and Shastri, N. (1994). LacZ inducible, antigen/MHC-specific T cell hybrids. *Int. Immunol.* *6*, 369–376.
- Schaeffer, M., Han, S.J., Chtanova, T., van Dooren, G.G., Herzmark, P., Chen, Y., Roysam, B., Striepen, B., and Robey, E.A. (2009). Dynamic imaging of T cell-parasite interactions in the brains of mice chronically infected with *Toxoplasma gondii*. *J. Immunol.* *182*, 6379–6393.
- Schwab, S.R., Li, K.C., Kang, C., and Shastri, N. (2003). Constitutive display of cryptic translation products by MHC class I molecules. *Science* *301*, 1367–1371.
- Serwold, T., Gaw, S., and Shastri, N. (2001). ER aminopeptidases generate a unique pool of peptides for MHC class I molecules. *Nat. Immunol.* *2*, 644–651.
- Shen, H., Miller, J.F., Fan, X., Kolwyck, D., Ahmed, R., and Harty, J.T. (1998). Compartmentalization of bacterial antigens: differential effects on priming of CD8 T cells and protective immunity. *Cell* *92*, 535–545.
- Suzuki, Y., Joh, K., Orellana, M.A., Conley, F.K., and Remington, J.S. (1991). A gene(s) within the H-2D region determines the development of toxoplasmic encephalitis in mice. *Immunology* *74*, 732–739.
- Suzuki, Y., Joh, K., Kwon, O.C., Yang, Q., Conley, F.K., and Remington, J.S. (1994). MHC class I gene(s) in the D/L region but not the TNF-alpha gene determines development of toxoplasmic encephalitis in mice. *J. Immunol.* *153*, 4649–4654.
- Vaughan, A.M., and Kappe, S.H. (2012). Malaria vaccine development: persistent challenges. *Curr. Opin. Immunol.* *24*, 324–331.
- Wilson, D.C., Grotenbreg, G.M., Liu, K., Zhao, Y., Frickel, E.M., Gubbels, M.J., Ploegh, H.L., and Yap, G.S. (2010). Differential regulation of effector- and central-memory responses to *Toxoplasma gondii* infection by IL-12 revealed by tracking of Tgd057-specific CD8+ T cells. *PLoS Pathog.* *6*, e1000815.