

NMR studies of [$U\text{-}^{13}\text{C}$]cyclosporin A bound to human cyclophilin B

Placido Neri¹, Gerd Gemmecker¹, Lynne D. Zydowsky², Christopher T. Walsh² and Stephen W. Fesik¹

¹Pharmaceutical Discovery Division, Abbott Laboratories, Abbott Park, IL 60064, USA and ²Department of Biological Chemistry and Molecular Pharmacology, Harvard Medical School, Boston, MA 02115, USA

Received 19 July 1991

NMR data (^1H and ^{13}C chemical shifts, NOEs) on [$U\text{-}^{13}\text{C}$]cyclosporin A bound to cyclophilin B were compared to previously published data on the [$U\text{-}^{13}\text{C}$]CsA/CyPA complex [Fesik et al., (1991) *Biochemistry* 30, 6574-6583]. Despite only 64% sequence identity between CyPA and CyPB, the conformation and active site environment of CsA when bound to CyPA and CyPB are nearly identical as judged by the similarity of the NMR data.

Cyclophilin B; Cyclosporin A; NMR; Conformation

1. INTRODUCTION

The cyclophilins (CyPs) are a family of proteins found in mammalian, fungal, and prokaryotic organisms [1-12] that catalyze *cis-trans* peptidyl-prolyl isomerization [13,14]. Cyclophilin from human T cells (CyPA) as well as other cyclophilins, bind to and are inhibited by the immunosuppressive drug cyclosporin A (CsA) (Fig. 1) [1-12]. For most CsA analogs, the relative affinity for cyclophilin correlates with their immunosuppressive activity, suggesting that the immunosuppressive effects of CsA involve binding to cyclophilin [1,15-18]. However, for a few CsA analogs cyclophilin binding does not correlate with immunosuppressive activity (e.g. [MeAla⁶] and [MeBm₂t¹CsA]), questioning the role of cyclophilin in CsA-mediated immunosuppression [17,18].

One possible explanation for these apparent discrepancies is that CsA exerts its immunosuppressive effects by binding to a different, yet structurally related, cyclophilin. Recently, a second human cyclophilin, cyclophilin B (CyPB) [19], has been cloned and was found to be 64% identical to human CyPA. Unlike CyPA which is cytosolic, CyPB contains a hydrophobic N-terminal signal sequence which is removed upon expression in *E. coli* [19]. Recent localization studies have shown CyPB to be located in the endoplasmic reticulum, Golgi and vesicles in the cytoplasm (Jin, et al.,

unpublished). CyPB possesses *cis-trans* peptidyl-prolyl isomerase (PPIase) activity which is inhibited by CsA [19] at three-fold higher concentration than CyPA. Thus, CyPB may also play a significant role in T-cell activation.

In this report we describe NMR studies on the binding of [$U\text{-}^{13}\text{C}$]CsA to human cyclophilin B and compare the results to previously described NMR studies on the [$U\text{-}^{13}\text{C}$]CsA/CyPA complex [20].

2. MATERIALS AND METHODS

2.1. Materials

[$U\text{-}^{13}\text{C}$]CsA was isolated from *Beauveria nivea* ATCC 34921 (American Type Culture Collection; Rockville, MD) culture containing [$U\text{-}^{13}\text{C}$]Celtone medium (Marterk Corp., Columbia, MD) [20]. Recombinant human cyclophilin A was prepared as recently described [21].

A genetically truncated human cyclophilin B (residues 24-208) was overexpressed in *E. coli* and purified to homogeneity as previously described [19].

2.2. NMR experiments

The cyclophilin samples (A and B) were exchanged into a D₂O solution (pH 6.5) containing phosphate buffer (50 mM), NaCl (100 mM), and dithiothreitol (5 mM) and concentrated to ~1.0 mM with a centricon YM-10 (Amicon).

The [$U\text{-}^{13}\text{C}$]CsA/cyclophilin (A and B) complexes were prepared by gently shaking a suspension of ~1.5 mol equivalents of [$U\text{-}^{13}\text{C}$]CsA in a D₂O solution of cyclophilin at 6°C for 12 h under argon. The excess CsA was removed by centrifugation.

All NMR spectra were acquired on a Bruker AMX500 NMR spectrometer at 20°C. The spectra were processed using in-house written software on Silicon Graphics computers or the Bruker UXNMR software package on a Bruker X32.

The heteronuclear single quantum correlation (HSQC) spectra [22] were acquired using 2048 complex points in F₂ and 512 complex points in F₁, with 96 scans. The sweep width was 7812.5 Hz in F₂ and 7462.7 Hz in F₁.

The 3D HMQC-NOESY experiments [23] were acquired as 48(t₁)

Abbreviations: CyP, cyclophilin (A or B); CsA, cyclosporin A; NOE, nuclear Overhauser effect; HSQC, heteronuclear single quantum correlation; HMQC-NOESY, heteronuclear multiple quantum correlation-nuclear Overhauser effect spectroscopy.

Correspondence address: S.W. Fesik, Pharmaceutical Discovery Division, D-47G, AP9, Abbott Laboratories, Abbott Park, Illinois 60064, USA. Fax: (1) (708) 938-2478.




Fig. 1. (a) Structure of CsA. (b) Conformation of CsA when bound to CyPA [20].

$\times 128 (t_2) \times 2048 (t_3)$ complex points using a mixing time of 50 ms and spectral widths of 7462.7 Hz in ω_1 and 7812.5 Hz in ω_2 and ω_3 . 32 acquisitions were collected per t_3 experiment for a total experimental time of approximately 6 days.

3. RESULTS

As a first step in probing for any structural differences between the $[U-^{13}C]CsA/CyPA$ and $[U-^{13}C]CsA/CyPB$ complexes, we compared their proton NMR spectra. As expected for two proteins with 64% sequence identity, the proton NMR spectra of the two CsA/CyP complexes, which mainly consisted of protein signals (data not shown), were markedly different. In contrast, as shown in the two-dimensional heteronuclear single quantum correlation (HSQC) spectra [22] of $[U-^{13}C]CsA$ bound to CyPA (Fig. 2a) and CyPB (Fig. 2b), the 1H and ^{13}C chemical shifts of $[U-^{13}C]CsA$ when bound to either protein are nearly identical. The assignment of the signals was easily deduced by a direct comparison of the two HSQC spectra and confirmed in the 3D HMQC-NOESY spectra [20]. The observed chemical shift differences in ppm ($\delta_{CyPB} - \delta_{CyPA}$) were: $1H^\beta$ (-0.04), $5H^\alpha$ (-0.07), $8H^\alpha$ (-0.05), $9H^\alpha$ (-0.10), $9H^\beta$

(-0.16, -0.06), $9C^\beta$ (-0.3), $11NCH_3$ (+0.05), $11H^\alpha$ (-0.04), $11C^\alpha$ (+0.4). All other signals resonate at the same frequencies within the limits of experimental error [20]. In Fig. 2 some signals are invisible at the level chosen for the plot, but can be observed with lower contouring. The residual protein cross-peaks that are detected in the HSQC spectra (indicated by P in Fig. 2) are different in the two cases, reflecting the sequence differences in the two proteins.

In Fig. 3 three-dimensional HMQC-NOESY spectra [23] are compared for $[U-^{13}C]CsA$ bound to CyPA (top of each panel a-e) and CyPB (bottom of each panel a-e). 1H , 1H planes were extracted at the same ^{13}C frequencies in the two data sets. As shown by the comparison of the spectra obtained for the two complexes, similar NOEs, with the same relative intensities, were observed. Some weak NOEs present in the CypA dataset could not be observed in the 3D NOE spectrum of CypB due to a lower signal-to-noise ratio.

In Fig. 1b the conformation of CsA when bound to CyPA is shown that had been previously determined from 3D NOE data [20]. The bound conformation of CsA [20,24,25] was found to be very different from the previously determined NMR-derived or X-ray crystal structure of cyclosporin A determined in the absence of cyclophilin A [26]. When bound to cyclophilin, CsA adopts a *trans* 9,10 peptide bond [24] in contrast to the 9,10 *cis* peptide bond found in free CsA [26]. In addition, the backbone conformation and the orientations of some of the side chains greatly differ between the structures [20,25].

It is important to note key long range NOEs important in defining the overall backbone conformation of CsA when bound to CyPA (Fig. 1b) [20] were also observed in the 3D NOE spectrum of the CsA/CyPB complex. Some of these include NOEs from $6NCH_3$ to $5H^\alpha$, $1NCH_3$ and $10NCH_3$ (Fig. 3a) and NOEs from $5H\gamma^2$ to $1NCH_3$ and $3NCH_3$ (Fig. 3b). NOEs between CsA protons indicative of a particular side chain conformation were also found to be the same in the CsA/CyPA [20] and CsA/CyPB complexes. For example, the α -proton of MeBmt¹ is close to its own β and δCH_3 side chain protons (Fig. 3c) and MeLeu⁶(H ^{α}) is in close proximity to MeLeu⁶(H ^{α}) (Fig. 3b) in both complexes. NOEs between CsA and the protein were also found to be similar in the two complexes (e.g. Fig. 3d). An exception is shown in Fig. 3e in which an additional NOE between MeVal¹¹ (NCH₃) and an aromatic proton of the protein (arrow) was observed in the $[U-^{13}C]CsA/CyPB$ complex. Another difference is shown in Fig. 2d in which a small change in chemical shift of one of the NOE peaks involving the protein and MeVal¹¹(H γ^2) was observed, suggesting the CyP binding pocket in the vicinity of the MeVal¹¹ CsA residue is slightly different in the two complexes. This is supported by the differences in chemical shift of the $11NCH_3$, $11H^\alpha$, and $11C^\alpha$ signals of CsA when bound to CyPA or CyPB.




Fig. 2. Contour plots of the HSQC spectra of [U-¹³C]CsA bound to (a) CyPA and (b) CyPB. The 'x' refers to experimental artifacts. *The MeBmt¹(H^c) signals were not fully ¹³C-decoupled at the power level employed.




Fig. 3. Comparison of corresponding cross-sections (ω_2 , vertical axis; ω_3 , horizontal axis) from 3D-HMQC-NOESY spectra of $[U-^{13}C]CsA$ bound to CyPA (top of each panel) and CyPB (bottom of each panel), acquired with a mixing time of 50 ms. In both spectra the planes were taken at the same ^{13}C chemical shifts (ω_1) of (a) 31.8, (b) 19.4, (c) 58.1, (d) 16.3, (e) 33.2 ppm. The skewed solid lines indicate the diagonal peaks on the $^1H-^1H$ NOESY planes.

4. DISCUSSION

Despite only 64% sequence identity between CyPA and CyPB, the chemical shift and NOE data suggest that the conformation and active site environment of CsA when bound to CyPA and CyPB is very similar. Even for CsA residues (e.g. residue 1 and 6) in which structural changes produced analogs in which cyclophilin binding and immunosuppressive activity did not correlate [17,18], the chemical shifts and NOEs involving these residues were found to be identical. These results suggest that the apparent discrepancies in CyP binding and immunosuppressive activity displayed by some CsA analogs cannot simply be explained by invoking CyPB as the biologically relevant cyclophilin. Indeed, the relative amounts of PPIase inhibition for CyPA and CyPB with [MeAla⁶] and [MeBm₂t¹]CsA show little difference with the two cyclophilin isoforms (Anderson et al., unpublished). It may be that the complexes of cyclophilins (A and/or B) with cyclosporin or CsA analogs may exhibit differential affinities with partner proteins immediately downstream in the signal transduction cascade and that is the relevant immunosuppressive readout.

Acknowledgements: We thank M. Jackson and J. Hochlowski for the preparation of [U-¹³C]CsA. P. Neri acknowledges support from a fellowship from the CNR, Italy (Bando no. 203.03.22, Com. Scienze Chimiche). This work was supported in part by National Institutes of Health Grant GM20011 to C.T.W. and National Institutes of Health NSRA Grant ES05459 to L.D.Z.

REFERENCES

- [1] Handschumacher, R.E., Harding, M.W., Rice, J., Drugge, R.J. and Speicher, D.W. (1984) *Science* 226, 544-547.
- [2] Harding, M.W., Handschumacher, R.E. and Speicher, D.W. (1986) *J. Biol. Chem.* 261, 8547-8555.
- [3] Koletsky, A.J., Harding, M.W. and Handschumacher, R.E. (1986) *J. Immunol.* 137, 1054-1059.
- [4] Tropschug, M., Nicholson, D.W., Hartl, F.-U., Kohler, H., Pfanner, N., Wachter, E. and Neupert, W. (1988) *J. Biol. Chem.* 263, 14433-14440.
- [5] Shieh, B.-H., Stammes, M.A., Seavello, S., Harris, G.L. and Zuker, C.S. (1989) *Nature* 338, 67-70.
- [6] Liu, J. and Walsh, C.T. (1990) *Proc. Natl. Acad. Sci. USA* 87, 4028-4032.
- [7] Dietmeier, L. and Tropschug, M. (1990) *Nucleic Acids Res.* 18, 373.
- [8] Haendler, B., Keller, R., Heistand, P.C., Kocher, H.P., Wegmann, G. and Movva, R.N. (1989) *Gene* 83, 39-46.
- [9] Koser, P.L., Sylvester, D., Livi, G.P. and Bergsma, D.J. (1990) *Nucleic Acids Res.* 18, 1643.
- [10] Gasser, C.S., Gunning, D.A., Budelier, K.A. and Brown, S.M. (1990) *Proc. Natl. Acad. Sci. USA* 87, 9519-9523.
- [11] Hayano, T., Takahashi, N., Kato, S., Maki, N. and Suzuki, M. (1991) *Biochemistry* 30, 3041-3048.
- [12] Liu, J., Albers, M., Chen, C.-M., Schreiber, S.L. and Walsh, C.T. (1990) *Proc. Natl. Acad. Sci. USA* 87, 2304-2308.
- [13] Takahashi, N., Hayano, T. and Suzuki, M. (1989) *Nature* 337, 473-475.
- [14] Fischer, G., Wittman-Liebold, B., Lang, K., Kiefhaber, T. and Schmid, F.X. (1989) *Nature* 337, 476-478.
- [15] Quesniaux, V.F.J., Schreier, M.H., Wenger, R.M., Hiestand, P.C., Harding, M.W. and Van Regenmortel, M.H.V. (1987) *Eur. J. Immunol.* 17, 1359-1365.
- [16] Quesniaux, V.F.J., Schreier, M.H., Wenger, R.M., Hiestand, P.C., Harding, M.W. and Van Regenmortel, M.H.V. (1988) *Transplantation* 46, 23S-28S.
- [17] Durette, P.L., Boger, J., Dumont, F., Firestone, R., Frankshun, R.A., Koprak, S.L., Lin, C.S., Melino, M.R., Pessolano, A.A., Pisano, J., Schmidt, J.A., Sigal, N.H., Staruch, M.J. and Witzel, B.E. (1988) *Transplant. Proc.* 20, 51-57.
- [18] Sigal, N.H., Dumont, F., Durette, P., Siekierka, J.J., Peterson, L., Rich, D.H., Dunlap, B.E., Staruch, M.J., Melino, M.R., Koprak, S.L., Williams, D., Witzel, B. and Pisano, J.M. (1991) *J. Exp. Med.* 173, 619-627.
- [19] Price, E.R., Zydowsky, L.D., Jin, M., Baker, C.H., McKeon, F.D. and Walsh, C.T. (1991) *Proc. Natl. Acad. Sci. USA* 88, 1903-1907.
- [20] Fesik, S.W., Gampe, Jr. R.T., Eaton, H.L., Gemmecker, G., Olejniczak, E.T., Neri, P., Holzmann, T.F., Egan, D.A., Edalji, R., Simmer, R., Helfrich, R., Hochlowski, J. and Jackson, M. (1991) *Biochemistry* 30, 6574-6583.
- [21] Holzman, T.F., Egan, D.A., Edalji, R., Simmer, R.S., Helfrich, R., Taylor, A. and Burres, N.S. (1991) *J. Biol. Chem.* 266, 2474-2479.
- [22] Bodenhausen, G. and Ruben, D.J. (1980) *Chem. Phys. Lett.* 69, 185-189.
- [23] Fesik, S.W. and Zuiderweg, E.R.P. (1988) *J. Magn. Reson.* 78, 588-593.
- [24] Fesik, S.W., Gampe, Jr. R.T., Holzman, T.F., Egan, D.A., Edalji, R., Luly, J.R., Simmer, R., Helfrich, R., Kishore, V. and Rich, D.H. (1990) *Science* 250, 1406-1409.
- [25] Weber, C., Wider, G., von Freyberg, B., Traber, R., Braun, W., Widmer, H. and Wüthrich, K. (1991) *Biochemistry* 30, 6563-6574.
- [26] Loosli, H.R., Kessler, H., Oschkinat, H., Weber, H.-P., Petcher, T.J. and Widmer, T.J. (1985) *Helv. Chim. Acta* 68, 682-704.