

Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

## Genomics Data

journal homepage: <http://www.journals.elsevier.com/genomics-data/>

# Computational identification of microRNAs and their targets in *Catharanthus roseus* expressed sequence tags

Alok Pani, Rajani Kanta Mahapatra \*

School of Biotechnology, Campus-11, KIIT University, Bhubaneswar, Odisha 751024, India

## ARTICLE INFO

## Article history:

Received 31 March 2013

Received in revised form 27 June 2013

Accepted 28 June 2013

Available online 14 August 2013

## Keywords:

Vincristine

microRNAs

ESTs

Computational prediction

*C. roseus*

## ABSTRACT

No study has been performed on identifying microRNAs (miRNAs) and their targets in the medicinal plant, *Catharanthus roseus*. In the present study, using the comparative genomics approach, we have predicted two potential *C. roseus* miRNAs. Furthermore, twelve potential mRNA targets were identified in *C. roseus* genome based on the characteristics that miRNAs exhibit perfect or nearly perfect complementarity with their targeted mRNA sequences. Among them many of the targets were predicted to encode enzymes that regulate the biosynthesis of terpenoid indole alkaloids (TIA). In addition, most of the predicted targets were the gene coding for transcription factors which are mainly involved in cell growth and development, signaling and metabolism. This is the first in silico study to indicate that miRNA target gene encoding enzymes involved in vinblastine and vincristine biosynthesis, which may help to understand the miRNA-mediated regulation of TIA alkaloid biosynthesis in *C. roseus*.

© 2013 The Authors. Published by Elsevier Inc. Open access under [CC BY-NC-ND license](http://creativecommons.org/licenses/by-nc-nd/4.0/).

## 1. Introduction

The medicinal plant *Catharanthus roseus* L. G. Don is of enormous pharmaceutical potential because of the presence of >120 terpenoid indole alkaloids (TIAs), some of which are known to exhibit strong pharmacological activities [1]. Vinblastine and vincristine, the antineoplastic bisindole alkaloids are produced only in trace amounts; vinblastine (0.01%) and vincristine (0.003%). Furthermore, with significant international efforts cell cultures are also not yet a valid alternative for production for these low yielding secondary metabolites [2]. Therefore, a deeper understanding of the regulatory system governing TIA metabolism is of particular interest and could eventually make successful metabolic engineering of alkaloid biosynthesis possible.

Several hundred miRNAs have been identified in plants by computational and experimental approaches [3,4]. However, a little is known about experimental or computational identification of miRNA in *C. roseus* species. *C. roseus* belongs to Apocyanaceae family, reflecting a disparity between the important values of this plant family and insufficient molecular and genetic studies, including small RNA mediated gene regulation. So to gain insight into miRNAs and their important regulatory functions in terpenoid indole alkaloid biosynthetic pathway, we studied

miRNA and their targets in *C. roseus* genome using computational approach.

## 2. Materials & methods

### 2.1. miRNA reference set

To search potential miRNAs, a total of previously known 328 miRNAs from *Arabidopsis thaliana* were obtained from miRNA Registry Database (Release 18.0, November 2011). Although some of these *A. thaliana* miRNAs were initially identified by computational approaches, a majority of them have been validated by experimental approaches including direct cloning, PCR, Northern blotting, and/or 5' rapid amplification of cDNA end (5'RACE) [3]. To avoid the overlap of miRNAs, the redundant miRNA sequences were removed manually and the remaining sequences were used as a reference miRNA for homologous prediction in *C. roseus*. We have referred to the previous work on computational prediction of miRNAs [5].

### 2.2. Availability of software

Comparative software BLAST-2.2.14 was used from NCBI Genbank. MFOLD 3.1 was used online to analyze secondary structure of RNAs. MirEval (<http://tagc.univ-mrs.fr/mireval>) was used to predict miRNA precursor [6]. These precursor sequences were used for BLASTx analysis for removing the protein-coding sequences and retained only non-protein encoding sequences. BLASTn from NCBI (<http://www.ncbi.nlm.nih.gov>) was used to analyze potential targets of miRNAs.

\* Corresponding author. Tel.: +91 9337627961.

E-mail address: [rmahapatra@kiitbiotech.ac.in](mailto:rmahapatra@kiitbiotech.ac.in) (R.K. Mahapatra).

### 2.3. Prediction of miRNAs

Procedure for searching potential miRNAs in *C. roseus* is shown in Fig. 1. The non-redundant miRNA sequences after initial screening were subjected to BLASTn search for *C. roseus* miRNA homolog against EST database. The adjusted BLASTn parameters setting were as follows: expect value was set at 1000 to increase the hit chance for more potential sequences; low complexity was chosen as the sequence filter; the number of descriptions and alignments were raised to 1000. The default word match size between the query and a database sequence was 7. RNA sequences were considered as miRNA candidates only if they fit the following criteria: (1) at least 18 nt length were adopted between the predicted and mature miRNAs and (2) allowed to have 0–3 nt mismatches in sequence with all previously known plant mature miRNAs. The ESTs that closely match the previously known plant mature miRNAs were included in the set of miRNA candidates and used for additional characterization by subjected to evaluation for miRNA precursor prediction properties using mirEval software. These precursor sequences were used for BLASTX analysis for removing the protein-coding sequences and retaining only the non-protein-coding sequences.

### 2.4. Prediction of secondary structure

Precursor sequences of these potential miRNA homologs were subjected to hairpin structure prediction using the Zuker folding algorithm with MFOLD 3.1 [7]. In brief, the following criteria were applied in designating the RNA sequence as a miRNA homolog described by (1) the sequence could fold into an appropriate stem-loop hairpin secondary structure; (2) the small RNA sits in one arm of the hairpin structure; (3) no more than 6 mismatches are between the predicted mature miRNA sequence and its opposite miRNA (miRNA\*) sequence

in the secondary structure; (4) no loop or break is in the miRNA or miRNA\* sequences, and (5) predicted secondary structure has higher MFEI and negative MFE [8].

The MFEI was calculated using the following equation:

$$\text{MFEI} = [(\text{MFE}/\text{length of the RNA sequence}) \times 100]/(\text{G} + \text{C}) \%$$

MFE denotes the negative folding free energies ( $\Delta G$ ).

### 2.5. Prediction of mRNA targets of miRNAs

In brief, we used the potential *C. roseus* miRNA BLAST analysis against the *C. roseus* mRNA database to search sequences conforming to the following standards as the *C. roseus* candidate targets gene. (1) The maximum number of mismatched nucleotides between the mature miRNA and its potential target genes was four; (2) the maximum number of mismatched nucleotides at positions 1–9 was one; (3) no mismatches were allowed at positions 10–11; (4) more than two continuous mismatches at any position were not allowed [9].

### 2.6. Analysis of GO and KEGG pathway

To better understand the function of *C. roseus* miRNAs, Blast2go [10,11] was employed to investigate the predicted target genes. First, the identified miRNA targeted mRNAs were used to BLASTX against NR database. Second, the best hits identified by BLASTX were further searched against the GO and KEGG databases using default settings.




Fig. 1. Procedure of potential *C. roseus* miRNA gene search by identifying homologs of previously known *A. thaliana* miRNAs.

### 3. Results

#### 3.1. Potential miRNAs in *C. roseus*


Following the procedure depicted in Fig. 1, 19910 ESTs from *C. roseus* were searched against 321 mature miRNA sequences of *Arabidopsis*. In total, two potential miRNAs were predicted in *C. roseus*. The two identified *C. roseus* candidate miRNAs belong to miR5021 family. The two predicted microRNAs are having maximal mismatches of 2 & 3 respectively against its homolog (Fig. 2). The length of the two EST is 611 nt and 592 nt while the precursor's length is 80 nt and 100 nt respectively as identified by MirEVal software. The mature miRNA sequences are located at the 3' end of the miRNA precursors. Minimal folding free energy (MFE) is an important characteristic that determines the secondary structure of nucleic acids (DNA and RNA). The lower the MFE is, the higher the thermodynamically stable secondary structure of the corresponding sequence is [12]. The MFE value of the structures is estimated to  $-17.90$  kcal/mol and  $-34.70$  kcal/mol respectively. We have also calculated the MFE index (MFEI) for each sequence, to avoid false calling of other RNAs as miRNA candidates. The MFE index (MFEI) for each sequence was calculated as previously reported [13]. In this study, the MFEI value is 0.57 and 0.83 respectively. During the screening of the potential miRNAs, candidate miRNAs were evaluated for A + U content. The sequences of the miRNA precursors have A + U content of 61.25% and 58% respectively (Table 1), which is in agreement with the previous results [5]. It is estimated that in plants, approximately 10,000 ESTs contain 1 miRNA [14]. Therefore, the total of 19,910 ESTs in *C. roseus* examined in this study may contain 1–2 miRNAs. *C. roseus* belongs to Apocyanaceae family. Unfortunately, not a single miRNA from Apocyanaceae family has been deposited in the MiRbase [15]. We

expect that as more miRNAs of this plant family are publicly available, more miRNAs will be identified in *C. roseus*. ESTs are partially transcribed gene sequences, which have been used to confirm the existence and expression of potential miRNAs predicted by computational approaches in *Arabidopsis*, rice and maize [16,17]. In this study, we also tested the predicted *C. roseus* miRNAs individually against the EST databases of GenBank. Our BLASTn search results indicated that several predicted periwinkle miRNAs exist in *C. roseus* EST databases, suggesting that these miRNAs were expressed in the *Catharanthus* genome. Moreover, many of the reference set miRNAs from *Arabidopsis* which are used for homology search in *C. roseus* have been validated by experimental approaches. So the result from the computational prediction will be useful to guide experimental design for biological verification.

### 4. Discussion

#### 4.1. *C. roseus* miRNA targets and their functions

The miRNA-regulated genes control a variety of biological and metabolic processes. Gaining insight into the miRNA targets will help us to understand the spectrum of miRNA regulation and elucidate the functional importance of miRNAs. Increasing evidences have demonstrated that most plant miRNAs bind to their target mRNA sequences with perfect or near-perfect sequence complementarity [18,19]. This provides a powerful strategy for discovering potential miRNA targets by comparing and aligning miRNAs with mRNAs sequences. Here, we performed more stringent criterion [19] to identify potential *C. roseus* targets. After a set of screening criteria as described in the method, we achieved 12 target genes. Out of these 12 target genes many they have



**Fig. 2.** Mature and precursor sequences and the predicted stem and loop structures of newly identified miRNAs in *C. roseus*. The mature miRNAs are highlighted in cyan. A. miR5021 (EG560894; homolog of *A. thaliana*); B. miR5021 (EG558543; homolog of *A. thaliana*).

**Table 1**Newly identified miRNAs from ESTs of *C. roseus*.

miRNA	Reference species	Gene ID	EST length (nt)	NM (nt)	LM (nt)	LP (nt)	Side	A + U (%)	MFE	MFEI
miR-5021	<i>A. thaliana</i>	EG560894	697	2	20	80	3'	61.25	17.90	0.57
	<i>A. thaliana</i>	EG558543	795	3	20	100	3'	58	34.70	0.83

Note: NM, number of mismatch; LM, length of mature miRNAs; LP, length of precursor; MFE, minimal folding free energy; MFEI, minimal folding free energy index.

**Table 2**Potential targets of the identified miRNAs in *C. roseus*.

miRNA family	Target accession ID	Target description	Function
mir-5021 (EG558543)	EF625593	MYB transcription factor	Transcription factor
	EF625552	ADP-ribosylation factor 1	Protein transport
	EF625539	Geranylgeranyl diphosphate synthase	Isoprenoid biosynthetic process
	EF625523	Gamma-tocopherol methyltransferase related protein	Methyltransferase activity
	EF625416	GCPE protein	Terpenoid biosynthetic process
mir-5021 (EG560894)	EF625531	UDP-glucose glucosyltransferase	Transferase activity
	EF625513	Cytochrome c oxidase subunit I	Oxidative phosphorylation
	EF625489	Type-A response regulator	Regulation of Transcription
	EF625469	UDP-glucose iridoid glucosyltransferase	Transferase activity
	EF625448	Magnesium chelatase subunit H	Biological process
	EF625362	Chloroplast terpenoid cyclase	Terpene synthase activity
	EF661875	Putative secretory peroxidase	Response to oxidative stress

different biological functions, including the terpenoid biosynthesis, transcription regulation, cell growth and development (Table 2). These predicted mRNA targets are statistically significant in compare to the other mRNA targets with least Expect (E) value.

Out of several mRNA targets predicted; one target gene identified for *C. roseus* miRNA (EG560894) is UDP-glucose iridoid glucosyltransferase. This enzyme is responsible for glucosylation step in the biosynthetic pathway of the secondary metabolite, iridoid in higher plants [20]. They are pharmacologically active principles in various medicinal plants and key intermediates in the biosynthesis of monoterpenoid indole alkaloids as well as quinoline alkaloids. In addition, strictosidine synthase is another potential of target of miR5021 in *C. roseus*. The transcriptionally regulated strictosidine synthase condenses tryptamine and the iridoid secologanin to yield strictosidine, the universal precursor of TIA [21]. Interestingly, another *Arabidopsis* miR5021 homolog in *C. roseus* is predicted to target GCPE protein. GCPE protein is involved in the terpenoid biosynthesis process [22].

It is expected that freeing the expression of key enzymatic activities form the strict regulation to which they are normally subjected is expected to increase the flux through the pathway and product formation. Therefore, the next major steps are to successfully apply metabolic engineering approach, to regulate the enzymes through miRNA mediated regulation. This will help to explore unconventional alternate strategies that are economically viable for the commercial production of indole alkaloids.

In addition EG560894 was also predicted to target type-A response regulator (EF625489), which is involved in the final steps of a histidine-to-aspartate phosphorelay in cytokinin (CK) signaling in *C. roseus*. CKs are plant growth regulators with pleiotropic functions in plants. They can also control some secondary metabolite biosynthetic pathways in *Arabidopsis* [23] and terpenoid indole alkaloids (TIAs) in periwinkle cells [24].

Recent studies have shown that miRNAs are also involved in plant adaptation to environmental stresses, such as cold [25,26], salt [27], drought [28], and nutrient deficiency [29,30]. Interestingly, we identified one target of miR-5021 is putative secretory peroxidase [EF661875] which response to oxidative stress. Further analysis of Gene ontology (GO) suggested that they are involved in peroxidase activity.

*C. roseus* miRNA also target a gene chloroplast geranylgeranyl diphosphate synthase (EF625539) that is involved in isoprenoid

biosynthesis. Isoprenoids are a large and highly diverse group of natural products with many functions in plant primary and secondary metabolism [31]. Another *Arabidopsis* miR-5021 homolog of *C. roseus*, EG558543 targets ADP-ribosylation factor 1. It is a GTP binding protein involved in the regulation of vesicle-mediated protein transport through the secretory pathway [32].

Many miRNA targets identified by bioinformatics and/or experimental methods were transcription factors that help control plant growth and development. In this study we found a target of miR-5021 called MYB transcription factor. MYB proteins are key factors in regulatory networks controlling development, metabolism and responses to biotic and abiotic stresses [33]. In *Arabidopsis*, the other functions of MYB transcription factor are (i) primary and secondary metabolism, (ii) cell fate and identity, (iii) developmental processes and (iv) responses to biotic and abiotic stresses. Magnesium chelatase subunit H is also predicted to be a target for mir5021 homolog in *C. roseus*. Previous studies have shown that magnesium-protoporphyrin IX (ProtoIX) chelatase large subunit (Mg chelatase H subunit; CHLH) is an ABA receptor. It mediates ABA responses in seed germination, post germination growth, and stomatal movement [34]. CHLH has multiple functions in plant cells. As a subunit of the Mg-chelatase, CHLH catalyzes the introduction of Mg to ProtoIX, a key regulatory step of chlorophyll biosynthesis. In addition, CHLH plays a key role in mediating plastid-to-nucleus retrograde signaling [35].

**Table 3**GO analysis of miRNA targets in *C. roseus*.

miRNAs	Biological Process	Accession IDs for the targets	GOs
mir-5021 (EG558543)	DNA binding	EF625593	GO: 0003677
	GTP binding	EF625552	GO: 0005525
	Geranyltranstransferase activity	EF625539	GO: 0004337
	Methyltransferase activity	EF625523	GO: 0008168
	Iron ion binding	EF625416	GO: 0005506
mir-5021 (EG560894)	Transferase activity	EF625531	GO: 0016758
	Oxidative phosphorylation	EF625513	GO: 0004129
	Regulation of transcription	EF625489	GO: 0000156
	Transferase activity	EF625469	GO: 0016758
	Biological process	EF625448	GO: 0009058
	Terpene synthase activity	EF625362	GO: 0010333
	Response to oxidative stress	EF661875	GO: 0004601



**Table 4**  
KEGG analysis of miRNA targets in *C. roseus* (Homology Search in *A. thaliana*).

miRNAs	Accession ID for targets	Target description	Enzyme	Pathways
mir-5021 (EG558543)	EF625416	GCPE protein	EC: 1.17.7.1	Terpenoid backbone biosynthesis
	EF625539	Geranylgeranyl diphosphate synthase, type II	EC: 2.5.1.1	Terpenoid backbone biosynthesis
mir-5021 (EG560894)	EF625531	UDP-glucose: glycoprotein glucosyltransferase	EC: 2.4.1.-	Protein processing in ER
	EF625513	Cytochrome c oxidase subunit 1	EC: 1.9.3.1	Oxidative phosphorylation
	EF625448	Magnesium chelatase subunit H	EC: 6.6.1.1	Porphyryn and chlorophyll metabolism

To further understand the function of *C. roseus* miRNAs, the predicted target mRNAs were subjected to analysis by GO and Kyoto Encyclopedia of Genes and Genomes (KEGG), a database for analyzing gene functions systematically [36]. The result suggested that *C. roseus* miRNAs were involved in various biological processes such as oxidation–reduction process, regulation of transcription, transport, growth and development, metabolism and translation (Table 3). Pathway enrichment analysis, based on the KEGG database, demonstrates that the identified miRNAs participated in four metabolic networks. These networks were involved in terpenoid backbone biosynthesis, protein processing, porphyrin and chlorophyll metabolism, oxidative phosphorylation, and other secondary metabolites biosynthesis process (Table 4). Obviously; our study will help further understanding of the important regulatory roles of miRNAs in *C. roseus* growth and development, stress response, and likewise in research and development to augment the production of vinblastine and vincristine biosynthesis.

## 5. Conclusion

*C. roseus* (periwinkle) is the sole source of anticancerous alkaloids vinblastine and vincristine which today is widely used for treatment of cancer. Ajmalicine is used as an antihypertensive alkaloid. Recent report demonstrates that, in addition to its anticancerous properties, the extracts from the leaves of this plant can be used as prophylactic agent in many infectious diseases. The unfortunately low yield of the antineoplastic bisindole alkaloids and with the failure of the alternative production systems, for example by in vitro culture of *C. roseus* cells, an attempt can be made for the genetic manipulation of the alkaloid biosynthesis pathway for higher production levels. Our study is the first in silico study to identify miRNAs and their targets in *C. roseus*, which we hope could help to better understand miRNA-mediated regulation of genes related to terpenoid indole alkaloids biosynthesis.

## Competing interests

The authors have declared that no competing interests exist.

## Acknowledgments

AP and RKM thank Dr. Mrutyunjay Suar, Director of School of Biotechnology in KIIT University for his encouragement and support and the Bioinformatics lab facility of School of Biotechnology, KIIT University during the course of work.

## References

- R. Van der Heijden, D.I. Jacobs, W. Snoeijer, D. Hallard, R. Verpoorte, The *Catharanthus* alkaloids: pharmacognosy and biotechnology. *Curr. Med. Chem.* 11 (2004) 607–628.
- R.L. Noble, The discovery of the vinca alkaloids—chemotherapeutic agents against cancer. *Biochem. Cell Biol.* 68 (1990) 1344–1351.
- S. Griffiths-Jones, R.J. Grocock, S. van Dongen, A. Bateman, A.J. Enright, miRBase: microRNA sequences, targets and gene nomenclature. *Nucleic Acids Res.* 34 (2006) D140–D144.
- B.H. Zhang, X.P. Pan, C.H. Cannon, G.P. Cobb, T.A. Anderson, Conservation and divergence of plant microRNAs genes. *Plant J.* 46 (2006) 243–259.
- B. Zhang, Q. Wang, K. Wang, X. Pan, F. Liu, T. Guo, G.P. Cobb, T.A. Anderson, Identification of cotton microRNAs and their targets. *Gene* 397 (2007) 26–37.
- W. Ritchie, F.X. Théodule, D. Gautheret, Mireval: a web tool for simple microRNA prediction in genome sequences. *Bioinformatics* 24 (11) (2008) 1394–1396.
- M. Zuker, Mfold web server for nucleic acid folding and hybridization prediction. *Nucleic Acids Res.* 13 (2003) 3406–3415.
- L. Wang, H. Liu, D. Li, H. Chen, Identification and characterization of maize microRNAs involved in the very early stage of seed germination. *BMC Genomics* 12 (18 2011) 154.
- F.Xie, T.P.Frazier, B.Zhang, Identification and characterization of micro rRNA and their targets in the bioenergy plant switchgrass (*panicum virgatum*). *Planta* 232 (2010) 417–434.
- A.Conesa, S.Gotz, J.M.Garcia-Gomez, J.Terol, M.Talon, M.Robles, Blast2go: a universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics* 21 (2005) 3674–3676.
- A.Conesa, S.Gotz, Blast2go: a comprehensive suite for functional analysis in plant genomics. *Int. J. Plant Genomics* (2008) 619832.
- G.R.Prabu, A.K.A.Mandal, Computational identification of miRNAs and their target genes from expressed sequence tags of tea (*Camellia sinensis*). *Genom. Proteomics Bioinforma.* 8 (2010) 113–121.
- Z.Yin, C.Li, X.Han, F.Shen, Identification of conserved microRNAs and their target genes in tomato (*Lycopersicon esculentum*). *Gene* 414 (2008) 60–66.
- B.H.Zhang, X.P.Pan, Q.L.Wang, G.P.Cobb, T.A.Anderson, Identification and characterization of new plant microRNAs using EST analysis. *Cell Res.* 15 (2005) 336–360.
- S.Griffiths-Jones, H.K.Saini, S.van Dongen, E.J.Enright, miRBase: tools for microRNAs genomics. *Nucleic Acids Res.* 36 (2008) D154–D158.
- E.Bonnet, J.Wuyts, P.Rouzé, Y.Van de Peer, Detection of 91 potential conserved plant microRNAs in *Arabidopsis thaliana* and *Oryza sativa* identifies important target genes. *Proc. Natl. Acad. Sci. U. S. A.* 101 (2004) 11511–11516.
- B.Zhang, X.Pan, T.A.Anderson, Identification of 188 conserved maize microRNAs and their targets. *FEBS Lett.* 580 (2006) 3753–3762.
- X.J.Wang, J.L.Reyes, N.H.Chua, T.Gaasterland, Prediction and identification of arabidopsis thaliana microRNAs and their mRNA targets. *Genome Biol.* 5 (9) (2004).
- R.Schwab, J.F.Palatnik, M.Riester, C.Schommer, M.Schmid, D.Weigel, Specific effects of microRNAs on the plant transcriptome. *Dev. Cell* 8 (2005) 517–527.
- M.Nagatoshi, K.Terasaka, A.Nagatsuo, H.Mizukami, An iridoid-specific glucosyltransferase from *Gardenia jasminoides*. *J. Biol. Chem.* 286 (2011) 32866–32874.
- A.De Waal, A.H.Meijer, R.Verpoorte, Strictosidine synthase from *Catharanthus roseus*: purification and characterization of multiple forms. *Biochem. J.* 306 (1995) 571–580.
- A.Oudin, S.Mahroug, V.Courdavault, N.Hervouet, C.Zelwer, M.Rodríguez-Concepción, B.St-Pierre, V.Burlat, Spatial distribution and hormonal regulation of gene products from methyl erythritol phosphate and monoterpenes-secoiridoid pathways in *Catharanthus roseus*. *Plant Mol. Biol.* 65 (2007) 13–30.
- J.Deikman, P.E.Hammer, Induction of anthocyanin accumulation by cytokinins in *Arabidopsis thaliana*. *Plant Physiol.* 108 (1995) 47–57.
- N.Papon, A.Oudin, A.Vansiri, M.Rideau, J.C.Chénieux, J.Crêche, Differential expression of two type-A response regulators in plants and cell cultures of *Catharanthus roseus* (L.) G. Don. *J. Exp. Bot.* 2003 (2003) 1793–1795.
- J.Zhang, Y.Xu, Q.Huan, K.Chong, Deep sequencing of brachypodium small rRNA at the global genome level identifies microRNAs involved in cold stress response. *BMC Genomics* 10 (2009) 449.
- F.Thiebaut, C.A.Rojas, K.L.Almeida, C.Gratiwo, G.C.Domiciano, C.R.Lamb, A.Engler Jde, A.C.Hemerly, P.C.Ferreira, Regulation of mir319 during cold stress in sugarcane. *Plant Cell Environ.* 35 (2012) 502–512.
- D.Ding, L.Zhang, H.Wang, Z.Liu, Z.Zhang, Y.Zheng, Differential expression of miRNAs in response to salt stress in maize roots. *Ann. Bot.* 103 (2009) 29–38.
- B.Li, Y.Qin, H.Duan, W.Yin, X.Xia, Genome-wide characterization of new and drought stress responsive microRNAs in populus euphratica. *J. Exp. Bot.* 62 (2011) 3765–3779.
- R.Sunkar, V.Chinnusamy, J.Zhu, J.K.Zhu, Small rRNA as big players in plant abiotic stress responses and nutrient deprivation. *Trends Plant Sci.* 12 (2007) 301–309.
- M.Zhao, H.Tai, S.Sun, F.Zhang, Y.Xu, W.X.Li, Cloning and characterization of maize miRNAs involved in responses to nitrogen deficiency. *PLoS One* 7 (2012) e29669.
- J.Kirby, J.D.Keasling, Biosynthesis of plant isoprenoids: perspectives for microbial engineering. *Annu. Rev. Plant Biol.* 60 (2009) 335–355.
- F.Regad, C.Bardet, D.Tremousaygue, A.Moisan, B.Lescure, cDNA cloning and expression of an Arabidopsis GTP-binding protein of the ARF family. *FEBS Lett.* 316 (1993) 133–136.
- C.Dubos, R.Stracke, E.Grotewold, B.Weisshaar, C.Martin, L.Lepiniec, MYB transcription factors in Arabidopsis. *Trends Plant Sci.* 15 (2010) 573–581.
- Y.Y.Shen, X.F.Wang, F.Q.Wu, S.Y.Du, Z.Cao, Y.Shang, X.L.Wang, C.C.Peng, X.C.Yu, S.Y.Zhu, R.C.Fan, Y.H.Xu, D.P.Zhang, The Mg-chelatase H subunit is an abscisic acid receptor. *Nature* 443 (2006) 823–826.
- A.Nott, H.S.Jung, S.Koussevitzky, J.Chory, Plastid-to-nucleus retrograde signaling. *Annu. Rev. Plant Biol.* 57 (2006) 739–759.
- M.Kanehisa, S.Goto, Kegg, Kegg encyclopedia of genes and genomes. *Nucleic Acids Res.* 28 (2000) 27–30.