


Cell Host & Microbe

The Cytosolic Sensor cGAS Detects *Mycobacterium tuberculosis* DNA to Induce Type I Interferons and Activate Autophagy

Graphical Abstract



Authors

Robert O. Watson, Samantha L. Bell, Donna A. MacDuff, ..., Christina L. Stallings, Herbert W. Virgin, Jeffery S. Cox

Correspondence

jeffery.cox@ucsf.edu

In Brief

During *Mycobacterium tuberculosis* infection, cytosolic bacterial DNA triggers cell-intrinsic innate immune responses. Watson et al. show that the cytosolic DNA sensor cGAS recognizes *M. tuberculosis* DNA. This sensing is required for activating pro-bacterial type I interferon expression as well as targeting *M. tuberculosis* to the anti-bacterial selective autophagy pathway.

Highlights

- cGAS is essential for type I IFN production during *M. tuberculosis* infection
- cGAS activation targets *M. tuberculosis* to the selective autophagy pathway
- cGAS deficiency leads to increased intracellular *M. tuberculosis* growth
- cGAS directly binds to *M. tuberculosis* genomic DNA during infection



The Cytosolic Sensor cGAS Detects *Mycobacterium tuberculosis* DNA to Induce Type I Interferons and Activate Autophagy

Robert O. Watson,^{1,6,7} Samantha L. Bell,^{1,7} Donna A. MacDuff,² Jacqueline M. Kimmey,³ Elie J. Diner,⁴ Joanna Olivas,¹ Russell E. Vance,^{4,5} Christina L. Stallings,³ Herbert W. Virgin,² and Jeffery S. Cox^{1,*}

¹Department of Microbiology and Immunology, Program in Microbial Pathogenesis and Host Defense, University of California, San Francisco, San Francisco, CA 94158, USA

²Department of Pathology and Immunology, Washington University School of Medicine, Saint Louis, MO 63110, USA

³Department of Molecular Microbiology, Washington University School of Medicine, Saint Louis, MO 63110, USA

⁴Department of Molecular & Cell Biology, University of California, Berkeley, Berkeley, CA 94720, USA

⁵Howard Hughes Medical Institute, University of California, Berkeley, Berkeley, CA 94720, USA

⁶Present address: Department of Microbiology and Immunology, Texas A&M Health Science Center, Bryan, TX 77807, USA

⁷Co-first author

*Correspondence: jeffery.cox@ucsf.edu

<http://dx.doi.org/10.1016/j.chom.2015.05.004>

SUMMARY

Type I interferons (IFNs) are critical mediators of antiviral defense, but their elicitation by bacterial pathogens can be detrimental to hosts. Many intracellular bacterial pathogens, including *Mycobacterium tuberculosis*, induce type I IFNs following phagosomal membrane perturbations. Cytosolic *M. tuberculosis* DNA has been implicated as a trigger for IFN production, but the mechanisms remain obscure. We report that the cytosolic DNA sensor, cyclic GMP-AMP synthase (cGAS), is required for activating IFN production via the STING/TBK1/IRF3 pathway during *M. tuberculosis* and *L. pneumophila* infection of macrophages, whereas *L. monocytogenes* short-circuits this pathway by producing the STING agonist, c-di-AMP. Upon sensing cytosolic DNA, cGAS also activates cell-intrinsic antibacterial defenses, promoting autophagic targeting of *M. tuberculosis*. Importantly, we show that cGAS binds *M. tuberculosis* DNA during infection, providing direct evidence that this unique host-pathogen interaction occurs in vivo. These data uncover a mechanism by which IFN is likely elicited during active human infections.

INTRODUCTION

Innate immune cells discriminate pathogens from non-pathogens at the earliest stages of infection and tailor their responses to match the level of the threat (Vance et al., 2009). A fundamental way this is achieved is through sensing membrane perturbations mediated by bacterial virulence factors, either directly or via the recognition of specific bacterial molecules in the cytosol (Manzanillo et al., 2012; Thurston et al., 2012; Vance et al., 2009). Cytosolic detection leads to activation of three

potent antimicrobial effector pathways—the inflammasome (Lamkanfi and Dixit, 2011), autophagy (Watson et al., 2012), and the cytosolic surveillance pathway (CSP) characterized by elicitation of type I interferons (IFNs) (Monroe et al., 2010; O’Riordan et al., 2002). While type I IFNs (IFN- α , IFN- β) are potent antiviral signaling molecules, they inhibit antibacterial signaling pathways (Mayer-Barber et al., 2011; Teles et al., 2013) and promote infection of many intracellular bacteria including *Mycobacterium tuberculosis* (Manca et al., 2001; Manzanillo et al., 2012) and *Listeria monocytogenes* (Auerbuch et al., 2004; Carrero et al., 2004; O’Connell et al., 2004), suggesting that bacterial pathogens have evolved mechanisms to activate this antiviral pathway for their benefit. Likewise, clinical studies have shown that elevated levels of type I IFNs are a biomarker of active TB disease in humans (Berry et al., 2010), indicating that this pathway is engaged during bacterial replication in vivo. Elicitation of IFN- β requires the pore-forming toxin listeriolysin O during *L. monocytogenes* infection (O’Riordan et al., 2002) and the membrane-disrupting activity of the ESX-1 secretion system during *M. tuberculosis* infection (Manzanillo et al., 2012). *L. monocytogenes* actively secretes the bacterial second messenger cyclic diadenylate monophosphate (c-di-AMP) that binds to the host protein STING and activates the STING/TBK1/IRF3 signaling axis to promote a signature transcriptional response that includes IFN- β (Burdette et al., 2011; Sauer et al., 2011). In contrast to *L. monocytogenes*, *M. tuberculosis* and another intracellular bacterial pathogen, *Legionella pneumophila*, appear to activate this same STING-dependent pathway (Lippmann et al., 2008; Manzanillo et al., 2012; Monroe et al., 2010) via recognition of pathogen-derived nucleic acids, although the evidence for this is indirect.

Surprisingly, the STING pathway also potentially activates autophagy, a degradative pathway implicated in resistance to intracellular pathogens (Birmingham et al., 2006; Deretic and Levine, 2009; Zhao et al., 2008). Activation of STING and the kinase TBK1 leads to targeting of bacteria and cytosolic DNA to the ubiquitin-mediated selective autophagy pathway in macrophages (Watson et al., 2012), and ATG5, a core autophagy

protein, is crucial for limiting *M. tuberculosis* growth during infection. Despite our growing understanding of the links between the CSP and selective autophagy, the nature of the nucleic acid ligand and the host receptor proteins involved remain major unanswered questions.

Recent studies have identified a DNA sensor, cyclic GMP-AMP synthase (cGAS), as the central cytoplasmic DNA sensor upstream of STING during viral infection (Gao et al., 2013; Schoggins et al., 2014; Sun et al., 2013). Upon binding dsDNA, cGAS synthesizes the secondary messenger cGAMP, which in turn binds to and activates STING, leading to the production of IFN- β through IRF3. Despite some controversy over the involvement of a variety of other DNA receptors (Burdette and Vance, 2013), cGAS is absolutely required for IFN induction upon both DNA transfection and viral infection and is likely the major cytosolic DNA receptor (Li et al., 2013). cGAS is required for IFN- β induction during *Chlamydia trachomatis*, but its functional role during bacterial infections is largely unknown (Zhang et al., 2014).

Here, we report two different mechanisms by which bacterial pathogens activate the CSP: *M. tuberculosis* and *L. pneumophila* activate cGAS during infection, while *L. monocytogenes* bypasses cGAS and directly activates STING via c-di-AMP. Furthermore, we show that recognition of *M. tuberculosis* by cGAS is a critical host-pathogen interaction, as it promotes the delivery of bacilli to the ubiquitin-mediated selective autophagy pathway during macrophage infection, and has an unexpected role in cell-autonomous bacterial control. Importantly, we also show that cGAS binds *M. tuberculosis* genomic DNA during macrophage infection, providing evidence of direct interactions between an *M. tuberculosis* ligand and a host sensor in vivo.

RESULTS

cGAS Is Required to Induce the CSP during Intracellular Bacterial Infection

To test the role of cGAS during *M. tuberculosis* infection, primary murine bone marrow-derived macrophages (BMDMs) were isolated from *cGas*^{-/-} and *Sting*^{-/-} mice. These cells were unresponsive to transfection with interferon-stimulatory DNA (ISD), a 45 bp dsDNA sequence sufficient for activating cGAS (Stetson and Medzhitov, 2006), as measured by qPCR of IRF3 targets (IFN- β and IFIT1 mRNAs, Figure S1A) and IRF3 phosphorylation (Figure S1G). Delivery of the STING ligand cGAMP via transfection bypassed the cGAS requirement (Figure S1B), in agreement with previous reports in other cell types establishing that cGAS functions upstream of STING (Sun et al., 2013).

Importantly, infection with the Erdman strain of *M. tuberculosis* led to robust CSP activation in wild-type BMDMs as measured by both protein and transcript levels. This response was completely blocked in *cGas*^{-/-} and *Sting*^{-/-} macrophages (Figures 1A and 1B), similar to infection of wild-type macrophages with an ESX-1 mutant, which fails to induce the CSP (Manzanillo et al., 2012). We observed similar results using a clinical isolate of *M. tuberculosis* (CDC1551), which, unlike the Erdman strain, has the capacity to produce the STING agonist c-di-GMP (Figure 1C) (Manzanillo et al., 2012). This response is specific to

the STING/TBK1/IRF3 pathway, as the mRNA level of TNF α , which is not regulated by IRF3, was generally unaffected by these mutations (Figure 1D). Independent shRNA knockdown of cGAS in the RAW 264.7 murine macrophage cell line (Figure S1C), and in the U937 human macrophage cell line (Figure S1D), confirmed the key role of this receptor in responding to cytosolic DNA (Figures S1E and S1F) and *M. tuberculosis* infection (Figures 1E and S1H). These data demonstrate that cGAS is the major sensor that activates the CSP during *M. tuberculosis* infection of macrophages. These results also provide strong evidence that CSP activation by wild-type *M. tuberculosis* is due to exposure of DNA in the cytosol and support previous findings that endogenous levels of bacterial-produced cyclic dinucleotides are not a major contributor in triggering this response during macrophage infection (Dey et al., 2015; Manzanillo et al., 2012).

We next tested the role of cGAS during infection with another intracellular bacterial pathogen, *L. pneumophila*, which elicits type I IFNs by accessing the cytosol through its type IV secretion system (T4SS) (Stetson and Medzhitov, 2006). *L. pneumophila* infection failed to induce the CSP in *cGas*^{-/-} BMDMs (Figure 1F). In addition, *sdhA* mutant *L. pneumophila* cells, which enter into the cytoplasm more readily than wild-type bacteria (Creasey and Isberg, 2012), induced significantly higher levels of IFN- β mRNA (Monroe et al., 2009; Vance et al., 2009), and this too was dependent on cGAS (Figure 1F). The absolute requirement for cGAS is consistent with the notion that DNA is exposed to the cytosol during *L. pneumophila* infection via vacuolar perforations from its T4SS (Monroe et al., 2010; Stetson and Medzhitov, 2006).

In contrast with *M. tuberculosis*, CSP activation by *L. monocytogenes* infection was mostly independent of cGAS (Figure 1G). In *cGas*^{-/-} BMDMs there was only a slight reduction in IFN- β and no decrease in IFIT1 mRNA levels compared to infected wild-type BMDMs, whereas the response was completely blocked in *Sting*^{-/-} BMDMs (Sauer et al., 2011). The absolute requirement of STING, but not cGAS, indicates that *L. monocytogenes* successfully short-circuits the CSP pathway by providing its own second messenger, whereas *M. tuberculosis* and *L. pneumophila* generate the signal via DNA binding to cGAS.

Finally, infection with *Salmonella enterica* serovar Typhimurium led to normal levels of IFN- β and IFIT1 mRNAs in both *cGas*^{-/-} and *Sting*^{-/-} BMDMs, indicating that this pathogen predominantly activates type I IFNs via a CSP-independent mechanism, likely via the TLR4/TRIF pathway (Figure 1H) (Kawai and Akira, 2010; Zughraier et al., 2005). Taken together, these data suggest that intracellular bacterial pathogens have evolved the ability to activate type I IFNs via different mechanisms.

cGAS Targets Cytosolic DNA and *M. tuberculosis* to the Selective Autophagy Pathway

Newly-recognized connections between DNA sensing and autophagy suggests that cGAS activation may also promote bacterial clearance (Liang et al., 2014; Manca et al., 2001; Manzanillo et al., 2012; Watson et al., 2012). To test the role of cGAS in autophagic targeting, we transfected murine embryonic fibroblasts (MEFs) with Cy3-labeled dsDNA and observed that a similar percentage of DNA puncta colocalized with cGAS as with the selective autophagy markers LC3, ubiquitin, NDP52,




Figure 1. cGAS Is Essential for Induction of the Cytosolic Surveillance Pathway during Intracellular Bacterial Infection

(A) BMDMs were infected with WT or ΔESX-1 *M. tuberculosis* (Erdman) for 4 hr, and IFN-β and IFIT1 transcripts were measured by RT-qPCR. mRNA levels are expressed as percentages relative to infected WT cells.

(B) Same as (A), but IFN-β protein was measured by ELISA (left) or with ISRE-luciferase cells (right) 24 hr post-infection.

(C) Same as (A), but with *M. tuberculosis* strain CDC1551.

(D) Same as (A), but with TNFα.

(E) Same as (A), but with U937 knockdown cell lines. mRNA levels are expressed as percentage relative to infected scramble control cells.

(F–H) BMDMs were infected for 4 hr, and transcript levels were measured during infection with *L. pneumophila* ΔflaA (*L.p.*) or ΔflaAΔsdhA (*L.p.*ΔsdhA) (F), *L. monocytogenes* (*L.m.*) (G), or *S. Typhimurium* (*S. T.*) (H).

n.s., not significant; **p < 0.005 by two-tailed t test comparing to WT or SCR infected with WT bacteria. Error bars indicate SD. See also Figure S1.

did not observe the cGAS-independent contribution to this effect. As an additional method of testing the role of cGAS in targeting dsDNA to the selective autophagy pathway, we measured the levels of LC3-II, the lipidated form of LC3 generated during activation of autophagy, in wild-type, *cGas*^{-/-}, and *Sting*^{-/-} BMDMs after transfection with dsDNA. Western blot analysis revealed that while wild-type BMDMs had increased LC3-II levels after dsDNA transfection, *cGas*^{-/-} and *Sting*^{-/-} BMDMs failed to induce LC3-II conversion

and activated phospho-TBK1 (pTBK1) (Figures S2A and S2B). Multicolor fluorescence microscopy revealed that the large majority of cGAS+ dsDNA structures contained all of these selective autophagy markers in both transfected MEFs and RAW 264.7 cells (Figures 2A–2C). Importantly, in *cGas*^{-/-} macrophages, colocalization of the autophagy-targeting components, ubiquitin and LC3, with transfected DNA was reduced by approximately 50% (Figure 2D). Curiously, we consistently observed a stronger reduction in selective autophagy marker colocalization with DNA in *Sting*^{-/-} macrophages compared to *cGas*^{-/-} cells, suggesting that other factors, perhaps additional DNA sensors including IFI204 (Manzanillo et al., 2012), work upstream of STING to target cytosolic DNA to autophagy. The requirement for STING in selective autophagic targeting is consistent with our previous finding that TBK1 is also required for this effect, whereas IRF3 is only required for the transcriptional output of this pathway (Watson et al., 2012). ShRNA-mediated knockdown of cGAS or STING in human or murine macrophage cell lines also led to decreased recruitment of selective autophagy markers to dsDNA 4 hr post-transfection (Figures 2E and 2F), though we

did not observe the cGAS-independent contribution to this effect. As an additional method of testing the role of cGAS in targeting dsDNA to the selective autophagy pathway, we measured the levels of LC3-II, the lipidated form of LC3 generated during activation of autophagy, in wild-type, *cGas*^{-/-}, and *Sting*^{-/-} BMDMs after transfection with dsDNA. Western blot analysis revealed that while wild-type BMDMs had increased LC3-II levels after dsDNA transfection, *cGas*^{-/-} and *Sting*^{-/-} BMDMs failed to induce LC3-II conversion

Importantly, the effect of cGAS on autophagy is not due to general defects in bulk autophagy, as starvation of macrophages led to LC3-II conversion and degradation regardless of genotype (Figure S2C). To begin to test the role of cGAS in autophagic targeting of *M. tuberculosis*, we infected RAW 264.7 cells expressing epitope-tagged cGAS with mCherry-expressing *M. tuberculosis* and monitored cGAS localization. Early in infection, cGAS colocalized with approximately 20% of wild-type *M. tuberculosis*, and this colocalization decreased at later time points (Figures 3A and 3B). Importantly, we observed significantly decreased colocalization of cGAS with ESX-1 mutant bacteria at all time points, confirming that recognition requires ESX-1-mediated cytosolic access. As observed with transfected DNA, the majority of cGAS+ bacilli also colocalized with pTBK1 and LC3, suggesting that cGAS is important for recruiting these selective autophagy markers to intracellular *M. tuberculosis* (Figures 3C and 3D). Time course studies showed that the appearance of cGAS on bacilli coincided with pTBK1 recruitment but preceded the recruitment of the terminal autophagy marker LC3 (Figure 3E),




Figure 2. cGAS Is Required to Target Cytosolic DNA to the Ubiquitin-Mediated Selective Autophagy Pathway

(A) MEFs expressing 3xFLAG-tagged cGAS transfected with Cy3-labeled plasmid DNA for 4 hr and immunostained for 3xFLAG or indicated markers.

(B) Quantification of cGAS+ Cy3-DNA co-stained with indicated marker from (A). Differences are not statistically significant.

(C) RAW 264.7 cells stably expressing 3xFLAG-cGAS transfected with Cy3-DNA for 4 hr and immunostained for 3xFLAG and indicated markers.

(D) Quantification of ubiquitin and LC3 colocalization with Cy3-DNA 4 hr post-transfection in BMDMs.

(E) Same as (D), but with RAW 264.7 knockdown cell lines.

(F) Same as (D), but with U937 knockdown cell lines.

(G) BMDMs were transfected with interferon-stimulatory DNA (ISD) for 2 hr, and LC3-II conversion was analyzed by quantitative western blot (left) and expressed as a fold increase in the ratio of LC3-II/Actin. Results are representative of at least three independent experiments.

* $p < 0.05$, ** $p < 0.005$ by two-tailed t test comparing to SCR unless otherwise indicated. Error bars indicate SD. See also Figure S2.

which is consistent with the notion that cGAS is a proximal sensor of bacteria that leads to subsequent targeting.

We next determined the requirement of cGAS for targeting *M. tuberculosis* to the selective autophagy pathway. Consistent with our previous observations, *Sting*^{-/-} BMDMs were defective in recruiting ubiquitin, LC3, NDP52, pTBK1, and Atg12, giving rise to an approximately 75%–80% decrease in colocalization of any of these markers with bacteria compared with wild-type (Figures 3F, 3G, and S3A–S3C) (Leber et al., 2008; Watson et al., 2012; Woodward et al., 2010). Importantly, *cGas*^{-/-} BMDMs were also defective for autophagic targeting of *M. tuberculosis*, with a reduction in colocalization of ~50%, which mirrors our observations with transfected cytosolic dsDNA. Stable shRNA knockdowns of cGAS or STING in human and mouse macrophage cell lines led to similar results (Figures 3H and 3I). Consistent with the recruitment of selective autophagy markers to *M. tuberculosis*, wild-type BMDMs had increased levels of LC3-II conversion,

while *cGas*^{-/-} and *Sting*^{-/-} BMDMs were partially defective for LC3-II conversion after infection (Figure 3J). Together, these data indicate that DNA sensing by cGAS is required for full targeting of *M. tuberculosis* to the ubiquitin-mediated selective autophagy pathway and that a cGAS-independent pathway is also at play.

Previous work demonstrated that ATG5 is important for controlling intracellular *M. tuberculosis* replication in vivo (Gutierrez et al., 2004; Watson et al., 2012). In contrast, IRF3-driven type I IFNs appear to work in a cell-extrinsic fashion, promoting infection in vivo (Manca et al., 2001; Manzanillo et al., 2012). Given the potential positive and negative roles of these two outputs of STING/TBK1 activation during infection, we sought to determine the overall role of cGAS in *M. tuberculosis* pathogenesis. Infection of *cGas*^{-/-} mice led to partially decreased type I IFN levels when compared with wild-type mice, indicating that while this pathway is activated by *M. tuberculosis* during

infection, other pathways can stimulate IFN in vivo (Figures 3L and 3M). Moreover, these animals had no overt defect in overall resistance to *M. tuberculosis*, as we observed similar bacterial burdens in the tissues of these mice at both early and late time points after infection (Figures 3N and 3O), and none of the *cGas*^{-/-} succumbed to infection during the 100-day experiment (Figure 3P). Consistent with these observations, a parallel study by Collins et al. (2015) using the same *M. tuberculosis* strain but different *cGas*^{-/-} mice (Li et al., 2013) also found that *M. tuberculosis*-infected *cGas*^{-/-} mice have similar bacterial loads and cytokine levels compared to wild-type mice through an extended time course of infection. In these studies, Collins et al. (2015) report a small increase in susceptibility of *cGas*^{-/-} mice to *M. tuberculosis* manifest after the 100-day time point, a phenotype that was not observed in *Sting*^{-/-} mice. Although we do not understand the basis for these small discrepancies, both studies support the overall conclusion that removing cGAS from the context of an intact immune system is insufficient to dramatically alter host resistance to *M. tuberculosis*. This is in stark contrast to both *Irf3*^{-/-} and myeloid-specific *Atg5*^{-/-} mice (Manzanillo et al., 2012; Watson et al., 2012). Since we observed a significant reduction in autophagic targeting in *cGas*^{-/-} macrophages, we sought to determine the cell-intrinsic contribution of cGAS in controlling *M. tuberculosis* replication in macrophages. Consistent with cGAS targeting *M. tuberculosis* to the selective autophagy pathway for destruction by lysosomes, *cGas*^{-/-} and *Sting*^{-/-} BMDMs were permissive for *M. tuberculosis* growth, resulting in bacterial numbers 3-fold higher than those of wild-type macrophages 5 days post-infection (Figure 3K). Thus, while cGAS appears to be dispensable for overt bacterial resistance in a mouse model, this receptor is required for cell-intrinsic bacterial killing in macrophages. While we do not understand why the susceptibility of *cGas*^{-/-} macrophages does not translate to a similar phenotype during infection of the entire mouse, we suspect that compensatory factors manifest only in vivo, such as the involvement of additional host DNA sensors or functionally redundant autophagic targeting pathways, may be at play. Alternatively, the decrease of pro-bacterial type I IFNs in *cGas*^{-/-} mice during *M. tuberculosis* infection may be balanced by the reduced antibacterial capacity of selective autophagy in the macrophages of *cGas*^{-/-} mice. Finally, while cGAS activation leads to both interferon and autophagic targeting in macrophages, specific functions of IRF3 and ATG5 (Choi et al., 2014; Zhao et al., 2008) independent of their roles in interferon production and autophagic targeting, respectively, may still be active in *cGas*^{-/-} mice.

cGAS Binds *M. tuberculosis* Genomic DNA In Vivo in an ESX-1-Dependent Manner

Our finding that cGAS is activated by *M. tuberculosis* supports an interaction between a bacterial-derived molecule (DNA) and a host cytosolic sensor in vivo. However, surprisingly few bona fide physical interactions have been identified between *M. tuberculosis* and macrophages in the context of infection. This prompted us to explore the molecular mechanism by which *M. tuberculosis* activates cGAS by detecting a physical association of cGAS and DNA within macrophages. Consistent with previous findings, we were able to efficiently co-precipitate cGAS following transfection with biotin-labeled ISD in macro-

phages (Figure 4A) (Liang et al., 2014; Sun et al., 2013). To perform the reciprocal experiment using unlabeled DNA, we adapted a chromatin-IP protocol to capture DNA bound by cGAS in live cells. We first transfected macrophages with ISD, and after formaldehyde crosslinking and immunoprecipitating with α -FLAG beads (Figure S4A) we measured the abundance of ISD by qPCR. Importantly, we identified an enrichment of ISD only in immunoprecipitates from FLAG-cGAS-expressing cells transfected with ISD, but not in untransfected cells or cells expressing cGAS with a different epitope tag (Strep-cGAS, Figure S4B).

We next utilized this ChIP-like methodology to determine if cGAS binds directly to *M. tuberculosis* genomic DNA during infection. We infected cGAS-expressing macrophages with *M. tuberculosis*, crosslinked and immunoprecipitated cGAS (Figure 4B), and measured the abundance of several *M. tuberculosis* DNA sequences, including the IS6110 transposon and the CRISPR repeat, two repetitive elements in the *M. tuberculosis* genome (Figure 4C). These *M. tuberculosis*-derived sequences were significantly enriched in FLAG-cGAS immunoprecipitates as compared to controls (FLAG-GFP and Strep-cGAS). Importantly, we observed an enrichment of mycobacterial DNA bound by cGAS in cells infected with wild-type *M. tuberculosis* compared to cells infected with ESX-1 mutant bacteria, which cannot actively access the cytosol (Figure 4C). The *M. tuberculosis* DNA sequences that were detected in the ESX-1 mutant-infected cells likely arise from nonspecific phagosomal rupture that can occur when macrophages undergo necrotic cell death during infections with high bacterial burdens (Lee et al., 2006). Furthermore, we did not observe any change in the abundance of host-derived DNA sequences in the immunoprecipitates after *M. tuberculosis* infection (Figure 4D), although the low levels of host sequences that were detected may be due to low levels of nonspecific binding of cGAS to nuclear DNA during normal cell processes (Schoggins et al., 2014). Together, these data show that bacterial-derived DNA exposed to the cytosol is bound by cGAS during infection.

DISCUSSION

Our data, as well as the work presented in the accompanying papers (Collins et al., 2015; Ablasser et al., 2015), support a model in which *M. tuberculosis* triggers the STING/TBK1 pathway using the ESX-1 secretion system to disrupt phagosomal membranes, allowing bacterial DNA access to cGAS in the cytosol. Subsequent activation of TBK1 likely couples a number of cellular events, including the IRF3-dependent IFN response and, as described here, antibacterial autophagic targeting. Although indirect evidence had implicated the role of cytosolic bacterial DNA in triggering these pathways (Manzanillo et al., 2012; Watson et al., 2012), here we provide direct evidence that bacterial DNA is a bona fide pathogen-associated molecular pattern sensed by *M. tuberculosis*-infected cells. The mechanism by which DNA is liberated from *M. tuberculosis* and how ESX-1 permeabilizes phagosomal membranes to allow cytosolic access are currently unknown. However, we suspect that the nucleic acid may be sensed by cGAS at the bacterial cell surface, providing important spatial cues for the selective recruitment of autophagic vesicles.




Figure 3. cGAS Is Required to Target *Mycobacterium tuberculosis* to the Ubiquitin-Mediated Selective Autophagy Pathway

(A) RAW 264.7 cells stably expressing 3xFLAG-cGAS infected with mCherry *M. tuberculosis* for 45 min and immunostained for 3xFLAG.
 (B) Quantification of (A) during the indicated time course.
 (C) RAW 264.7 cells stably expressing 3xFLAG-cGAS infected with mCherry *M. tuberculosis* for 45 min and immunostained for 3xFLAG and indicated markers.
 (D) Quantification of (C).
 (E) Quantification of (C) during the indicated time course.
 (F) BMDMs infected with mCherry *M. tuberculosis* and immunostained for ubiquitin and LC3 4 hr post-infection.
 (G) Quantification of (F).
 (H) Quantification of ubiquitin- and LC3-positive *M. tuberculosis* in RAW 264.7 knockdown cell lines.
 (I) Same as (H), but with U937 knockdown cell lines.
 (J) BMDMs were infected with *M. tuberculosis* for 2 hr, and LC3-II conversion was analyzed by quantitative western blot (left) and expressed as a fold change in the ratio of LC3-II/Actin. Results are representative of at least three independent experiments.
 (K) RAW 264.7 cells infected with *M. tuberculosis* for 120 hr. CFU counts were determined.
 (L) RAW 264.7 cells infected with *M. tuberculosis* for 120 hr. Relative transcript levels (IFIT1/Actin) were determined.
 (M) RAW 264.7 cells infected with *M. tuberculosis* for 120 hr. RLU levels were determined.
 (N) RAW 264.7 cells infected with *M. tuberculosis* for 120 hr. Lung CFU counts were determined.
 (O) RAW 264.7 cells infected with *M. tuberculosis* for 120 hr. Spleen CFU counts were determined.
 (P) Survival curves for RAW 264.7 cells infected with *M. tuberculosis* for 120 hr.

(legend continued on next page)




Figure 4. cGAS Binds to *M. tuberculosis*-Derived DNA during Infection

(A) Western blot of 3xFLAG-cGAS after streptavidin pull down of cell lysates from 3xFLAG-cGAS RAW 264.7 cells transfected with interferon-stimulatory DNA (ISD) or biotinylated-ISD.

(B) Western blot of inputs and FLAG IPs from WT or ΔESX-1 *M. tuberculosis*-infected RAW 264.7 cells stably expressing 3xFLAG-GFP or 3xFLAG-cGAS (left panel), or Strep-cGAS as a negative control (right panel). Whole-cell lysates (Input) or IPs (Flag-IP) were visualized by western blot using anti-FLAG (left) or anti-Strep (right) antibodies.

(C) qPCR of *M. tuberculosis*-derived sequences from DNA isolated from IPs in (B). Quantities were normalized to inputs.

(D) Same as (C), but with mouse *Actin*.

n.s., not significant; **p < 0.005 by two-tailed t test. Error bars indicate SD. See also Figure S4.

It is tempting to speculate that *M. tuberculosis* has evolved to use its own chromosomal DNA to elicit type I IFNs, as these cytokines are produced throughout in vivo mouse infections (Manzanillo et al., 2012) as well as in active human disease (Berry et al., 2010). While the modest decreases in type I IFNs and similar colony-forming unit (CFU) counts we observed during *cGas*^{-/-} mouse infection implicate a role for other host sensor molecules, this work provides a detailed view of one of the molecular interactions that occurs during *M. tuberculosis* infection of macrophages. Indeed, the situation in vivo is certainly much more complicated, and we cannot exclude the possible role of additional DNA sensors that may work in concert with cGAS to coordinate an innate immune response, or the potentially crucial role of signaling in other cell types at different stages of infection. Although these other factors remain uncharacterized, we suspect that cGAS contributes to the robust type I IFN signature associated with active disease in humans (Berry et al., 2010; Novikov et al., 2011).

The cGAS/STING/TBK1 pathway is a major pathway by which innate immune cells recognize both viral and bacterial pathogens that access the cytosol, but different bacteria have evolved diverse mechanisms to initiate this response. We suspect that this primarily antiviral pathway has been co-opted by bacterial pathogens, perhaps primarily to elicit type I IFNs. Indeed, this pathway may be an easy target for bacterial pathogens, as the potential for evolution of bacteria to elicit this response would, in theory, be relatively straightforward as it simply requires exposure of existing metabolites such as nucleic acids or cyclic dinucleotides to the inside of infected cells. However, activation also comes at a cost to bacterial replication as cGAS/STING/TBK1 activation simultaneously induces autophagic targeting. The ability of some pathogens, like *L. pneumophila* (Choy et al., 2012) and *L. monocytogenes* (Gao et al., 2013; Sun

et al., 2013; Tattoli et al., 2013), to inhibit autophagy may be a common mechanism by which pathogens shift the balance of cytosolic detection to promote infection. Understanding how to shift this balance instead toward autophagic targeting of bacterial pathogens represents an attractive host-directed therapeutic strategy to combat infection in humans.

EXPERIMENTAL PROCEDURES

Cell Lines and Cell Culture

BMDMs were generated from wild-type C57BL/6 (The Jackson Laboratory), *cGas*^{-/-} (Schoggins et al., 2014) (or Li et al., 2013 for experiments with *L. pneumophila*), and *Sting*^{gt/gt} (referred to as *Sting*^{-/-}) (Sauer et al., 2011) mice that were 8–12 weeks old. All mice were housed and treated using procedures described in animal care protocols approved by the Institutional Animal Care and Use Committee of UC San Francisco and Washington University St. Louis. Knockdown cell lines were generated using targeting sequences listed in Table S1.

Bacterial Strains

The wild-type *M. tuberculosis* strain used in these studies was the Erdman strain, except when CDC1551 was used as mentioned. The 10403S strain of *L. monocytogenes*, the SL1344 strain of *S. enterica* serovar Typhimurium, and the LPO2 *ΔflaA* strain of *L. pneumophila* were used as wild-type strains.

Bacterial Infections

For *M. tuberculosis* infections, cells were infected at an moi of 10 for RNA and cytokine analysis and 1 for CFUs and microscopy studies. For *L. monocytogenes*, *S. Typhimurium*, and *L. pneumophila* infections, BMDMs were infected at an moi of 10, 5, and 1, respectively. Cells were harvested 4 or 6 hr post-infection.

RNA Isolation and qPCR Analysis

For RNA analysis, cells were harvested in Trizol 4 hr post-treatment, and RNA was isolated using PureLink RNA Mini Kits (Invitrogen). cDNA was synthesized with iScript cDNA Synthesis Kit (Bio-Rad). qPCR was performed using Taq DNA Polymerase (NEB), SYBR Green I (Sigma), and primers listed in Table S2.

(K) CFUs from BMDMs infected with *M. tuberculosis* at 6, 24, 72, and 120 hr (left) and normalized to CFUs at 0 hr (right).

(L–P) WT and *cGas*^{-/-} mice were infected with ~100 aerosolized *M. tuberculosis* CFUs (n = 4 or 6 per group). IFIT1 transcripts in lungs of uninfected and infected mice (L). Serum IFN-β levels 21 days post-infection as measured by ISRE-luciferase reporter cells (M). CFUs in lungs (N) and spleens (O) 21 and 80 days post-infection. Survival of mice monitored for 102 days (n = 6 WT and 3 *cGas*^{-/-}) (P).

*p < 0.05; **p < 0.005; n.s., not significant by two-tailed t test comparing to SCR or WT unless otherwise indicated. Error bars indicate SD. See also Figure S3.

Both the averages and the standard deviations of the raw values were normalized to the average of the treated wild-type sample, which was set at 100%.

Colocalization of Markers with *M. tuberculosis* and Cytosolic DNA

Cytosolic DNA puncta or bacterium-containing phagosomes were visualized directly by fluorescence microscopy. A series of images were captured and analyzed by counting the number of cell-associated DNA puncta or bacteria that colocalized with the corresponding marker. At least one hundred events were analyzed per coverslip, and each condition was performed with triplicate coverslips. For experiments in knockout or knockdown cells, genotypes were blinded throughout.

Mouse Infection

All procedures involving animals were conducted following NIH guidelines in accordance with an approved IACUC protocol at Washington University in St. Louis School of Medicine. *cGas*^{+/+} and *cGas*^{-/-} mice (Schoggins et al., 2014) (C57BL/6 background, male and female littermates, approximately 8–10 weeks old; see Supplemental Experimental Procedures) were infected with ~100 *M. tuberculosis* (Erdman strain) using an inhalation exposure system (Glas-Col). Samples were collected for CFUs, RNA, or serum analysis at the indicated time points.

cGAS Pull Down and DNA Analysis

RAW 267.4 cells were infected at an moi of 20 or transfected with 25 µg ISD and fixed in 4% PFA after 2 hr. Tagged constructs were immunoprecipitated using M2 FLAG Magnetic Beads (Sigma) eluted with FLAG peptide (synthesized by Bioneer). IP efficiency was confirmed by western blot analysis. Samples were analyzed by qPCR to measure the abundance of fragments from the *M. tuberculosis* or mouse genome (see Table S3).

Statistics

Statistical analysis of data was performed using GraphPad Prism software (GraphPad). Two-tailed unpaired Student's *t* tests were used for statistical analyses, and unless otherwise noted, all results are representative of at least two independent biological experiments and are reported as the mean ± SD (*n* = 3 per group).

SUPPLEMENTAL INFORMATION

Supplemental Information includes Supplemental Experimental Procedures, four figures, and three tables and can be found with this article online at <http://dx.doi.org/10.1016/j.chom.2015.05.004>.

AUTHOR CONTRIBUTIONS

R.O.W. and S.L.B. contributed equally to this work. Experiments were designed, performed, and analyzed primarily by R.O.W., S.L.B., and J.S.C., with contributions from D.A.M., E.J.D., and J.O. Mouse infection was performed by J.M.K. The manuscript was written by R.O.W., S.L.B., and J.S.C., and figures were prepared by R.O.W. and S.L.B. All authors read and commented on the manuscript during preparation.

ACKNOWLEDGMENTS

The authors thank Z. Chen for providing femurs from *cGas*^{-/-} mice, T. Parry and B. Penn for sharing reagents, and J. Woodward, K. Patrick, C. Homer, and M. Bell for technical assistance. We are grateful to S. Johnson, K. Thorn, and the Nikon Imaging Center staff at UCSF for training and use of their microscopes. This work was supported by NIH grant PO1 AI063302, NIH grant U19 AI109725, Crohn's and Colitis Foundation of America grant 274415, the Arnold and Mabel Beckman Foundation, HHMI, The Damon Runyon Cancer Research Foundation, NSF Graduate Research Fellowships, and NIGMS Cell and Molecular Biology Training Grant GM007067.

Received: December 5, 2014

Revised: February 4, 2015

Accepted: April 29, 2015

Published: June 2, 2015

REFERENCES

- Ablasser, A., Wassermann, R., Gülen, M.F., Sala, C., Garcia Perin, S., Lou, Y., Rybniker, J., Schmid-Burgk, J.L., Schmidt, T., Hornung, V., and Cole, S.T. (2015). The ESX-1 secretion system of *Mycobacterium tuberculosis* differentially regulates cGAS- and inflammasome-dependent intracellular immune responses. *Cell Host Microbe* 17, this issue, 799–810.
- Auerbuch, V., Brockstedt, D.G., Meyer-Morse, N., O'Riordan, M., and Portnoy, D.A. (2004). Mice lacking the type I interferon receptor are resistant to *Listeria monocytogenes*. *J. Exp. Med.* 200, 527–533.
- Berry, M.P.R., Graham, C.M., McNab, F.W., Xu, Z., Bloch, S.A.A., Oni, T., Wilkinson, K.A., Banichereau, R., Skinner, J., Wilkinson, R.J., et al. (2010). An interferon-inducible neutrophil-driven blood transcriptional signature in human tuberculosis. *Nature* 466, 973–977.
- Birmingham, C.L., Smith, A.C., Bakowski, M.A., Yoshimori, T., and Brummell, J.H. (2006). Autophagy controls *Salmonella* infection in response to damage to the *Salmonella*-containing vacuole. *J. Biol. Chem.* 281, 11374–11383.
- Burdette, D.L., and Vance, R.E. (2013). STING and the innate immune response to nucleic acids in the cytosol. *Nat. Immunol.* 14, 19–26.
- Burdette, D.L., Monroe, K.M., Sotelo-Troha, K., Iwig, J.S., Eckert, B., Hyodo, M., Hayakawa, Y., and Vance, R.E. (2011). STING is a direct innate immune sensor of cyclic di-GMP. *Nature* 478, 515–518.
- Carrero, J.A., Calderon, B., and Unanue, E.R. (2004). Type I interferon sensitizes lymphocytes to apoptosis and reduces resistance to *Listeria* infection. *J. Exp. Med.* 200, 535–540.
- Choi, J., Park, S., Biering, S.B., Selleck, E., Liu, C.Y., Zhang, X., Fujita, N., Saitoh, T., Akira, S., Yoshimori, T., et al. (2014). The parasitophorous vacuole membrane of *Toxoplasma gondii* is targeted for disruption by ubiquitin-like conjugation systems of autophagy. *Immunity* 40, 924–935.
- Choy, A., Dancourt, J., Mugo, B., O'Connor, T.J., Isberg, R.R., Melia, T.J., and Roy, C.R. (2012). The *Legionella* effector RavZ inhibits host autophagy through irreversible Atg8 deconjugation. *Science* 338, 1072–1076.
- Collins, A., Cai, H., Li, T., Franco, L., Li, X.-D., Nair, V., Scharn, C., Stamm, C., Levine, B., Chen, Z., et al. (2015). Cyclic GMP-AMP Synthase Is an Innate Immune Sensor of *Mycobacterium tuberculosis* DNA. *Cell Host Microbe* 17, this issue, 820–828.
- Creasey, E.A., and Isberg, R.R. (2012). The protein SdhA maintains the integrity of the *Legionella*-containing vacuole. *Proc. Natl. Acad. Sci. USA* 109, 3481–3486.
- Deretic, V., and Levine, B. (2009). Autophagy, immunity, and microbial adaptations. *Cell Host Microbe* 5, 527–549.
- Dey, B., Dey, R.J., Cheung, L.S., Pokkali, S., Guo, H., Lee, J.-H., and Bishai, W.R. (2015). A bacterial cyclic dinucleotide activates the cytosolic surveillance pathway and mediates innate resistance to tuberculosis. *Nat. Med.* 21, 401–406.
- Gao, D., Wu, J., Wu, Y.T., Du, F., Aroh, C., Yan, N., Sun, L., and Chen, Z.J. (2013). Cyclic GMP-AMP synthase is an innate immune sensor of HIV and other retroviruses. *Science* 341, 903–906.
- Gutierrez, M.G., Master, S.S., Singh, S.B., Taylor, G.A., Colombo, M.I., and Deretic, V. (2004). Autophagy is a defense mechanism inhibiting BCG and *Mycobacterium tuberculosis* survival in infected macrophages. *Cell* 119, 753–766.
- Kawai, T., and Akira, S. (2010). The role of pattern-recognition receptors in innate immunity: update on Toll-like receptors. *Nat. Immunol.* 11, 373–384.
- Lamkanfi, M., and Dixit, V.M. (2011). Modulation of inflammasome pathways by bacterial and viral pathogens. *J. Immunol.* 187, 597–602.
- Leber, J.H., Crimmins, G.T., Raghavan, S., Meyer-Morse, N.P., Cox, J.S., and Portnoy, D.A. (2008). Distinct TLR- and NLR-mediated transcriptional responses to an intracellular pathogen. *PLoS Pathog.* 4, e6.
- Lee, J., Remold, H.G., Jeong, M.H., and Kornfeld, H. (2006). Macrophage apoptosis in response to high intracellular burden of *Mycobacterium tuberculosis* is mediated by a novel caspase-independent pathway. *J. Immunol.* 176, 4267–4274.

- Li, X.-D., Wu, J., Gao, D., Wang, H., Sun, L., and Chen, Z.J. (2013). Pivotal roles of cGAS-cGAMP signaling in antiviral defense and immune adjuvant effects. *Science* *341*, 1390–1394.
- Liang, Q., Seo, G.J., Choi, Y.J., Kwak, M.-J., Ge, J., Rodgers, M.A., Shi, M., Leslie, B.J., Hopfner, K.-P., Ha, T., et al. (2014). Crosstalk between the cGAS DNA sensor and Beclin-1 autophagy protein shapes innate antimicrobial immune responses. *Cell Host Microbe* *15*, 228–238.
- Lippmann, J., Rothenburg, S., Deigendesch, N., Eitel, J., Meixenberger, K., van Laak, V., Slevogt, H., N'guessan, P.D., Hippenstiel, S., Chakraborty, T., et al. (2008). IFN β responses induced by intracellular bacteria or cytosolic DNA in different human cells do not require ZBP1 (DLM-1/DAI). *Cell Microbiol.* *10*, 2579–2588.
- Manca, C., Tsenova, L., Bergtold, A., Freeman, S., Tovey, M., Musser, J.M., Barry, C.E., 3rd, Freedman, V.H., and Kaplan, G. (2001). Virulence of a *Mycobacterium tuberculosis* clinical isolate in mice is determined by failure to induce Th1 type immunity and is associated with induction of IFN- α / β . *Proc. Natl. Acad. Sci. USA* *98*, 5752–5757.
- Manzanillo, P.S., Shiloh, M.U., Portnoy, D.A., and Cox, J.S. (2012). *Mycobacterium tuberculosis* activates the DNA-dependent cytosolic surveillance pathway within macrophages. *Cell Host Microbe* *11*, 469–480.
- Mayer-Barber, K.D., Andrade, B.B., Barber, D.L., Hieny, S., Feng, C.G., Caspar, P., Oland, S., Gordon, S., and Sher, A. (2011). Innate and adaptive interferons suppress IL-1 α and IL-1 β production by distinct pulmonary myeloid subsets during *Mycobacterium tuberculosis* infection. *Immunity* *35*, 1023–1034.
- Monroe, K.M., McWhirter, S.M., and Vance, R.E. (2009). Identification of host cytosolic sensors and bacterial factors regulating the type I interferon response to *Legionella pneumophila*. *PLoS Pathog.* *5*, e1000665.
- Monroe, K.M., McWhirter, S.M., and Vance, R.E. (2010). Induction of type I interferons by bacteria. *Cell Microbiol.* *12*, 881–890.
- Novikov, A., Cardone, M., Thompson, R., Shenderov, K., Kirschman, K.D., Mayer-Barber, K.D., Myers, T.G., Rabin, R.L., Trinchieri, G., Sher, A., and Feng, C.G. (2011). *Mycobacterium tuberculosis* triggers host type I IFN signaling to regulate IL-1 β production in human macrophages. *J. Immunol.* *187*, 2540–2547.
- O'Connell, R.M., Saha, S.K., Vaidya, S.A., Bruhn, K.W., Miranda, G.A., Zarnegar, B., Perry, A.K., Nguyen, B.O., Lane, T.F., Taniguchi, T., et al. (2004). Type I interferon production enhances susceptibility to *Listeria monocytogenes* infection. *J. Exp. Med.* *200*, 437–445.
- O'Riordan, M., Yi, C.H., Gonzales, R., Lee, K.-D., and Portnoy, D.A. (2002). Innate recognition of bacteria by a macrophage cytosolic surveillance pathway. *Proc. Natl. Acad. Sci. USA* *99*, 13861–13866.
- Sauer, J.-D., Sotelo-Troha, K., von Moltke, J., Monroe, K.M., Rae, C.S., Brubaker, S.W., Hyodo, M., Hayakawa, Y., Woodward, J.J., Portnoy, D.A., and Vance, R.E. (2011). The N-ethyl-N-nitrosourea-induced Goldenticket mouse mutant reveals an essential function of Sting in the in vivo interferon response to *Listeria monocytogenes* and cyclic dinucleotides. *Infect. Immun.* *79*, 688–694.
- Schoggins, J.W., MacDuff, D.A., Imanaka, N., Gainey, M.D., Shrestha, B., Eitson, J.L., Mar, K.B., Richardson, R.B., Ratushny, A.V., Litvak, V., et al. (2014). Pan-viral specificity of IFN-induced genes reveals new roles for cGAS in innate immunity. *Nature* *505*, 691–695.
- Stetson, D.B., and Medzhitov, R. (2006). Recognition of cytosolic DNA activates an IRF3-dependent innate immune response. *Immunity* *24*, 93–103.
- Sun, L., Wu, J., Du, F., Chen, X., and Chen, Z.J. (2013). Cyclic GMP-AMP synthase is a cytosolic DNA sensor that activates the type I interferon pathway. *Science* *339*, 786–791.
- Tattoli, I., Sorbara, M.T., Yang, C., Tooze, S.A., Philpott, D.J., and Girardin, S.E. (2013). *Listeria* phospholipases subvert host autophagic defenses by stalling pre-autophagosomal structures. *EMBO J.* *32*, 3066–3078.
- Teles, R.M.B., Graeber, T.G., Krutzik, S.R., Montoya, D., Schenk, M., Lee, D.J., Komisopoulou, E., Kelly-Scumpia, K., Chun, R., Iyer, S.S., et al. (2013). Type I interferon suppresses type II interferon-triggered human anti-mycobacterial responses. *Science* *339*, 1448–1453.
- Thurston, T.L.M., Wandel, M.P., von Muhlinen, N., Foeglein, A., and Randow, F. (2012). Galectin 8 targets damaged vesicles for autophagy to defend cells against bacterial invasion. *Nature* *482*, 414–418.
- Vance, R.E., Isberg, R.R., and Portnoy, D.A. (2009). Patterns of pathogenesis: discrimination of pathogenic and nonpathogenic microbes by the innate immune system. *Cell Host Microbe* *6*, 10–21.
- Watson, R.O., Manzanillo, P.S., and Cox, J.S. (2012). Extracellular *M. tuberculosis* DNA targets bacteria for autophagy by activating the host DNA-sensing pathway. *Cell* *150*, 803–815.
- Woodward, J.J., Iavarone, A.T., and Portnoy, D.A. (2010). c-di-AMP secreted by intracellular *Listeria monocytogenes* activates a host type I interferon response. *Science* *328*, 1703–1705.
- Zhang, Y., Yeruva, L., Marinov, A., Prantner, D., Wyrick, P.B., Lupashin, V., and Nagarajan, U.M. (2014). The DNA sensor, cyclic GMP-AMP synthase, is essential for induction of IFN- β during *Chlamydia trachomatis* infection. *J. Immunol.* *193*, 2394–2404.
- Zhao, Z., Fux, B., Goodwin, M., Dunay, I.R., Strong, D., Miller, B.C., Cadwell, K., Delgado, M.A., Ponpuak, M., Green, K.G., et al. (2008). Autophagosome-independent essential function for the autophagy protein Atg5 in cellular immunity to intracellular pathogens. *Cell Host Microbe* *4*, 458–469.
- Zughaier, S.M., Zimmer, S.M., Datta, A., Carlson, R.W., and Stephens, D.S. (2005). Differential induction of the toll-like receptor 4-MyD88-dependent and -independent signaling pathways by endotoxins. *Infect. Immun.* *73*, 2940–2950.