

Errata

The following corrections may be observed in the articles listed below:

Grewal, K.K., P.J. Warner and P.H. Williams, An inducible outer membrane protein involved in aerobactin-mediated iron transport by ColV strains of *Escherichia coli* (1982) FEBS Letters 140, 27–30.

page 28 column 2 and page 29 column 2: figures 1 and 2 should be interchanged

Flechner, C.R., A.H. Pershadsingh, M.L. Vorbeck, J.W. Long jr and A.P. Martin, Phosphate-dependent trifluoperazine-sensitive Ca^{2+} efflux from rat liver mitochondria: Modulation by a cytosol factor (1982) FEBS Letters 141, 45–48.

page 45, column 2, line 13 *should read:*
centrifugation at $9500 \times g$. The cytosol was obtained

instead of:
centrifugation at $95\,000 \times g$. The cytosol was obtained

page 47, column 1, table 1 footnote ^a *should read:*
^aProtein was 3.0 ± 0.51 mg

instead of:
^aProtein was 3.0 ± 0.51 mg/ml

Capiod, T., B. Berthon, J. Poggioli, G.M. Burgess and M. Claret, The effect of Ca^{2+} -mobilising hormones on the $\text{Na}^+ - \text{K}^+$ pump in isolated rat liver hepatocytes (1982) FEBS Letters 141, 49–52.

page 50, fig. 1, legend line 2 *should read:*
ence of the β -blocker propranolol at $5 \mu\text{M}$ ATP

instead of:
ence of the β -blocker propranolol (POB) at $5 \mu\text{M}$ ATP

page 50, table 1, column 3 *should read:*
Noradrenaline
+ phenoxybenzine

instead of:
Noradrenaline
+ propranolol

page 51, column 2, line 1 *should read:*
similar time courses [8,9]. In contrast, the β -adreno-

instead of:
similar time courses [8,9]. In contrast, the α -adreno-

Praaning-van Dalen, D.P. and D.L. Knook, Quantitative determination of in vivo endocytosis by rat liver Kupffer and endothelial cells facilitated by an improved cell isolation method (1982) FEBS Letters 141, 229–232.

page 230, column 1, lines 10 and 11 *should read:*
fied by centrifugal elutriation in a Beckman JE-6 elutriator rotor at 4°C with GBSS–BSA as the elutriation fluid

instead of:
fied by centrifugal elution in a Beckman JE-6 elution rotor at 4°C with GBSS–BSA as the elution fluid

page 230, column 1, lines 14 to 9 from below

should read:

before elutriation. The amount of sinusoidal cells recovered from a liver of ~5 g varied from 160–200 × 10⁶ cells, of which ~20% were Kupffer cells, 27% lymphocytes and 53% endothelial cells. After elutriation >90% of the cells could be covered. The viability of the elutriated cells was >90%. The average

instead of:

before elution. The amount of sinusoidal cells recovered from a liver of ~5 g varied from 160–200 × 10⁶ cells, of which ~20% were Kupffer cells, 27% lymphocytes and 53% endothelial cells. After elution, >90% of the cells could be recovered. The viability of the eluted cells was >90%. The average

page 231, column 2, line 6 *should read:*

centrifuged elutriation. Table 3 shows the amount of

instead of:

centrifuged elution. Table 3 shows the amount of

De Mattheis, F., A.H. Jackson, A.H. Gildos, K.R.N. Rao, J. Atton, S. Weerasinghe and C. Hollands, Structural isomerism and chirality of *N*-monosubstituted protoporphyrins (1982) FEBS Letters 142, 44–48.

page 47, column 1, line 23 *should read:*

itory *N*-ethyl protoporphyrin [18], the loss of

instead of:

itory *N*-methyl protoporphyrin [18], the loss of

Vogt, W., B. Hinsch, G. Schmidt and I. von Zabern, Function of the activated fourth component of complement (C4b) in activation of C2 (1982) FEBS Letters 144, 195–198.

page 197, column 1, line 10 from below *should read:*

cleaving C42 complex AgAb was treated as above

instead of:

cleaving C22 complex AgAb was treated as above
