



Packing disjoint cycles over vertex cuts

Jochen Harant^a, Dieter Rautenbach^{a,*}, Peter Recht^b, Ingo Schiermeyer^c, Eva-Maria Sprengel^b

^a Institut für Mathematik, TU Ilmenau, D-98684 Ilmenau, Germany

^b Lehrstuhl für Operations Research und Wirtschaftsinformatik, Universität Dortmund, D-44227 Dortmund, Germany

^c Institut für Diskrete Mathematik und Algebra, Technische Universität Bergakademie Freiberg, D-09596 Freiberg, Germany

ARTICLE INFO

Article history:

Received 2 April 2009

Received in revised form 26 February 2010

Accepted 7 March 2010

Available online 2 April 2010

Keywords:

Connectivity

Cycle

Edge-disjoint cycles

Graph

Packings of cycles

Vertex cut

Vertex-disjoint cycles

ABSTRACT

For a graph G , let $\nu(G)$ and $\nu'(G)$ denote the maximum cardinalities of packings of vertex-disjoint and edge-disjoint cycles of G , respectively. We study the interplay of these two parameters and vertex cuts in graphs. If G is a graph whose vertex set can be partitioned into three non-empty sets S , V_1 , and V_2 such that there is no edge between V_1 and V_2 , and $k = |S|$, then our results imply that $\nu(G)$ is uniquely determined by the values $\nu(H)$ for at most $2^{k+1}k!^2$ graphs H of order at most $\max\{|V_1|, |V_2|\} + k$, and $\nu'(G)$ is uniquely determined by the values $\nu'(H)$ for at most $2^{\binom{k}{2}+1}$ graphs H of order at most $\max\{|V_1|, |V_2|\} + k$.

© 2010 Elsevier B.V. All rights reserved.

1. Introduction

Packing vertex- or edge-disjoint cycles in graphs is a very well-studied and classical graph-theoretical problem. There is a large amount of literature concerning conditions in terms of, for instance, order, size, vertex degrees, degree sums, independence number, chromatic number, and feedback vertex sets that are sufficient for the existence of some number of disjoint cycles which may satisfy further restrictive conditions. We refer the reader to [3,6,9,10,8,11–13,15,17,20,22,25–30], which is just a small selection. The algorithmic problems concerning cycle packings are typically hard [1,2,14,16,18,19,22,24], and approximation algorithms were described [14,19,24]. Several authors mention practical applications in computational biology such as reconstruction of evolutionary trees or genomic analysis.

The starting point for the research presented here is the simple observation that for graphs that contain cutvertices, the problems to find optimal packings of vertex- or edge-disjoint cycles essentially reduce to the blocks of the graph. Here we extend this observation and study the behaviour of these packing problems with respect to larger vertex cuts. Related results concerning vertex cuts of order 2 were used in [7,21].

We consider a finite and undirected graph G with vertex set $V(G)$ and edge set $E(G)$, which may contain parallel edges but no loops. The *neighbourhood* of a vertex u in a graph G is denoted by $N_G(u)$, and the *degree* of u in G is the number of incident edges. For a vertex x and a set of vertices Y in G , let $E_G(x, Y)$ denote the set of all edges of G between x and a vertex in Y . A *vertex cut* in G is a set of vertices S whose removal disconnects G . A *cycle* in G is a connected subgraph of G in which all vertices have degree 2. A *packing of vertex-disjoint cycles (edge-disjoint cycles)* in G is a set of cycles in G that are pairwise vertex-disjoint (edge-disjoint). Let $\nu(G)$ and $\nu'(G)$ denote the maximum cardinalities of packings of vertex-disjoint

* Corresponding author.

E-mail addresses: jochen.harant@tu-ilmenau.de (J. Harant), dieter.rautenbach@tu-ilmenau.de, dieter.rautenbach@gmx.de (D. Rautenbach), peter.recht@tu-dortmund.de (P. Recht), eva-maria.schulte-loh@tu-dortmund.de (I. Schiermeyer), schierme@math.tu-freiberg.de (E.-M. Sprengel).

cycles and edge-disjoint cycles in G , respectively. While parallel edges occur naturally in our constructions and proofs, it is a reasonable restriction to consider graphs without loops; that is, graphs that do not contain cycles of length 1, because every loop is contained in every maximum packing of edge-disjoint cycles and in some maximum packing of vertex-disjoint cycles.

In Sections 2 and 3 we consider packings of vertex-disjoint cycles and edge-disjoint cycles in graphs G that contain a vertex cut S . In both cases we prove that $\nu(G)$ and $\nu'(G)$ are uniquely determined by the values $\nu(H)$ and $\nu'(H)$ for graphs H that arise from G by some simple modifications and contain a vertex cut of cardinality strictly less than $|S|$. In Section 4 we discuss some algorithmical consequences of these results. In this context, we observe that for fixed l the graph properties “ $\nu(G) \geq l$ ” and “ $\nu'(G) \geq l$ ” can be decided in linear time for graphs of bounded clique-width.

2. Vertex-disjoint cycles

Throughout this section, let G be a graph and let $S \cup V_1 \cup V_2$ be a partition of the vertex set of G into three non-empty sets such that there is no edge between V_1 and V_2 , i.e. if G is connected, then S is a vertex cut of G .

For $x \in S$ and $i \in \{1, 2\}$, let $G_{x,i}$ arise from G by deleting all edges in $E_G(x, V_{3-i})$.

For $x, y \in S$, let $G_{x,y}$ arise from G by deleting x and y , adding two new vertices z_1 and z_2 , adding a new edge between z_1 and u for every edge between x or y and a vertex u in V_1 , and adding a new edge between z_2 and v for every edge between x or y and a vertex v in V_2 . Note that there are parallel edges between z_i and the vertices in $N_G(x) \cap N_G(y) \cap V_i$ for $i \in \{1, 2\}$.

Clearly, for $x \in S$ and $i \in \{1, 2\}$, $G - x$ is a subgraph of $G_{x,i}$ and $G_{x,i}$ is a subgraph of G . Furthermore, deleting x from G , can only reduce $\nu(G)$ by 1. Therefore, for $x \in S$ and $i \in \{1, 2\}$,

$$\nu(G - x) \leq \nu(G_{x,i}) \leq \nu(G) \leq \nu(G - x) + 1. \tag{1}$$

We proceed to our main result in this section.

Theorem 1. *Let G, S, V_1 , and V_2 be as above.*

- (i) *If $\nu(G - x) < \nu(G_{x,i})$ for some $x \in S$ and $i \in \{1, 2\}$, then $\nu(G) = \nu(G - x) + 1$.*
- (ii) *If $\nu(G - x) < \nu(G - y)$ for some x and y in S , then $\nu(G) = \nu(G - x) + 1$.*
- (iii) *If $\nu(G - x) = \nu(G_{y,i})$ for all x and y in S and $i \in \{1, 2\}$, then $\nu(G) = \nu(G - x) + 1$ if and only if for every $x \in S$, there is some $y \in S$ such that x and y are non-adjacent and $\nu(G_{x,y}) \geq \nu(G - x) + 2$.*

Proof. (i) By (1),

$$\nu(G - x) + 1 \leq \nu(G_{x,i}) \leq \nu(G) \leq \nu(G - x) + 1,$$

which implies $\nu(G) = \nu(G - x) + 1$.

(ii) By (1),

$$\nu(G - y) \leq \nu(G) \leq \nu(G - x) + 1 \leq \nu(G - y),$$

which implies $\nu(G) = \nu(G - x) + 1$.

(iii) Let $\nu(G - x) = \nu(G_{y,i})$ for all x and y in S and $i \in \{1, 2\}$. Let $\nu^- = \nu(G - x)$ for some $x \in S$. Note that ν^- is independent of the choice of x .

If $\nu(G) = \nu^- + 1$, then for every packing \mathcal{C} of $\nu(G)$ vertex-disjoint cycles in G and for every $x \in S$, there is a unique cycle $C_x \in \mathcal{C}$ such that C_x contains an edge from $E_G(x, V_1)$ and an edge from $E_G(x, V_2)$. Let $x \in S$. Clearly, the cycle C_x contains at least two vertices from S . Let $y \in S$ be a vertex on C_x such that x and y are consecutive vertices from S on C_x . Note that $C_x = C_y$. If x and y are adjacent, then either $\nu(G_{x,1}) \geq \nu^- + 1$ or $\nu(G_{x,2}) \geq \nu^- + 1$, which is a contradiction. Hence x and y are non-adjacent. Since $G_{x,y}$ contains two vertex-disjoint cycles whose vertex set is contained in $\{z_1, z_2\} \cup V(C_x)$, we have $\nu(G_{x,y}) \geq \nu(G) + 1 = \nu^- + 2$.

Conversely, let $x, y \in S$ be such that x and y are non-adjacent and $\nu(G_{x,y}) \geq \nu^- + 2$. Let $\mathcal{C}_{x,y}$ be a packing of $\nu(G_{x,y})$ vertex-disjoint cycles in $G_{x,y}$. Since

$$\nu(G_{x,y} - \{z_1, z_2\}) \geq \nu(G_{x,y}) - 2 \geq \nu^-$$

and

$$\nu(G_{x,y} - \{z_1, z_2\}) \leq \nu(G - x) = \nu^-,$$

we have

$$\nu(G_{x,y} - \{z_1, z_2\}) = \nu(G_{x,y}) - 2 = \nu^-,$$

and the vertices z_1 and z_2 are contained in distinct cycles C_{z_1} and C_{z_2} of $\mathcal{C}_{x,y}$, respectively. Clearly, the subgraph induced in G by the vertex set $(\{x, y\} \cup V(C_{z_1}) \cup V(C_{z_2})) \setminus \{z_1, z_2\}$ contains a subgraph H in which all vertices except for x and y are of degree 2 and $d_H(x) + d_H(y) = 4$. Since H contains at least one cycle, $\nu(G) \geq \nu(G_{x,y}) - 1 = \nu^- + 1$. By (1), $\nu(G) = \nu^- + 1$, which completes the proof. \square

3. Edge-disjoint cycles

Throughout this section, let G be a graph and let $S \cup V_1 \cup V_2$ be a partition of the vertex set of G into three non-empty sets such that there is no edge between V_1 and V_2 . Let $x \in S$ and let $E_1 \cup E_2$ be a partition of $E_G(x, S)$.

For a set $T \subseteq S \setminus \{x\}$, let $G(T)$ be the graph that arises from G by deleting x , adding two new vertices x_1 and x_2 , adding a new edge between x_1 and y for all edges in $E_1 \cup E_G(x, V_1)$ between x and a vertex y , adding a new edge between x_2 and y for all edges in $E_2 \cup E_G(x, V_2)$ between x and a vertex y , and adding two new edges $e_{1,y} = x_1y$ and $e_{2,y} = x_2y$ for all $y \in T$. Note that $d_{G(T)}(x_1) + d_{G(T)}(x_2) = d_G(x) + 2|T|$.

We proceed to our main result in this section.

Theorem 2. *If $G, S, V_1, V_2, x, E_1,$ and E_2 are as above, then*

$$v'(G) = \max \{v'(G(T)) - |T| \mid T \subseteq S \setminus \{x\}\}.$$

Theorem 2 follows immediately from the next two lemmas.

Lemma 3. *If $G, S, V_1, V_2, x, E_1,$ and E_2 are as above, then $v'(G) \geq v'(G(T)) - |T|$ for all sets $T \subseteq S \setminus \{x\}$.*

Proof. We prove the result by induction on $|T|$. For $T = \emptyset$, every cycle in $G(\emptyset)$ corresponds to a subgraph of G of minimum degree at least 2. This immediately implies $v'(G) \geq v'(G(\emptyset))$. Now let $|T| \geq 1$. By induction, it suffices to determine a non-empty set $\partial T \subseteq T$ with

$$v'(G(T \setminus \partial T)) \geq v'(G(T)) - |\partial T|. \tag{2}$$

Let \mathcal{C} be a packing of $v'(G(T))$ edge-disjoint cycles in $G(T)$.

If there is some $y \in T$ such that \mathcal{C} contains at most one cycle whose edge set intersects $\{e_{1,y}, e_{2,y}\}$, then $\partial T = \{y\}$ satisfies (2). Hence we may assume that for every $y \in T$, the two edges $e_{1,y}$ and $e_{2,y}$ are contained in two different cycles in \mathcal{C} .

Next, we assume that there are two distinct vertices $y, z \in T$ and indices $i, j \in \{1, 2\}$ such that $e_{i,y}$ and $e_{j,z}$ are both contained in one cycle $C \in \mathcal{C}$. Let $e_{3-i,y}$ be contained in $C' \in \mathcal{C}$ and let $e_{3-j,z}$ be contained in $C'' \in \mathcal{C}$. If $|\{C, C', C''\}| = 2$, then $\partial T = \{y, z\}$ clearly satisfies (2). Hence, we may assume that C, C' and C'' are three distinct cycles. Since $(E(C) \cup E(C') \cup E(C'')) \setminus \{e_{1,y}, e_{2,y}, e_{1,z}, e_{2,z}\}$ contains the edge set of a cycle, $\partial T = \{y, z\}$ satisfies (2). Hence, we may assume that no cycle in \mathcal{C} contains two of the edges in $E_T = \{e_{1,y}, e_{2,y} \mid y \in T\}$.

Now, for every $i \in \{1, 2\}$ and every $y \in T$, there is a cycle $C_{i,y}$ in \mathcal{C} such that $\{e_{i,y}\} = E(C_{i,y}) \cap E_T$. The edge set in G corresponding to $(E(C_{1,y}) \cup E(C_{2,y})) \setminus \{e_{1,y}, e_{2,y}\}$ contains a cycle C_y for every $y \in T$. Furthermore, the edge set in G corresponding to $E(C)$ contains a cycle for every $C \in \mathcal{C} \setminus \{C_{i,y} \mid i \in \{1, 2\}, y \in T\}$. Altogether, this implies

$$\begin{aligned} v'(G) &\geq |T| + |\mathcal{C} \setminus \{C_{i,y} \mid i \in \{1, 2\}, y \in T\}| \\ &= |T| + (v'(G(T)) - 2|T|) \\ &= v'(G(T)) - |T|. \end{aligned}$$

This completes the proof. \square

Lemma 4. *If $G, S, V_1, V_2, x, E_1,$ and E_2 are as above, then there is a set $T \subseteq S \setminus \{x\}$ such that $v'(G) = v'(G(T)) - |T|$.*

Proof. A cycle C in G is a *crossing x -cycle*, if it intersects $E_G(x, V_1) \cup E_1$ and $E_G(x, V_2) \cup E_2$. Let \mathcal{C} be a packing of $v'(G)$ edge-disjoint cycles in G with the minimum possible number of crossing x -cycles. Since the union of two edge-disjoint crossing x -cycles that contain a common vertex other than x contains two edge-disjoint cycles that are not crossing x -cycles, the choice of \mathcal{C} implies that no two crossing x -cycles in \mathcal{C} contain a common vertex apart from x .

For every crossing x -cycle $C \in \mathcal{C}$ choose a vertex $y_C \in S \setminus \{x\}$ such that C passes through y_C . Let $T = \{y_C \mid C \in \mathcal{C} \text{ is a crossing } x\text{-cycle}\}$.

Clearly, by the definition of y_C and T , there are two edge-disjoint cycles in $G(T)$ corresponding to each crossing x -cycle in \mathcal{C} . Furthermore, there is a cycle in $G(T)$ corresponding to each cycle in \mathcal{C} that contains x but is not a crossing- x -cycle. Finally, there is a cycle in $G(T)$ corresponding to each cycle in \mathcal{C} that does not contain x . Since all these cycles are edge-disjoint, we obtain $v'(G(T)) \geq v'(G) + |T|$. By Lemma 3, $v'(G) = v'(G(T)) - |T|$, which completes the proof. \square

4. Algorithmic consequences

Theorems 1 and 2 are clearly suitable for an inductive argument with respect to the cardinality of the vertex cut.

Theorem 5. *Let G be a graph and let $S \cup V_1 \cup V_2$ be a partition of the vertex set of G into three non-empty sets such that there is no edge between V_1 and V_2 . Let $k = |S|$.*

- (i) $v(G)$ is uniquely determined by the values $v(H)$ for at most $2^{k+1}k!^2$ graphs H of order at most $\max\{|V_1|, |V_2|\} + k$.
- (ii) $v'(G)$ is uniquely determined by the values $v'(H)$ for at most $2^{\binom{k}{2}+1}$ graphs H of order at most $\max\{|V_1|, |V_2|\} + k$.

Proof. (i) We prove the statement by induction over k .

For $k = 1$, S contains exactly one vertex and no cycle of G intersects V_1 and V_2 . Therefore, $\nu(G)$ equals $\max\{\nu(G[V_1 \cup S]) + \nu(G[V_2]), \nu(G[V_1]) + \nu(G[V_2 \cup S])\}$, i.e. $\nu(G)$ is uniquely determined by the values $\nu(H)$ for 4 graphs H of order at most $\max\{|V_1|, |V_2|\} + k$.

For $k \geq 2$, **Theorem 1** implies that $\nu(G)$ is uniquely determined by the values $\nu(H)$ for $k + 2k + \binom{k}{2}$ graphs H whose vertex set $V(H)$ can be partitioned into three non-empty sets S' , V'_1 , and V'_2 such that there is no edge between V'_1 and V'_2 , and $(|S'|, |V'_1|, |V'_2|) = (|S|, |V_1|, |V_2|) + (a, b, c)$ for some $(a, b, c) \in \{(-1, 0, 0), (-1, 1, 0), (-1, 0, 1), (-2, 1, 1)\}$. Note that $k + 2k + \binom{k}{2} \leq 2k^2$ for $k \geq 2$. By induction, we obtain that $\nu(G)$ is uniquely determined by the values $\nu(H)$ for at most $2k^2 2^k (k-1)!$ graphs H of order at most $\max\{|V_1|, |V_2|\} + k$. Since $2k^2 2^k (k-1)! = 2^{k+1} k!^2$, this completes the proof of (i).

(ii) We prove the statement by induction on k .

For $k = 1$, S contains exactly one vertex, and no cycle of G intersects V_1 and V_2 . Therefore, $\nu'(G)$ equals $\nu'(G[V_1 \cup S]) + \nu'(G[V_2 \cup S])$; that is, $\nu'(G)$ is uniquely determined by the values $\nu'(H)$ for 2 graphs H of order at most $\max\{|V_1|, |V_2|\} + k$.

For $k \geq 2$, **Theorem 2** implies that $\nu'(G)$ is uniquely determined by the values $\nu'(H)$ for 2^{k-1} graphs H whose vertex set $V(H)$ can be partitioned into three non-empty sets S' , V'_1 , and V'_2 such that there is no edge between V'_1 and V'_2 , and $(|S'|, |V'_1|, |V'_2|) = (|S|, |V_1|, |V_2|) + (-1, +1, +1)$. By induction, we obtain that $\nu'(G)$ is uniquely determined by the values $\nu'(H)$ for $2^{k-1} 2^{\binom{k-1}{2}+1}$ graphs H of order at most $\max\{|V_1|, |V_2|\} + k$. Since $2^{k-1} 2^{\binom{k-1}{2}+1} = 2^{\binom{k}{2}+1}$, this completes the proof of (ii). \square

Considering the proofs of **Theorems 1** and **2**, it is not difficult to see that optimal packings of cycles in G can also be derived efficiently from optimal packings of cycles in the graphs H from **Theorem 5**.

Since graphs of bounded tree-width and order n have vertex cuts of bounded order whose removal results in components of order at most $2n/3$ [23], such graphs seem to be a natural choice for an algorithmic application of **Theorem 5**. In view of [4,5], our next and final result implies that for fixed $l \in \mathbb{N}$, the graph properties “ $\nu(G) \geq l$ ” and “ $\nu'(G) \geq l$ ” can be decided in linear time for the even larger class of graphs of bounded clique-width. Furthermore, the corresponding cycle packings can be found efficiently, too.

Theorem 6. For fixed $l \in \mathbb{N}$, the two graph properties “ $\nu(G) \geq l$ ” and “ $\nu'(G) \geq l$ ” can be expressed in monadic second order logic [4] avoiding quantification over sets of edges (MSO₁-logic).

Proof. We only give details for the property “ $\nu'(G) \geq l$ ”, which is more difficult to express in monadic second order logic avoiding quantification over sets of edges.

Let G be a graph with $\nu'(G) \geq l$. If \mathcal{C} is a packing of l edge-disjoint cycles of G with the smallest total size $\sum_{C \in \mathcal{C}} |E(C)|$, then every two cycles in \mathcal{C} intersect in at most two vertices. This implies that every cycle in \mathcal{C} contains at most $2(l-1)$ vertices that belong also to other cycles in \mathcal{C} . Therefore, it is easy to see that a graph G satisfies $\nu'(G) \geq l$ if and only if there are (not necessarily distinct) vertices v_j^i for $1 \leq i \leq l$ and $1 \leq j \leq 2(l-1)$, edges e_j^i for $1 \leq i \leq l$ and $1 \leq j \leq 4(l-1)$, and sets U_i of vertices for $1 \leq i \leq l$ such that

- (i) $v_j^{i_1} \notin U_{i_2}$ for $1 \leq i_1, i_2 \leq l$ and $1 \leq j \leq 2(l-1)$,
- (ii) $U_{i_1} \cap U_{i_2} = \emptyset$ for $1 \leq i_1, i_2 \leq l$ with $i_1 \neq i_2$,
- (iii) e_{2j-1}^i and e_{2j}^i are distinct edges incident with v_j^i whose other endvertex lies in

$$V_i := U_i \cup \left\{ v_{j'}^i \mid 1 \leq j' \leq 2(k-1) \right\}$$

for $1 \leq i \leq l$ and $1 \leq j \leq 2(l-1)$,

- (iv) every vertex in U_i has two distinct neighbours in V_i for $1 \leq i \leq l$, and
- (v) $e_{j_1}^{i_1} \neq e_{j_2}^{i_2}$ for $1 \leq i_1, i_2 \leq l$ and $1 \leq j_1, j_2 \leq 4(l-1)$ with $i_1 \neq i_2$.

The vertices v_j^i correspond to the vertices of the i th cycle C_i of a packing \mathcal{C} of l edge-disjoint cycles which may belong to more than one cycle of \mathcal{C} . Note that the vertices v_j^i are allowed to coincide. Therefore, if C_i contains no vertex that belongs to another cycle of \mathcal{C} , then one can choose $v_1^i = v_2^i = \dots = v_{2(l-1)}^i$ equal to an arbitrary vertex of C_i . The set U_i corresponds to the set of the remaining vertices of C_i . Since the sets U_i are disjoint, edges incident with vertices in distinct sets U_i are necessarily distinct.

Since the existence of the vertices v_j^i , the edges e_j^i , and the sets U_i and also the conditions (i)–(v) can clearly be expressed in monadic second order logic avoiding quantification over sets of edges, the proof is complete. \square

References

[1] A. Caprara, A. Panconesi, R. Rizzi, Packing cycles in undirected graphs, J. Algorithms 48 (2003) 239–256.
 [2] A. Caprara, R. Rizzi, Packing triangles in bounded degree graphs, Inf. Process. Lett. 84 (2002) 175–180.
 [3] K. Corradi, A. Hajnal, On the maximal number of independent circuits in a graph, Acta Math. Acad. Sci. Hung. 14 (1963) 423–439.

- [4] B. Courcelle, The expression of graph properties and graph transformations in monadic second-order logic, in: G. Rozenberg (Ed.), Handbook of Graph Grammars and Computing by Graph Transformations, in: Foundations, vol. 1, World Scientific, 1997, pp. 313–400.
- [5] B. Courcelle, J.A. Makowsky, U. Rotics, Linear time solvable optimization problems on graphs of bounded clique-width, *Theory Comput. Syst.* 33 (2000) 125–150.
- [6] J. Degenhardt, P. Recht, On a relation between the cycle packing number and the cyclomatic number of a graph, manuscript, 2008.
- [7] G. Ding, W. Zang, Packing Cycles in Graphs, *J. Combin. Theory, Ser. B* 86 (2002) 381–407.
- [8] Y. Egawa, H. Enomoto, S. Jendrol', K. Ota, I. Schiermeyer, Independence number and vertex-disjoint cycles, *Discrete Math.* 307 (2007) 1493–1498.
- [9] Y. Egawa, R.J. Faudree, E. Györi, Y. Ishigami, R.H. Schelp, H. Wang, Vertex-disjoint cycles containing specified edges, *Graphs Combin.* 16 (2000) 81–92.
- [10] Y. Egawa, M. Hagita, K. Kawarabayashi, H. Wang, Covering vertices of a graph by k disjoint cycles, *Discrete Math.* 270 (2003) 115–125.
- [11] H. Enomoto, On the existence of disjoint cycles in a graph, *Combinatorica* 18 (1998) 487–492.
- [12] P. Erdős, L. Pósa, On the maximal number of disjoint circuits of a graph, *Publ. Math. Debrecen* 9 (1962) 3–12.
- [13] P. Erdős, L. Pósa, On independent circuits contained in a graph, *Can. J. Math.* 17 (1965) 347–352.
- [14] Z. Friggstad, M.R. Salavatipour, Approximability of packing disjoint cycles, in: Proceedings of ISAAC 2007, *Lecture Notes in Comput. Sci.* 4835 (2007), 304–315.
- [15] S. Fujita, Recent results on disjoint cycles in graphs, *Electronic Notes Discrete Math.* 22 (2005) 409–412.
- [16] M.R. Garey, D.S. Johnson, *Computers and Intractability: A Guide to the Theory of Incompleteness*, Freeman, San Francisco, 1979.
- [17] J. Harant, D. Rautenbach, P. Recht, F. Regen, Packing Edge-Disjoint Cycles in Graphs and the Cyclomatic Number, manuscript, 2008.
- [18] I. Holyer, The NP-completeness of some edge-partition problems, *SIAM J. Comput.* 10 (1981) 713–717.
- [19] M. Krivelevich, Z. Nutov, M.R. Salavatipour, J. Yuster, R. Yuster, Approximation algorithms and hardness results for cycle packing problems, *ACM Trans. Algorithms* 3 (2007) Article No. 48.
- [20] J.W. Moon, On edge-disjoint cycles in a graph, *Can. Math. Bull.* 7 (1964) 519–523.
- [21] Z. Nutov, M. Penn, On integrality, stability and composition of dicycle packings and covers, *J. Combin. Optim.* 4 (2000) 235–251.
- [22] D. Rautenbach, F. Regen, On packing shortest cycles in graphs, *Inf. Process. Lett.* 109 (2009) 816–821.
- [23] N. Robertson, P. Seymour, Graph minors II. Algorithmic aspects of tree width, *J. Algorithms* 7 (1986) 309–322.
- [24] M.R. Salavatipour, J. Verstraëte, Disjoint cycles: integrality gap, hardness, and approximation, in: Proceedings of IPCO 2005, *Lecture Notes in Comput. Sci.* 3509 (2005), 51–65.
- [25] M. Simonovits, A new proof and generalizations of a theorem of Erdős and Posa on graphs without $k + 1$ independent circuits, *Acta Math. Acad. Sci. Hung.* 18 (1967) 191–206.
- [26] J. Verstraëte, Vertex-disjoint cycles of the same length, *J. Combin. Theory, Ser. B* 88 (2003) 45–52.
- [27] J. Verstraëte, A note on vertex-disjoint cycles, *Comb. Probab. Comput.* 11 (2002) 97–102.
- [28] H. Wang, Large vertex-disjoint cycles in a bipartite graph, *Graphs Combin.* 16 (2000) 359–366.
- [29] H. Wang, On independent cycles in a bipartite graph, *Graphs Combin.* 17 (2001) 177–183.
- [30] H. Wang, Maximal total length of k disjoint cycles in bipartite graphs, *Combinatorica* 25 (2005) 367–377.