



Discrete Mathematics 131 (1994) 345–350

DISCRETE
MATHEMATICS

Note

R-sequenceability and *R**-sequenceability of abelian 2-groups

Patrick Headley

Department of Mathematics, University of Michigan, Ann Arbor, MI 48109-4903, USA

Received 20 December 1991; revised 15 June 1992

Abstract

A group of order n is said to be *R*-sequenceable if the nonidentity elements of the group can be listed in a sequence a_1, a_2, \dots, a_{n-1} such that the quotients $a_1^{-1}a_2, a_2^{-1}a_3, \dots, a_{n-2}^{-1}a_{n-1}, a_{n-1}^{-1}a_1$ are distinct. An abelian group is *R**-sequenceable if it has an *R*-sequencing a_1, a_2, \dots, a_{n-1} such that $a_{i-1}a_{i+1} = a_i$ for some i (subscripts are read modulo $n-1$). Friedlander, Gordon and Miller (1978) showed that an *R**-sequenceable Sylow 2-subgroup is a sufficient condition for a group to be *R*-sequenceable. In this paper we also show that all noncyclic abelian 2-groups are *R**-sequenceable except for $\mathcal{L}_2 \times \mathcal{L}_4$ and $\mathcal{L}_2 \times \mathcal{L}_2 \times \mathcal{L}_2$.

A group of order n is said to be *R*-sequenceable if the nonidentity elements of the group can be listed in a sequence a_1, a_2, \dots, a_{n-1} such that the quotients $a_1^{-1}a_2, a_2^{-1}a_3, \dots, a_{n-2}^{-1}a_{n-1}, a_{n-1}^{-1}a_1$ are distinct. The concept of *R*-sequenceability has been around for more than 40 years in one form or another. In 1951 Paige observed that it is a sufficient condition for a group to have a complete mapping. In 1955 Hall and Paige [3] showed that a solvable group has a complete mapping if and only if its Sylow 2-subgroup is either trivial or noncyclic. In 1974 Ringel [5] was led to the concept of *R*-sequenceability in his solution of the map coloring problem for all compact two-dimensional manifolds except the sphere. In their book [1] Dénes and Keedwell used an alternative definition of *R*-sequenceable and discussed the topic in great depth. They also showed that an abelian group is a super *P*-group if and only if it is either *R*-sequenceable or sequenceable. Friedlander et al. [2] showed that the following types of abelian groups are *R*-sequenceable: cyclic groups of odd order greater than 1; groups of odd order whose Sylow 3-subgroup is cyclic; groups whose orders are relatively prime to 6; elementary abelian p -groups, except the group of order 2; groups of type $\mathcal{L}_2 \times \mathcal{L}_{4k}$, $k \geq 1$; groups whose Sylow p -subgroup has the form \mathcal{L}_2^m , $m > 1$ but $m \neq 3$; groups G whose Sylow p -subgroup has the form $S = \mathcal{L}_2 \times \mathcal{L}_n$

where $n = 2^k$ and either k is odd or $k \geq 2$ is even and G/S has a direct cyclic factor of order congruent to 2 modulo 3. Ringel [1] has claimed that abelian groups of the form $\mathcal{L}_2 \times \mathcal{L}_{6k+2}$ are R -sequenceable.

Friedlander et al. [2] conjectured that an abelian group is R -sequenceable if and only if its Sylow 2-subgroup is either trivial or noncyclic. This paper proves the conjecture for abelian 2-groups.

The following types of nonabelian groups are known to be R -sequenceable: groups of order pq where p and q are odd primes, $p < q$, and p has 2 as a primitive root [4]; dihedral groups of order $2n$ where n is even [4]; dicyclic groups of order $4n$ where n is divisible by 4 [6].

An abelian group is R^* -sequenceable if it has an R -sequencing a_1, a_2, \dots, a_{n-1} such that $a_{i-1}a_{i+1} = a_i$ for some i (subscripts are read modulo $n-1$). The term was introduced by Friedlander et al. [2], who showed that the existence of an R^* -sequenceable Sylow 2-subgroup is a sufficient condition for a group to be R -sequenceable. In this paper we also show that all noncyclic abelian 2-groups are R^* -sequenceable except for $\mathcal{L}_2 \times \mathcal{L}_4$ and $\mathcal{L}_2 \times \mathcal{L}_2 \times \mathcal{L}_2$.

We begin with two results of Friedlander et al. concerning abelian 2-groups.

Lemma 1 (Friedlander et al. [2]). $(\mathcal{L}_2)^m$ is R^* -sequenceable for $m > 1$, $m \neq 3$, $\mathcal{L}_2 \times \mathcal{L}_{2^k}$ is R^* -sequenceable for k odd, and R -sequenceable for all k .

Lemma 2 (Friedlander et al. [2]). $\mathcal{L}_2 \times \mathcal{L}_2 \times \mathcal{L}_2$ and $\mathcal{L}_2 \times \mathcal{L}_4$ are R -sequenceable but not R^* -sequenceable.

Lemma 3. *If an abelian group G is an extension of $\mathcal{L}_2 \times \mathcal{L}_2$ by an R^* -sequenceable group H , then G is R^* -sequenceable.*

Proof of Lemma 3. Let $n = |H|$. Since H is R^* -sequenceable, the cosets of $\mathcal{L}_2 \times \mathcal{L}_2$, excluding $\mathcal{L}_2 \times \mathcal{L}_2$ itself, have an ordering K_1, \dots, K_{n-1} that is an R -sequence with $K_{n-1}K_2 = K_1$. Choose k_i , $1 \leq i \leq n-1$, such that $k_i \in K_i$ and $k_{n-1}k_2 = k_1$. Then any element in G can be uniquely expressed as a product of an element in $\mathcal{L}_2 \times \mathcal{L}_2$ and an element in $\{k_1, \dots, k_{n-1}, e\}$. Let $\{y_i\}_{i=1}^{4n-1}$ be the sequence $k_1, k_2, \dots, k_{n-1}, e, k_2, k_3, \dots, k_{n-1}, k_1, k_1, k_1, k_2, \dots, k_{n-1}, e, e, k_2, k_3, \dots, k_{n-1}$. Let a and b be generators of the $\mathcal{L}_2 \times \mathcal{L}_2$ subgroup of G . Define $\{x_i\}_{i=1}^{4n-1}$ as follows.

Case 1: $|H| \bmod 3 \equiv 0$. Let $3k = |H|$, $\{x_i\}$ is given by the successive rows of the $4 \times n$ matrix

$$\begin{pmatrix} e & e & \dots & b & a \\ ab & k-2 \text{ copies of } \{a, b, ab\} & a & ab & ab & b & a \\ ab & b & k-2 \text{ copies of } \{ab, a, b\} & ab & b & a & ab \\ b & a & k-2 \text{ copies of } \{b, ab, a\} & b & a & e \end{pmatrix}.$$

If $k=1$, then $H = \mathcal{L}_3$, so $G = \mathcal{L}_2 \times \mathcal{L}_6$, which is R^* -sequenceable since its Sylow 2-subgroup is R^* -sequenceable.

Case 2: $|H| \bmod 3 \equiv 1$. Let $3k+1 = |H|$. $\{x_i\}$ is read from the successive rows of the $4 \times n$ matrix.

$$\begin{pmatrix} e & e & \dots & b & a \\ ab & k-1 \text{ copies of } \{b, a, ab\} & ab & b & a \\ ab & b & k-1 \text{ copies of } \{a, ab, b\} & a & ab \\ b & a & k-1 \text{ copies of } \{ab, b, a\} & e & \end{pmatrix}.$$

Case 3: $|H| \bmod 3 \equiv 2$. Let $3k+2 = |H|$. $\{x_i\}$ is read from the successive rows of the $4 \times n$ matrix

$$\begin{pmatrix} e & e & \dots & b & a \\ ab & k-1 \text{ copies of } \{b, a, ab\} & b & ab & b & a \\ ab & b & k-1 \text{ copies of } \{a, ab, b\} & a & a & ab \\ b & a & k-1 \text{ copies of } \{ab, b, a\} & ab & e & \end{pmatrix}.$$

Then $\{x_i y_i\}$ is an R^* -sequence. Clearly $(x_{4n-1} y_{4n-1})(x_2 y_2) = x_1 y_1$. Verifying that $\{x_i y_i\}$ is an R -sequence is straightforward with the following observations:

(i) $k_n^{-1} e = k_1^{-1} k_2$ and $e^{-1} k_2 = k_{n-1}^{-1} k_1$, so $\{y_i^{-1} y_{i+1}\}_{i=1}^{4n-1}$ (with $y_{4n} = y_1$) is the sequence $k_1^{-1} k_2, k_2^{-1} k_3, \dots, k_{n-2}^{-1} k_{n-1}, k_1^{-1} k_2, k_{n-1}^{-1} k_1, k_2^{-1} k_3, k_3^{-1} k_4, \dots, k_{n-2}^{-1} k_{n-1}, k_{n-1}^{-1} k_1, e, e, k_1^{-1} k_2, k_2^{-1} k_3, \dots, k_{n-2}^{-1} k_{n-1}, k_1^{-1} k_2, e, k_{n-1}^{-1} k_1, k_2^{-1} k_3, k_3^{-1} k_4, \dots, k_{n-2}^{-1} k_{n-1}, k_{n-1}^{-1} k_1$.

(ii) If x_m is the first element of the first copy of one of the repeated 3-element sequences in $\{x_i\}$, then $y_m = k_3$, and the sequence $\{a, b, ab\}$ is itself an R -sequence. \square

Lemma 4. $\mathcal{L}_2 \times \mathcal{L}_{2^n}$ is R^* -sequenceable for $n \geq 1, n \neq 2$.

Proof of Lemma 4. Any sequence of the nonidentity elements of $\mathcal{L}_2 \times \mathcal{L}_2$ is an R^* -sequence. $\mathcal{L}_2 \times \mathcal{L}_8 \cong \langle a, b \mid a^8 = b^2 = e, ab = ba \rangle$ has the R^* -sequence $ba^7, b, a^5, a^3, ba^6, ba, a^2, a^6, ba^5, ba^2, a^4, ba^4, ba^3, a^7, a$. The relevant triple is ba^4, ba^3 and a^7 .

For $n \geq 4, \mathcal{L}_2 \times \mathcal{L}_{2^n} \cong \langle a, b \mid a^{2^n} = b^2 = e, ab = ba \rangle$, an R^* -sequence can be read from the successive rows of this $2m \times 8$ matrix, where $m = 2^{n-3}$:

ba^{8m-1}	b	a^{3m}	a^{5m}	ba^{8m-2}	ba	a^{m-2}	a^{7m+2}
ba^{8m-3}	ba^2	a^{3m-2}	a^{5m+2}	ba^{8m-4}	ba^3	a^{m-4}	a^{7m+4}
			\vdots				
ba^{7m+3}	ba^{m-4}	a^{2m+4}	a^{6m-4}	ba^{7m+2}	ba^{m-3}	a^2	a^{8m-2}
ba^{7m+1}	ba^{m-2}	a^{2m+2}	a^{6m-2}	ba^{7m}	ba^{m-1}	a^{8m-1}	a
ba^{7m-1}	ba^m	a^{6m-1}	a^{2m+1}	ba^{7m-2}	ba^{m+1}	a^{8m-3}	a^3
ba^{7m-3}	ba^{m+2}	a^{6m-3}	a^{2m+3}	ba^{7m-4}	ba^{m+3}	a^{8m-5}	a^5
			\vdots				
ba^{5m+3}	ba^{3m-4}	a^{4m+3}	a^{4m-3}	ba^{5m+2}	ba^{3m-3}	a^{6m+1}	a^{2m-1}
ba^{5m+1}	ba^{3m-2}	a^{4m+1}	a^{4m-1}	ba^{5m}	ba^{3m-1}	a^{2m}	a^{6m}
ba^{5m-1}	ba^{3m}	a^{4m}	—	ba^{5m-2}	ba^{3m+1}	a^{2m-2}	a^{6m+2}
ba^{5m-3}	ba^{3m+2}	a^{4m-2}	a^{4m+2}	ba^{5m-4}	ba^{3m+3}	a^{2m-4}	a^{6m+4}
			\vdots				
ba^{4m+1}	ba^{4m-2}	a^{3m+2}	a^{5m-2}	ba^{4m}	ba^{4m-1}	a^m	a^{7m}

To see that the sequence is an R -sequence, the successive quotients are listed in the successive rows of this matrix:

a	ba^{3m}	a^{2m}	ba^{3m-2}	a^3	ba^{m-3}	a^{6m+4}	ba^{m-5}
a^5	ba^{3m-4}	a^{2m+4}	ba^{3m-6}	a^7	ba^{m-7}	a^{6m+8}	ba^{m-9}
			\vdots				
a^{2m-7}	ba^{m+8}	a^{4m-8}	ba^{m+6}	a^{2m-5}	ba^{7m+5}	a^{8m-4}	ba^{7m+3}
a^{2m-3}	ba^{m+4}	a^{4m-4}	ba^{m+2}	a^{2m-1}	ba^{7m}	a^2	ba^{7m-2}
a^{2m+1}	ba^{5m-1}	a^{4m+2}	ba^{5m-3}	a^{2m+3}	ba^{7m-4}	a^6	ba^{7m-6}
a^{2m+5}	ba^{5m-5}	a^{4m+6}	ba^{5m-7}	a^{2m+7}	ba^{7m-8}	a^{10}	ba^{7m-10}
			\vdots				
a^{6m-7}	ba^{m+7}	a^{8m-6}	ba^{m+5}	a^{6m-5}	ba^{3m+4}	a^{4m-2}	ba^{3m+2}
a^{6m-3}	ba^{m+3}	a^{8m-2}	ba^{m+1}	a^{6m-1}	ba^{7m+1}	a^{4m}	ba^{7m-1}
a^{6m+1}	ba^m	—	ba^{m-2}	a^{6m+3}	ba^{7m-3}	a^{4m+4}	ba^{7m-5}
a^{6m+5}	ba^{m-4}	a^4	ba^{m-6}	a^{6m+7}	ba^{7m-7}	a^{4m+8}	ba^{7m-9}
			\vdots				
a^{8m-7}	ba^{7m+8}	a^{2m-8}	ba^{7m+6}	a^{8m-5}	ba^{5m+5}	a^{6m-4}	ba^{5m+3}
a^{8m-3}	ba^{7m+4}	a^{2m-4}	ba^{7m+2}	a^{8m-1}	ba^{5m+1}	a^{6m}	ba^{m-1}

If $m=6k+2$, we have ..., $ba^{7m-1-2(4k+1)}$, $ba^{m+2(4k+1)}$, $a^{6m-1-2(4k+1)}$, ... and $(ba^{7m-1-2(4k+1)})(a^{6m-1-2(4k+1)})=ba^{14k+4}=ba^{m+2(4k+1)}$. If $m=6k+4$, we have ..., $a^{2m+1+2(4k+2)}$, $ba^{7m-2-2(4k+2)}$, $ba^{m+1+2(4k+2)}$, ... and $(a^{2m+1+2(4k+2)})(ba^{m+1+2(4k+2)})=ba^{34k+22}=ba^{7m-2-2(4k+2)}$. Thus, the sequence is an R^* -sequence for all $n \geq 4$. \square

Theorem. If G is a non-cyclic abelian 2-group, then G is R -sequenceable. Moreover, if $|G| \neq 8$, then G is R^* -sequenceable.

Proof. If $|G|=8$, the result follows from Lemma 2. Otherwise, we use induction on n , where $|G|=2^n$. For n even, the base of the induction is $n=2$, so that $G \cong \mathcal{L}_2 \times \mathcal{L}_2$, which is R^* -sequenceable by Lemma 1. For n odd, the base of the induction is $n=5$, so that either $G \cong \mathcal{L}_2 \times \mathcal{L}_2 \times \mathcal{L}_2 \times \mathcal{L}_2 \times \mathcal{L}_2$, $G \cong \mathcal{L}_2 \times \mathcal{L}_2 \times \mathcal{L}_2 \times \mathcal{L}_4$, $G \cong \mathcal{L}_2 \times \mathcal{L}_2 \times \mathcal{L}_8$, $G \cong \mathcal{L}_2 \times \mathcal{L}_4 \times \mathcal{L}_4$, $G \cong \mathcal{L}_2 \times \mathcal{L}_{16}$ or $G \cong \mathcal{L}_4 \times \mathcal{L}_8$, $\mathcal{L}_2 \times \mathcal{L}_2 \times \mathcal{L}_2 \times \mathcal{L}_2 \times \mathcal{L}_2$ is R^* -sequenceable by Lemma 1. $\mathcal{L}_2 \times \mathcal{L}_{16}$ is R^* -sequenceable by Lemma 4. The other groups are extensions of $\mathcal{L}_2 \times \mathcal{L}_4$ by $\mathcal{L}_2 \times \mathcal{L}_2$. Let H_1, H_2, H_3 be the cosets, other than $\mathcal{L}_2 \times \mathcal{L}_4$ itself, of $\mathcal{L}_2 \times \mathcal{L}_4$, and let h_1, h_2, h_3 be elements of H_1, H_2, H_3 , respectively, such that $h_1 h_3 = h_2$. This is possible since H_1, H_2, H_3 must be an R^* -sequence of $G/(\mathcal{L}_2 \times \mathcal{L}_4)$. Let the subgroup of G isomorphic to $\mathcal{L}_2 \times \mathcal{L}_4$ be generated by a and b with $a^4 = b^2 = e, ab = ba$. Then the following is an R^* -sequence: $h_1, h_2, ba^2 h_3, a, bah_2, bah_3, a^2 h_1, ah_1, bh_1, ba^2 h_2, ba^3 h_3, ba^2, b, ba^3 h_2, a^2 h_3, a^3 h_1, bah_1, a^2 h_2, ah_3, ba^3, a^3, ah_2, a^3 h_3, ba^2 h_1, ba^3 h_1, a^3 h_2, bh_3, ba, a^2, bh_2, h_3$. The relevant triple is $ba^3 h_1, a^3 h_2$ and bh_3 .

To complete the induction, we assume the result is true for n . Let $|G|=2^{n+2}$. If $G \cong \mathcal{L}_2 \times \mathcal{L}_{2^{n+1}}$, G is R^* -sequenceable by Lemma 4. Otherwise, G is an extension of $\mathcal{L}_2 \times \mathcal{L}_2$ by a noncyclic abelian 2-group H , and $|H|=2^n$. Since H is R^* -sequenceable by assumption, G is R^* -sequenceable by Lemma 3. \square

Since Friedlander et al. [2] have shown that an abelian group whose Sylow 2-subgroup is R^* -sequenceable is itself R^* -sequenceable, we have the following corollary.

Corollary. An abelian group whose Sylow 2-subgroup is noncyclic and not of order 8 is R^* -sequenceable.

Acknowledgment

This paper was written at the University of Minnesota, Duluth, under the direction of Professor Joseph Gallian. Donald Keedwell read a preliminary version of the paper and made several helpful suggestions that are incorporated in this version. The research was supported by NSF grant DMS 8709428.

References

- [1] J. Dénes and A.D. Keedwell, Latin squares: new developments in the theory and applications, Ann. Discrete Math. 46 (1991).
- [2] R.J. Friedlander, B. Gordon and M.D. Miller, On a group sequencing problem of Ringel, Proc. 9th S-E Conf. Combinatorics, Graph Theory and Computing, Congr. Numer. XXI (1978) 307–321.

- [3] M. Hall and L.J. Paige, Complete mappings of finite groups, *Pacific J. Math.* 5 (1955) 541–549.
- [4] A.D. Keedwell, On R -sequenceability and R_n -sequenceability of groups, *Ann. Discrete Math.* 18 (1983) 535–548.
- [5] G. Ringel, Cyclic arrangements of the elements of a group, *Notices Amer. Math. Soc.* 21 (1974) A95–96.
- [6] C. Wang, On the R -sequenceability of dicyclic groups, preprint.