



## Review

Biogenesis of *cbb*<sub>3</sub>-type cytochrome *c* oxidase in *Rhodobacter capsulatus*☆Seda Ekici <sup>a,1</sup>, Grzegorz Pawlik <sup>b,1</sup>, Eva Lohmeyer <sup>b</sup>, Hans-Georg Koch <sup>b,\*</sup>, Fevzi Daldal <sup>a,\*\*</sup><sup>a</sup> University of Pennsylvania, Department of Biology, Philadelphia, PA 19104, USA<sup>b</sup> Albert-Ludwigs University of Freiburg, Institute of Biochemistry and Molecular Biology, ZBMZ, Freiburg, 79110 Germany

## ARTICLE INFO

## Article history:

Received 24 September 2011

Accepted 31 October 2011

Available online 4 November 2011

## Keywords:

*cbb*<sub>3</sub>-type cytochrome *c* oxidase  
Co- or post-translational cofactor insertion  
Assembly of membrane proteins  
Copper acquisition  
Photosynthesis and respiration  
Energy transduction

## ABSTRACT

The *cbb*<sub>3</sub>-type cytochrome *c* oxidases (*cbb*<sub>3</sub>-Cox) constitute the second most abundant cytochrome *c* oxidase (Cox) group after the mitochondrial-like *aa*<sub>3</sub>-type Cox. They are present in bacteria only, and are considered to represent a primordial innovation in the domain of Eubacteria due to their phylogenetic distribution and their similarity to nitric oxide (NO) reductases. They are crucial for the onset of many anaerobic biological processes, such as anoxygenic photosynthesis or nitrogen fixation. In addition, they are prevalent in many pathogenic bacteria, and important for colonizing low oxygen tissues. Studies related to *cbb*<sub>3</sub>-Cox provide a fascinating paradigm for the biogenesis of sophisticated oligomeric membrane proteins. Complex subunit maturation and assembly machineries, producing the *c*-type cytochromes and the binuclear heme *b*<sub>3</sub>-Cu<sub>B</sub> center, have to be coordinated precisely both temporally and spatially to yield a functional *cbb*<sub>3</sub>-Cox enzyme. In this review we summarize our current knowledge on the structure, regulation and assembly of *cbb*<sub>3</sub>-Cox, and provide a highly tentative model for *cbb*<sub>3</sub>-Cox assembly and formation of its heme *b*<sub>3</sub>-Cu<sub>B</sub> binuclear center. This article is part of a Special Issue entitled: Biogenesis/Assembly of Respiratory Enzyme Complexes.

© 2011 Elsevier B.V. All rights reserved.

## 1. Introduction

## 1.1. Generalities

The emerging ease of whole genome sequencing approaches has revealed the existence of multiple terminal oxidases in bacteria, which allow them to utilize efficiently varying oxygen (O<sub>2</sub>) concentrations in their environments. Most of the terminal oxidases belong to the heme-Cu: O<sub>2</sub> reductase superfamily (heme-Cu: O<sub>2</sub> reductases); they reduce O<sub>2</sub> to H<sub>2</sub>O and couple electron transfer with vectorial proton translocation across the membrane [1]. Members of this superfamily are present in eukarya, archaea and eubacteria. They are thought to be of monophyletic origin, possibly related to nitric oxide reductases (NOR) [2]. NOR catalyze reduction of NO to N<sub>2</sub>O during denitrification. Although they do not reduce O<sub>2</sub> and do not generate a proton gradient [3], their structural organization is similar to heme-Cu: O<sub>2</sub> reductases.

The heme-Cu: O<sub>2</sub> reductases can be divided into different subgroups in respect to their electron donors: those that use a *c*-type cytochrome

(*i. e.*, cytochrome *c* oxidase, Cox) and those that use a quinol (*i. e.*, quinol oxidase, Qox). Cox are universally conserved oligomeric membrane proteins that terminate the respiratory chains of aerobic and facultative aerobic organisms. The structure and function of Cox enzymes have been studied intensely because of their central role in energy metabolism. Given the complexity of these multi-subunit, multi-cofactor enzymes, their assembly also attracted much attention. Indeed, assembly defects in the human Cox are major causes of mitochondrial disorders, and are crucial for neurodegenerative diseases, cancer and aging [4–7]. Among the Cox enzymes, the *aa*<sub>3</sub>-type Cox (*aa*<sub>3</sub>-Cox) of mitochondria and many bacterial species are highly abundant and the best-studied group. However, the heme-Cu: O<sub>2</sub> reductases represent a diverse ensemble of enzymes with significantly different subunit compositions and cofactors [8]. In this review, we focus on the structure, subunit maturation and assembly of the *cbb*<sub>3</sub>-type Cox (*cbb*<sub>3</sub>-Cox) that represent the second most abundant group after the *aa*<sub>3</sub>-Cox. Most of the studies on *cbb*<sub>3</sub>-Cox biogenesis were performed with the facultative phototrophic  $\alpha$ -proteobacterium *Rhodobacter capsulatus*. This species is used as a model organism for these studies as it contains only one heme-Cu: O<sub>2</sub> reductase, *cbb*<sub>3</sub>-Cox. *R. capsulatus* is still capable of aerobic respiration in the absence of it because it also contains an unrelated terminal oxidase (*bd*-type quinol oxidase) (Fig. 1) [9,10].

1.2. Classification of heme-Cu: O<sub>2</sub> reductases

Members of heme-Cu: O<sub>2</sub> reductases are diverse in terms of their subunit composition, heme cofactor content, electron donor, and O<sub>2</sub> affinity (Table 1) [8,11,12]. All members have a conserved core subunit

Abbreviations: Cox, cytochrome *c* oxidase; Qox, quinol oxidase; Ccm, cytochrome *c* maturation; NOR, nitric oxide reductase; BN-PAGE, blue-native polyacrylamide gel electrophoresis; Cu, Copper

☆ This article is part of a Special Issue entitled: Biogenesis/Assembly of Respiratory Enzyme Complexes.


\* Corresponding author. Tel.: +49 761 203 5250; fax: +49 761 203 5289.

\*\* Corresponding author. Tel.: +1 215 898 4394; fax: +1 215 898 8780.

E-mail addresses: [Hans-Georg.Koch@biochemie.uni-freiburg.de](mailto:Hans-Georg.Koch@biochemie.uni-freiburg.de) (H.-G. Koch),

[fdaldal@sas.upenn.edu](mailto:fdaldal@sas.upenn.edu) (F. Daldal).

<sup>1</sup> S. Ekici and G. Pawlik are first co-authors for this work.



**Fig. 1.** The respiratory chain of *R. capsulatus* and its regulation in response to oxygen availability. The protein complexes and electron carrier of the respiratory chain are shown in yellow. UQ and UQH<sub>2</sub> correspond to the oxidized and reduced form of quinones. Soluble cytochrome c<sub>2</sub> and the membrane-bound cytochrome c<sub>v</sub> (indicated as c in the figure) are electron donors for the photosynthetic reaction center and *cbb*<sub>3</sub>-Cox. Metabolic processes which also consume redox equivalents are shown in blue. Regulatory proteins are shown in gray. RegB is a sensor kinase that phosphorylates the response regulator RegA under low oxygen concentrations. Phosphorylated RegA (RegA-P) inhibits (shown as the red square) expression of *ccoNOQP* (*cbb*<sub>3</sub>-Cox) and *hupSLC* ([NiFe] hydrogenase), but stimulates the expression of the *bd*-type quinol oxidase and of genes required for photosynthesis. In addition, RegA-P stimulates (shown as an arrow) the expression of *nifHDK* (nitrogenase) and the *cbb*<sub>1</sub> and *cbb*<sub>2</sub> operons (CO<sub>2</sub> fixation) both directly and indirectly [154]. Under microaerobic and anaerobic conditions HrvA further represses *ccoNOQP* expression but stimulates *bd*-type quinol oxidase expression. Finally, FnrL probably activates *ccoNOQP* expression in the transition phase between aerobic and anaerobic conditions. *R. capsulatus* SenC and its *R. sphaeroides* homologue PrfC were suggested to be involved in oxygen-dependent expression of *ccoNOQP* and photosynthetic genes. FixIJ-K regulatory system discussed in the text is not shown in the figure.

(Subunit I), containing at least 12 transmembrane helices. This subunit contains a low spin heme (of *a*- or *b*-type), a binuclear metal center composed of a high spin heme (of *a*-, *o*-, or *b*-type heme, referred to as *a*<sub>3</sub>, *o*<sub>3</sub> or *b*<sub>3</sub>)-iron, and a Cu atom (Cu<sub>B</sub>), as well as a tyrosine residue covalently linked to a histidine ligand of Cu<sub>B</sub> [1,8]. Besides subunit I, which is the catalytic heart where O<sub>2</sub> is reduced to H<sub>2</sub>O, heme-Cu: O<sub>2</sub> reductases have at least two other core subunits: Subunit II is the primary electron acceptor, and in some cases, it harbors extra cofactors like a binuclear Cu center (Cu<sub>A</sub>), or is a *c*-type cytochrome. Subunit III in many cases does not contain any cofactor except in *cbb*<sub>3</sub>-Cox where it is a diheme *c*-type cytochrome (Table 1) [1]. Most bacterial Cox are composed of these three core subunits, but mitochondrial Cox are more complex with 13 subunits of which only three (subunits I, II, and III) are encoded in the mitochondrial genome. An additional fourth subunit with a single transmembrane helix is also present in some bacterial Cox. However, this subunit is unlike any of the ten nuclear encoded mitochondrial subunits [13].

A classification, based on common features of the core subunits, and key residues in proton transfer pathways, defines three (A, B and C) types of heme-Cu: O<sub>2</sub> reductases (Table 1) (Fig. 2) [1]. Mitochondrial-*aa*<sub>3</sub>-Cox (Type A), *ba*<sub>3</sub>-Cox of *Thermus thermophilus* (Type B) and *cbb*<sub>3</sub>-Cox of *R. capsulatus* (Type C) are the most representative members of these types.

Available three-dimensional (3D) structures of the heme-Cu: O<sub>2</sub> reductase family members (*i. e.*, the *aa*<sub>3</sub>-Cox from *Paracoccus denitrificans*, *Rhodobacter sphaeroides* and bovine heart mitochondria; *ba*<sub>3</sub>-Cox from *T. thermophilus*, *bo*<sub>3</sub>-Qox from *Escherichia coli* and the *cbb*<sub>3</sub>-Cox from *Pseudomonas stutzeri*) (Fig. 2A) [14–19] reveal a remarkably

conserved Cu and heme cofactor arrangement in these enzymes. Subunit I of type A enzymes contain two proton-transfer pathways referred to as K- and D- channels (Table 1). The four electrons required for O<sub>2</sub> reduction at the binuclear heme-Cu<sub>B</sub> center are conveyed sequentially via a non-covalently attached low spin heme, which itself receives electrons from the Cu<sub>A</sub> center in subunit II [15,19]. Type B enzymes are present only in bacteria and archaea, but not in eukaryotes, and constitute the least abundant group [20]. The catalytic center of type B enzymes is similar to that of type A, except that they lack the D-channel for proton transfer [21]. The prototypes of type C are the *cbb*<sub>3</sub>-Cox, which are present only in bacteria, and considered to represent the most distant members of heme-Cu: O<sub>2</sub> reductases. They also lack the D-channel for proton transfer [22–24], their subunits II and III are *c*-type cytochromes (Table 1), and they exhibit higher O<sub>2</sub> affinity as compared with other types of Cox enzymes.

## 2. The *cbb*<sub>3</sub>-Cox

### 2.1. Distribution of *cbb*<sub>3</sub>-Cox in bacteria and its role in pathogenesis

The *cbb*<sub>3</sub>-Cox enzymes are common to proteobacteria and also found in the *cytophaga*, *flexibacter* and *bacteriodes* (CFB) group [12]. They were first described in the facultative symbiotic N<sub>2</sub>-fixing *rhizobiacaea* [25,26]. In *Bradyrhizobium japonicum* expression of *cbb*<sub>3</sub>-Cox is required for symbiotic N<sub>2</sub> fixation under very low O<sub>2</sub> conditions in soybean root nodules [26]. Due to their importance for symbiotic N<sub>2</sub> fixation, the four structural genes of *cbb*<sub>3</sub>-Cox were initially termed *fix-NOQP* [26]. Subsequent studies identified these genes and their

**Table 1**  
Characteristic features of different Cox and NOR enzymes.

	<i>cbb</i> <sub>3</sub> -Cox	NOR	<i>ba</i> <sub>3</sub> -Cox	<i>aa</i> <sub>3</sub> -Cox
Subunit composition	<ul style="list-style-type: none"> <li>• CcoN<sup>a</sup></li> <li>• Two <i>c</i>-type cytochromes: CcoP and CcoO</li> <li>• Two putative small subunits: CcoQ, CcoH</li> </ul>	<ul style="list-style-type: none"> <li>• NorB<sup>a</sup></li> <li>• <i>c</i>-type cytochrome: NorC</li> </ul>	<ul style="list-style-type: none"> <li>• Subunit I<sup>a</sup></li> <li>• Subunit II with or without Cu<sub>A</sub></li> </ul>	<ul style="list-style-type: none"> <li>• Subunit I<sup>a</sup></li> <li>• Subunit II with Cu<sub>A</sub></li> <li>• Subunit III</li> <li>• Subunit IV</li> <li>• Other subunits</li> </ul>
Catalytic binuclear center	<ul style="list-style-type: none"> <li>• Heme <i>b</i><sub>3</sub></li> <li>• Heme <i>b</i></li> <li>• Cu<sub>B</sub></li> <li>• Covalent His-Tyr</li> <li>• Ca<sup>2+</sup> coordinating heme <i>b</i> and <i>b</i><sub>3</sub> propionates</li> </ul>	<ul style="list-style-type: none"> <li>• Heme <i>b</i><sub>3</sub></li> <li>• Heme <i>b</i></li> <li>• Non-heme iron Fe<sub>B</sub></li> <li>• No covalent His-Tyr</li> <li>• Ca<sup>2+</sup> coordinating heme <i>b</i> and <i>b</i><sub>3</sub> propionates</li> </ul>	<ul style="list-style-type: none"> <li>• Heme <i>a</i><sub>3</sub></li> <li>• Cu<sub>B</sub></li> <li>• Covalent His-Tyr</li> </ul>	<ul style="list-style-type: none"> <li>• Heme <i>a</i><sub>3</sub></li> <li>• Heme <i>a</i></li> <li>• Cu<sub>B</sub></li> <li>• Covalent His-Tyr</li> </ul>
Enzymatic properties	<ul style="list-style-type: none"> <li>K<sub>m</sub> ~ 7 nM (for O<sub>2</sub>)</li> <li>K<sub>m</sub> ~ 12 mM (for NO)</li> </ul>		K <sub>m</sub> ~ 0,1 μM	K <sub>m</sub> 0,7–2 μM
Proton pathway	'K' pathway	Protons from periplasm	K pathway	K and D pathways

<sup>a</sup> Catalytic subunit of the enzyme

products also in non-symbiotic bacteria such as *R. capsulatus*, and referred to them as *ccoNOQP* [27–29]. A recent bioinformatics study, based on the occurrence of *ccoN* and *ccoO* genes, identified the *ccoNOQP* cluster in all bacterial species, except the *Thermotogales*, *Deinococcales* and *Firmicutes* [12]. So far, no *cbb*<sub>3</sub>-Cox coding sequences were identified in archaea [30]. Interestingly, *ccoNO* genes were found recently in the last mitochondrial ancestor bacterium *Midichloria mitochondrii* [31].

The *cbb*<sub>3</sub>-Cox is present in many microaerophilic pathogenic bacteria and implicated in host colonization (Table 2). In some species including *Campylobacter jejuni* [32], *Helicobacter pylori* [33,34], *Neisseria meningitidis* and *Neisseria gonorrhoeae* [35] *cbb*<sub>3</sub>-Cox is the sole Cox, and it is suggested that its high O<sub>2</sub> affinity allows these pathogens to colonize low O<sub>2</sub> containing tissues [11]. Correlations between *cbb*<sub>3</sub>-Cox and host colonization have so far been examined mainly in *C. jejuni*, which is a non-fermenting microaerophile that colonizes the small and large intestines of humans and animals [36], causing acute gastroenteritis in humans. *C. jejuni* has a branched aerobic electron transport chain, with a *cbb*<sub>3</sub>-Cox of high O<sub>2</sub> affinity (K<sub>m</sub> = 40 nM) and a quinol oxidase (CioAB) of low (K<sub>m</sub> = 800 nM) O<sub>2</sub> affinity. Gene expression studies have shown that CioAB was mainly expressed at high O<sub>2</sub> conditions [37], while *cbb*<sub>3</sub>-Cox was induced significantly in *C. jejuni* cells colonizing the chick caecum, which is a low O<sub>2</sub> environment [38]. *C. jejuni* mutants lacking *cbb*<sub>3</sub>-Cox were unable to colonize the chick caecum, whereas mutants lacking CioAB were slightly impaired in colonization [38]. The caecum is primarily an anaerobic environment, hence *C. jejuni* is likely to be exposed transiently to anaerobic conditions. Probably, it tolerates anaerobiosis by using alternative electron acceptors like nitrate, nitrite, DMSO or TMAO [39], while its *cbb*<sub>3</sub>-Cox would keep cellular O<sub>2</sub> concentrations low to protect its crucial O<sub>2</sub>-labile enzymes and allow establishment of anaerobic respiration [40,41]. Indeed, *C. jejuni* mutants lacking nitrate or nitrite reductases are significantly attenuated for colonizing the chick caecum [39]. Thus, anaerobic respiration together with *cbb*<sub>3</sub>-Cox seems to allow *C. jejuni* to adapt successfully to environmental changes at their ecological niches.


A *cbb*<sub>3</sub>-Cox is also present in *Brucella suis*, which is an intracellular Gram-negative pathogen and the causative agent of brucellosis [42]. Under microaerobic conditions in liquid cultures, *B. suis* expresses mainly *cbb*<sub>3</sub>-Cox. However inside macrophages, it relies almost exclusively on a *bd*-type quinol oxidase [42], which apparently has a higher O<sub>2</sub> affinity than *cbb*<sub>3</sub>-Cox in this species. In *H. pylori* which is responsible for stomach ulcers, *cbb*<sub>3</sub>-Cox is the sole respiratory terminal oxidase recognizable in its genome (Table 2). Transposon insertion mutants in *ccoN*, *ccoO* and *ccoP* genes of this species were obtained under microaerobiosis, indicating that *cbb*<sub>3</sub>-Cox is not essential under these conditions [43]. Whether *H. pylori* needs *cbb*<sub>3</sub>-Cox to colonize the intestine is unknown. The fact that *H. pylori* mutants that lack fumarate reductase fail to colonize the mouse stomach [44] indicates that anaerobic respiration is crucial for virulence of this and other microaerophilic pathogens.

Although not yet studied extensively, the role of *cbb*<sub>3</sub>-Cox during pathogenesis might provide a potential antibacterial target for therapeutic interventions. Chemicals that interfere with *cbb*<sub>3</sub>-Cox might act as powerful specific antibiotics with minimal side effects due to the absence of this type of Cox in mammals [45].

## 2.2. Regulation of *cbb*<sub>3</sub>-Cox expression

In *R. capsulatus* and most other bacteria the steady-state concentration and activity of *cbb*<sub>3</sub>-Cox is dependent on the environmental O<sub>2</sub> concentrations. This activity is high under microaerobic, low under fully aerobic, and even lower under fully anaerobic growth conditions [46]. Several components, including RegA (or PrrA), RegB (or PrrB), FnrL, HvrA, and FixLJ-K that respond to different O<sub>2</sub> and redox conditions, are thought to coordinately regulate this process (Fig. 1) [47]. RegA and RegB constitute a global two-component regulatory system that controls the expression of many genes involved in energy metabolism, including *cbb*<sub>3</sub>-Cox, *bd*-quinol oxidase, and various genes related to photosynthesis, hydrogen utilization, nitrogen fixation and carbon assimilation [46,48–51]. RegA is a response regulator with a conserved helix–turn–helix DNA binding motif, whereas RegB is a sensor kinase proposed to contain a conserved region that monitors redox changes inside the cells. Under microaerobic and aerobic conditions the dephosphorylated form of RegA activates *cbb*<sub>3</sub>-Cox expression. Conversely, under anaerobic conditions RegA is phosphorylated by the sensor kinase RegB, and phosphorylated RegA represses *cbb*<sub>3</sub>-Cox and enhances photosynthesis genes expression [47]. In *R. sphaeroides*, it was proposed that *cbb*<sub>3</sub>-Cox monitors electron flow through its CcoN subunit [52]. When this electron flow is high, it sends an inhibitory signal to PrrA and PrrB (homologues of RegA and RegB) to repress photosynthesis gene expression [52]. Thus, in this species abolishing *cbb*<sub>3</sub>-Cox results in activation of photosynthetic genes under aerobic conditions [53].

Besides RegA and RegB, some other genes also regulate *cbb*<sub>3</sub>-Cox expression. FnrL-like genes are found upstream of *ccoNOQP* in *R. capsulatus* [54], and upstream of both *ccoNOQP* and *ccoGHIS* (involved in the assembly of *cbb*<sub>3</sub>-Cox, see below) in other organisms including *R. sphaeroides*, *P. denitrificans*, *B. japonicum*. Studies performed in *E. coli* established that FnrL controls switching from aerobic to anaerobic respiration [55]. Inactivation of FnrL in *R. capsulatus* decreases *cbb*<sub>3</sub>-Cox activity by about 80% under anaerobic growth conditions, suggesting that it acts as an activator in the absence of O<sub>2</sub> [47]. This effect is counter balanced by the trans-acting regulatory protein HvrA, which functions as a repressor of *ccoNOQP* under microaerobic and anaerobic conditions (Fig. 1) [47]. Another major set of regulators of microaerophilic growth as well as photosynthesis is the FixLJ-K regulatory system. FixL is a histidine kinase that can bind O<sub>2</sub> via a heme *b* group, whereas FixJ is a transcription factor that undergoes phosphorylation by FixL under microaerophilic



**Fig. 2.** A. 3D structures of different types of heme-Cu: O<sub>2</sub> reductases and NO reductase: Catalytic subunit I – blue (*cbb*<sub>3</sub>-Cox CcoN), subunit II (*cbb*<sub>3</sub>-Cox CcoO) – magenta, subunit III (*cbb*<sub>3</sub>-Cox CcoP) – green, additional subunit – orange. B. Architecture of the catalytic binuclear center of the different types of heme-Cu:O<sub>2</sub> reductases and NO reductase. Heme – red, His – green, Tyr – violet, Glu – orange. Spheres: copper – blue, heme iron – dark gray, non-heme iron – light gray, calcium – yellow. The structures depicted are taken from protein database (PDB) entries PDB ID: 3MK7 (*cbb*<sub>3</sub>-Cox) [14], PDB ID: 3OOR (NOR) [60], PDB ID: 1M56 (*aa*<sub>3</sub>-Cox) [19], PDB ID: 1EHK (*ba*<sub>3</sub>-Cox) [17] using Jmol software (<http://www.jmol.org/>).

conditions. Phosphorylated FixL then activates the expression of FixK, a downstream transcription regulator that binds specifically upstream of *ccoN* gene to activate its expression [56–58]. The FixLJ-K system regulating expression of *cbb*<sub>3</sub>-Cox is also found in many other bacteria, including *B. japonicum*, *Caulobacter crescentus* and *Novosphingobium aromativorans* [59].

### 2.3. Subunit composition and structure of *cbb*<sub>3</sub>-Cox

Usually, *cbb*<sub>3</sub>-Cox enzymes are composed of CcoN (subunit I), CcoO (subunit II), CcoQ (subunit IV) and CcoP (subunit III) proteins (Table 1) [12]. Excitingly, the 3D structure of *P. stutzeri cbb*<sub>3</sub>-Cox was solved recently at a resolution of 3.2 Å [14]. The structure revealed that the overall shape and size of the membrane-embedded part of *cbb*<sub>3</sub>-Cox is similar to those of *R. sphaeroides aa*<sub>3</sub>-Cox [19] and *T. thermophilus ba*<sub>3</sub>-Cox, [17] whereas its periplasmic part, mainly formed by the

heme containing domains of CcoO and CcoP, is more surface-exposed than in other Cox enzymes (Fig. 2A). Moreover, highlighting the close evolutionary relationship between the *cbb*<sub>3</sub>-Cox and NOR enzymes, the 3D structure of CcoN was found very similar to that of *Pseudomonas aeruginosa* NOR [60], even though the amino acid sequence conservation between these two proteins is below 40%.

CcoN forms a “clamshell”-like structure, with its amino (N-) and carboxyl (C-) termini located close to each other on the cytoplasmic side. CcoN also has two β strands that form a hairpin loop between the helices V and VI. Two calcium atoms with a proposed stabilizing role are found in the structure (Fig. 2B). One of these atoms interacts directly with the heme *b*<sub>3</sub>-Cu<sub>B</sub> catalytic center, while the second one is located between the loops IV and V at the edge of the structure. The clamshell structure surrounds the catalytic center located near the outer membrane surface and consists of a low spin heme *b*, a high spin heme *b*<sub>3</sub> and a Cu (Cu<sub>B</sub>) atom. The open edge of the clamshell is close to heme *b*, whereas heme *b*<sub>3</sub> and Cu<sub>B</sub> are at its distal end. Like in NOR, the hemes *b* and *b*<sub>3</sub> are linked together by a calcium atom, which is coordinated to the carboxyl groups of pyrrole D rings of both hemes. Like other Cox enzymes, three histidine residues coordinate the Cu<sub>B</sub> atom, and one of them is covalently ligated to a tyrosine residue on helix VII (Fig. 2B). This linkage is absent in NOR, and in *aa*<sub>3</sub>-Cox the ligating tyrosine residue is located on helix VI (Table 1) [19]. Like the type B enzymes, *cbb*<sub>3</sub>-Cox contains only one proton pathway that is positioned similar to the K-channel in the type A enzymes. In *cbb*<sub>3</sub>-Cox, hydrophobic residues block the equivalent of the D-channel, and moreover, none

**Table 2**  
Various terminal oxidases and NOR in some pathogenic bacteria.

	<i>cbb</i> <sub>3</sub> -Cox	NOR	<i>aa</i> <sub>3</sub> -Cox	<i>bd</i> -type Qox
<i>Campylobacter jejuni</i>	✓	✓		✓
<i>Helicobacter pylori</i> J99	✓			
<i>Neisseria gonorrhoeae</i>	✓	✓		
<i>Neisseria meningitidis</i>	✓	✓		
<i>Pseudomonas aeruginosa</i> PAO1	✓	✓	✓	✓
<i>Vibrio cholera</i>	✓			✓

of the residues of the proposed proton pathway of *cbb*<sub>3</sub>-Cox (Ser240, Tyr223, His243, Tyr317, Thr215, Tyr251 of CcoN and Glu49 of CcoP) is conserved in type A heme-Cu: O<sub>2</sub> reductases. However, Tyr223 and Tyr317 can be found in the predicted proton channel of type B enzyme *ba*<sub>3</sub>-Cox of *T. thermophilus* [17]. Similar proton pathways have not been identified in NOR which does not pump protons, but it is noteworthy that some *cbb*<sub>3</sub>-Cox also exhibit low NOR activity (Table 1) [61].

CcoO is a mono-heme *c*-type cytochrome, and has an N-terminally located transmembrane helix as an anchor (Table 1). It is thought to convey electrons to heme *b* of CcoN, and makes strong contacts with the  $\alpha$ -helices of CcoN. In particular, CcoO contacts the calcium atom that coordinates heme *b* and heme *b*<sub>3</sub> at the catalytic center via its Ser102 residue. Of the two cavities that are visible on the 3D structure of *cbb*<sub>3</sub>-Cox, one is close to the periplasmic face of the membrane, between CcoO and CcoN. This cavity is located at a position equivalent to the end of the D-channel of type A enzymes, and is proposed to provide an exit path for protons and water molecules from the catalytic site to the periplasm. The second cavity is membrane-embedded and connected by narrow and hydrophobic channels to the catalytic site, possibly providing an O<sub>2</sub> access pathway. An equivalent hydrophobic channel is also present in the NOR structure, and is suggested to allow access of NO to the catalytic site of this enzyme [60].

CcoP is a di-heme *c*-type cytochrome with both of its heme-groups solvent exposed. In the case of *P. stutzeri*, CcoP has two transmembrane helices connected via a long linker that makes multiple contacts to the cytoplasmic part of CcoN (Table 1). In some other organisms, including *R. capsulatus*, CcoP probably contains only one transmembrane helix as its anchor. The distal heme group of CcoP is thought to accept electrons from a soluble or membrane bound electron donor (e.g., cytochrome *c*<sub>2</sub> or cytochrome *c*<sub>1</sub> in *R. capsulatus*) and transfers them to the proximal heme of this subunit. This latter heme then conveys electrons to the heme group of CcoO, which transfers them to the catalytic center in CcoN.

CcoQ is a small subunit formed by a single transmembrane helix, and is not present in all *cbb*<sub>3</sub>-Cox. It does not contain any cofactor, and its elimination does not completely abolish *cbb*<sub>3</sub>-Cox activity. CcoQ is absent in *P. stutzeri* 3D structure, which instead contains an unassigned  $\alpha$ -helix, located close to helices IX and XI of CcoN. The location of CcoQ in *R. capsulatus cbb*<sub>3</sub>-Cox is unknown, but it can be cross-linked to CcoP, suggesting that it is associated with this subunit [62].

### 3. Assembly of the *cbb*<sub>3</sub>-Cox

In general, Cox assembly is an intrinsically complex process because this oligomeric enzyme is membrane embedded and contains multiple cofactors [63]. Membrane insertion and maturation of individual subunits, insertion of cofactors, and assembly of cognate partners have to be coordinated to produce an active enzyme [63,64]. In eukaryotes, *aa*<sub>3</sub>-Cox assembly involves more than 30 factors, which provide timely availability of mitochondrially-encoded subunits, or which associate transiently with various assembly intermediates [13,65]. Assembly of bacterial *aa*<sub>3</sub>-Cox is possibly less complex, but still relies on specific components that mediate heme and Cu insertion into the subunits. Assembly of *cbb*<sub>3</sub>-Cox is particularly challenging because the maturation of CcoO and CcoP require the *c*-type cytochrome maturation (Ccm) machinery, whereas CcoN relies on heme *b* and Cu atom insertion processes. A set of genes, *ccoGHIS* are located immediately downstream of *ccoNOQP* operon in most *cbb*<sub>3</sub>-Cox containing bacteria [54,66–68]. The roles of *ccoGHIS* products in the assembly of *cbb*<sub>3</sub>-Cox are described below.

#### 3.1. An assembly pathway for the subunits of *cbb*<sub>3</sub>-Cox and role of CcoH

Assembly of *cbb*<sub>3</sub>-Cox has been studied in *R. capsulatus* membranes using blue-native polyacrylamide gel electrophoresis (BN-PAGE) analyses (Fig. 3A) [62,67,69]. This technique allows identification of

membrane protein complexes in their native state, and is particularly suitable for examining abundant and stable respiratory chain complexes [70]. In *R. capsulatus* membranes, BN-PAGE, activity staining and immunodetection analyses showed that *cbb*<sub>3</sub>-Cox forms an active complex of 230 kDa, which contains all four (CcoNOQP) structural subunits [67]. However, BN-PAGE does not necessarily reflect the correct molecular mass of a protein complex, as binding of lipids, detergent and Coomassie blue can interfere with the migration on BN-PAGE [71]. Indeed *R. capsulatus cbb*<sub>3</sub>-Cox purified after detergent-solubilization migrates as an active 160 kDa complex on BN-PAGE (Pawlik et al., unpublished data).

Later on, it was also found that the putative assembly factor CcoH is a part of the 230 kDa complex (Fig. 3A) [69]. CcoH encodes a single-spanning membrane protein with an extended periplasmic domain that is suggested to serve as a dimerization domain [69]. CcoH was also found in the purified *R. capsulatus cbb*<sub>3</sub>-Cox (Pawlik et al., unpublished results). Unlike the mitochondrial *aa*<sub>3</sub>-Cox, which forms together with other respiratory complexes a network of supercomplexes [70], no similar large macromolecular assemblies were so far observed in the case of *cbb*<sub>3</sub>-Cox in *R. capsulatus* membranes [67]. In addition to the active 230 kDa complex, using BN-PAGE one large (210 kDa) and one small (~40 kDa) inactive assembly intermediates were also detected in *R. capsulatus* membranes (Fig. 3A). The 210 kDa complex contained CcoN, CcoO and CcoH, but lacked CcoP [67,69], and the ~40 kDa complex had CcoP [67], CcoQ and CcoH [62,69]. The CcoO subunit in the 210 kDa, and the CcoP subunit in the ~40 kDa complexes contained their covalently attached heme groups as indicated by their peroxidase activity [67]. Thus, maturation of the *c*-type cytochrome subunits occurred prior to their assembly into *cbb*<sub>3</sub>-Cox. Genetic data supported these findings, as mutants expressing truncated CcoP derivatives contained only the 210 kDa, and not the 230 kDa complex [67], and mutations abolishing the *c*-type cytochrome maturation process also prevented *cbb*<sub>3</sub>-Cox assembly [72]. Although not essential, CcoQ seemed to improve *cbb*<sub>3</sub>-Cox assembly as in its absence active enzymes were produced at reduced amounts. More recent chemical cross-linking data suggest that interactions between CcoQ and CcoP favor assembly of the ~40 kDa complex with the 210 kDa complex [62]. The surprising finding that CcoH is associated with the assembly intermediates of *cbb*<sub>3</sub>-Cox [69] suggested that this protein behaved more like a *bona fide* subunit of *R. capsulatus* enzyme rather than an assembly factor. This is further supported by the observation that the steady-state stability of CcoH is strictly dependent on the presence of CcoNOQP [69]. Available data suggest that *cbb*<sub>3</sub>-Cox assembly might occur via a fusion between the CcoNOH and CcoQPH subcomplexes mediated by CcoH (Fig. 3A) [67,69]. The 3D structure of *cbb*<sub>3</sub>-Cox indicates that CcoO is sandwiched between CcoN and CcoP [14]. A similar assembly pathway has also been proposed for *B. japonicum cbb*<sub>3</sub>-Cox, although CcoH was not included in this model [73].

#### 3.2. Maturation of *cbb*<sub>3</sub>-Cox subunits

##### 3.2.1. CcoN: Roles of CcoS, CcoI and CcoG in hemes *b* and Cu incorporation

CcoN is an integral membrane protein and is most likely co-translationally inserted into *R. capsulatus* cytoplasmic membrane [74,75]. The two heme *b* and the Cu cofactor are deeply buried within CcoN, and their incorporation into this subunit might be critical for proper folding and stabilization of *cbb*<sub>3</sub>-Cox. However, whether these cofactors are inserted co-translationally into CcoN, and if so, in which order, is unknown. A co-translational insertion process for hemes *a* and Cu cofactors for *aa*<sub>3</sub>-Cox has been proposed [76], but not yet established.

In *R. capsulatus*, CcoS is a small membrane protein of 56 amino acids with no putative heme or Cu binding motifs, which is highly tolerant towards mutations of its conserved residues (Pawlik et al., unpublished results). A mutant lacking CcoS assembles an inactive *cbb*<sub>3</sub>-Cox variant of 230 kDa that lacks both of the heme *b* cofactors

of CcoN, but contains properly matured CcoO and CcoP subunits [66]. This finding indicates that *c*-type cytochrome maturation is not affected, and further supports that the CcoN and *c*-type cytochrome maturation processes occur independently of each other [66]. Whether the inactive *cbb*<sub>3</sub>-Cox variant present in a *R. capsulatus* *ccoS* knock out mutant contains Cu<sub>B</sub> is unknown. Clearly, CcoS is required for proper maturation of CcoN, but defining its precise role in *cbb*<sub>3</sub>-Cox assembly deserves further studies.

In both mitochondria and bacteria, heme insertion into *aa*<sub>3</sub>-Cox is mediated by Surf1 (called Shy1 in yeast), which accepts heme *a* directly from heme *a* synthase (CtaA) and is thought to transfer it to subunit I [77]. In the absence of Surf1, *aa*<sub>3</sub>-Cox assembly is significantly reduced, but not completely abolished [78–80] implying that Surf1 is important, but not essential for heme *a* insertion into the subunit I. Moreover, Surf1 homologues are not present in all species that contain heme *a* containing Cox (e. g. they are missing in *Bacillus subtilis* and *T. thermophilus*). In the case of *T. thermophilus*, a protein called CbaX, which has no homology to Surf1, was implicated in heme *a* insertion into *ba*<sub>3</sub>-Cox [81]. A Surf1 homologue is absent in *R. capsulatus* genome, and a *R. sphaeroides* mutant that lacks Surf1 is impaired only in the assembly of *aa*<sub>3</sub>- but not *cbb*<sub>3</sub>-Cox [78]. Available data indicate that Surf1 is dedicated to heme *a*, and not involved in heme *b* insertion. In fact, how heme *b* is inserted into any membrane protein is still enigmatic.

Copper is an essential cofactor of many enzymes such as heme-Cu: O<sub>2</sub> reductases, Cu-Zn superoxide dismutase, multicopper oxidase (or laccase) and tyrosinase. Like many metals, free Cu is basically undetectable in cells, minimizing the risk of reactive oxygen species production. Hence, Cu acquisition, trafficking, storage, and delivery to the target sites are strictly controlled processes [82]. Cu can be found in two redox states: the reduced Cu<sup>+</sup> is considered to be the transported form while the oxidized Cu<sup>2+</sup> is probably the catalytically active form. In eukaryotes, the Cu transporter (Ctr) proteins represent the main Cu importers, and they have been studied intensely [83,84]. How Cu is imported into the bacterial cytoplasm is not clear. In the Gram-positive *Enterococcus hirae* a universally conserved P-type ATPase [84], and in *B. subtilis* the membrane protein YcnJ [85] was suggested to import Cu. In addition, two P-type ATPases are required to import Cu into cyanobacteria (i. e., CtaA and PacS) [86,87] and into the thylakoids of plant chloroplasts (i. e., PAA1/HMA6 and PAA2/HMA8) [86,87]. Very recently, CcoA, a novel member of the major facilitator superfamily (MFS) of membrane proteins, was implicated in Cu import in *R. capsulatus* (Ekici et al., unpublished results, see below) and *Schizosaccharomyces pombe* [88]. The MFS-type transporters are structurally and functionally distinct from the Ctr-type Cu transporters, and might illustrate a novel Cu uptake system.

Different to Cu import pathways, bacterial Cu efflux pathways are intensely studied, mainly motivated by the toxicity of Cu for all organisms. A major Cu efflux mechanism is provided by P-type ATPases (P1B subgroup). P-type ATPases are integral membrane pumps that hydrolyse ATP for maintaining ion homeostasis, electrochemical gradients and lipid asymmetry [89]. The P-type ATPase superfamily is composed of 11 distinct subgroups of which the P1B is one of the largest and most widespread. P1B-type ATPases use ATP hydrolysis for Cu extrusion across the cytoplasmic membrane [90–93], and contain the characteristic motifs of P-type ATPases, including the ATP binding domain, the phosphorylation domain, and the phosphatase domain [94]. They also contain an N-terminal heavy metal binding domain (HMBD) with the conserved Cu-binding motif (MXCXXC), the membrane embedded ion translocation (CPX)-motif and the conserved histidine-proline (HP)-motif [94]. The mechanism of loading cytosolic Cu to Cu-ATPases is not well understood, but HMBD domains are thought to interact with specific metallochaperones to deliver Cu atoms to these transporters [95,96]. The recent resolution of the first 3D structure of a P1B-type ATPase from *Legionella pneumophila* revealed a putative docking platform for Cu chaperones [97].

The P1B-type ATPases pump Cu<sup>+</sup> out of the bacterial cytoplasm, but their specific roles seem to be determined by their rate of Cu efflux [98]. For example two of them, CopA1 and CopA2 of *P. aeruginosa*, are highly homologous to each other (35% identity and 50% similarity), yet their functions differ. CopA1 has a faster Cu efflux rate and is required for Cu detoxification as its absence renders cells highly sensitive to Cu. CopA2 has a much slower Cu efflux rate and its absence does not affect cellular Cu sensitivity, but leads to the loss of *cbb*<sub>3</sub>-Cox activity [99].


CcoI is a homologue of CopA2, and is the product of *ccoI* gene located in the *ccoGHIS* cluster in many species [54,68]. In *B. japonicum* and *R. capsulatus*, CcoI appears to be specifically required for *cbb*<sub>3</sub>-Cox assembly, but a *Rubrivivax gelatinosus* mutant lacking the CcoI homologue CtpA seems to be defective in other Cu containing enzymes such as *caa*<sub>3</sub>-Cox and N<sub>2</sub>O reductase as well [100]. In *R. capsulatus*, absence of CcoI [66] or mutations in its N-terminal HMBD domain or CPC motifs, drastically reduce the steady-state amounts of *cbb*<sub>3</sub>-Cox subunits [66] (Pawlik et al., unpublished results). Cu supplementation does not suppress the *cbb*<sub>3</sub>-Cox defect of a mutant lacking CcoI, which is also not sensitive to Cu [66]. Whether Cu<sub>B</sub> is inserted into CcoN before or after heme *b* is not known. In case of *aa*<sub>3</sub>-Cox, the availability of enzyme variants containing heme *a*, but lacking Cu<sub>B</sub> suggests that heme *a* is likely to be inserted before Cu<sub>B</sub> [78].

The essential role of CcoI for *cbb*<sub>3</sub>-Cox assembly suggests a model that involves transport of Cu<sup>+</sup> from the cytoplasm to the periplasm, then its concomitant or subsequent insertion into CcoN (Fig. 3B). However it is intriguing that a mutant lacking CcoI is not rescued by exogenous Cu addition, because Cu is thought to be diffused freely across the outer membrane into the periplasm. A possibility is that either Cu<sup>+</sup> transported by CcoI is inserted into CcoN without its full transport across membrane, or that it is delivered upon transport to specific periplasmic Cu chaperones (perhaps SenC or PCu<sub>A</sub>C, see below), which then supplies it to CcoN (Fig. 3B). This model would rationalize the cytoplasmic origin of Cu, and suggests that in the absence of CcoI, its loading onto the specific chaperone might be inefficient.

Among the *R. capsulatus* mutants lacking any one of the *ccoGHIS* products, those devoid of CcoG exhibited the mildest effect on *cbb*<sub>3</sub>-Cox assembly. In the absence of CcoG, all subunits were present at quasi wild type levels, and the enzyme activity was not significantly reduced [66]. CcoG is an integral membrane protein, which has five predicted transmembrane helices and two putative [4Fe-4S] cluster-binding motifs, but whether it contains an iron-sulfur cluster has not been shown experimentally. The exact function of CcoG remains unknown. It has been proposed that it might be involved in intracellular oxidation of Cu<sup>+</sup> to Cu<sup>2+</sup> [68]. Another possibility is that CcoG might reduce Cu<sup>2+</sup> to Cu<sup>+</sup> to provide the substrate for CcoI. If so, then CcoG might work together with the putative MFS-type Cu importer CcoA, provided that CcoA imports Cu<sup>2+</sup> into the cytoplasm (as suggested by the insensitivity of *ccoA* mutant to silver which mimics Cu<sup>+</sup>) (Ekici et al., unpublished results) (Fig. 3B). Clearly, both the structure and function of CcoG are poorly defined and warrant further studies.

### 3.2.2. CcoO and CcoP: *c*-type cytochrome maturation (Ccm) process

The CcoO and CcoP subunits of *cbb*<sub>3</sub>-Cox are matured by a *c*-type cytochrome maturation (Ccm) process, which operates independently from CcoN maturation. Ccm is a post-translation and post-translocation process that occurs on the periplasmic face of the cytoplasmic membrane (Fig. 4). During this process, heme *b* groups are covalently and stereospecifically attached by thioether bonds formed between their vinyl groups and the cysteine thiols of conserved CXXCH motifs of apocytochromes [101,102]. Several Ccm systems are encountered in nature, and in  $\alpha$ - and  $\gamma$ -proteobacteria and *Deinococcus* species maturation of the CcoO and CcoP subunits relies on Ccm-system I, which was reviewed




**Fig. 3.** A. An assembly scheme for *cbb3*-Cox in *R. capsulatus*. Using BN-PAGE, active *cbb3*-Cox is detectable in *R. capsulatus* membranes as a 230 kDa complex. Assembly of the 230 kDa complex proceeds via two inactive assembly intermediates: a 210 kDa subcomplex that contains the catalytic subunit CcoN, the mono-heme cytochrome *c* subunit CcoO and the single-spanning membrane protein CcoH, and a 40 kDa subcomplex composed of the di-heme cytochrome *c* subunit CcoP, the small subunit CcoQ and a second copy of CcoH. Full assembly of *cbb3*-Cox is probably achieved by the dimerization of the cytoplasmic domains of the two CcoH. CcoH was originally considered to function as an assembly factor, but recent data demonstrate that it is a stable component of the fully assembled *cbb3*-Cox. The *c*-type cytochromes in both assembly subcomplexes have their heme groups attached indicating that *c*-type cytochrome maturation preceded the formation of the assembly intermediates. B. A tentative model for the maturation of CcoN subunit. The maturation of CcoN requires Cu and heme *b* insertions. Cu is probably imported into the cytoplasm of *R. capsulatus* by the major facilitator superfamily (MFS) protein CcoA as oxidized  $\text{Cu}^{2+}$ . On the cytoplasmic face of the membrane  $\text{Cu}^{2+}$  is reduced to  $\text{Cu}^{+}$  by the ferredoxin-like protein CcoG and transported back to the periplasmic face of the membrane by the P1B-type ATPase CcoI. CcoI then supplies  $\text{Cu}^{+}$  to CcoN either directly within the membrane or from the periplasm, or via some putative periplasmic Cu binding proteins such as SenC and PCu<sub>A</sub>C. The insertion of heme *b* and heme *b*<sub>3</sub> into CcoN is probably mediated by the assembly factor CcoS.

in detail recently [103,104]. Ccm-system I can be divided into three operation modules that are described briefly below.

**3.2.2.1. Module 1: Transport and relay of heme.** The function of this module is to translocate cytoplasmically synthesized heme *b* to the periplasm, and to prepare it for ligation to apo-cytochromes (Fig. 4, right). This module consists of five proteins, named CcmABCDE. CcmABCD is suggested to form an ATP binding cassette (ABC)-type transporter involved in loading heme *b* to CcmE. CcmE has a single membrane-anchoring helix and a conserved histidine residue that covalently binds heme, which then acts as a heme donor to apocytochromes. CcmA has an ATP binding domain and Walker A and B motifs needed for ATPase activity. CcmB is an integral membrane protein with six transmembrane helices, required for membrane localization of CcmA. CcmC contains a conserved tryptophan-rich WWD motif and is involved in loading heme to CcmE, and apparently CcmD enhances this process. A recent bioinformatics study grouped CcmC and its homologues as the “heme handling proteins” (HHP) [105]. To what extent CcmAB and CcmCD function together or separately for loading heme onto CcmE and delivering heme-loaded CcmE to the heme ligation complex is unclear. A possibility is that CcmC and CcmD are necessary for attachment of heme to CcmE

[106–108], and that CcmAB is required for ATP dependent release of heme-loaded CcmE from CcmCD [109]. On the other hand, whether CcmAB has another, yet to be defined, role in this process is unknown [110,111].

**3.2.2.2. Module 2: Apocytochrome thio-oxidoreduction and chaperoning.** Like most *c*-type apocytochromes, CcoO and CcoP are thought to be secreted via the SecYEG secretory pathway (Fig. 4, left). Their signal-anchor sequences are not processed, and serve as their N-terminal membrane-anchors. Upon translocation across the cytoplasmic membrane, the DsbA-DsbB dependent oxidative protein-folding pathway [112,113] is thought to rapidly oxidize the thiol groups of apocytochromes to possibly avoid their proteolytic degradation. The disulfide bonds formed between the cysteines at the conserved heme binding (CXXCH) motifs need to be reduced prior to heme *b* attachment. Thus, a Ccm-specific thioreduction pathway involving CcdA, CcmG and CcmH proteins has been proposed [114,115]. CcdA is an integral membrane protein with six transmembrane helices and is responsible for conveying from the cytoplasm to the periplasm reducing equivalents for this process. CcmG and CcmH are membrane-anchored thioredoxin-like proteins that contain a single CXXC domain facing the periplasm [115], and thought to reduce the



**Fig. 4.** Cytochrome *c* maturation (Ccm) system in *R. capsulatus*. The cytochrome *c* maturation components can be divided into three modules. First module (right) is the transport of heme and its preparation for ligation to apocytochrome that involves CcmABCD proteins which forms a ABC-type transporter that has a role in the delivery of heme to the heme chaperone CcmE before the ligation. It is unknown whether heme is transported via CcmABCD complex or by another unknown protein. CcmA and CcmB are required for the release of heme-bound CcmE from CcmC and CcmD. CcmC and CcmD are involved in the heme attachment to CcmE. Second module (left) involves the apocytochrome thiooxidation and chaperoning processes. After the translocation of apocytochrome into the periplasm, the cysteine thiols are first oxidized by DsbA-DsbB, then reduced by CcdA, CcmG and/or CcmH. Third module (middle) consists of CcmHIF which is the heme ligation core complex. CcmI binds to C-terminal part of apocytochrome to deliver it to the core complex for the catalysis of the thioether bond formation between reduced apocytochrome and heme vinyl groups forming the mature cytochrome *c* (adapted from [155]).

disulfide bonds at the heme binding sites. Of the three components, CcdA and CcmG are not required in the absence of DsbA-DsbB dependent thio-oxidation pathway, but CcmH is still required for *c*-type cytochrome production, suggesting that it has an additional role. Moreover, the 3D structure of CcmG showed that, in addition to its canonical thioredoxin fold, it has a cavity that might bind apocytochromes which is in accordance with a putative holdase role of CcmG as shown in *R. capsulatus* [116].

**3.2.2.3. Module 3: Apocytochrome and heme *b* ligation.** In *R. capsulatus*, CcmF, CcmH and CcmI proteins have been proposed to form a heme ligation core complex (Fig. 4, middle), based on reciprocal copurification experiments documenting protein-protein interactions between these components [117]. Of these proteins, CcmF is a large integral membrane protein with 11 transmembrane helices, with a tryptophan-rich (WWD) signature motif and four conserved histidine residues facing the periplasm proposed to bind heme *b* before its ligation to apocytochromes [106]. It interacts with heme-loaded CcmE [118], is a heme handling protein (HHP) like CcmC [106]. CcmI is a bipartite protein that contains a membrane embedded N-terminal region with two transmembrane helices and a cytoplasmic leucine-zipper-like motif containing loop (CcmI-1 domain) and a large periplasmic C-terminal extension with tetratricopeptide repeat (TPR)-like motifs (CcmI-2 domain) [119–121]. Unlike most other bacteria, in *E. coli* the homologue of CcmI-2 domain is fused to the C-terminal end of CcmH, leaving only CcmF and a modified “CcmH” as components of heme ligation core complex [122]. *R. capsulatus*, mutants lacking CcmI can be suppressed by overproduction of CcmF and CcmH or CcmG. Complementation studies with two distinct domains of CcmI indicated that CcmF and CcmH overproduction relates to the

functional role of CcmI-1 domain [123], whereas CcmG overproduction to that of CcmI-2 [120]. These findings suggested that CcmI might be a junction point between the CcdA-CcmG dependent thio-reduction and CcmF-CcmH dependent heme ligation processes. Indeed, CcmI was initially proposed to chaperone apocytochromes to the heme ligation complex [121], and very recently *in vitro* studies provided strong biochemical evidence to support this proposal [156]. Protein-protein interaction studies conducted using purified CcmI and purified apocytochrome *c*<sub>2</sub> indicated that the C-terminal CcmI-2 domain of CcmI recognizes and binds tightly the most C-terminal helix of apocytochrome *c*<sub>2</sub> in the absence of heme *b*. The folding process of cytochrome *c* (and cytochrome *c*<sub>2</sub>) [124] indicates that their most C-terminal helix interacts with their most N-terminal heme binding helix to form a stable folding intermediate that can trap heme *b* non-covalently [125]. Altogether, these observations led to the proposal that at least some *c*-type apocytochromes are first recognized via their C-terminal helices by the periplasmic CcmI-2 domain of the CcmFHI core complex (Fig. 4, middle), and then released from this domain upon transfer of heme *b* from CcmF to apocytochrome, and subsequently, the thioether bonds are formed. Once the thioether bonds are formed, cytochrome *c* folds into its final structure.

Recently, another type of post-translational modification was seen in CcoP of pathogenic *Neisseria* species [126]. The CcoP subunit of *N. gonorrhoeae* contains a O-linked glycosylation in its periplasmic moiety [126]. The role of this glycosylation, perhaps important for pathogenesis, remains to be determined. It is noteworthy that O-linked glycosylation was also observed for ScoI (see below SenC) and cytochrome CysB, which are involved in *cbb*<sub>3</sub>-Cox assembly and electron transfer to *cbb*<sub>3</sub>-Cox, respectively [126]. Whether this or other kinds of modification of *cbb*<sub>3</sub>-Cox subunits also occurs in other species,



and whether this affects biogenesis of this enzyme deserves further studies.

### 3.2.3. Additional proteins involved in *cbb*<sub>3</sub>-Cox biogenesis

Several additional proteins of not yet fully defined roles, including SenC [127], PCu<sub>A</sub>C [128,129], DsbA [130] and CcoA (Ekici et al., unpublished results) might be involved in *cbb*<sub>3</sub>-Cox biogenesis, and described below.

**3.2.3.1. SenC (Scol).** SenC is a membrane-anchored protein with a single transmembrane helix, and in its absence, *cbb*<sub>3</sub>-Cox biogenesis in *R. capsulatus* [127] and in *P. aeruginosa* [131] is drastically decreased and *cbb*<sub>3</sub>-Cox activity is regained upon addition of exogenous Cu. Its periplasmic domain contains a thioredoxin fold and a conserved Cu binding motif comprising CxxxC and a His ligand [127]. SenC is a close homologue of the universally conserved Scol protein, which has been implicated mainly into the assembly of the Cu<sub>A</sub> center of subunit II of *aa*<sub>3</sub>-Cox. Whether Scol acts as a direct donor of Cu to subunit II [132], or is a thiol:disulfide oxidoreductase reducing appropriate cysteines of subunit II for subsequent Cu delivery by another Cu chaperone [129] is not yet clear [133]. *B. japonicum* mutants lacking SenC were impaired in the assembly of *aa*<sub>3</sub>- but not *cbb*<sub>3</sub>-Cox and showed reduced symbiotic N<sub>2</sub> fixation [134]. Similarly, *R. sphaeroides* mutants lacking PrrC (a SenC homologue) had no effect on *cbb*<sub>3</sub>-Cox assembly although they were defective in photosynthetic growth [52]. Like *R. capsulatus* *senC*, *R. sphaeroides* *prcC* is located next to *prxAB* (homologues of *R. capsulatus* *regAB*) genes that encode a two component regulatory system controlling energy processes, including photosynthesis [46,135]. Whether *prcC* mutations have any polar effect on downstream *prxB* gene, indirectly interfering with cellular amounts of PrrB, is not known (Fig. 1).

Several studies have shown that the role of prokaryotic Scol homologues is not restricted to their involvement in *aa*<sub>3</sub>-Cox assembly. A role of these proteins in oxidative stress response [136] or photosynthetic gene regulation [52] has been proposed. Bioinformatics based genome surveys indicated that Scol homologues are also present in many bacterial species, like *R. capsulatus*, which do not have a Cu<sub>A</sub> containing Cox [133]. *R. capsulatus* mutants lacking SenC produce very low amounts of *cbb*<sub>3</sub>-type Cox [127,137], a phenotype which was rescued by addition of exogenous Cu [127]. *In vitro* experiments showed that in a mutant lacking SenC *cbb*<sub>3</sub>-Cox assembly proceeded at a significantly reduced level (Lohmeyer et al., unpublished results). Very recent chemical cross-linking studies indicated that SenC interacts with CcoP and CcoH *in vivo*, suggesting that it is involved directly in *cbb*<sub>3</sub>-Cox assembly (Lohmeyer et al., unpublished results) (Fig. 3B).

**3.2.3.2. PCu<sub>A</sub>C (DR1885).** The periplasmic Cu<sub>A</sub>-chaperone (PCu<sub>A</sub>C) was first identified in *Deinococcus radiodurans* and its structural gene is located near Scol [128]. In this species, PCu<sub>A</sub>C is thought to provide Cu to the Cu<sub>A</sub> center after Scol reduces the appropriate disulfide bond at subunit II of *aa*<sub>3</sub>-Cox. In the case of *T. thermophilus* PCu<sub>A</sub>C is involved in Cu transfer to Cu<sub>A</sub> of *ba*<sub>3</sub>-Cox [129]. Homologues of PCu<sub>A</sub>C are also found in many species that lack *aa*<sub>3</sub>-Cox such as *R. capsulatus*. Unlike *D. radiodurans*, in *R. capsulatus* the structural gene of PCu<sub>A</sub>C is not located near that of SenC, but its product contains similar conserved metal binding motifs. In this species whether PCu<sub>A</sub>C affects *cbb*<sub>3</sub>-Cox Cu<sub>B</sub> center assembly is unknown as no chromosomal PCu<sub>A</sub>C knockout mutant could be obtained so far (Fig. 3B).

**3.2.3.3. DsbA.** *R. capsulatus* mutants lacking the periplasmic thiol:disulfide oxidoreductase DsbA, primarily involved in oxidative protein folding pathway [112], overproduce the periplasmic protease DegP [138]. They have pleiotropic phenotypes, including temperature sensitivity for growth (35 °C), osmosensitivity, filamentation and decreased respiratory capabilities. Even at permissive growth

temperature (25 °C), they exhibit reduced *cbb*<sub>3</sub>-Cox activity. Supplementation of the growth medium with redox active chemicals like cysteine/cystine or Cu<sup>2+</sup>, restores both temperature sensitive growth and *cbb*<sub>3</sub>-Cox defects of mutants lacking DsbA. Remarkably, DsbA knock out mutants revert frequently to regain their growth ability at 35 °C without any need for redox active supplements, and concomitantly, they restore their *cbb*<sub>3</sub>-Cox production. These revertants acquire mutations in *degP* that decrease drastically the protease activity of DegP [138]. These findings point to important links between the formation of disulfide bond, degradation of misfolded periplasmic proteins, and production of active *cbb*<sub>3</sub>-Cox. In the absence of DsbA, whether exogenously supplied Cu<sup>2+</sup> acts exclusively as a source of oxidant, or also eases acquisition of Cu as a missing cofactor for *cbb*<sub>3</sub>-Cox production, or both is unknown. Future studies might clarify these emerging links between major cellular processes and *cbb*<sub>3</sub>-Cox biogenesis.

**3.2.3.4. CcoA.** Very recent studies of *R. capsulatus* mutants that exhibit “Cu supplement dependent *cbb*<sub>3</sub>-Cox production” phenotype yielded a novel component, CcoA involved in *cbb*<sub>3</sub>-Cox biogenesis (Ekici et al., unpublished results). These mutants lacked *cbb*<sub>3</sub>-Cox activity on Cu-depleted media, but regained it upon exogenous Cu<sup>2+</sup> supplementation. Molecular genetic studies established that these mutants were defective in an open reading frame (ORF) of previously unknown function, now named *ccoA* due to its role in *cbb*<sub>3</sub>-Cox biogenesis. CcoA is an integral membrane protein formed by two subdomains of six transmembrane helices each, which are separated by a large cytoplasmic loop. Its sequence is highly homologous to the Major Facilitator Superfamily (MFS) type transporters [139]. It contains the AVYGRR and ARFGRE between its second-third and eighth-ninth transmembrane helices, respectively. These motifs are highly similar to DRXGRR motifs that characterize members of the MFS family. In addition, unlike many other MFS members, CcoA contains a putative metal (Cu) binding (MX<sub>x</sub>M) motif as found in Ctr type Cu importers [140], suggesting a role in metal transport. In *R. capsulatus* mutants lacking CcoA, transcription and translation initiation of *cco-NOQP* occur, but the steady-state amounts of *cbb*<sub>3</sub>-Cox subunits are quasi-absent (Ekici et al., unpublished results). Thus, absence of CcoA perturbs a co- or post-translational step(s) of *cbb*<sub>3</sub>-Cox biogenesis. Remarkably, supplementation with Cu<sup>2+</sup>, but not with other metals like Mn<sup>2+</sup>, Zn<sup>2+</sup> or Fe<sup>3+</sup>, palliates the *cbb*<sub>3</sub>-Cox production defect. CcoA knock out mutants are not sensitive to Cu<sup>2+</sup>, but their cellular Cu content is lower than in normal cells, and they revert very frequently to restore their *cbb*<sub>3</sub>-Cox activity. These by-pass suppressors restore normal cellular Cu content and production of *cbb*<sub>3</sub>-Cox. Surprisingly, they also become hypersensitive to Cu<sup>2+</sup> but not to other metals, including Ag<sup>+</sup>. These findings suggest that the MFS-type transporter CcoA may be involved in Cu acquisition by a hitherto unknown pathway, which seems needed for normal biogenesis of *cbb*<sub>3</sub>-Cox. Very recently, the Mfc1 protein of *Schizosaccharomyces pombe*, which is also a MFS-type transporter with a MX<sub>x</sub>M motif and homologous to CcoA, was shown to import Cu across the plasma membrane [88]. In *S. pombe*, the Ctr-type transporters are thought to mediate high affinity Cu uptake, and Mfc1 might provide an additional low affinity Cu import route. Clearly, future studies of this novel family of MFS-type putative Cu importers might provide invaluable information about bacterial Cu acquisition pathways and their link to *cbb*<sub>3</sub>-Cox biogenesis.

### 3.2.4. Role of membrane lipids for *cbb*<sub>3</sub>-Cox biogenesis

Membrane lipids, especially those that act as “lipochaperones”, are important determinants for membrane protein structure and activity [141]. These lipids bind to specific locations of proteins in stoichiometric amounts, and influence their folding, stability, steady-state amounts and activity [142]. Phospholipids were shown to co-crystallize with heme-Cu: O<sub>2</sub> reductases from many different species [143–145], and

the stability and organization of respiratory chain super complexes were significantly influenced by some phospholipids like cardiolipin [146,147]. A specific role of lipids on biogenesis and catalytic activity of *cbb*<sub>3</sub>-Cox is also likely, as changes in membrane lipid composition seem to affect the amounts of *cbb*<sub>3</sub>-Cox found in *R. capsulatus* membranes [148]. Moreover, different lipid environments were also shown to affect carbon monoxide binding to CcoP [149]. *R. capsulatus* mutants that were defective in ornithine lipid (OL) biosynthesis lacked *cbb*<sub>3</sub>-Cox [148]. Studies of these mutants defined two genes, *olsA* and *olsB* coding for N-acyltransferase and O-acyltransferase, respectively, which are both required for OL biosynthesis [148,150]. OL is a non-phosphorus membrane lipid that usually accounts only for a small fraction of the total lipids found in bacteria. In some species, OL are synthesized under phosphate limiting conditions, and are used as replacement for phosphate containing lipids [151–153]. In *R. capsulatus* OL biosynthesis is not regulated by phosphate availability, but absence of OL affects the steady-amounts of a group of membrane proteins, including most of the c-type cytochromes. Mutants unable to synthesize OL have very low amounts of *cbb*<sub>3</sub>-Cox and the electron carrier cytochromes *c*<sub>2</sub> and *c*<sub>γ</sub>. They contain a small amount of cytochrome *c*<sub>1</sub>, and consequently, very low ubiquinol: cytochrome *c*<sub>2</sub> oxidoreductase (cytochrome *bc*<sub>1</sub>) activity. Interestingly, these defects are both temperature- and growth medium-dependent. Especially on enriched media at regular growth temperature (35 °C), mutants lacking OL are photosynthesis-incompetent and have no *cbb*<sub>3</sub>-Cox. Pulse-chase studies conducted using cytochrome *c*<sub>γ</sub> indicated that absence of OL decreased drastically its cellular amount [148], suggesting that in the absence of OL, a group of proteins including *cbb*<sub>3</sub>-Cox, may be more prone to misfolding and enhanced degradation.

#### 4. Summary and future prospects

In recent years, enormous amount of progress has been accomplished in studies aimed at understanding the biogenesis of *cbb*<sub>3</sub>-Cox. This second most common class of heme-Cu: O<sub>2</sub> reductases is becoming attractive for future studies due to the availability of an excellent model system like *R. capsulatus* providing molecular genetic analyses, a 3D structure at high resolution allowing refined structural and functional inquiries, and exquisite biochemical approaches developed and practiced in various laboratories. Salient recent accomplishments on this topic include an emerging subunit assembly process highlighting the indispensable role of CcoH as a “subcomplex assembler”, the better defined role of CcoI subgroup among the P1B-type ATPases as a Cu exporter to *cbb*<sub>3</sub>-Cox, and possibly to a few other Cu containing enzymes. Furthermore, the discovery of a periplasmic Cu chaperone PCu<sub>A</sub>C, the emerging interactions between the ScoI homologue SenC/PrrC and *cbb*<sub>3</sub>-Cox subunits and its assembly factors, as well as the novel MFS-type putative Cu importer CcoA, whose Cu acquisition functions correlate with the production of *cbb*<sub>3</sub>-Cox, provide exciting novel insight into the complexity of binuclear center assembly. An attempt was made to organize all known components into a highly speculative working scheme that outlines an emerging path of Cu delivery to the heme-Cu<sub>B</sub> center of *cbb*<sub>3</sub>-Cox (Fig. 3B). Indeed, much work is needed to probe experimentally the validity of the various facets of this working model under construction. Currently, we are reaching a stage where the crucial players required for *cbb*<sub>3</sub>-Cox assembly are becoming identified, and ongoing molecular genetic and biochemical studies are now initiating research aimed to untangle the sophisticated and unique biogenesis process of *cbb*<sub>3</sub>-Cox.

#### Acknowledgments

This work was supported by grants DOE 91ER20052 and NIH GM38239 to F.D., and by grants from the German Science Foundation

(DFG-GRK1478) and from the German-French PhD college (DFH/UFA) on “Membranes and Membrane Proteins” to H.G.K.

#### References

- [1] M.M. Pereira, M. Santana, M. Teixeira, A novel scenario for the evolution of haem-copper oxygen reductases, *Biochim. Biophys. Acta* 1505 (2001) 185–208.
- [2] J. Castresana, M. Lubben, M. Saraste, D.G. Higgins, Evolution of cytochrome oxidase, an enzyme older than atmospheric oxygen, *EMBO J.* 13 (1994) 2516–2525.
- [3] U. Flock, J. Reimann, P. Adelroth, Proton transfer in bacterial nitric oxide reductase, *Biochem. Soc. Trans.* 34 (2006) 188–190.
- [4] S. Matoba, J.G. Kang, W.D. Patino, A. Wragg, M. Boehm, O. Gavrilova, P.J. Hurley, F. Bunz, P.M. Hwang, p53 regulates mitochondrial respiration, *Science* 312 (2006) 1650–1653.
- [5] K.Y. Won, S.J. Lim, G.Y. Kim, Y.W. Kim, S.A. Han, J.Y. Song, D.K. Lee, Regulatory role of p53 in cancer metabolism via SCO2 and TIGAR in human breast cancer, *Hum. Pathol.* (2011).
- [6] K. Vesela, H. Hulkova, H. Hansikova, J. Zeman, M. Elleder, Structural analysis of tissues affected by cytochrome *c* oxidase deficiency due to mutations in the SCO2 gene, *APMIS* 116 (2008) 41–49.
- [7] C. Bernstein, A. Facista, H. Nguyen, B. Zaitlin, N. Hassounah, C. Loustaunau, C.M. Payne, B. Banerjee, S. Goldschmid, V.L. Tsikitis, R. Krouse, H. Bernstein, Cancer and age related colonic crypt deficiencies in cytochrome *c* oxidase I, *World J. Gastrointest. Oncol.* 2 (2010) 429–442.
- [8] J.A. Garcia-Horsman, B. Barquera, J. Rumbley, J. Ma, R.B. Gennis, The superfamily of heme-copper respiratory oxidases, *J. Bacteriol.* 176 (1994) 5587–5600.
- [9] B. Marrs, H. Gest, Genetic mutations affecting the respiratory electron-transport system of the photosynthetic bacterium *Rhodospseudomonas capsulata*, *J. Bacteriol.* 114 (1973) 1045–1051.
- [10] F. Daldal, S. Mandaci, C. Winterstein, H. Myllykallio, K. Duyck, D. Zannoni, Mobile cytochrome *c*<sub>2</sub> and membrane-anchored cytochrome *c*<sub>γ</sub> are both efficient electron donors to the *cbb*<sub>3</sub>- and *aa*<sub>3</sub>-type cytochrome *c* oxidases during respiratory growth of *Rhodobacter sphaeroides*, *J. Bacteriol.* 183 (2001) 2013–2024.
- [11] R.S. Pitcher, N.J. Watmough, The bacterial cytochrome *cbb*<sub>3</sub> oxidases, *Biochim. Biophys. Acta* 1655 (2004) 388–399.
- [12] A.L. Ducluzeau, S. Ouchane, W. Nitschke, The *cbb*<sub>3</sub> oxidases are an ancient innovation of the domain bacteria, *Mol. Biol. Evol.* 25 (2008) 1158–1166.
- [13] J.M. Herrmann, S. Funes, Biogenesis of cytochrome oxidase-sophisticated assembly lines in the mitochondrial inner membrane, *Gene* 354 (2005) 43–52.
- [14] S. Buschmann, E. Warkentin, H. Xie, J.D. Langer, U. Ermler, H. Michel, The structure of *cbb*<sub>3</sub> cytochrome oxidase provides insights into proton pumping, *Science* 329 (2010) 327–330.
- [15] S. Iwata, C. Ostermeier, B. Ludwig, H. Michel, Structure at 2.8 Å resolution of cytochrome *c* oxidase from *Paracoccus denitrificans*, *Nature* 376 (1995) 660–669.
- [16] J. Abramson, S. Riistama, G. Larsson, A. Jasaitis, M. Svensson-Ek, L. Laakkonen, A. Puustinen, S. Iwata, M. Wikstrom, The structure of the ubiquinol oxidase from *Escherichia coli* and its ubiquinone binding site, *Nat. Struct. Biol.* 7 (2000) 910–917.
- [17] T. Soulimane, G. Buse, G.P. Bourenkov, H.D. Bartunik, R. Huber, M.E. Than, Structure and mechanism of the aberrant ba(3)-cytochrome *c* oxidase from *Thermus thermophilus*, *EMBO J.* 19 (2000) 1766–1776.
- [18] T. Tsukihara, H. Aoyama, E. Yamashita, T. Tomizaki, H. Yamaguchi, K. Shinzawa-Ittoh, N. Nakashima, R. Yaono, S. Yoshikawa, The whole structure of the 13-subunit oxidized cytochrome *c* oxidase at 2.8 Å, *Science* 272 (1996) 1136–1144.
- [19] M. Svensson-Ek, J. Abramson, G. Larsson, S. Tornroth, P. Brzezinski, S. Iwata, The X-ray crystal structures of wild-type and EQ(I-286) mutant cytochrome *c* oxidases from *Rhodobacter sphaeroides*, *J. Mol. Biol.* 321 (2002) 329–339.
- [20] B.H. Zimmermann, C.I. Nitsche, J.A. Fee, F. Rusnak, E. Munck, Properties of a copper-containing cytochrome *ba*<sub>3</sub>: a second terminal oxidase from the extreme thermophile *Thermus thermophilus*, *Proc. Natl. Acad. Sci. U. S. A.* 85 (1988) 5779–5783.
- [21] H.Y. Chang, J. Hemp, Y. Chen, J.A. Fee, R.B. Gennis, The cytochrome *ba*<sub>3</sub> oxygen reductase from *Thermus thermophilus* uses a single input channel for proton delivery to the active site and for proton pumping, *Proc. Natl. Acad. Sci. U. S. A.* 106 (2009) 16169–16173.
- [22] E. Arslan, A. Kannt, L. Thony-Meyer, H. Hennecke, The symbiotically essential *cbb*<sub>3</sub>-type oxidase of *Bradyrhizobium japonicum* is a proton pump, *FEBS Lett.* 470 (2000) 7–10.
- [23] O. Preisig, R. Zufferey, L. Thony-Meyer, C.A. Appleby, H. Hennecke, A high-affinity *cbb*<sub>3</sub>-type cytochrome oxidase terminates the symbiosis-specific respiratory chain of *Bradyrhizobium japonicum*, *J. Bacteriol.* 178 (1996) 1532–1538.
- [24] J. Hemp, H. Han, J.H. Roh, S. Kaplan, T.J. Martinez, R.B. Gennis, Comparative genomics and site-directed mutagenesis support the existence of only one input channel for protons in the C-family (*cbb*<sub>3</sub> oxidase) of heme-copper oxygen reductases, *Biochemistry* 46 (2007) 9963–9972.
- [25] D. Kahn, M. David, O. Domergue, M.L. Daveran, J. Ghai, P.R. Hirsch, J. Batut, *Rhizobium meliloti* fixGHI sequence predicts involvement of a specific cation pump in symbiotic nitrogen fixation, *J. Bacteriol.* 171 (1989) 929–939.
- [26] O. Preisig, D. Anthamatten, H. Hennecke, Genes for a microaerobically induced oxidase complex in *Bradyrhizobium japonicum* are essential for a nitrogen-fixing endosymbiosis, *Proc. Natl. Acad. Sci. U. S. A.* 90 (1993) 3309–3313.
- [27] L. Thony-Meyer, C. Beck, O. Preisig, H. Hennecke, The ccoNOQP gene cluster codes for a *cb*-type cytochrome oxidase that functions in aerobic respiration of *Rhodobacter capsulatus*, *Mol. Microbiol.* 14 (1994) 705–716.

- [28] K.A. Gray, M. Grooms, H. Myllykallio, C. Moomaw, C. Slaughter, F. Daldal, *Rhodobacter capsulatus* contains a novel *cb*-type cytochrome c oxidase without a CuA center, *Biochemistry* 33 (1994) 3120–3127.
- [29] J.A. Garcia-Horsman, E. Berry, J.P. Shapleigh, J.O. Alben, R.B. Gennis, A novel cytochrome c oxidase from *Rhodobacter sphaeroides* that lacks CuA, *Biochemistry* 33 (1994) 3113–3119.
- [30] J. Hemp, R.B. Gennis, Diversity of the heme-copper superfamily in archaea: insights from genomics and structural modeling, *Results Probl. Cell Differ.* 45 (2008) 1–31.
- [31] D. Sasser, N. Lo, S. Epis, G. D'Auria, M. Montagna, F. Comandatore, D. Horner, J. Pereto, A.M. Luciano, F. Franciosi, E. Ferri, E. Crotti, C. Bazzocchi, D. Daffonchio, L. Sacchi, A. Moya, A. Latorre, C. Bandi, Phylogenomic evidence for the presence of a flagellum and *cbb3* oxidase in the free-living mitochondrial ancestor, *Mol. Biol. Evol.* (2011).
- [32] J. Parkhill, B.W. Wren, K. Mungall, J.M. Kettle, C. Churcher, D. Basham, T. Chillingworth, R.M. Davies, T. Feltham, S. Holroyd, K. Jagels, A.V. Karlyshev, S. Moule, M.J. Pallen, C.W. Penn, M.A. Quail, M.A. Rajandream, K.M. Rutherford, A.H. van Vliet, S. Whitehead, B.G. Barrell, The genome sequence of the food-borne pathogen *Campylobacter jejuni* reveals hypervariable sequences, *Nature* 403 (2000) 665–668.
- [33] J.F. Tomb, O. White, A.R. Kerlavage, R.A. Clayton, G.G. Sutton, R.D. Fleischmann, K.A. Ketchum, H.P. Klenk, S. Gill, B.A. Dougherty, K. Nelson, J. Quackenbush, L. Zhou, E.F. Kirkness, S. Peterson, B. Loftus, D. Richardson, R. Dodson, H.G. Khalak, A. Glodek, K. McKenney, L.M. Fitzgerald, N. Lee, M.D. Adams, E.K. Hickey, D.E. Berg, J.D. Gocayne, T.R. Utterback, J.D. Peterson, J.M. Kelley, M.D. Cotton, J.M. Weidman, C. Fujii, C. Bowman, L. Watthey, E. Wallin, W.S. Hayes, M. Borodovsky, P.D. Karp, H.O. Smith, C.M. Fraser, J.C. Venter, The complete genome sequence of the gastric pathogen *Helicobacter pylori*, *Nature* 388 (1997) 539–547.
- [34] R.A. Alm, L.S. Ling, D.T. Moir, B.L. King, E.D. Brown, P.C. Doig, D.R. Smith, B. Noonan, B.C. Guild, B.L. deJonge, G. Carmel, P.J. Tummino, A. Caruso, M. Uria-Nickelsen, D.M. Mills, C. Ives, R. Gibson, D. Merberg, S.D. Mills, Q. Jiang, D.E. Taylor, G.F. Vovis, T.J. Trust, Genomic-sequence comparison of two unrelated isolates of the human gastric pathogen *Helicobacter pylori*, *Nature* 397 (1999) 176–180.
- [35] Y. Li, A. Hopper, T. Overton, D.J. Squire, J. Cole, N. Tovell, Organization of the electron transfer chain to oxygen in the obligate human pathogen *Neisseria gonorrhoeae*: roles for cytochromes c4 and c5, but not cytochrome c2, in oxygen reduction, *J. Bacteriol.* 192 (2010) 2395–2406.
- [36] G.I. Perez-Perez, M.J. Blaser, *Campylobacter* and *Helicobacter* 2011/03/18 ed., University of Texas Medical Branch at Galveston, Galveston (TX), 1996.
- [37] R.J. Jackson, K.T. Elvers, L.J. Lee, M.D. Gidley, L.M. Wainwright, J. Lightfoot, S.F. Park, R.K. Poole, Oxygen reactivity of both respiratory oxidases in *Campylobacter jejuni*: the *cydAB* genes encode a cyanide-resistant, low-affinity oxidase that is not of the cytochrome *bd* type, *J. Bacteriol.* 189 (2007) 1604–1615.
- [38] C.A. Woodall, M.A. Jones, P.A. Barrow, J. Hinds, G.L. Marsden, D.J. Kelly, N. Dorrell, B.W. Wren, D.J. Maskell, *Campylobacter jejuni* gene expression in the chick cecum: evidence for adaptation to a low-oxygen environment, *Infect. Immun.* 73 (2005) 5278–5285.
- [39] R.A. Weingarten, J.L. Grimes, J.W. Olson, Role of *Campylobacter jejuni* respiratory oxidases and reductases in host colonization, *Appl. Environ. Microbiol.* 74 (2008) 1367–1375.
- [40] C.W. Jones, J.M. Brice, V. Wright, B.A. Ackrell, Respiratory protection of nitrogenase in *Azotobacter vinelandii*, *FEBS Lett.* 29 (1973) 77–81.
- [41] J. Oelze, Respiratory protection of nitrogenase in *Azotobacter* species: is a widely held hypothesis unequivocally supported by experimental evidence? *FEMS Microbiol. Rev.* 24 (2000) 321–333.
- [42] S. Loisel-Meyer, M.P. Jimenez de Bagues, S. Kohler, J.P. Liautard, V. Jubier-Maurin, Differential use of the two high-oxygen-affinity terminal oxidases of *Brucella suis* for in vitro and intramacrophagic multiplication, *Infect. Immun.* 73 (2005) 7768–7771.
- [43] N.R. Salama, B. Shepherd, S. Falkow, Global transposon mutagenesis and essential gene analysis of *Helicobacter pylori*, *J. Bacteriol.* 186 (2004) 7926–7935.
- [44] Z. Ge, Y. Feng, C.A. Dangler, S. Xu, N.S. Taylor, J.G. Fox, Fumarate reductase is essential for *Helicobacter pylori* colonization of the mouse stomach, *Microb. Pathog.* 29 (2000) 279–287.
- [45] M.A. Smith, M. Finel, V. Korolik, G.L. Mendz, Characteristics of the aerobic respiratory chains of the microaerophiles *Campylobacter jejuni* and *Helicobacter pylori*, *Arch. Microbiol.* 174 (2000) 1–10.
- [46] L.R. Swem, S. Elsen, T.H. Bird, D.L. Swem, H.G. Koch, H. Myllykallio, F. Daldal, C.E. Bauer, The RegB/RegA two-component regulatory system controls synthesis of photosynthesis and respiratory electron transfer components in *Rhodobacter capsulatus*, *J. Mol. Biol.* 309 (2001) 121–138.
- [47] D.L. Swem, C.E. Bauer, Coordination of ubiquinol oxidase and cytochrome *cbb(3)* oxidase expression by multiple regulators in *Rhodobacter capsulatus*, *J. Bacteriol.* 184 (2002) 2815–2820.
- [48] S. Elsen, W. Dischert, A. Colbeau, C.E. Bauer, Expression of uptake hydrogenase and molybdenum nitrogenase in *Rhodobacter capsulatus* is coregulated by the RegB-RegA two-component regulatory system, *J. Bacteriol.* 182 (2000) 2831–2837.
- [49] P. Vichivanives, T.H. Bird, C.E. Bauer, F. Robert Tabita, Multiple regulators and their interactions in vivo and in vitro with the *cbb* regulons of *Rhodobacter capsulatus*, *J. Mol. Biol.* 300 (2000) 1079–1099.
- [50] C.S. Mosley, J.Y. Suzuki, C.E. Bauer, Identification and molecular genetic characterization of a sensor kinase responsible for coordinately regulating light harvesting and reaction center gene expression in response to anaerobiosis, *J. Bacteriol.* 176 (1994) 7566–7573.
- [51] M.W. Sganga, C.E. Bauer, Regulatory factors controlling photosynthetic reaction center and light-harvesting gene expression in *Rhodobacter capsulatus*, *Cell* 68 (1992) 945–954.
- [52] J.M. Eraso, S. Kaplan, From redox flow to gene regulation: role of the PrrC protein of *Rhodobacter sphaeroides* 2.4.1, *Biochemistry* 39 (2000) 2052–2062.
- [53] J.P. O'Gara, J.M. Eraso, S. Kaplan, A redox-responsive pathway for aerobic regulation of photosynthesis gene expression in *Rhodobacter sphaeroides* 2.4.1, *J. Bacteriol.* 180 (1998) 4044–4050.
- [54] H.G. Koch, O. Hwang, F. Daldal, Isolation and characterization of *Rhodobacter capsulatus* mutants affected in cytochrome *cbb<sub>3</sub>* oxidase activity, *J. Bacteriol.* 180 (1998) 969–978.
- [55] P.A. Cotter, S. Darie, R.P. Gunsalus, The effect of iron limitation on expression of the aerobic and anaerobic electron transport pathway genes in *Escherichia coli*, *FEMS Microbiol. Lett.* 79 (1992) 227–232.
- [56] F.E. Rey, C.S. Harwood, FixK, a global regulator of microaerobic growth, controls photosynthesis in *Rhodospseudomonas palustris*, *Mol. Microbiol.* 75 (2010) 1007–1020.
- [57] H.M. Fischer, Environmental regulation of rhizobial symbiotic nitrogen fixation genes, *Trends Microbiol.* 4 (1996) 317–320.
- [58] M.A. Gilles-Gonzalez, G. Gonzalez, Heme-based sensors: defining characteristics, recent developments, and regulatory hypotheses, *J. Inorg. Biochem.* 99 (2005) 1–22.
- [59] C. Cosseau, J. Batut, Genomics of the *ccoNOQP*-encoded *cbb<sub>3</sub>* oxidase complex in bacteria, *Arch. Microbiol.* 181 (2004) 89–96.
- [60] T. Hino, Y. Matsumoto, S. Nagano, H. Sugimoto, Y. Fukumori, T. Murata, S. Iwata, Y. Shiro, Structural basis of biological N<sub>2</sub>O generation by bacterial nitric oxide reductase, *Science* 330 (2010) 1666–1670.
- [61] E. Forte, A. Urbani, M. Saraste, P. Sarti, M. Brunori, A. Giuffrè, The cytochrome *cbb<sub>3</sub>* from *Pseudomonas stutzeri* displays nitric oxide reductase activity, *Eur. J. Biochem.* 268 (2001) 6486–6491.
- [62] A. Peters, C. Kulajta, G. Pawlik, F. Daldal, H.G. Koch, Stability of the *cbb<sub>3</sub>*-type cytochrome oxidase requires specific CcoQ-CcoP interactions, *J. Bacteriol.* 190 (2008) 5576–5586.
- [63] P.A. Cobine, F. Pierrel, D.R. Winge, Copper trafficking to the mitochondrion and assembly of copper metalloenzymes, *Biochim. Biophys. Acta* 1763 (2006) 759–772.
- [64] P. Greiner, A. Hannappel, C. Werner, B. Ludwig, Biogenesis of cytochrome c oxidase—in vitro approaches to study cofactor insertion into a bacterial subunit I, *Biochim. Biophys. Acta* 1777 (2008) 904–911.
- [65] D.U. Mick, T.D. Fox, P. Rehling, Inventory control: cytochrome c oxidase assembly regulates mitochondrial translation, *Nat. Rev. Mol. Cell Biol.* 12 (2011) 14–20.
- [66] H.G. Koch, C. Winterstein, A.S. Saribas, J.O. Alben, F. Daldal, Roles of the *ccoGHIS* gene products in the biogenesis of the *cbb<sub>3</sub>*-type cytochrome c oxidase, *J. Mol. Biol.* 297 (2000) 49–65.
- [67] C. Kulajta, J.O. Thumfart, S. Haid, F. Daldal, H.G. Koch, Multi-step assembly pathway of the *cbb<sub>3</sub>*-type cytochrome c oxidase complex, *J. Mol. Biol.* 355 (2006) 989–1004.
- [68] O. Preisig, R. Zufferey, H. Hennecke, The *Bradyrhizobium japonicum* fixGHIS genes are required for the formation of the high-affinity *cbb<sub>3</sub>*-type cytochrome oxidase, *Arch. Microbiol.* 165 (1996) 297–305.
- [69] G. Pawlik, C. Kulajta, I. Sachelaru, S. Schroder, B. Waidner, P. Hellwig, F. Daldal, H.G. Koch, The putative assembly factor CcoH is stably associated with the *cbb<sub>3</sub>*-type cytochrome oxidase, *J. Bacteriol.* 192 (2010) 6378–6389.
- [70] H. Schagger, G. von Jagow, Blue native electrophoresis for isolation of membrane protein complexes in enzymatically active form, *Anal. Biochem.* 199 (1991) 223–231.
- [71] E.H. Heuberger, L.M. Veenhoff, R.H. Duurkens, R.H. Friesen, B. Poolman, Oligomeric state of membrane transport proteins analyzed with blue native electrophoresis and analytical ultracentrifugation, *J. Mol. Biol.* 317 (2002) 591–600.
- [72] M. Deshmukh, G. Brasseur, F. Daldal, Novel *Rhodobacter capsulatus* genes required for the biogenesis of various c-type cytochromes, *Mol. Microbiol.* 35 (2000) 123–138.
- [73] R. Zufferey, O. Preisig, H. Hennecke, L. Thony-Meyer, Assembly and function of the cytochrome *cbb<sub>3</sub>* oxidase subunits in *Bradyrhizobium japonicum*, *J. Biol. Chem.* 271 (1996) 9114–9119.
- [74] H.G. Koch, M. Moser, M. Muller, Signal recognition particle-dependent protein targeting, universal to all kingdoms of life, *Rev. Physiol. Biochem. Pharmacol.* 146 (2003) 55–94.
- [75] H.G. Koch, Folding, assembly, and stability of transmembrane cytochromes, *Curr. Chem. Biol.* 1 (2007) 59–74.
- [76] O. Khalimonchuk, K. Ostermann, G. Rodel, Evidence for the association of yeast mitochondrial ribosomes with Cox11p, a protein required for the Cu(B) site formation of cytochrome c oxidase, *Curr. Genet.* 47 (2005) 223–233.
- [77] A. Hannappel, F.A. Bundschuh, B. Ludwig, Characterization of heme-binding properties of *Paracoccus denitrificans* Surf1 proteins, *FEBS J.* 278 (2011) 1769–1778.
- [78] D. Smith, J. Gray, L. Mitchell, W.E. Antholine, J.P. Hosler, Assembly of cytochrome-c oxidase in the absence of assembly protein Surf1p leads to loss of the active site heme, *J. Biol. Chem.* 280 (2005) 17652–17656.
- [79] F.A. Bundschuh, K. Hoffmeier, B. Ludwig, Two variants of the assembly factor Surf1 target specific terminal oxidases in *Paracoccus denitrificans*, *Biochim. Biophys. Acta* 1777 (2008) 1336–1343.
- [80] V. Tiranti, K. Hoertmagel, R. Carozzo, C. Galimberti, M. Munaro, M. Granatiero, L. Zelante, P. Gasparini, R. Marzella, M. Rocchi, M.P. Bayona-Bafaluy, J.A. Enriquez, G. Uziel, E. Bertini, C. Dionisi-Vici, B. Franco, T. Meitinger, M. Zeviani, Mutations

- of SURF-1 in Leigh disease associated with cytochrome c oxidase deficiency, *Am. J. Hum. Genet.* 63 (1998) 1609–1621.
- [81] J. Werner, O.M. Richter, B. Ludwig, A novel heme a insertion factor gene cotranscribes with the *Thermus thermophilus* cytochrome ba3 oxidase locus, *J. Bacteriol.* 192 (2010) 4712–4719.
- [82] L. Banci, I. Bertini, F. Cantini, S. Ciofi-Baffoni, Cellular copper distribution: a mechanistic systems biology approach, *Cell. Mol. Life Sci.* 67 (2011) 2563–2589.
- [83] S. Lutsenko, Human copper homeostasis: a network of interconnected pathways, *Curr. Opin. Chem. Biol.* 14 (2010) 211–217.
- [84] A. Odermatt, H. Suter, R. Krapf, M. Solioz, Primary structure of two P-type ATPases involved in copper homeostasis in *Enterococcus hirae*, *J. Biol. Chem.* 268 (1993) 12775–12779.
- [85] S. Chillappagari, M. Miethke, H. Trip, O.P. Kuipers, M.A. Marahiel, Copper acquisition is mediated by YcnJ and regulated by YcnK and CsoR in *Bacillus subtilis*, *J. Bacteriol.* 191 (2009) 2362–2370.
- [86] S. Tottey, P.R. Rich, S.A. Rondet, N.J. Robinson, Two Menkes-type atpases supply copper for photosynthesis in *Synechocystis* PCC 6803, *J. Biol. Chem.* 276 (2001) 19999–20004.
- [87] C. Nouet, P. Motte, M. Hanikenne, Chloroplastic and mitochondrial metal homeostasis, *Trends Plant Sci.* 16 (2011) 395–404.
- [88] J. Beaudoin, R. Ioannoni, D. Lopez-Maury, J. Bahler, S. Ait-Mohand, B. Guerin, S.C. Dodani, C.J. Chang, S. Labbe, Mfc1 is a novel forespore membrane copper transporter in meiotic and sporulating cells, *J. Biol. Chem.* (2011).
- [89] W. Kuhlbrandt, Biology, structure and mechanism of P-type ATPases, *Nat. Rev. Mol. Cell Biol.* 5 (2004) 282–295.
- [90] R. Linz, S. Lutsenko, Copper-transporting ATPases ATP7A and ATP7B: cousins, not twins, *J. Bioenerg. Biomembr.* 39 (2007) 403–407.
- [91] S. Lutsenko, N.L. Barnes, M.Y. Bartee, O.Y. Dmitriev, Function and regulation of human copper-transporting ATPases, *Physiol. Rev.* 87 (2007) 1011–1046.
- [92] J.M. Arguello, E. Eren, M. Gonzalez-Guerrero, The structure and function of heavy metal transport PIB-ATPases, *Biomaterials* 20 (2007) 233–248.
- [93] M.G. Palmgren, K.B. Axelsen, Evolution of P-type ATPases, *Biochim. Biophys. Acta* 1365 (1998) 37–45.
- [94] M. Solioz, C. Vulpe, CPx-type ATPases: a class of P-type ATPases that pump heavy metals, *Trends Biochem. Sci.* 21 (1996) 237–241.
- [95] M. Gonzalez-Guerrero, D. Hong, J.M. Arguello, Chaperone-mediated Cu<sup>+</sup> delivery to Cu<sup>+</sup> transport ATPases: requirement of nucleotide binding, *J. Biol. Chem.* 284 (2009) 20804–20811.
- [96] M. Gonzalez-Guerrero, J.M. Arguello, Mechanism of Cu<sup>+</sup>-transporting ATPases: soluble Cu<sup>+</sup> chaperones directly transfer Cu<sup>+</sup> to transmembrane transport sites, *Proc. Natl. Acad. Sci. U. S. A.* 105 (2008) 5992–5997.
- [97] P. Gourdon, X.Y. Liu, T. Skjorringe, J.P. Morth, L.B. Moller, B.P. Pedersen, P. Nissen, Crystal structure of a copper-transporting PIB-type ATPase, *Nature* 475 (2011) 59–64.
- [98] D. Raimunda, M. Gonzalez-Guerrero, B.W. Leeber III, J.M. Arguello, The transport mechanism of bacterial Cu<sup>+</sup>-ATPases: distinct efflux rates adapted to different function, *Biomaterials* 24 (2011) 467–475.
- [99] M. Gonzalez-Guerrero, D. Raimunda, X. Cheng, J.M. Arguello, Distinct functional roles of homologous Cu<sup>+</sup> efflux ATPases in *Pseudomonas aeruginosa*, *Mol. Microbiol.* 78 (2010) 1246–1258.
- [100] B.K. Hassani, C. Astier, W. Nitschke, S. Ouchane, CtpA, a copper-translocating P-type ATPase involved in the biogenesis of multiple copper-requiring enzymes, *J. Biol. Chem.* 285 (2010) 19330–19337.
- [101] P.D. Barker, S.J. Ferguson, Still a puzzle: why is haem covalently attached in c-type cytochromes? *Structure* 7 (1999) R281–R290.
- [102] L. Thony-Meyer, Haem-polypeptide interactions during cytochrome c maturation, *Biochim. Biophys. Acta* 1459 (2000) 316–324.
- [103] C. Sanders, S. Turkarslan, D.W. Lee, F. Daldal, Cytochrome c biogenesis: the Ccm system, *Trends Microbiol.* 18 (2010) 266–274.
- [104] R.G. Kranz, C. Richard-Fogal, J.S. Taylor, E.R. Frawley, Cytochrome c biogenesis: mechanisms for covalent modifications and trafficking of heme and for heme-iron redox control, *Microbiol. Mol. Biol. Rev.* 73 (2009) 510–528 Table of Contents.
- [105] J.H. Lee, E.M. Harvat, J.M. Stevens, S.J. Ferguson, M.H. Saier Jr., Evolutionary origins of members of a superfamily of integral membrane cytochrome c biogenesis proteins, *Biochim. Biophys. Acta* 1768 (2007) 2164–2181.
- [106] B.S. Goldman, D.L. Beck, E.M. Monika, R.G. Kranz, Transmembrane heme delivery systems, *Proc. Natl. Acad. Sci. U. S. A.* 95 (1998) 5003–5008.
- [107] B.S. Goldman, D.L. Beckman, A. Bali, E.M. Monika, K.K. Gabbert, R.G. Kranz, Molecular and immunological analysis of an ABC transporter complex required for cytochrome c biogenesis, *J. Mol. Biol.* 268 (1997) 724–738.
- [108] L. Thony-Meyer, A heme chaperone for cytochrome c biosynthesis, *Biochemistry* 42 (2003) 13099–13105.
- [109] R.E. Feissner, C.L. Richard-Fogal, E.R. Frawley, R.G. Kranz, ABC transporter-mediated release of a haem chaperone allows cytochrome c biogenesis, *Mol. Microbiol.* 61 (2006) 219–231.
- [110] O. Christensen, E.M. Harvat, L. Thony-Meyer, S.J. Ferguson, J.M. Stevens, Loss of ATP hydrolysis activity by CcmAB results in loss of c-type cytochrome synthesis and incomplete processing of CcmE, *FEBS J.* 274 (2007) 2322–2332.
- [111] H. Schulz, R.A. Fabianek, E.C. Pelliccioli, H. Hennecke, L. Thony-Meyer, Heme transfer to the heme chaperone CcmE during cytochrome c maturation requires the CcmC protein, which may function independently of the ABC-transporter CcmAB, *Proc. Natl. Acad. Sci. U. S. A.* 96 (1999) 6462–6467.
- [112] H. Kadokura, F. Katzen, J. Beckwith, Protein disulfide bond formation in prokaryotes, *Annu. Rev. Biochem.* 72 (2003) 111–135.
- [113] H. Nakamoto, J.C. Bardwell, Catalysis of disulfide bond formation and isomerization in the *Escherichia coli* periplasm, *Biochim. Biophys. Acta* 1694 (2004) 111–119.
- [114] D.L. Beckman, R.G. Kranz, Cytochromes c biogenesis in a photosynthetic bacterium requires a periplasmic thioredoxin-like protein, *Proc. Natl. Acad. Sci. U. S. A.* 90 (1993) 2179–2183.
- [115] E.M. Monika, B.S. Goldman, D.L. Beckman, R.G. Kranz, A thioreduction pathway tethered to the membrane for periplasmic cytochromes c biogenesis; in vitro and in vivo studies, *J. Mol. Biol.* 271 (1997) 679–692.
- [116] S. Turkarslan, C. Sanders, S. Ekici, F. Daldal, Compensatory thio-redox interactions between DsbA, CcdA and CcmG unveil the apocytochrome c holdase role of CcmG during cytochrome c maturation, *Mol. Microbiol.* 70 (2008) 652–666.
- [117] C. Sanders, S. Turkarslan, D.W. Lee, O. Onder, R.G. Kranz, F. Daldal, The cytochrome c maturation components CcmF, CcmH, and CcmI form a membrane-integral multi-subunit heme ligation complex, *J. Biol. Chem.* 283 (2008) 29715–29722.
- [118] Q. Ren, U. Ahuja, L. Thony-Meyer, A bacterial cytochrome c heme lyase. CcmF forms a complex with the heme chaperone CcmE and CcmH but not with apocytochrome c, *J. Biol. Chem.* 277 (2002) 7657–7663.
- [119] C. Sanders, C. Boulay, F. Daldal, Membrane-spanning and periplasmic segments of CcmI have distinct functions during cytochrome c biogenesis in *Rhodobacter capsulatus*, *J. Bacteriol.* 189 (2007) 789–800.
- [120] C. Sanders, M. Deshmukh, D. Astor, R.G. Kranz, F. Daldal, Overproduction of CcmG and CcmFH(Rc) fully suppresses the c-type cytochrome biogenesis defect of *Rhodobacter capsulatus* CcmI-null mutants, *J. Bacteriol.* 187 (2005) 4245–4256.
- [121] S.E. Lang, F.E. Jenney Jr., F. Daldal, *Rhodobacter capsulatus* CycH: a bipartite gene product with pleiotropic effects on the biogenesis of structurally different c-type cytochromes, *J. Bacteriol.* 178 (1996) 5279–5290.
- [122] R.A. Fabianek, T. Hofer, L. Thony-Meyer, Characterization of the *Escherichia coli* CcmH protein reveals new insights into the redox pathway required for cytochrome c maturation, *Arch. Microbiol.* 171 (1999) 92–100.
- [123] M. Deshmukh, M. May, Y. Zhang, K.K. Gabbert, K.A. Karber, R.G. Kranz, F. Daldal, Overexpression of ccl1-2 can bypass the need for the putative apocytochrome chaperone CycH during the biogenesis of c-type cytochromes, *Mol. Microbiol.* 46 (2002) 1069–1080.
- [124] S. Akiyama, S. Takahashi, K. Ishimori, I. Morishima, Stepwise formation of alpha-helices during cytochrome c folding, *Nat. Struct. Biol.* 7 (2000) 514–520.
- [125] W. Colon, G.A. Elove, L.P. Wakem, F. Sherman, H. Roder, Side chain packing of the N- and C-terminal helices plays a critical role in the kinetics of cytochrome c folding, *Biochemistry* 35 (1996) 5538–5549.
- [126] A. Vik, F.E. Aas, J.H. Anonsen, S. Bilsborough, A. Schneider, W. Egge-Jacobsen, M. Koomey, Broad spectrum O-linked protein glycosylation in the human pathogen *Neisseria gonorrhoeae*, *Proc. Natl. Acad. Sci. U. S. A.* 106 (2009) 4447–4452.
- [127] D.L. Swem, L.R. Swem, A. Setterdahl, C.E. Bauer, Involvement of SenC in assembly of cytochrome c oxidase in *Rhodobacter capsulatus*, *J. Bacteriol.* 187 (2005) 8081–8087.
- [128] L. Banci, I. Bertini, S. Ciofi-Baffoni, E. Katsari, N. Katsaros, K. Kubicek, S. Mangani, A copper(I) protein possibly involved in the assembly of CuA center of bacterial cytochrome c oxidase, *Proc. Natl. Acad. Sci. U. S. A.* 102 (2005) 3994–3999.
- [129] L.A. Briata, L. Banci, I. Bertini, S. Ciofi-Baffoni, P. Gkazonis, G.A. Spyroulias, A.J. Vila, S. Wang, Mechanism of Cu(A) assembly, *Nat. Chem. Biol.* 4 (2008) 599–601.
- [130] M. Deshmukh, S. Turkarslan, D. Astor, M. Valkova-Valchanova, F. Daldal, The dithiol:disulfide oxidoreductases DsbA and DsbB of *Rhodobacter capsulatus* are not directly involved in cytochrome c biogenesis, but their inactivation restores the cytochrome c biogenesis defect of CcdA-null mutants, *J. Bacteriol.* 185 (2003) 3361–3372.
- [131] E. Frangipani, D. Haas, Copper acquisition by the SenC protein regulates aerobic respiration in *Pseudomonas aeruginosa* PAO1, *FEMS Microbiol. Lett.* 298 (2009) 234–240.
- [132] D.M. Glerum, A. Shtanko, A. Tzagoloff, SCO1 and SCO2 act as high copy suppressors of a mitochondrial copper recruitment defect in *Saccharomyces cerevisiae*, *J. Biol. Chem.* 271 (1996) 20531–20535.
- [133] L. Banci, I. Bertini, S. Ciofi-Baffoni, T. Kozyreva, M. Mori, S. Wang, Sco proteins are involved in electron transfer processes, *J. Biol. Inorg. Chem.* 16 (2011) 391–403.
- [134] D. Buhler, R. Rossmann, S. Landolt, S. Balsiger, H.M. Fischer, H. Hennecke, Disparate pathways for the biogenesis of cytochrome oxidases in *Bradyrhizobium japonicum*, *J. Biol. Chem.* 285 (2011) 15704–15713.
- [135] S. Masuda, Y. Matsumoto, K.V. Nagashima, K. Shimada, K. Inoue, C.E. Bauer, K. Matsuura, Structural and functional analyses of photosynthetic regulatory genes regA and regB from *Rhodovulum sulfidophilum*, *Roseobacter denitrificans*, and *Rhodobacter capsulatus*, *J. Bacteriol.* 181 (1999) 4205–4215.
- [136] K.L. Seib, M.P. Jennings, A.G. McEwan, A Sco homologue plays a role in defence against oxidative stress in pathogenic *Neisseria*, *FEBS Lett.* 546 (2003) 411–415.
- [137] F. Borsetti, V. Tremaroli, F. Michelacci, R. Borghese, C. Winterstein, F. Daldal, D. Zannoni, Tellurite effects on *Rhodobacter capsulatus* cell viability and superoxide dismutase activity under oxidative stress conditions, *Res. Microbiol.* 156 (2005) 807–813.
- [138] O. Onder, S. Turkarslan, D. Sun, F. Daldal, Overproduction or absence of the periplasmic protease DegP severely compromises bacterial growth in the absence of the dithiol: disulfide oxidoreductase DsbA, *Mol. Cell. Proteomics* 7 (2008) 875–890.
- [139] M.H. Saier Jr., J.T. Beatty, A. Goffeau, K.T. Harley, W.H. Heijne, S.C. Huang, D.L. Jack, P.S. Jahn, K. Lew, J. Liu, S.S. Pao, I.T. Paulsen, T.T. Tseng, P.S. Virk, The major facilitator superfamily, *J. Mol. Microbiol. Biotechnol.* 1 (1999) 257–279.
- [140] S. Puig, J. Lee, M. Lau, D.J. Thiele, Biochemical and genetic analyses of yeast and human high affinity copper transporters suggest a conserved mechanism for copper uptake, *J. Biol. Chem.* 277 (2002) 26021–26030.

- [141] M. Bogdanov, E. Mileykovskaya, W. Dowhan, Lipids in the assembly of membrane proteins and organization of protein supercomplexes: implications for lipid-linked disorders, *Subcell. Biochem.* 49 (2008) 197–239.
- [142] W. Dowhan, M. Bogdanov, Lipid-protein interactions as determinants of membrane protein structure and function, *Biochem. Soc. Trans.* 39 (2011) 767–774.
- [143] T. Tiefenbrunn, W. Liu, Y. Chen, V. Katritch, C.D. Stout, J.A. Fee, V. Cherezov, High resolution structure of the ba3 cytochrome c oxidase from *Thermus thermophilus* in a lipidic environment, *PLoS One* 6 (2011) e22348.
- [144] K. Shinzawa-Itoh, H. Aoyama, K. Muramoto, H. Terada, T. Kurauchi, Y. Tadehara, A. Yamasaki, T. Sugimura, S. Kurono, K. Tsujimoto, T. Mizushima, E. Yamashita, T. Tsukihara, S. Yoshikawa, Structures and physiological roles of 13 integral lipids of bovine heart cytochrome c oxidase, *EMBO J.* 26 (2007) 1713–1725.
- [145] L. Qin, C. Hiser, A. Mulichak, R.M. Garavito, S. Ferguson-Miller, Identification of conserved lipid/detergent-binding sites in a high-resolution structure of the membrane protein cytochrome c oxidase, *Proc. Natl. Acad. Sci. U. S. A.* 103 (2006) 16117–16122.
- [146] K. Pfeiffer, V. Gohil, R.A. Stuart, C. Hunte, U. Brandt, M.L. Greenberg, H. Schagger, Cardiolipin stabilizes respiratory chain supercomplexes, *J. Biol. Chem.* 278 (2003) 52873–52880.
- [147] G. Paradies, F.M. Ruggiero, G. Petrosillo, E. Quagliariello, Age-dependent decline in the cytochrome c oxidase activity in rat heart mitochondria: role of cardiolipin, *FEBS Lett.* 406 (1997) 136–138.
- [148] S. Aygun-Sunar, S. Mandaci, H.G. Koch, I.V. Murray, H. Goldfine, F. Daldal, Ornithine lipid is required for optimal steady-state amounts of c-type cytochromes in *Rhodobacter capsulatus*, *Mol. Microbiol.* 61 (2006) 418–435.
- [149] Y. Huang, J. Reimann, L.M. Singh, P. Adelroth, Substrate binding and the catalytic reactions in *cbb<sub>3</sub>*-type oxidases: the lipid membrane modulates ligand binding, *Biochim. Biophys. Acta* 1797 (2010) 724–731.
- [150] S. Aygun-Sunar, R. Bilaloglu, H. Goldfine, F. Daldal, *Rhodobacter capsulatus* OlsA is a bifunctional enzyme active in both ornithine lipid and phosphatidic acid biosynthesis, *J. Bacteriol.* 189 (2007) 8564–8574.
- [151] D.E. Minnikin, H. Abdolrahimzadeh, The replacement of phosphatidylethanolamine and acidic phospholipids by an ornithine-amide lipid and a minor phosphorus-free lipid in *Pseudomonas fluorescens* NCMB 129, *FEBS Lett.* 43 (1974) 257–260.
- [152] C. Benning, Z.H. Huang, D.A. Gage, Accumulation of a novel glycolipid and a betaine lipid in cells of *Rhodobacter sphaeroides* grown under phosphate limitation, *Arch. Biochem. Biophys.* 317 (1995) 103–111.
- [153] O. Geiger, V. Rohrs, B. Weissenmayer, T.M. Finan, J.E. Thomas-Oates, The regulator gene *phoB* mediates phosphate stress-controlled synthesis of the membrane lipid diacylglycerol-N, N, N-trimethylhomoserine in *Rhizobium* (*Sinorhizobium*) *meliloti*, *Mol. Microbiol.* 32 (1999) 63–73.
- [154] S. Elsen, L.R. Swem, D.L. Swem, C.E. Bauer, RegB/RegA, a highly conserved redox-responding global two-component regulatory system, *Microbiol. Mol. Biol. Rev.* 68 (2004) 263–279.
- [155] N.C. Hunter, F. Daldal, M. Thurnauer, J.T. Beaty, *The Purple Phototrophic Bacteria*, Springer, Dordrecht, 2008.
- [156] A.F. Verissimo, H. Yang, X. Wu, C. Sanders, F. Daldal, CcmI subunit of CcmFHI heme ligation complex functions as an apocytochrome c chaperone during c-type cytochrome maturation, *J. Biol. Chem.* (2011 Sep 28) [Electronic publication ahead of print].