

effects of chronic oral administration of quinapril (40 mg/day), captopril (150 mg/day) and placebo on vascular angiotensin (Ang) I conversion in humans.

**Methods:** Patients (n = 187) scheduled for coronary bypass surgery were randomized at least 7 days before surgery. Segments of the internal mammary artery were obtained and contracted to increasing doses of Ang I and II (0.1 nM to 1  $\mu$ M) in organ baths. Ang I conversion was defined as (1) the difference between pEC50 of the dose-response curves to Ang I and II and (2) as the area between curves (ABC) of Ang I and II.

**Results:** Baseline characteristics were comparable between groups. Quinapril and captopril treated patients had a comparable and significant reduction in mean blood pressure compared to placebo (p = 0.04).

	Quinapril	Captopril	Placebo
delta pEC50	1.0 $\pm$ 0.12*	0.86 $\pm$ 0.1	0.64 $\pm$ 0.1
ABC	81 $\pm$ 8*	84 $\pm$ 4	67 $\pm$ 8

Mean  $\pm$  SEM, \* p < 0.05 (quinapril compared to placebo)

**Conclusion:** This is the first randomized study in patients showing tissue effects of ACE-inhibitors after treatment with adequate clinical doses. The results suggest functional differences between quinapril and captopril.

### 1090 Coronary Artery Disease II

Tuesday, March 31, 1998, 9:00 a.m.-11:00 a.m.

Georgia World Congress Center, West Exhibit Hall Level  
Presentation Hour: 9:00 a.m.-10:00 a.m.

### 1090-6 Segmental Analysis of Coronary Artery Lesions Using 3-dimensional Echocardiography in Comparison With Coronary Angiography: A Clinical Semiquantitative Study

J. Yao, F.J. de Feijter, J.D. Kasprzak, M.A. Taams, J.R.T.C. Roelandt.  
Thoraxcenter, Erasmus University Rotterdam, The Netherlands

Stenotic lesions of coronary artery (CA) are of clinical predictive value for ischemic heart events. The accuracy of transoesophageal 3-dimensional echocardiography (3DTEE) in defining the severity of CA stenosis is not known.

**Purpose:** To analyze the quantitative potential of 3DTEE in defining moderate to severe CA stenosis.

**Methods:** A group of 20 subjects with CA angiography (A) less than 3 months scheduled for routine or intra-operative TEE was selected. 3DTEE was performed using a 5 MHz transducer with an integrated on-line 3DE acquisition system. The proximal CA was divided into 5 segments including left main (1), anterior descending (1), circumflex (2) and right (1) CA. CAA and 3DTEE were analyzed segment-by-segment by independent observers and a score of 0-3 was given to each segment according to none, mild (< 50%), moderate (50-90%) and severe (> 90%) stenosis located. A score was given to each patient according to distribution pattern of the stenotic lesions defined as none (0), localized (1) or diffusid (2).

**Results:** 11 of 20 pts had CA stenosis of proximal segments by CAA (3 localized, 8 diffusid). A total of 85 segments of CA were observed by 3DTEE in all 20 subjects. Mild, moderate and severe stenotic segments were 15, 9 and 5 by 3DTEE and 14, 13 and 5 by CAA. Agreement (4 x 4 table) between these two methods was 80%. The sensitivity, specificity, positive and negative predictive value and general accuracy of 3DTEE for detecting moderate to severe CA stenosis were 61%, 95%, 79%, 90% and 88%, respectively.

**Conclusion:** 3DTEE provides an alternative specific method for detecting moderate to severe stenosis of proximal CA segments. In addition, 3DTEE can provide both longitudinal and cross-sectional views of the CA and plaques.

### 1090-7 Dynamic Systolic Coronary Blood Flow Reversal in Patients With Hypertrophic Cardiomyopathy: Relationship to Provokable Outflow Tract Obstruction

R.G. Bach, M.J. Kern, T.J. Donohue, T. Wolford, E. Caracciolo. St. Louis University Health Sciences Center, St. Louis, MO, USA

Hypertrophic cardiomyopathy (HCM) is characterized by variable physiologic phenotypes, with a subset of patients showing only provokable left ventricular outflow tract (LVOT) obstruction. Unusual and variably abnormal patterns of resting coronary blood flow have been reported in patients with HCM. The relationship between dynamic LVOT obstruction and phasic coronary blood flow has not been examined. Therefore, we studied coronary flow velocity (0.014" Doppler guidewire) in 13 arteries (7 LAD, 4 CX, 2 RCA) in 8 patients with HCM before and during Valsalva maneuver at catheterization. Six patients had no resting systolic LVOT gradient (mean resting gradient =

5  $\pm$  14, range = 0-50 mmHg). At peak Valsalva strain, the LVOT systolic gradient increased to 56  $\pm$  23 mmHg (range 18-100 mmHg). Phasic coronary flow velocity pattern was normal in all arteries at rest. At peak Valsalva, there was reversal of systolic flow and a significant decrease in diastolic flow velocity in all arteries (see Table;  $\Delta$ P = gradient, mmHg; SPV = systolic peak velocity, cm/s; DPV = diastolic peak velocity, cm/s; SPVI = SPV integral, units; DPVI = DPV integral, units; values are mean  $\pm$  SD).

	LVOT $\Delta$ P	SPV	SPVI	DPV	DPVI
Rest	5 $\pm$ 14	25 $\pm$ 17	8 $\pm$ 4	40 $\pm$ 15	18 $\pm$ 6
Valsalva	56 $\pm$ 23	-24 $\pm$ 13	-3 $\pm$ 1	34 $\pm$ 12	13 $\pm$ 5
p	<0.0001	<0.0001	<0.0001	0.05	0.01

Systolic coronary flow velocity reversal was related to the dynamic LVOT gradient with Valsalva (r = 0.65, p = 0.01).

**Conclusion:** Dynamic pathophysiologic alterations in left ventricular hemodynamics seen in HCM are associated with dynamic changes in coronary blood flow.

### 1090-8 Injection Rate Minimally Affects the TIMI Frame Count

T. Dodge, M. Rizzo, W. Daley, M. Nykiel, J. Altmann, K. Hokirk, M. Brennan, J. Larson, M. Gibson. Brigham and Women's Hospital & West Roxbury Veterans Affairs Medical Center, Boston, MA, USA

**Background:** The Corrected TIMI Frame Count (CTFC) is a clinically important index of coronary artery blood flow. The goal of this study was to determine whether injection rate affects CTFC.

**Methods:** Dye injection rates were recorded during 361 hand injections to characterize the 10th, 50th, and 90th %-tiles for human angiographers. 273 ECG-gated power injections were then performed using a MedRad V power injector to deliver dye randomized to these 10th, 50th, or 90th %-tiles. The injection rate group and actual injection rate delivered (ml/sec) were recorded. Cineangiograms were analyzed for CTFC, proximal normal reference lumen diameter, and the presence of >50% stenoses (disease).

**Results:** In univariate analyses, CTFC was significantly affected by diameter (p < 0.0001), and disease (p < 0.001), but not injection rate group (p = 0.43) or actual injection rate (p = 0.94). Multivariate analysis revealed: CTFC = 20.9 frames + 4.1 frames x diameter (mm), (p < 0.001) + 4.6 frames if diseased, (p < 0.001) - 1.6 frames if 50th %-tile injection rate, (p = 0.31) - 1.9 frames if 90th %-tile injection rate, (p = 0.23) (p-values for 50th and 90th %-tiles represent comparison with 10th %-tile injection rate group, the dummy variable) (p < 0.0001 for model, r = 0.39).

**Conclusion:** Proximal lumen diameter and presence of >50% stenoses independently and significantly affect the Corrected TIMI Frame Count. Injection technique minimally impacts the Corrected TIMI Frame Count: an increase in injection rate from the 10th to 90th %-tile tends to decrease the CTFC by less than two frames.

### 1090-9 Coronary Flow Reserve in the Contralateral Artery Increases After Successful Coronary Angioplasty

Z.S. Kyriakides, A. Antoniadis, T.M. Kolettis, D.Th. Kremastinos. 2<sup>nd</sup> Dept. of Cardiology, Onassis Cardiac Surgery Center, Athens, Greece

**Background:** Coronary flow reserve (CFR) is reduced in remote regions supplied by angiographically normal arteries. Recent studies have found that myocardial flow reserve is impaired in the myocardium in the perfusion area of a vessel with non-critical stenosis supplying collaterals to other segments. We, therefore, tested the hypothesis that CFR in the grossly normal contralateral artery could increase after successful angioplasty (PTCA).

**Methods:** We studied 31 patients (pts), 55  $\pm$  2 years old, having stable angina and single artery disease, undergoing PTCA of the right coronary or the left anterior descending artery. CFR of the diseased and the contralateral artery was estimated using a Doppler flow guidewire, by giving ic adenosine, before and 15 min after the end of PTCA. All the procedures were successful and in 19 pts intracoronary stent was placed.

**Results:** (mean  $\pm$  standard error)

	Before PTCA	After PTCA
Blood pressure (mmHg)	98 $\pm$ 3	95 $\pm$ 3
Heart rate (beats/min)	74 $\pm$ 1	76 $\pm$ 2
CFR of the contralateral artery	2.4 $\pm$ 0.1	2.8 $\pm$ 0.1*
Baseline APV (cm/sec)	21 $\pm$ 2	12 $\pm$ 1*
Maximum APV (cm/sec)	47 $\pm$ 3	34 $\pm$ 3*

\* p < 0.05; APV: average peak velocity;

The contralateral CFR after PTCA increased by 0.8  $\pm$  0.1 in 11 pts with visible collaterals before PTCA and by 0.3  $\pm$  0.1 in the remaining pts (p < 0.05). CFR of the diseased artery after PTCA was 2.2  $\pm$  0.1.

**Conclusion:** CFR in the contralateral artery increases after successful PTCA. This increase is more exaggerated in pts with more well developed collaterals.

**1090-10 Clinical and Angiographic Predictors of New Total Occlusions in Patients Awaiting Coronary Angioplasty**

C.G. de Cock, A.J. vd Berg, N. vd Nieuwenhuizen, J.G. Bronzwaer, C.A. Visser. *Free University Hospital, Amsterdam, The Netherlands*

**Background:** To study predictive indices to identify patients (pts) awaiting coronary angioplasty who progress to total occlusion, 84 consecutive (pts) who developed new total occlusion were compared with 158 pts with similar luminal diameter stenoses on quantitative coronary angiography (CAAS). The time interval between the 2 angiograms was not significantly different (7 ± 0.8 weeks vs 7 ± 0.5 weeks).

**Results:**

	Total occlusion (n = 84)	No total occlusion (n = 158)	p
Age	57 ± 10	58 ± 9	ns
Men	68%	79%	0.01
Smoking	62%	59%	ns
Diabetes mellitus	15%	11%	ns
Angina class ≥ III	32%	38%	ns
Complex lesion	61%	38%	0.005
Long > 10 mm	37%	11%	0.005
Diameter Stenosis	64 ± 10	67 ± 9	ns
MLD	1.06 ± 0.31	1.07 ± 0.32	ns

MLD = minimal luminal diameter

**Conclusion:** Female pts with long complex lesions should be considered at high risk for subsequent total occlusions and should therefore be candidates for early angioplasty.

**1091 Coronary Collaterals**

Tuesday, March 31, 1998, 9:00 a.m.–11:00 a.m.  
Georgia World Congress Center, West Exhibit Hall Level  
Presentation Hour: 9:00 a.m.–10:00 a.m.

**1091-11 Recruited Collateral Channels Are Not Responsible for Ischemic Preconditioning During Single Vessel Angioplasty**

M.M. Mason, D.J. Patel, N.S. Jeyson, S. Brant, V. Paul, C.D.J. Ilesley. *Harefield Hospital, Harefield, Middlesex, UK*

**Background:** Collateral channel recruitment during balloon occlusion varies between patients, but their timing and contribution to myocardial protection in this setting are not well established. This study assessed the degree and timing of recruitment of collaterals and their possible functional contribution during single vessel angioplasty in patients with normal LV function and no spontaneous collaterals.

**Methods:** Patients (n = 17) had four 90s balloon inflations performed. Collaterals were assessed by contralateral injection at 30, 60 and 90s into each inflation and graded using 'Rentrop' criteria, and by 0.014" doppler guide wire across the lesion.

**Results:** Collaterals were recruited in 12/17 patients (7 grade I, 3 grade II and 2 grade III). This was maximal by 30s of inflation (1) in 11 and by 60s in 1 patient, with no progressive recruitment during inflations (2)–(4). Only 4 patients had significant doppler flow during balloon occlusion (all 4 had recruitable collaterals - two grade I; one grade II, one grade III), though all patients had good antegrade flow pre- and post-inflation. Average (APV), integral (PVI) and maximum peak velocity (MPV) during inflations expressed as a percentage of their baseline value (%BF), were compared. No differences were observed during successive inflations-see table (mean (SD)), or comparing those with grade 0/I and grade II/III collaterals.

	30s inf (1)	90s inf (1)	90s inf (4)	p
APV	0.14 (0.13)	0.13 (0.17)	0.13 (0.13)	ns
PVI	0.13 (0.17)	0.11 (0.15)	0.12 (0.14)	ns
MPV	0.36 (0.24)	0.34 (0.23)	0.32 (0.19)	ns

**Conclusions:** Collateralisation varies between patients, but is maximal early in the first inflation. The low level of measurable Doppler flow, together with the lack of incremental recruitment of collaterals, suggests these channels do not make a major contribution to myocardial protection in this setting.


**1091-12 Chronic Physical Exercise and the Quantitatively Assessed Human Coronary Collateral Circulation**

C. Seiler, S. Senti, M. Fleisch, B. Meier. *University Hospital, Cardiology, Bern, Switzerland*

**Background:** There have been no studies in humans assessing the effect of chronic physical exercise on quantitative, directly determined measures of the coronary collateral circulation. This cross-sectional study tested the hypothesis that the degree of exercise (Ex) is directly related to an intracoronary (ic) flow velocity-based index for collateral flow ( $V_{ic,occl}/V_{ic,pre-occl}$ )

**Methods:** In 84 patients (pts, age 57 ± 9 years, yrs) with a coronary artery stenosis to be dilated, a simple, structured interview for the determination of the level of chronic (≥ 1 yr) physical activity (score 1–4) during leisure time and during work hours was performed (J Chron Dis 1986; 18: 527-). The validated collateral flow index:  $V_{ic,occl}/V_{ic,pre-occl}$  (Eur Heart J 1997; 18: 239), was determined using an ic 0.014" Doppler guidewire positioned distal to the stenosis to be dilated.  $V_{ic,occl}/V_{ic,pre-occl}$  was calculated as the flow velocity time integral distal to the stenosis during ( $V_{ic,pre-occl}$ , cm) divided by that obtained at the same location after ( $V_{ic,occl}$ , cm) balloon occlusion (post-hyperemia).

**Results:**



**Conclusions:** 1) In humans, the degree of chronic physical activity during leisure time but not during work hours is directly associated with a quantitative index of coronary collateral flow to a vascular area in need (2) Corroboration of a causal relation among the two factors requires a longitudinal study, and a more refined, objective measure of physical activity (i.e. physical fitness).

**1091-13 Asymmetrical Effects of Angiographically Assessed Collateral Flow on Vasodilator and Exercise Stress-induced Ischemia**

C. Lu, A. D'Alonte, M. Marzilli, M. De Nes, Y. Wang, P. Marraccini, A. L'Abbate. *Institute of Clinical Physiology, CNR, Pisa, Italy*

**Background:** isolated, totally occluded coronary vessel subtending a non-interacted, entirely collateral filling-dependent myocardial region provides a unique opportunity to assess the role of angiographically assessed coronary collateral circulation (ACC) in modulating myocardial vulnerability to pharmacological or physiologic stress testing.

**Objective:** To assess whether a high-grade ACC has similar protective role in induced collateral-dependent ischemia during coronary vasodilator and exercise stress tests.

**Methods and Results:** The ischemia in collateral-dependent myocardium was determined by analysis of dipyridamole echocardiography test (0.84 mg/kg over 10', DET) and bicycle exercise ECG test (EET) data from 53 infarction patients with isolated coronary occlusion, and no significant stenosis in other coronary arteries. ACC was scored from 0: absent to 3: complete and rapid filling of distal portion. There were 34 patients with high-grade collateral flow (score ≥ 2, Group I) and 19 patients with poor or absent collateral flow (score < 2, Group II). During the stress tests, ischemia in the collateral-dependent myocardium was detected by DET in 91% patients in Group I and 74% in Group II (p = 0.09). EET could be completed in 32 patients of Group I and in all patients of Group II. EET was positive in 69% of Group I and in 68% of Group II patients (p = ns). Ischemia detected in Group I was higher by DET than by EET (91 vs 69%, p < 0.05) and did not differ in Group II patients (74 vs 68%, p = ns)

**Conclusions:** ACC has asymmetrical effects on induced ischemia depending on the employed stress. Collateral-dependent ischemia is more frequent during dipyridamole vasodilator stimulation than during physiologic exercise testing in patients with high-grade collateral flow.