

- Shprintzen RJ, Goldberg RB, Lewin ML, Sidoti EJ, Berkman MD, Argamaso RV, Young D (1978) A new syndrome involving cleft palate, cardiac anomalies, typical facies, and learning disabilities: velo-cardio-facial syndrome. *Cleft Palate J* 15:56–62
- Shprintzen RJ, Goldberg RB, Young D, Wolford L (1981) The velo-cardio-facial syndrome: a clinical and genetic analysis. *Pediatrics* 67:167–172
- Tezenas Du Montcel S, Mendizabal H, Ayme S, Levy A, Philip N (1996) Prevalence of 22q11 microdeletion. *J Med Genet* 33:719
- Wilson DI, Britton SB, McKeown C, Kelly D, Cross IE, Strobel S, Scambler PJ (1993) Noonan's and DiGeorge syndrome with monosomy 22q11. *Arch Dis Child* 68:187–189
- Wilson DI, Cross IE, Wren C, Scambler PJ, Burn J, Goodship J (1994) Minimum prevalence of chromosome 22q11 deletions. *Am J Hum Genet Suppl* 55:A169

Address for correspondence and reprints: Dr. Anita Rauch, Institut für Humangenetik der Friedrich-Alexander-Universität Erlangen-Nürnberg, Schwabachanlage 10, D-91054 Erlangen, Germany. E-mail: arauch@humgenet.uni-erlangen.de

© 1999 by The American Society of Human Genetics. All rights reserved. 0002-9297/99/6402-0040\$02.00

Am. J. Hum. Genet. 64:667–668, 1999

RB1 Gene Mutations in Peripheral Blood DNA of Patients with Isolated Unilateral Retinoblastoma

To the Editor:

Two recent reports in this *Journal* (Lohmann et al. 1997; Sippel et al. 1998) indicated that the proportion of patients with isolated unilateral retinoblastoma who carry RB1 gene mutations in constitutional cells is higher than estimated previously (Vogel 1979; Draper et al. 1992). Mutation analysis in patients with unilateral tumors is important because it helps to significantly reduce the number of infant relatives who require clinical surveillance for retinoblastoma (Gallie 1997). Moreover, molecular investigation of these patients can identify carriers of mutations associated with incomplete penetrance and reduced expressivity and thus can extend our knowledge of the genotype-phenotype correlation (Gallie 1997; Lohmann et al. 1997). We have now analyzed additional tumors and have found that the frequency of constitutional mutations in patients with isolated unilateral retinoblastoma is not as high as indicated by our previous study (Lohmann et al. 1997).

Forty-two retinoblastomas that showed loss of constitutional heterozygosity (LOH) at the intragenic loci RBi2 (Toguchida et al. 1993) or RB1.20 (Yandell et al. 1989) were available for mutation analysis. Twenty-one of these tumors had been part of a previous study but

were not analyzed for small mutations at that time (Lohmann et al. 1997). We analyzed the methylation status at the 5' end of the RB1 gene by Southern blot analysis, using the methylation-sensitive enzymes *Bss*HII and *Sac*II as described elsewhere (Greger et al. 1994). Hypermethylation was identified in tumors from six patients. We performed SSCP to screen for small mutations, using a method reported elsewhere (Lohmann et al. 1996). Single base substitutions, including 17 transitions at CpG-dinucleotides, and small length alterations were identified in 24 and 3 tumors, respectively. To identify mutations in the remaining 10 tumors, we sequenced all 27 exons and the promoter region of the RB1 gene. However, no small mutation was identified.

In all, mutations were identified in tumors from 32 (76%) of 42 patients (RB1 gene mutation database). In the tumor of one patient (M6485), a missense base change (c.929G→A, E310G) in exon 9 and a nonsense mutation in exon 15 (c.1399C→T, R467X) were identified in addition to LOH. The missense base change was also present in peripheral blood DNA of this patient. Further investigation showed that this variant RB1 allele was inherited from the father and is, at least, carried by four adult relatives who are unaffected by retinoblastoma. The sequence variant, which, to our knowledge, has not been reported before, is expected to alter an amino acid located N-terminal of the pocket domains A and B (Hu et al. 1990). Only a few reported missense mutations with putative oncogenic effect are located outside the regions that code for these domains (RB1 gene mutation database). Considering that the tumor of patient M6485 also shows a somatic nonsense mutation and LOH, the missense base change is probably a neutral polymorphism and has not contributed to tumorigenesis. However, detailed analyses are required, to demonstrate that this sequence variant does not change the functional properties of the Rb protein (Bremner et al. 1997; Otterson et al. 1997).

In our previous study, we detected small RB1 gene mutations in leukocyte DNA from 6 (17%) of 36 patients with isolated unilateral tumors (Lohmann et al. 1997). In the present study, none of the bona fide oncogenic mutations identified in tumors was also detected in corresponding peripheral blood DNA by direct sequencing of PCR products. Therefore, when the data presented here are included, the proportion of patients with mutations in leukocyte DNA drops to 6 (9%) of 68. Because of mutational mosaicism (Lohmann et al. 1997; Sippel et al. 1998), this figure underestimates the true prevalence of constitutional RB1 gene mutations in patients with isolated unilateral retinoblastoma. However, in almost all patients with isolated unilateral retinoblastoma who have affected children, the mutation is readily detectable in peripheral blood DNA (Sippel et al. 1998; authors' unpublished data). It is reasonable to

assume, therefore, that the proportion of unilaterally affected patients with mutations in leukocyte DNA approximates the prevalence of hereditary retinoblastoma among patients with isolated unilateral tumors. The percentage obtained from our conjoint studies is now in accord with a previous estimate by Vogel (1979) that was based on epidemiological data.

Acknowledgments

We thank N. Bornfeld, B. Lorenz, U. Oehlschläger, and S. Schüler, for referral of patients, and E. Passarge for continuous support.

MARTINA KLUTZ, BERNHARD HORSTHEMKE, AND
DIETMAR R. LOHMANN

*Institut für Humangenetik,
Universitätsklinikum Essen,
Essen, Germany*

Electronic-Database Information

URL for data in this article is as follows:

RB1 gene mutation database, <http://home.kamp.net/home/dr.lohmann>

References

- Bremner R, Du DC, Connolly-Wilson MJ, Bridge P, Ahmad KF, Mostachfi H, Rushlow D, et al (1997) Deletion of RB exons 24 and 25 causes low-penetrance retinoblastoma. *Am J Hum Genet* 61:556–70
- Draper GJ, Sanders BM, Brownbill PA, Hawkins MM (1992) Patterns of risk of hereditary retinoblastoma and applications to genetic counselling. *Br J Cancer* 66:211–219
- Gallie BL (1997) Predictive testing for retinoblastoma comes of age. *Am J Hum Genet* 61:279–281
- Greger V, Debus N, Lohmann D, Höpping W, Passarge E, Horsthemke B (1994) Frequency and parental origin of hypermethylated RB1 alleles in retinoblastoma. *Hum Genet* 94:491–496
- Hu QJ, Dyson N, Harlow E (1990) The regions of the retinoblastoma protein needed for binding to adenovirus E1A or SV40 large T antigen are common sites for mutations. *EMBO J* 9:1147–1155
- Lohmann DR, Brandt B, Höpping W, Passarge E, Horsthemke B (1996) Spectrum of RB1 germ-line mutations in hereditary retinoblastoma. *Am J Hum Genet* 58:940–949
- Lohmann DR, Gerick M, Brandt B, Oehlschläger U, Lorenz B, Passarge E, Horsthemke B (1997) Constitutional RB1-gene mutations in patients with isolated unilateral retinoblastoma. *Am J Hum Genet* 61:282–294
- Otterson GA, Chen WD, Coxon AB, Khleif SN, Kaye FJ (1997) Incomplete penetrance of familial retinoblastoma linked to germ-line mutations that result in partial loss of RB function. *Proc Natl Acad Sci USA* 94:12036–12040
- Sippel KC, Fraioli RE, Smith GD, Schalkoff ME, Sutherland J, Gallie BL, Dryja TP (1998) Frequency of somatic and germ-line mosaicism in retinoblastoma: implications for genetic counseling. *Am J Hum Genet* 62:610–619
- Toguchida J, McGee TL, Paterson JC, Eagle JR, Tucker S, Yandell DW, Dryja TP (1993) Complete genomic sequence of the human retinoblastoma susceptibility gene. *Genomics* 17:535–543
- Vogel F (1979) Genetics of retinoblastoma. *Hum Genet* 52:1–54
- Yandell DW, Campbell TA, Dayton SH, Petersen R, Walton D, Little JB, McConkie-Rosell A, et al (1989) Oncogenic point mutations in the human retinoblastoma gene: their application to genetic counseling. *N Engl J Med* 321:1689–1695

Address for correspondence and reprints: Dr. Dietmar R. Lohmann, Institut für Humangenetik, Hufelandstrasse 55, D-45122 Essen, Germany. E-mail: dr.lohmann@uni-essen.de

© 1999 by The American Society of Human Genetics. All rights reserved.
0002-9297/99/6402-0041\$02.00

Am. J. Hum. Genet. 64:668–669, 1999

TDT Clarification

To the Editor:

A potentially misleading statement occurs in the invited editorial “The TDT and other family-based tests for linkage disequilibrium and association” by Spielman and Ewens (59:983–989), published in the November 1996 issue of the *Journal*. We wish to make the following clarification.

In discussing some issues that arise when the transmission/disequilibrium test (TDT) is used in families where genotype data are unavailable for one parent, we noted the finding of Curtis and Sham (1995) that, when there is a single affected offspring who is homozygous for an allele present in the available (heterozygous) parent, the TDT gives a biased result and should not be used. We then stated that “[w]hen there is more than one offspring in the sibship, it sometimes will be possible to deduce that the unavailable parent [is also heterozygous], and, in these cases, we may proceed as though this [reconstructed] genotype were known directly” (Spielman and Ewens 1996, p. 987).

We should have emphasized that this claim assumes that the reconstruction is done from the genotypes of unaffected offspring. A bias will usually arise in the TDT statistic if the reconstruction uses, in whole or in part, genotype data from affected offspring whose genotypes are then used in the TDT. A bias can also arise when both parental genotypes are reconstructed from the genotypes of the offspring.

The bias resulting from reconstruction occurs for a reason different from that noted by Curtis and Sham