

## REFERENCES

1. Miyamoto T, Neuman Y, Luo H, et al. Coronary vasodilation by noninvasive transcutaneous ultrasound. *J Am Coll Cardiol* 2003;41:1623–7.
2. Kipshidze NN, Chapidze G, Shperling LV, et al. USSR patent no. 1172554: method for the treatment of patients with acute myocardial ischemia. January 18, 1982.
3. McPherson DD, Holland CK. Seizing the science of ultrasound: beyond imaging and into physiology and therapeutics. *J Am Coll Cardiol* 2003;41:1628–30.
4. Kipshidze N, Keelan MH, Petersen JR, et al. Photoactivation of vascular iNOS and elevation of cGMP in vivo: possible mechanism for photovaso-relaxation and inhibition of restenosis in an atherosclerotic rabbit model. *Photochem Photobiol* 2000;72:579–82.
5. Kipshidze N, Nikolaychik V, Keelan MH, et al. Low-power helium: neon laser irradiation enhances production of vascular endothelial growth factor and promotes growth of endothelial cells in vitro. *Lasers Surg Med* 2001;28:355–64.

## REPLY

We believe the potential benefits of ultrasound include enhancement of thrombolysis, vasodilation, and increased tissue perfusion. Transcutaneous ultrasound in animals facilitates thrombolysis (1). A pilot trial of transcutaneous low-frequency ultrasound, in combination with tissue-type plasminogen activator (t-PA), has shown feasibility and safety to treat patients with acute myocardial infarction (2).

The study by Miyamoto et al. (3) found that the coronary arteries as well as the coronary veins dilate in response to transthoracic low-frequency ultrasound. Transcutaneous low-frequency ultrasound augments ST-segment elevation resolution in dogs with coronary occlusion treated with t-PA (4). Suchkova et al. (5) found that ultrasound improves tissue perfusion in ischemic limbs, which is associated with capillary dilation. Pretreatment with an inhibitor of nitric oxide synthase blocked ultrasound enhancement of tissue perfusion and capillary dilation. These findings indicate that the effects of ultrasound are mediated through the nitric oxide. Based on this work, it is likely that ultrasound promotes mechanical shear on the endothelial cells and, thus, release of nitric oxide. This thesis is in agreement with that of Drs. Kipshidze.

**Robert J. Siegel, MD, FACC**  
Cedars-Sinai Medical Center  
Division of Cardiology  
8700 Beverly Boulevard  
Room 5335  
Los Angeles, CA 90048  
E-mail: siegel@cshs.org

**Huai Luo, MD**  
**Yochai Birnbaum, MD, FACC**  
**Michael C. Fishbein, MD, FACC**  
**Takashi Miyamoto, MD**

doi:10.1016/j.jacc.2004.04.008

## REFERENCES

1. Siegel RJ, Atar S, Fishbein MC, et al. Noninvasive, transthoracic, low-frequency ultrasound augments thrombolysis in a canine model of acute myocardial infarction. *Circulation* 2000;101:2026–9.

2. Cohen MG, Tuero E, Blughermann J, et al. Transcutaneous ultrasound facilitated coronary thrombolysis during acute myocardial infarction. *Am J Cardiol* 2003;92:454–7.
3. Miyamoto T, Neuman Y, Luo H, et al. Coronary vasodilation by noninvasive transcutaneous ultrasound. *J Am Coll Cardiol* 2003;41:1623–7.
4. Birnbaum Y, Luo H, Atar S, et al. Noninvasive transthoracic low frequency ultrasound augments thrombolysis in a canine model of acute myocardial infarction—evaluation of the extent of ST-segment resolution. *J Thromb Thrombolysis* 2001;11:229–34.
5. Suchkova VN, Baggs RB, Sahni SK, Francis CW. Ultrasound improves tissue perfusion in ischemic tissue through a nitric oxide dependent mechanism. *Thromb Haemost* 2002;88:865–70.

## Transient Ischemic Dilation

Abidov et al. (1) note an increased cardiac event rate in follow-up of patients found to have transient ischemic dilation (TID) as the sole abnormality on stress myocardial perfusion single-photon emission computed tomography (SPECT). However, the finding of severe coronary disease in only a minority (5 of 20) of such patients is in contrast to previous reports (2,3), which conclude TID to be highly specific for severe coronary disease, and it suggests that there are alternate or additional mechanisms to that of balanced diffuse endocardial ischemia due to epicardial coronary stenosis.

Earlier experimental studies have demonstrated a decreased endocardial-to-epicardial flow ratio in response to adenosine (4,5), tachycardia (4), and hypotension (6), particularly in the presence of left ventricular hypertrophy (LVH) and elevated left ventricular end-diastolic pressure (LVEDP) (4–6), as well as coronary stenosis. In addition, redistribution of transmural perfusion would be expected to produce a more pronounced increase in the TID ratio in the presence of LVH, due to a greater subendocardial-to-subepicardial distance. One recent clinical report (7) associates TID, in the absence of focal perfusion defect, with hypertensive LVH.

The possibility that TID may represent a nonspecific marker of stress-induced subendocardial underperfusion by one or more of a number of possible mechanisms and may thus serve as a surrogate of various risk factors for a future cardiac event should be considered. As this phenomenon may not truly be “ischemic” nor represent genuine left ventricular dilation, the term “transient ischemic dilation” may be a misnomer.

**Charles J. Hardebeck, MD, FACC**  
Division of Cardiology  
University of Kentucky  
210 Thomas More Park  
Crestview Hills, KY 41017  
E-mail: chardebeck@hotmail.com

doi:10.1016/j.jacc.2004.04.003

## REFERENCES

1. Abidov A, Bax JJ, Hayes SW, et al. Transient ischemic dilation ratio of the left ventricle is a significant predictor of future cardiac events in patients with otherwise normal myocardial perfusion SPECT. *J Am Coll Cardiol* 2003;42:1818–25.
2. Weiss AT, Berman DS, Lew AS, et al. Transient ischemic dilation of the left ventricle on stress thallium-201 scintigraphy: a marker of severe and extensive coronary artery disease. *J Am Coll Cardiol* 1987;9:752–9.

3. Mazzanti M, Germano G, Kiat H, et al. Identification of severe and extensive coronary artery disease by automatic measurements of transient ischemic dilation of the left ventricle in dual-isotope myocardial perfusion SPECT. *J Am Coll Cardiol* 1996;27:1612–20.
4. Duncker DJ, Ishibashi Y, Bache RJ. Effect of treadmill exercise on transmural distribution of blood flow in hypertrophied left ventricle. *Am J Physiol* 1998;275:H1274–82.
5. Hittinger L, Mirsky I, Shen Y, et al. Hemodynamic mechanisms responsible for reduced subendocardial coronary reserve in dogs with severe left ventricular hypertrophy. *Circulation* 1995;92:978–86.
6. Harrison DG, Florentine MS, Brooks LA, et al. The effect of hypertension and left ventricular hypertrophy on the lower range of coronary autoregulation. *Circulation* 1988;77:1108–15.
7. Robinson VJ, Corley JH, Merks DS, et al. Causes of transient dilatation of the left ventricle during myocardial perfusion imaging. *AJR Am J Roentgenol* 2000;174:1349–52.

## REPLY

We thank Dr. Hardebeck for his comments about our study; his comments provide an opportunity to clarify our concepts regarding the finding of “transient ischemic dilation” or “TID” of the left ventricle.

First, research from our laboratory and others clearly established that the finding of TID is a worrisome finding on myocardial perfusion single-photon emission computed tomography (MPS), and one that is often associated with severe and extensive coronary artery disease (CAD) (1–4). After the initial case report by Stolzenberg (5), our first contribution on this subject with planar thallium studies (1), we reasoned that the finding of TID was likely to represent extensive ischemia, because: 1) it was reversible; 2) the amount of ischemia had to be large enough to cause apparent transient ventricular enlargement even in a planar study; and 3) such ischemia was likely to be severe because it lasted for at least one-half hour after stress, a time at which usual exercise-induced ischemia would be expected to have been resolved. This hypothesis was supported by our observations that TID is predictive of proximal left anterior descending (LAD) coronary artery or multiple vessels with >90% stenosis. That the finding represented ventricular dilation and was not due to extensive subendocardial ischemia was suggested by the fact that the regions of interest were drawn on the epicardial and not the endocardial boundaries (6). It has subsequently been suggested that TID can also be caused by extensive subendocardial ischemia, in the absence of true left ventricular (LV) dilation (2,3). The relative frequencies of these underlying processes are likely to vary with the population being studied. In either case, the finding is suggestive of extensive pathology. The relationship of TID to severe and extensive disease was subsequently investigated by our laboratory with planar dipyridamole thallium studies (7) and then with dual isotope MPS (3), with findings that supported our initial observations.

Second, our current study (8) describes the largest population of patients with perfectly normal rest/stress MPS that has ever been reported. Our primary goal was to evaluate the prognostic and not the diagnostic implications of TID (as measured by the TID ratio) in this population, which is at low risk, but not necessarily at very low risk, of subsequent cardiac events. Our results clearly demonstrated that TID has independent and incremental prognostic value over clinical variables in these patients with perfectly normal MPS after all clinical variables—hemodynamic data, historical variables (including history of hypertension), and resting electro-

cardiogram (ECG) abnormalities—were considered in the multi-variable analysis for predicting future cardiac events.

Third, the principal concern of Dr. Hardebeck is that severe CAD was seen in only a minority (5 of 20) of the patients with TID who were subsequently catheterized. Of importance, only 65 of 1,560 patients were catheterized, indicating that this was a very selected group. Our findings did indicate that severe and extensive CAD was found solely in the patients with TID; however, owing to the very small proportion of our population represented by these patients, it is difficult to draw conclusions from these findings. The diagnostic accuracy of TID is yet to be evaluated in this clinical setting.

Fourth, again, our intent was not to further assess the diagnostic value but rather the prognostic value of the finding of TID with perfectly normal MPS. Dr. Hardebeck’s comments that subendocardial hypoperfusion, left ventricular hypertrophy (LVH), and high left ventricular end-diastolic pressure (LVEDP) might represent a reason for the TID to be valid, but at the same time his comments do not detract from the prognostic value of the finding. All of these conditions are pathological and are by themselves likely to indicate a worse prognostic outcome compared to normal patients. The higher prevalence of TID in these patients (especially in those without apparent perfusion defects) only makes our point stronger.

Finally, we would like to note that the material in Robinson et al. (9) is a pictorial essay and not a clinical report, and that the material cited dealing with LVH represents only preliminary reports from 1997. The authors of these reports should be encouraged to submit full manuscripts for peer review so that their interesting preliminary observations can be more completely evaluated.

### Daniel S. Berman, MD, FACC

Cedars-Sinai Medical Center Imaging  
8700 Beverly Boulevard  
Room 1254  
Los Angeles, CA 90048  
E-mail: daniel.berman@cshs.org

### Guido Germano, PhD, FACC

Aiden Abidov, MD, PhD

doi:10.1016/j.jacc.2004.04.004

## REFERENCES

1. Weiss AT, Berman DS, Lew AS, et al. Transient ischemic dilation of the left ventricle on stress thallium-201 scintigraphy: a marker of severe and extensive coronary artery disease. *J Am Coll Cardiol* 1987;9:752–9.
2. Iskandrian AS, Heo J, Nguyen T, Lyons E, Paugh E. Left ventricular dilatation and pulmonary thallium uptake after single-photon emission computer tomography using thallium-201 during adenosine-induced coronary hyperemia. *Am J Cardiol* 1990;66:807–11.
3. Mazzanti M, Germano G, Kiat H, et al. Identification of severe and extensive coronary artery disease by automatic measurement of transient ischemic dilation of the left ventricle in dual-isotope myocardial perfusion SPECT. *J Am Coll Cardiol* 1996;27:1612–20.
4. Romanens M, Gradel C, Saner H, Pfisterer M. Comparison of 99mTc-sestamibi lung/heart ratio, transient ischaemic dilation and perfusion defect size for the identification of severe and extensive coronary artery disease. *Eur J Nucl Med* 2001;28:907–10.
5. Stolzenberg J. Dilatation of left ventricular cavity on stress thallium scan as an indicator of ischemic disease. *Clin Nucl Med* 1980;5:289–91.
6. Mazzanti M, Germano G, Kiat H, et al. Identification of severe and