



ELSEVIER

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

SCIENCE @ DIRECT®

Journal of Combinatorial Theory, Series A 109 (2005) 339–343

Journal of  
Combinatorial  
Theory

Series A

[www.elsevier.com/locate/jcta](http://www.elsevier.com/locate/jcta)

Note

## Partition analysis and symmetrizing operators

Amy M. Fu<sup>a</sup>, Alain Lascoux<sup>a, b</sup><sup>a</sup>Center for Combinatorics, LPMC, Nankai University, Tianjin 300071, P.R. China<sup>b</sup>CNRS, IGM Université de Marne-la-Vallée, 77454 Marne-la-Vallée Cedex, France

Received 25 November 2003

Available online 12 November 2004

**Abstract**

Using a symmetrizing operator, we give a new expression for the Omega operator used by MacMahon in Partition Analysis, and given a new life by Andrews, Paule and Riese. Our result is stated in terms of Schur functions.

© 2004 Elsevier Inc. All rights reserved.

In his book “Combinatory Analysis”, MacMahon introduced an Omega operator. This operator has been the subject of many recent articles, among which [1–4]. We show in Theorem 4 that the Omega operator can be expressed by a symmetrizing operator, due in fact to Cauchy and Jacobi [6]. As a consequence, we can formulate:

$$\Omega_{\geq} \lambda^k / \prod_{x \in \mathbb{X}} (1 - x\lambda) \prod_{y \in \mathbb{Y}} (1 - \frac{y}{\lambda})$$

in terms of Schur functions of  $\mathbb{X}$  and  $\mathbb{Y}$  (and therefore in terms of the elementary symmetric functions in  $\mathbb{X}$  and  $\mathbb{Y}$ ).

Recall the definitions of MacMahon’s Omega operator  $\Omega_{\geq}$  and of the symmetrizing operator  $\pi_{\omega}$ .

**Definition 1.**

$$\Omega_{\geq} \sum_{s_1=-\infty}^{\infty} \cdots \sum_{s_r=-\infty}^{\infty} A_{s_1, \dots, s_r} \lambda_1^{s_1} \cdots \lambda_r^{s_r} := \sum_{s_1=0}^{\infty} \cdots \sum_{s_r=0}^{\infty} A_{s_1, \dots, s_r},$$

E-mail addresses: [fmfu@eyou.com](mailto:fmfu@eyou.com) (A.M. Fu), [Alain.Lascoux@univ-mlv.fr](mailto:Alain.Lascoux@univ-mlv.fr) (A. Lascoux).

0097-3165/\$ - see front matter © 2004 Elsevier Inc. All rights reserved.

doi:10.1016/j.jcta.2004.08.005

where the domain of the  $A_{s_1, \dots, s_r}$  is the field of rational functions over  $\mathbb{C}$  in several complex variables and the  $\lambda_i$  are restricted to a neighborhood of the circle  $|\lambda_i| = 1$ .

By iteration, it is sufficient to treat the case of one variable  $\lambda$  only .

**Definition 2** (Lascoux [6]). Given  $\mathbb{X} = \{x_1, x_2, \dots, x_n\}$  of cardinality  $\text{Card}(\mathbb{X}) = n$ , the symmetrizing operator  $\pi_\omega$  is defined by:

$$\forall f(x_1, \dots, x_n), \pi_\omega f(x_1, \dots, x_n) = \sum_{\sigma \in \mathfrak{S}(\mathbb{X})} \sigma \left( \frac{f(x_1, \dots, x_n)}{\Delta(\mathbb{X})} x_1^{n-1} \cdots x_n^0 \right),$$

writing  $\Delta(\mathbb{X})$  for the Vandermonde  $\prod_{1 \leq i < j \leq n} (x_i - x_j)$ , the sum being over all permutations  $\sigma$  in the symmetric group  $\mathfrak{S}(\mathbb{X})$ .

Recall that complete symmetric functions  $S^j(\mathbb{X})$  are defined by the generating function:

$$\sum_{j=0}^{\infty} S^j(\mathbb{X}) \lambda^j = \frac{1}{\prod_{i=1}^n (1 - x_i \lambda)}.$$

Complete symmetric functions are compatible with union of alphabets (denoted ‘+’). Given  $\mathbb{Y} = \{y_1, y_2, \dots, y_m\}$ , we have:

$$S^n(\mathbb{X} + \mathbb{Y}) = \sum_{k=0}^n S^k(\mathbb{X}) S^{n-k}(\mathbb{Y}).$$

Schur functions have two classical expressions:

$$S_\mu(\mathbb{X}) = \left| x_i^{\mu_j + j - 1} \right|_{1 \leq i, j \leq n} / \Delta(\mathbb{X}) = \left| S^{\mu_i - i + j}(\mathbb{X}) \right|_{1 \leq i, j \leq n},$$

where  $\mu = [\mu_1, \dots, \mu_n]$  with  $\mu_1 \geq \mu_2 \geq \dots \geq \mu_n \geq 0$ . We denote by  $\mu \rightarrow \mu'$  the conjugation of partitions.

From the definition of  $\pi_\omega$ , we get [6] :

$$\pi_\omega x_1^{\mu_1} \cdots x_n^{\mu_n} = \left| x_i^{\mu_j + j - 1} \right|_{1 \leq i, j \leq n} / \Delta(\mathbb{X}) = S_\mu(\mathbb{X}). \tag{1}$$

This formula is still valid if  $\mu \in \mathbb{Z}^n$ ,  $\mu_1 > -n, \dots, \mu_n > -1$ :

$$\pi_\omega x_1^{\mu_1} \cdots x_n^{\mu_n} = S_\mu(\mathbb{X}), \tag{2}$$

the Schur function  $S_\mu$ , still defined as the determinant  $|S^{\mu_i - i + j}|_{1 \leq i, j \leq n}$ , being either null or equal to  $\pm$  a Schur function indexed by a partition.

Notice that by convention,  $S_i(\mathbb{X}) = 0, i < 0$ . However,  $S_{-1,2}(\mathbb{X}) = -S_{1,0}(\mathbb{X}) \neq 0$ , and indeed, in Theorem 4, we need to use vector-indexed Schur functions with possibly negative components.

Symmetrizing first in  $x_2, \dots, x_n$ , one also has, with the same hypotheses on  $\mu$ :

$$\pi_\omega x_1^{\mu_1} S_{\mu_2, \dots, \mu_n}(x_2, \dots, x_n) = S_\mu(\mathbb{X}). \tag{3}$$

**Lemma 3.** Given  $\mathbb{X}, \mathbb{Y}$  and  $k$  such that  $0 \leq k < \text{Card}(\mathbb{X})$ , then one has:

$$\pi_\omega \left( \sum_{j=0}^{\infty} x_1^{j-k} S^j(\mathbb{Y}) \right) = \sum_{j=0}^{\infty} S^{j-k}(\mathbb{X}) S^j(\mathbb{Y}). \tag{4}$$

**Proof.** Since powers of  $x_1$  range from  $-k$  to  $\infty$ , we can apply (2):

$$\pi_\omega \left( \sum_{j=0}^{\infty} x_1^{j-k} S^j(\mathbb{Y}) \right) = \sum_{j=0}^{\infty} S_{j-k, 0^{n-1}}(\mathbb{X}) S^j(\mathbb{Y}).$$

The terms such that  $j < k$  are all null, being determinants with two identical rows, and the sum reduces to the expression stated in the lemma.  $\square$

Let us remark that the action of the operator  $\Omega$  relative to  $x_1, \dots, x_n$  can be obtained from the action of the operator  $x_1, \dots, x_{n+r}, r \geq 0$  by specializing  $x_{n+1}, \dots, x_{n+r}$  to 0. Therefore we can suppose that  $n$  is bigger than any given integer. This allows us in the following theorem to suppose that  $n > k$ .

**Theorem 4.** Given two alphabets  $\mathbb{X} = \{x_1, x_2, \dots, x_n\}$  and  $\mathbb{Y} = \{y_1, y_2, \dots, y_m\}$  of cardinality  $n$  and  $m$ , let  $\mathbb{B} = 1 + \mathbb{Y} = \{1\} \cup \mathbb{Y}$ . If  $0 \leq k < n$ , then we have:

$$\begin{aligned} & \Omega \frac{\lambda^k}{(1-x_1\lambda) \cdots (1-x_n\lambda)(1-\frac{y_1}{\lambda}) \cdots (1-\frac{y_m}{\lambda})} \\ & \geq \pi_\omega \sum_{j=0}^{\infty} x_1^{j-k} S^j(\mathbb{B}) = \frac{\sum_{\mu} (-1)^{|\mu|} S_{\mu'}(\mathbb{B}) S_{-k, \mu}(\mathbb{X})}{R(1, \mathbb{X}\mathbb{B})}, \end{aligned} \tag{5}$$

where  $R(1, \mathbb{X}\mathbb{Y})$  is equal to  $\prod_{x \in \mathbb{X}, y \in \mathbb{Y}} (1 - xy)$ , and the sum is over all partitions  $\mu$  (the sum is in fact finite). The vector  $[-k, \mu_1, \dots, \mu_{n-1}]$  is denoted  $-k, \mu$ .

**Proof.** We first recall Cauchy’s formula [7, p. 65]:

$$\begin{aligned} R(1, \mathbb{X}\mathbb{Y}) &= \sum_{\mu} (-1)^{|\mu|} S_{\mu}(\mathbb{X}) S_{\mu'}(\mathbb{Y}), \\ & \geq \sum_{i,j=0}^{\infty} S^i(\mathbb{X}) S^j(\mathbb{Y}) \lambda^{i-j+k} = \Omega \frac{\lambda^k}{(1-x_1\lambda) \cdots (1-x_n\lambda)(1-\frac{y_1}{\lambda}) \cdots (1-\frac{y_m}{\lambda})} \\ & = \sum_{i=0}^{\infty} S^i(\mathbb{X}) \sum_{j=0}^{i+k} S^j(\mathbb{Y}) = \sum_{i=0}^{\infty} S^i(\mathbb{X}) S^{i+k}(\mathbb{B}) \\ & = \sum_{j=0}^{\infty} S^{j-k}(\mathbb{X}) S^j(\mathbb{B}). \end{aligned}$$

On the other hand, Lemma 3 allows us to write this last sum as  $\pi_\omega \left( \sum_{j=0}^{\infty} x_1^{j-k} S^j(\mathbb{B}) \right)$ . We shall now directly compute the action of  $\pi_\omega$  on  $\sum_{j=0}^{\infty} x_1^{j-k} S^j(\mathbb{B})$ , denoting

$$\mathbb{X} \setminus x_1 = \{x_2, \dots, x_n\}.$$

$$\begin{aligned} \pi_\omega \sum_{j=0}^\infty x_1^{j-k} S^j(\mathbb{B}) &= \pi_\omega x_1^{-k} \sum_{j=0}^\infty x_1^j S^j(\mathbb{B}) \\ &= \pi_\omega \frac{x_1^{-k}}{R(1, x_1 \mathbb{B})} = \pi_\omega \frac{x_1^{-k} R(1, (\mathbb{X} \setminus x_1) \mathbb{B})}{R(1, \mathbb{X} \mathbb{B})} \\ &= \frac{\pi_\omega \left( x_1^{-k} \sum_{\mu} (-1)^{|\mu|} S_{\mu'}(\mathbb{B}) S_{\mu}(\mathbb{X} \setminus x_1) \right)}{R(1, \mathbb{X} \mathbb{B})} \\ &= \frac{\sum_{\mu} (-1)^{|\mu|} S_{\mu'}(\mathbb{B}) S_{-k, \mu}(\mathbb{X})}{R(1, \mathbb{X} \mathbb{B})} \end{aligned}$$

and the theorem is proved.  $\square$

The result can be expressed in terms of elementary symmetric functions because  $e_i(\mathbb{B}) = e_i(\mathbb{Y}) + e_{i-1}(\mathbb{Y})$  and Schur functions are determinants in elementary symmetric functions.

Theorem 4 allows us to recover the “fundamental recurrence” given in [4, Theorem 2.1]. Let us remark that a different algorithm is provided in [1].

In [5, Theorem 1.4], Guo-Niu Han expresses the Omega operator in terms of Lagrange interpolation:

$$\Omega \frac{\lambda^k}{A(\lambda)B(\lambda^{-1})} = \sum_{i=1}^n \frac{x_i^{n-1-k}}{(1-x_i)B(x_i) \prod_{j \neq i} (x_i - x_j)}, \tag{6}$$

where

$$A(\lambda) = \prod_{i=1}^n (1 - x_i \lambda), \quad B(\lambda) = \prod_{j=1}^m (1 - y_j \lambda).$$

To relate his result to our expression, let us first recall the definition [6] of the Lagrange operator  $L_{\mathbb{X}}$ .

**Definition 5.**

$$\forall f \in \mathfrak{Sym}(1|n-1), \quad L_{\mathbb{X}} f(x_1, \dots, x_n) = \sum_{x \in \mathbb{X}} \frac{f(x, \mathbb{X} \setminus x)}{R(x, \mathbb{X} \setminus x)},$$

where  $\mathfrak{Sym}(1|n-1)$  is the space of polynomials in  $x_1, x_2, \dots, x_n$ , symmetrical in  $x_2, \dots, x_n$ , and  $R(x, \mathbb{X} \setminus x) = \prod_{x' \in \mathbb{X} \setminus x} (x - x')$ .

We can express the Lagrange operator in terms of  $\pi_\omega$ .

**Lemma 6.**  $\forall f \in \mathfrak{Sym}(1|n-1)$ , we have:

$$\pi_\omega f(x_1, \dots, x_n) = L_{\mathbb{X}} \left( f(x_1, \dots, x_n) x_1^{n-1} \right). \tag{7}$$

**Proof.**  $f(x_1, x_2, \dots, x_n)$  can be written as sums of powers of  $x_1$  [6], with coefficients symmetrical in  $x_1, \dots, x_n$ . Checking

$$L_{\mathbb{X}}(x_1^k x_1^{n-1}) = \pi_{\omega}(x_1^k) = S^k(\mathbb{X}),$$

is immediate.  $\square$

Formula (7) shows that the Lagrange operator in formula (6) can be replaced by  $\pi_{\omega}$ ; therefore [5, Theorem 1.4] is a consequence of Theorem 4.

One does not need to suppose that all the  $x_i$ 's are distinct. Indeed, in a Schur function, one may specialize  $x_1, \dots, x_k$  to the same value  $a$ . This is more of a problem in the Lagrange interpolation formula, where one has to use derivatives of different orders.

Let us finish with a small explicit example, for  $\mathbb{X} = \{x_1, x_2\}$ ,  $\mathbb{Y} = \{y\}$ , and  $k = 1$ .

$$\begin{aligned} \pi_{\omega} \left( \sum_{j=0}^{\infty} x_1^{j-1} S^j(\mathbb{B}) \right) &= \frac{\sum_{\mu} (-1)^{|\mu|} S_{\mu}(\mathbb{B}) S_{-1, \mu}(\mathbb{X})}{R(1, \mathbb{X}\mathbb{B})} \\ &= \frac{-S_1(\mathbb{B}) S_{-1, 1}(\mathbb{X}) + S_{1, 1}(\mathbb{B}) S_{-1, 2}(\mathbb{X})}{R(1, \mathbb{X}\mathbb{B})} \\ &= \frac{(1+y) - y(x_1 + x_2)}{(1-x_1)(1-x_2)(1-x_1y)(1-x_2y)} \\ &= \frac{\Omega}{\geq} \frac{\lambda}{(1-\lambda x_1)(1-\lambda x_2)(1-y/\lambda)}. \end{aligned}$$

### Acknowledgements

This work was done under the auspices of the 973 Project on Mathematical Mechanization of the Ministry of Science and Technology, and the National Science Foundation of China. A. Lascoux was partially supported by the EC's IHRP Program, within the Research Training Network "Algebraic Combinatorics in Europe", Grant HPRN-CT-2001-00272.

### References

[1] G.E. Andrews, MacMahon's Partition Analysis II: fundamental theorems, *Ann. Combin.* 4 (2000) 327–338.  
 [2] G.E. Andrews, P. Paule, A. Riese, MacMahon's Partition Analysis VI: a new reduction algorithm, *Ann. Combin.* 5 (2001) 251–270.  
 [3] G.E. Andrews, P. Paule, A. Riese, MacMahon's Partition Analysis IX:  $k$ -Gon Partitions, *Bull. Austral. Math. Soc.* 64 (2001) 321–329.  
 [4] G.E. Andrews, P. Paule, A. Riese, MacMahon's Analysis: The Omega Package, *Euro. J. Combin.* 22 (2001) 887–904.  
 [5] Guo-Niu Han, A general algorithm for the MacMahon Omega operator, *Ann. Combin.* 7 (2003) 467–480.  
 [6] A. Lascoux, *Symmetric Functions & Combinatorial Operators on Polynomials*, CBMS/AMS Lecture Notes, 2003.  
 [7] I.G. Macdonald, *Symmetric Functions and Hall Polynomials*, second ed., Oxford Mathematical Monographs, Oxford University Press, Oxford, 1995.