




# Revista Brasileira de Farmacognosia

## BRAZILIAN JOURNAL OF PHARMACOGNOSY

[www.journals.elsevier.com/revista-brasileira-de-farmacognosia](http://www.journals.elsevier.com/revista-brasileira-de-farmacognosia)



### Original article

## Pakistamide C, a new sphingolipid from *Abutilon pakistanicum*

Bakhat Ali<sup>a,b</sup>, Muhammad Ibrahim<sup>c,d</sup>, Iqbal Hussain<sup>e,\*</sup>, Nusrat Hussain<sup>d</sup>,  
Muhammad Imran<sup>d</sup>, Haq Nawaz<sup>b</sup>, Saleem Jan<sup>a</sup>, Muhammad Khalid<sup>b</sup>,  
Tahseen Ghous<sup>f</sup>, Muhammad Sajid Hameed Akash<sup>g</sup>

<sup>a</sup>University of Science & Technology, Bannu, Khyber Pakhtunkhawa, Pakistan

<sup>b</sup>Instituto de Química, Universidade de São Paulo, São Paulo, SP, Brazil

<sup>c</sup>Department of Applied Chemistry, Government College University, Faisalabad, Pakistan

<sup>d</sup>H. E. J. Research Institute of Chemistry, International Centre for Chemical and Biological Sciences University of Karachi, Karachi, Pakistan

<sup>e</sup>Department of Botany, Government College University, Faisalabad, Pakistan

<sup>f</sup>Department of Chemistry, A.J.K. University, Muzaffarabad, Pakistan.

<sup>g</sup>Institute of Pharmacology, Toxicology and Biochemical Pharmaceutics, College of Pharmaceutical Sciences, Zhejiang University, Hangzhou, P.R. China.

#### ARTICLE INFO

##### Article history:

Received 11 December 2013

Accepted 30 April 2014

##### Keywords:

*Abutilon pakistanicum*

Malvaceae

Sphingolipid

Pakistamide C

#### ABSTRACT

The *Abutilon* genus from the Malvaceae family is of medicinal importance, and members of this genus are distributed in the tropical and subtropical regions of Asia and other parts of the world. *Abutilon pakistanicum* Jafri & Ali is mainly found in Pakistan. It has been used by different systems of traditional medicines to treat different diseases. Pakistamide C, a new sphingolipid, has been isolated from the ethyl acetate soluble fraction of the methanolic extract of *A. pakistanicum*. Different spectroscopic techniques such as NMR (<sup>1</sup>H, <sup>13</sup>C, COSY, NOESY, HSQC, HMBC) and Mass spectrometry (EI-MS, and FAB-MS experiments) were used to elucidate the structure of pakistamide C.

© 2014 Sociedade Brasileira de Farmacognosia. Published by Elsevier Editora Ltda. All rights reserved.

### Introduction

The genus *Abutilon* consists of about 150 species, which are distributed in the tropical and subtropical regions of Asia and other parts of the world (Nasir and Ali, 1972; Baquar, 1989). The species are distributed among small trees, shrubs and perennial herbs. Generally, *Abutilon* species were used for various disorders, like rheumatism, and used as diuretic and a demulcent, due to their considerable amount of mucilage content (Nasir and Ali, 1972; 1979; Baquar, 1989). *Abutilon*

*pakistanicum* Jafri and Ali, Malvaceae, commonly grows in the southern parts of Pakistan. Different steroids (Ahmed et al., 1990; Hussain et al., 2008a), esters glycosides, flavonoids (Ahmed et al., 1991; Ali et al., 2010b), triterpenes, iridoids (Ali et al., 2010b), and sphingolipids (Ali et al., 2012) have been found in species from genus *Abutilon*. Sphingolipids are known to have a vast range of biological activities (Ahmed et al., 1990; Hussain et al., 2008b). Therefore, the chemotaxonomic and ethnopharmacological importance of this genus prompted us to carry out further phytochemical studies on *Abutilon pakistanicum*.

\* Corresponding author.

E-mail: [driqbal@gcuf.edu.pk](mailto:driqbal@gcuf.edu.pk) (I. Hussain).

0102-695X/\$ - see front matter © 2014 Sociedade Brasileira de Farmacognosia. Published by Elsevier Editora Ltda. All rights reserved.

<http://dx.doi.org/10.1016/j.bjp.2014.07.002>

As a continuation of our research (Ibrahim et al., 2013a; 2013b) on the phytochemical characterization of medicinal plants, we isolated pakistamide C (**1**) from *Abutilon pakistanicum*. The isolation and structural characterization by spectrometric data of this new sphingolipid, pakistamide C are presented here.

## Materials and methods

### Plant material

Whole specimen of *Abutilon pakistanicum* Jafri & Ali, Malvaceae (8 kg) was collected at Karachi, Pakistan in June 2004. Identification of material was done by plant taxonomist, Prof. Dr. Surraiya Khatoon, at the Department of Botany, University of Karachi. A specimen voucher was deposited in the herbarium of the Department of Botany (voucher no. 697 KUH).

### Extraction and isolation

The shade-dried plant material (8 kg) was grounded and soaked in MeOH (3 × 20 l) to get the crude methanolic extract (350 g). The extract (350 g) was then divided into *n*-hexane (65 g), chloroform (75 g), ethyl acetate (35 g), *n*-butanol (90 g), and water (60 g) soluble fractions using solvent-solvent extraction procedure.

### Ethyl acetate fraction (35 g)

Ethyl acetate soluble fraction was subjected to column chromatography (CC) with silica gel as stationary phase and *n*-hexane/AcOEt, AcOEt and AcOEt/MeOH (with polarity increase) as mobile phase. That yielded six major sub-fractions (A, B, C, D, E, F).

### Sub-fraction D

The fraction retrieved from the AcOEt layer (2.1 g), was subjected to chromatography on a silica gel column, and eluted with CHCl<sub>3</sub>-MeOH (8:2).

### Sub-fraction D2

The semipure compound was further purified by preparative TLC using the CHCl<sub>3</sub>-MeOH (8.5:1.5) as eluent to afford compound **1** (14 mg).

### Instruments and materials

The column chromatography (CC) was carried out using silica gel (230-400 mesh, E. Merck, Darmstadt/Germany). Thin layer chromatography (TLC) was performed on silica gel F254 precoated plates (E. Merck, Germany), and UV light detection was done at 254 and 366 nm, while spraying with the ceric sulfate in 10% H<sub>2</sub>SO<sub>4</sub> with gentle heating. Optical rotations were recorded on a Jasco P-2000 polarimeter. The UV spectra Hitachi UV-3200 spectrophotometer were used for the UV spectrum, while IR spectra were taken from the KBr pellets on a Jasco 302-A spectrometer. EI-MS were performed on a Finnigan MAT 312 mass spectrometer, while the FAB-MS were recorded on the JEOL JMS-HX-110 mass spectrometer with glycerol as a matrix. The Bruker AMX-500 spectrometer was used to get the <sup>1</sup>H and <sup>13</sup>C NMR spectra in the solvent C<sub>5</sub>D<sub>5</sub>N. The 2D NMR (COSY, HMQC, HMBC, NOESY) spectra were recorded on the same instrument. Chemical shifts (δ) were taken in ppm with reference to the tetramethylsilane as the internal standard, while the coupling constant (*J*) scalar were reported in Hertz.


## Results and discussion

MeOH extract of *Abutilon pakistanicum* Jafri & Ali, Malvaceae, was suspended in water and fractionated into *n*-hexane, CHCl<sub>3</sub>, AcOEt, *n*-BuOH, and H<sub>2</sub>O fractions. The AcOEt-soluble fraction was subjected to repeated silica gel column chromatography, which yielded compound **1**.

The compound pakistamide C (**1**) was a gummy colorless solid. Absorption peaks for UV spectrum were observed at 203 and 230 nm. The IR showed absorption bands at 3510-3345, 1543, 1640, and 1645 cm<sup>-1</sup> for the characteristic peaks of amine, hydroxyl and olefinic functional groups and secondary amide (Gao et al., 2001).

The molecular formula C<sub>42</sub>H<sub>83</sub>NO<sub>5</sub> was deduced from the EI-MS, FAB-MS, and high-resolution EI-MS spectra with a molecular ion peak at *m/z* 681.6289 (calculated for C<sub>42</sub>H<sub>83</sub>NO<sub>5</sub>, 681.6282). Further peaks appeared at *m/z* 663.6192, 645.6083 and 627.5979, yielded from consecutive water losses (Fig. 1).

The <sup>1</sup>H NMR spectrum showed (Table 1), a doublet at δ 8.48 (d, *J* = 8.9 Hz), assigned to the proton attached to nitrogen of the amide group. The disubstituted double bond signals were



**Figure 1** – Mass fragmentation pattern of **1**.

observed at  $\delta$  5.58 (dt,  $J = 16.0, 6.1$  Hz) and 5.50 (dt,  $J = 16.0, 5.9$  Hz). Protons of upfield signals determined four methylene groups at  $\delta$  1.90-2.24, while the rest of the methylene protons resonated at  $\delta$  1.21-1.33 (br, s,  $28 \times \text{CH}_2$ ). Methyl protons have a triplet at  $\delta$  0.85 (t,  $J = 7.0$  Hz, 6H), which confirms the defined sequence of terminal groups.

The two oxymethylene protons resonated at  $\delta$  4.42 (dd,  $J = 4.4, 10.6$  Hz) and  $\delta$  4.48 (dd,  $J = 7.8, 10.6$  Hz), while oxymethine protons resonated at  $\delta$  4.52 (dd,  $J = 4.0, 7.6$  Hz), 4.32 (dd,  $J = 5.0, 7.1$  Hz) and 4.24-4.26 (1H, m). These signals confirm the basic skeleton to be a sphingolipid (Gao et al., 2001; Muralidhar et al., 2003). The coupling constant of olefinic protons ( $J = 16.0$  Hz) confirmed the trans configuration of the double bond (Jung et al., 1996). The  $^{13}\text{C}$  NMR spectrum (BB and DEPT) (Table 1) showed signal for an amide carbonyl carbon at  $\delta$  175.1, whereas the olefinic methines resonated at  $\delta$  130.8 and  $\delta$  130.4. The four methine carbon signals appeared at  $\delta$  52.8 (CHNH),  $\delta$  77.0 (CHOH),  $\delta$  73.5 (CHOH) and  $\delta$  72.4 (CHOH), while an oxymethylene carbon resonated at  $\delta$  62.5. The aliphatic methylenic carbons resonated in the range of  $\delta$  22.8-35.8, while the two terminal methyl carbons were observed at  $\delta$  15.5.

The  $^1\text{H}$ - $^1\text{H}$  COSY spectrums showed the correlation between the oxygenated methylene hydrogens at  $\delta$  4.48 and 4.42 with the azomethine hydrogen at  $\delta$  5.07-5.09. This relates with the oxymethine proton H-3 at  $\delta$  4.32, which further cross peaks showed another oxymethine proton H-4 at  $\delta$  4.24-4.26 confirming the position of C-3 and C-4 for the two hydroxyl groups, respectively (Fig. 2). The third hydroxyl group position was assigned at C-2', in which H-2' at  $\delta$  4.52 showed HMBC correlation with amide carbonyl carbon ( $\delta$  175.1).

The COSY spectrum showed the correlations of olefinic protons at  $\delta$  5.58 and  $\delta$  5.50 with  $\delta$  2.00-2.02 and 1.90-1.92. Also, H-17 at  $\delta$  5.50 showed correlation with C-15 ( $\delta$  33.2) and C-19 ( $\delta$  22.8) in HMBC spectrum, thus confirming the position of the double bond at C-16. The fragmentation pattern and methanolysis confirmed the position of the double bond and the length of the fatty acid chain.

Methanolysis yielded the fatty acid methyl ester, characterized by mass spectrometry as methyl 2-hydroxydocosanoate ( $m/z$  370  $[\text{M}]^+$ ) (Inagaki et al., 2006). Thus confirming the sphingosine base length composed of 20 carbons with double bond located in the base chain.

The olefinic bond position was long-established by the fragmentation peak in the EI-MS at  $m/z$  613 due to the loss of  $\text{C}_5\text{H}_8$  by McLafferty rearrangement. The peaks at  $m/z$  339 and 342 result from the cleavage of the amide bond, further confirmation for the length of the fatty acid consisting of 20 and 22 carbons, respectively (Fig. 1). The  $^{13}\text{C}$  and  $^1\text{H}$  NMR data along with specific rotation of compound 1 were compared with previously reported data of sphingolipids, which confirmed the stereochemistry of the sphingosine base and the fatty acid. The stereochemistry was assigned to be (2'R) for 2'-hydroxy fatty acid and (2S, 3S, 4R) for phytosphingosines moiety (Loukaci et al., 2000; Zhang et al., 2007; Sandjo et al., 2010).

The stereochemistry at different stereocenters was further confirmed through NOESY spectrum, which showed correlation of the azomethine proton at  $\delta$  5.07-5.09 (H-2) with H-4 at  $\delta$  4.24-4.26 and H-2' at  $\delta$  4.52 (Fig. 3). While H-3 at  $\delta$  4.32 showed through space coupling with the amide proton




Figure 2 – COSY (↔) and HMBC (→) correlations of 1.





Figure 3 – Key NOESY (↔) of 1.

at  $\delta$  8.48 and H-1 $\alpha$  at  $\delta$  4.42. So, on the basis of different spectroscopic data as mentioned above the structure of pakistamide C (1) was unambiguously established as (R)-2-hydroxy-N-((2S,3S,4R,E)-1,3,4-trihydroxyicos-16-en-2-yl) docosanamide.

#### Pakistamide C (1)

(R)-2-hydroxy-N-((2S,3S,4R,E)-1,3,4-trihydroxyicos-16-en-2-yl) docosanamide, colorless gummy solid.  $[\alpha]_D^{25} = +25.0$  ( $c = 0.05$ , MeOH). UV  $\lambda_{\text{max}}$  (MeOH) nm: 202 (1.5), 230 (4.0). IR (KBr)  $\text{cm}^{-1}$ : 3510-3338 (OH/NH), 2920 (CH), 1647, 1545 (HN-C=O), 1636 (C=C); 730 (aliphatic Cs).  $^1\text{H}$  and  $^{13}\text{C}$  NMR: see Table 1. EI-MS:  $m/z$  (%) = 681 (7)  $[\text{M}]^+$ , 663 (10)  $[\text{M}-\text{H}_2\text{O}]^+$ , 645 (10)  $[\text{M}-2\text{H}_2\text{O}]^+$ , 627 (15)  $[\text{M}-3\text{H}_2\text{O}]^+$ , 613 (10), 342 (17), 339 (16), 283 (10), 281 (40), 223 (65), 68 (55), 43 (80). HREI-MS:  $m/z$  681.6266 (calcd. 681.6271 for  $\text{C}_{42}\text{H}_{83}\text{NO}_5$ ,  $[\text{M}]^+$ ). FAB-MS:  $m/z$  682.6338 (calcd. 682.6344 for  $\text{C}_{42}\text{H}_{83}\text{NO}_5 + \text{H}^+$ ).



#### Methanolysis of 1

Compound 1 (2 mg) was dissolved in MeOH (10 ml). This solution was taken in a 50 ml round bottom flask. After the addition of few drops of 5% HCl, the reaction mixture was refluxed for 2 h at 80°C. On completion of the reaction, the

**Table 1**  
 $^1\text{H-NMR}$  (500 MHz) and  $^{13}\text{C-NMR}$  (125 MHz) spectral data of compound **1** recorded in  $\text{C}_5\text{D}_5\text{N}$ .

Position	$^{13}\text{C-NMR}$	$^1\text{H-NMR}$
1	62.5	4.42 (dd, 4.4, 10.6) 4.48 (dd, 7.8, 10.6)
2	52.8	5.07-5.09 (m)
3	77.0	4.32 (dd, 5.0, 7.1)
4	73.5	4.24-4.26 (m)
5	34.2	1.92-1.93 (m) 2.21-2.24 (m)
6-14	29.4 – 30.4	1.21-1.33 (br, s)
15	33.2	2.00-2.02 (m)
16	130.8	5.58 (dt, 6.1, 16.0)
17	130.4	5.50 (dt, 5.9, 16.0)
18	33.5	1.90-1.91 (m)
19	22.8	1.21-1.33 (m)
20	15.5	0.85 (t, 7.0)
NH	-	8.48 (1H, d, $J = 8.9$ )
1'	175.1	-
2'	72.4	4.52 (dd, 4.0, 7.6)
3'	35.8	2.21-2.23 (m) 2.01-2.03 (m)
4'	27.0	1.21-1.33 (br, s)
5' – 20'	29.4 – 30.4	1.21-1.33 (br, s)
21'	22.8	1.21-1.33 (br, s)
Me 22'	15.5	0.85 (t, 7.0)

solution was diluted with water, and extracted with *n*-hexane. The *n*-hexane layer was then separated and concentrated for further analysis using GC-MS, which agrees with the data reported for methyl 2-hydroxydocosanoate ( $m/z$  370  $[\text{M}]^+$ ) [13].

### Authors' contributions

BA, HN, KH, IH, MSHA and MI contributed in collecting the plant sample, running the laboratory work, analysis and drafting of the data. MI, NH and IH contributed to the analysis of the spectroscopic data and writing of the manuscript. BA, TG and MI designed the study, supervised the laboratory work and contributed to critical reading of the manuscript. All the authors have read the final manuscript and approved the submission.

### Conflicts of interest

The authors declare no conflicts of interest.

### Acknowledgements

We are thankful to Higher Education Commission of Pakistan, for financial support. The authors are grateful to H. E. J. Research institute of chemistry, University of Karachi, for providing spectro-support in carrying out NMR studies.

### REFERENCES

- Ahmed, Z., Kazmi, S.N.H., Malik, A., 1990. Pakistanol, a new pentacyclic triterpene from *Abutilon pakistanicum*. *J. Nat. Prod.* 53, 1342-1344.
- Ahmed, Z., Kazmi, S.N.H., Malik, A., 1991. Phytochemical investigation of *Abutilon pakistanicum*. *Fitoterapia* 62, 349-352.
- Ali, B., Imran, M., Hussain, R., Ahmed, Z., Malik, A., 2010a. Structural determination of abutilins A and B, new flavonoids from *Abutilon pakistanicum*, by 1D and 2D NMR spectroscopy. *Mag. Res. Chem.* 48, 159-163.
- Ali, B., Fatima, I., Malik, A., Ahmed, Z., 2010b. New glycosidic constituents of *Abutilon pakistanicum*. *Helv. Chim. Acta.* 93, 2245-2250.
- Ali, B., Mehmood, R., Hussain, R., Malik, A., Imran, M., Nawaz, H., Hussain, A., 2012. New sphingolipids from *Abutilon pakistanicum*. *Naturforsch.* 67b, 433-437.
- Baqar, S.R., 1989. *Medicinal and Poisonous Plants of Pakistan*, Printas, Karachi, p. 506.
- Gao, J.M., Dong, Z.J., Liu, J.K., 2001. A new ceramide from the Basidiomycete *Russula cyanoxantha*. *Lipids* 36, 175-181.
- Hussain, M., Zahra, D.N., Hussain, S.M.S., Ahmed, E., Ahmad, I., Malik, A., Ahmed, Z., 2008a. Structure determination of new steroids from *Abutilon pakistanicum* by NMR techniques. *Mag. Res. Chem.* 46, 274-277.
- Hussain, M., Zahra, D.N., Malik, A., Ejaz, A., Siddiqui, H., Choudhary, M.I., Ahmed, Z., 2008b. Flavonoids C-glycosides from *Abutilon pakistanicum*. *Heterocycles* 75, 645-653.
- Ibrahim, M., Farooq, T., Hussain, N., Hussain, A., Gulzar, T., Hussain, I., Akash, M.S., Rehmani, F.S., 2013b. Acetyl and butyryl cholinesterase inhibitory sesquiterpene lactones from *Amberboa ramosa*. *Chem. Cent. J.* 7, 116, doi: 10.1186/1752-153X-7-116.
- Ibrahim, M., Hussain, I., Imran, M., Hussain, N., Hussain, A., Mahboob, T., 2013a. Corniculatin A, a new flavonoidal glucoside from *Oxalis corniculata*. *Rev. Bras. Farmacogn.* 23, 630-634.
- Inagaki, M., Ikeda, Y., Kawatake, S., Nakamura, K., Tanaka, M., Misawa, E., Yamada, M., Higuchi, R., 2006. Isolation and structure of four new ceramides from the starfish *Luidia maculata*. *Chem. Pharm. Bull.* 54, 1647-1649.
- Jung, J.H., Lee, C.O., Kim, Y.C., Kang, S.S., 1996. New bioactive cerebrosides from *Arisaema amurense*. *J. Nat. Prod.* 59, 319-322.
- Loukaci, A., Bultel-Poncé, V., Longeon, A., Guyot, M., 2000. New lipids from the tunicate *Cystodytes cf. dellechiaiei*, as PLA2 inhibitors. *J. Nat. Prod.* 63, 799-802.
- Muralidhar, P., Radhika, P., Krishna, N., Rao, D.V., Rao, C.B., 2003. Sphingolipids from marine organisms: a review. *J. Nat. Prod. Sci.* 9, 117-142.
- Nasir, E., Ali, S.I., 1972. *Flora of West Pakistan*, Fakhri Printing Press, Karachi, p. 383.
- Nasir, E., Ali, S.I., 1979. *Flora of West Pakistan*, Fakhri Printing Press, Karachi, p. 60.

Sandjo, L.P., Tchoukoua, A., Ntede, H.N., Yemoul, M., Perspicace, E., Keumedjio, F., Couty, F., Kirsch, G., Ngadjui, B.T., 2010. New nortriterpenoid and ceramides from stems and leaves of cultivated *Triumfetta cordifolia* A Rich (Tiliaceae). J. Am. Oil Chem. Soc. 87, 1167-1177.

Zhang, W.K., Xu, J.K., Zhang, X.Q., Yao, X.S., Ye, W.C., 2007. Sphingolipids with neuritogenic activity from *Euphorbia sororia*. Chem. Phys.Lipids 148, 77-83.