

Ribocation Transition State Capture and Rebound in Human Purine Nucleoside Phosphorylase

Mahmoud Ghanem,^{1,2,3,4} Andrew S. Murkin,^{1,3,5} and Vern L. Schramm^{1,*}¹Department of Biochemistry, Albert Einstein College of Medicine, Bronx, NY 10461, USA²Faculty of Veterinary Medicine, Cairo University, Giza, Egypt³These authors contributed equally to this work⁴Present address: Eli Lilly Pharmaceuticals, Inc., Indianapolis, IN 46285, USA⁵Present address: Department of Chemistry, University at Buffalo, SUNY, Buffalo, NY 14260, USA*Correspondence: vern@aecom.yu.edu

DOI 10.1016/j.chembiol.2009.07.012

SUMMARY

Purine nucleoside phosphorylase (PNP) catalyzes the phosphorolysis of 6-oxo-purine nucleosides to the corresponding purine base and α -D-ribose 1-phosphate. Its genetic loss causes a lethal T cell deficiency. The highly reactive ribocation transition state of human PNP is protected from solvent by hydrophobic residues that sequester the catalytic site. The catalytic site was enlarged by replacing individual catalytic site amino acids with glycine. Reactivity of the ribocation transition state was tested for capture by water and other nucleophiles. In the absence of phosphate, inosine is hydrolyzed by native, Y88G, F159G, H257G, and F200G enzymes. Phosphorolysis but not hydrolysis is detected when phosphate is bound. An unprecedented N9-to-N3 isomerization of inosine is catalyzed by H257G and F200G in the presence of phosphate and by all PNPs in the absence of phosphate. These results establish a ribocation lifetime too short to permit capture by water. An enlarged catalytic site permits ribocation formation with relaxed geometric constraints, permitting nucleophilic rebound and N3-inosine isomerization.

INTRODUCTION

Purine nucleoside phosphorylase (PNP) catalyzes the reversible phosphorolysis of 6-oxo-purine (deoxy)ribonucleosides to the corresponding purine base and α -D-ribose 1-phosphate (Figure 1, path b) (Schramm, 2005; Stoeckler, et al., 1978, 1980). This enzyme is a current target for clinical trials involving T cell cancers and autoimmune disease (BioCryst, 2009). Kinetic isotope effects have characterized the reaction catalyzed by human PNP as proceeding via a fully dissociated ribocation transition state (TS) with complete leaving-group departure prior to ribosyl migration to the phosphate nucleophile (Lewandowicz and Schramm, 2004). Despite the high amino acid sequence identity (87%) with human PNP, bovine PNP forms an earlier ribooxacarbenium-ion TS featuring a C1'–N9 bond length of

~ 1.8 Å (Kline and Schramm, 1993). Bovine PNP catalyzes the slow hydrolysis of inosine to hypoxanthine and ribose in the absence of phosphate with a burst of $2 \times 10^{-3} \text{ S}^{-1}$ turnover number of $1.3 \times 10^{-4} \text{ s}^{-1}$ (Figure 1, path c) (Kline and Schramm, 1992, 1995). Inosine hydrolysis has not been previously reported for human PNP.

Crystal structures of human PNP complexed with substrate and TS analogs have defined the active site residues in contact with the purine nucleoside and the phosphate nucleophile (de Azevedo, et al., 2003; Koellner, et al., 1997; Rinaldo-Matthis, et al., 2008; Shi, et al., 2004). Human PNP is a homotrimer, and the catalytic site of PNP is located near the subunit-subunit interfaces. The catalytic site conforms closely to bound inosine with residues N243, E201, H257, F200, Y88, M219, and F159' (from the adjacent subunit) as contacts for the nucleoside. The side chains of N243 and E201 are involved in leaving-group activation by forming hydrogen-bond contacts with the purine base (N7H and N1H, respectively) (Ghanem, et al., 2008b; Nunez, et al., 2006). Residues F159', H257, F200, and Y88 form a hydrophobic cover to the catalytic site and are proposed to shield reactants in the catalytic site from bulk solvent (Figure 2) (Saen-Oon, et al., 2008a). The phenyl group of F200 is perpendicular to the purine base, and residues Y88, F159', H257 and the bound phosphate nucleophile provide contacts to the ribose. Residue F159' from a mobile loop in the adjacent subunit covers the top surface of the ribosyl group of the purine nucleoside (Figure 2). Quantum mechanical simulations and laser-induced temperature-jump studies establish that dynamic motion of this residue is closely associated with TS formation (Ghanem, et al., 2009; Saen-Oon, et al., 2008a). H257 is a key residue in positioning the "oxygen stack" (H257:N δ –O5'–O4'–O ρ) that contributes to catalysis through a vibrational promoting mode of the three oxygen atoms to provide electron density changes to destabilize the ribosyl group, form the carbocation TS and thereby enhance departure of the purine-base leaving group (Murkin, et al., 2007; Saen-Oon, et al., 2008a). The crystal structure of human PNP in complex with a TS analog (Immucillin-H) and phosphate shows structural water molecules near O6 (2.6 Å) and another within 4.4 Å of the ribosyl anomeric carbon (Figure 2), and these structural waters are also found in the catalytic site of bovine PNP (Fedorov, et al., 2001).

We explored two possibilities for water capture of the highly reactive ribocation: (1) water molecules located within the catalytic site prior to ribocation formation, or (2) water molecules that




Figure 1. Reactions Catalyzed by Purine Nucleoside Phosphorylase

Depending on reaction conditions, the ribo-oxacarbenium-ion transition state can partition between (a) return to inosine [1], (b) reaction with phosphate to yield hypoxanthine [2] and α -D-ribose 1-phosphate [4], (c) reaction with water to yield hypoxanthine [2] and D-ribose [3], or (d) reaction with N3 of hypoxanthine to yield N3-inosine isomer [5].

diffuse from bulk solvent into the catalytic site as a consequence of catalytic-site mutations. Water capture of the ribocation in the presence of phosphate would partition the pentose between ribose and α -D-ribose 1-phosphate products. This approach provides a test of ribocation reactivity, its lifetime relative to water diffusion, and allows an estimate of the relative proclivity of the ribocation to react with water, phosphate, the hypoxanthine leaving group or adventitious catalytic site nucleophiles. This work contributes substantially to our understanding of the ribocation transition state.

^{13}C and ^1H nuclear magnetic resonance (NMR) spectroscopy were used to monitor the hydrolysis and phosphorolysis reac-

tions with [$1'$ - ^{13}C]inosine as substrate. Native PNP and catalytic-site glycine mutants of human PNP were analyzed. Four different glycine mutants of residues that are proposed to shield the catalytic site from bulk solvent were engineered in attempts to introduce a solvent leak. Partition of products between D-ribose and α -D-ribose 1-phosphate permit estimation of ribocation lifetime, based on the diffusion rate of water and the distance of the ribocation from bulk solvent in a leaky catalytic site.

RESULTS

Steady-State Kinetic Properties

The steady-state kinetic parameters of four catalytic-site glycine mutants were determined with inosine as substrate and compared with those for the native enzyme. All catalytic-site glycine mutants showed a decrease in their k_{cat} values ranging from 2-fold (Y88G) to 40-fold (F200G) compared with native PNP (see Table S1 available online). The catalytic efficiencies (k_{cat}/K_m) of those mutants were reduced by \sim 8-fold (Y88G) and up to 11,000-fold (F200G) relative to native PNP (Table S1). Purified PNPs showed sufficient catalytic stability in the reaction mixtures at 25°C to permit collection of ^{13}C or ^1H NMR spectra of actively equilibrating enzyme mixtures for 24 hr.

^{13}C and ^1H NMR Spectroscopy and Assignments

The ^{13}C and ^1H NMR spectra of reaction mixtures containing PNPs (native, Y88G, F159G, H257G and F200G) with [$1'$ - ^{13}C]inosine in the presence or absence of phosphate were acquired and compared with the reference spectra of free [$1'$ - ^{13}C]inosine and [$1'$ - ^{13}C]ribose (Figure 3 and Table S2). The ^{13}C NMR spectrum of free [$1'$ - ^{13}C]inosine [1] is a singlet with a chemical shift of 88.68 ppm (Chenon, et al., 1975a; Ebrahimi, et al., 2001).





Figure 2. Views of the Subunit-Subunit Interface of Human PNP in Complex with a TS Analog and Phosphate (PO_4)

(A) PNP catalytic-site contact residues surrounding Immucillin-H (ImmH) (PDB 1RR6). Adjacent subunits in the trimeric protein have been colored yellow and blue.

(B) Contact residues map, with distances in Å units. The two water molecules (W) closest to ImmH have been included.



The downfield region of the ^1H NMR spectrum of free $[1\text{-}^{13}\text{C}]$ inosine shows two singlets for H8 and H2 with chemical shifts of 8.33 and 8.21 ppm, respectively, and a doublet of doublets for H-1' ($J_{1',2'} = 5$ Hz, $J_{\text{H,C}} = 170$ Hz), with a chemical shift of 6.07 ppm (Figure 3A and Table S2). The peak assignments were based on the previously reported chemical shifts for purine nucleosides (Chenon, et al., 1975a, 1975b). The ^{13}C NMR spectrum of $[1\text{-}^{13}\text{C}]$ ribose [3], obtained through the irreversible arsenolysis of $[1\text{-}^{13}\text{C}]$ inosine in the presence of PNP (Kline and Schramm, 1993), shows four singlets with chemical shifts of 93.94, 94.22, 96.68 and 101.35 ppm (Figure 3B and Table S2), corresponding to the four major isomers of D-ribose. The chemical shifts and ratio of the four isomer singlets agree with previously determined values of 95.1 (20%), 95.4 (60%), 97.8 (7%), and 102.5 ppm (13%) for α -D-ribofuranose, β -D-ribofuranose, α -D-ribofuranose and β -D-ribofuranose, respectively (Kenneth et al., 1998; Ryu, et al., 2004). The ^1H NMR spectrum of free hypoxanthine [2] obtained through the irreversible arsenolysis of labeled inosine shows two downfield singlets for H-8 and H-2, with chemical shifts of 8.19 and 8.17 ppm, respectively (Figure 3B).

Phosphate-free PNP Reactions and Inosine Isomerization

In the absence of phosphate, the time course (to 24 hr) ^{13}C NMR spectra of the inosine reaction mixtures for F159G, Y88G, H257G and F200G showed no spectral changes (data not shown). Native PNP with $[1\text{-}^{13}\text{C}]$ inosine showed the formation of a new singlet with a chemical shift of 90.59 ppm (Figure 4-I).

Figure 3. ^{13}C NMR (I) and ^1H NMR (II) Spectra of $[1\text{-}^{13}\text{C}]$ inosine, $[1\text{-}^{13}\text{C}]$ ribose, and Hypoxanthine Spectra are of (A) 5.3 mM free $[1\text{-}^{13}\text{C}]$ inosine and (B) $[1\text{-}^{13}\text{C}]$ ribose and hypoxanthine obtained through the irreversible arsenolysis of 5.3 mM $[1\text{-}^{13}\text{C}]$ inosine in the presence of human PNP and 50 mM Na_2HAsO_4 (pH 7.4). Upfield signals have been omitted to highlight the downfield regions of interest. All spectra were acquired at 25°C in 10% D_2O . [1] Ino, inosine; [2] Hx, hypoxanthine.

This chemical shift does not match those for ribose isomers (Figure 3B). The time course (0–24 hr) ^1H NMR spectra of mixtures of PNPs (native, F159G, Y88G, H257G, and F200G) with $[1\text{-}^{13}\text{C}]$ inosine were also acquired. The ^1H NMR spectra of $[1\text{-}^{13}\text{C}]$ inosine with native PNP showed two new downfield singlets with chemical shifts of 8.55 and 8.18 ppm and another doublet of doublets with a chemical shift of 6.28 ppm between ~3–6 and 24 hr (Figure 4-II). The chemical shifts resemble those reported for the chemically synthesized N3-isomer of inosine (3- β -D-ribofuranosyl hypoxanthine or N3-isoinosine) [5] (s δ 8.67 or 8.59, s δ 8.32 or 8.25 and d δ 5.98 or 5.87 for H-8, H-2 and H-1', respectively, in d_6 , DMSO) (Lehikoinen, et al., 1989; Tindall, et al., 1972). The rate of isomerization was approximately $1 \times 10^{-3} \text{ s}^{-1}$ (Table 1). In comparison, the four catalytic-site glycine mutants did not form any detectable isomerized product during the time course of ^1H NMR observation (Figure 4-III, as an example for F200G).

The identity of the isomerized product as N3-isoinosine was determined by several independent criteria. Resolution from inosine and hypoxanthine by high-performance liquid chromatography (HPLC) established a distinct chemical entity. The electrospray ionization mass spectrometry (ESI-MS) mass value ($m/z = 270$, $\text{M}+\text{H}^+$ for the $1\text{-}^{13}\text{C}$ substituted species) corresponds to $[1\text{-}^{13}\text{C}]$ N3-isoinosine (but does not distinguish it from $[1\text{-}^{13}\text{C}]$ inosine or $[1\text{-}^{13}\text{C}]$ N7-isoinosine). UV-spectral properties at pH 1, 7, and 11 matched published findings for N3-isoinosine and are distinct from those reported for inosine and N7-isoinosine (Table S3) (Lehikoinen, et al., 1989; Montgomery and Thomas, 1969; Tindall, et al., 1972; Wolfenden, et al., 1966). The ^1H NMR spectrum of this isolated product agreed with the published chemical shift data for authentic N3-isoinosine but not for inosine or N7-isoinosine (Table S3) (Chenon, et al., 1975a; Lehikoinen, et al., 1989; Tindall, et al., 1972). The $1\text{-}^{13}\text{C}$ chemical shift is consistent with $[1\text{-}^{13}\text{C}]$ N3-isoinosine but not inosine or N7-isoinosine (Table S3) (Chenon, et al., 1975a).

PNP Reaction Fidelity with Phosphate

Conditions were selected to permit 2×10^7 catalytic turnovers of PNP at chemical equilibrium in the presence of phosphate and inosine. The ^{13}C and ^1H NMR spectra of reaction mixtures with native, Y88G and F159G PNPs showed only the formation of hypoxanthine [2] and α -D- $[1\text{-}^{13}\text{C}]$ ribose 1-phosphate [4] (Figure 5). The ^{13}C and ^1H NMR spectra for H257G and F200G




Figure 4. ^{13}C and ^1H NMR Spectra of $[1'-^{13}\text{C}]$ Inosine with Human PNP in the Absence of Phosphate

(Panel I) ^{13}C NMR spectra of $[1'-^{13}\text{C}]$ inosine with native human PNP after (A) 30 min and (B) 24 hr.

(Panel II) ^1H NMR spectra of $[1'-^{13}\text{C}]$ inosine with native human PNP after (A) 30 min, (B) 3 hr, (C) 6 hr, (D) 12 hr, and (E) 24 hr.

(Panel III) ^1H NMR spectra of $[1'-^{13}\text{C}]$ inosine with F200G after (A) 40 min and (B) 24 hr. All ^{13}C and ^1H spectra were acquired with 5.3 mM $[1'-^{13}\text{C}]$ inosine and 6 μM PNP in the absence of phosphate at 25°C in 10% D_2O . [1] Ino, inosine; [5] N3-Ino, N3-inosine.

Table 1. Comparison of the N3 Isomerization and Single-Turnover Hydrolysis Reactions for PNPs with Inosine as Substrate

PNP	Isomerization Kinetics ^a		Hydrolysis Kinetics ^b
	k_{iso} (s^{-1}), no phosphate ^c	k_{iso} (s^{-1}), with phosphate ^c	k_{hyd} (s^{-1}) ^c
BtPNP ^d	Not determined	Not determined	2.0×10^{-3}
Native	1×10^{-3}	Undetectable ^e	$(7.5 \pm 1.6) \times 10^{-4}$
Y88G	Undetectable ^e	Undetectable ^e	$(1.7 \pm 0.5) \times 10^{-3}$
F159G	Undetectable ^e	Undetectable ^e	$(3.0 \pm 0.8) \times 10^{-3}$
H257G	Undetectable ^e	7.0×10^{-5}	$(4.8 \pm 0.4) \times 10^{-4}$
F200G	Undetectable ^e	4.2×10^{-4}	$(3.6 \pm 1.2) \times 10^{-3}$

^a Isomerization assays were performed with 5.3 mM inosine in 20 mM Tris-HCl (pH 7.4) in the presence or absence of 50 mM KH_2PO_4 (pH 7.4). PNPs were added to give a phosphorolysis V_{max} of $2.6 \times 10^{-4} \text{ M s}^{-1}$ ($k_{\text{cat}} \times [E]_{\text{T}}$).

^b Hydrolysis assays were performed with equal amounts of inosine and PNP (80 μM) in 20 mM Tris-Cl (pH 7.4) at 25°C. Reactions were quenched with 1 N HCl, and the ratio between substrate (inosine) and product (hypoxanthine) was quantified by reverse-phase HPLC.

^c k_{iso} and k_{hyd} were calculated as indicated in Supplemental Data.

^d From Kline and Schramm (1992), at 30°C.

^e N3-Isoinosine was not detectable by ^1H or ^{13}C NMR during the time course; the limit for detection was estimated to be $\sim 25 \mu\text{M}$ (0.5% of the initial [inosine]).

PNPs also showed formation of N3-isoinosine [5] but not ribose (Figures 5-I and 5-II, 5E, and 5F). N3-Isoinosine is apparent after $\sim 3 \times 10^6$ catalytic turnovers in the phosphorolysis reactions catalyzed by both H257G and F200G. After the entire inosine sample had passed through the catalytic site approximately 4000 times (~ 24 hr), the N3-isoinosine peak was integrated to be $\sim 5\%$ and $\sim 10\%$ for H257G and F200G, respectively, relative to the initial $[1'-^{13}\text{C}]$ inosine peak (Figure 5); thus, the probability for N3 capture of the ribocation is 1:80,000 for H257G and 1:40,000 for F200G. In contrast, the phosphorolysis reaction of native human PNP showed no detectable N3-isoinosine or ribose formation after more than 10^7 transits through the catalytic site.

Single-Turnover Hydrolysis

Rates of single-turnover inosine hydrolysis catalyzed by native and catalytic-site glycine mutants of human PNP were compared with the previously reported values for bovine PNP (Table 1). The single-turnover hydrolysis rates for native human PNP were found to be ~ 3 -fold slower than those for the bovine enzyme (Table 1, Figure 6A). The introduction of solvent leaks into the active site of human PNP through the four engineered glycine mutants caused increased rates of inosine hydrolysis compared to native PNP (Table 1 and Figure 6A). The rate of hydrolysis for Y88G increased 2.3-fold, for F159G increased by 4-fold, for H257G decreased by 1.6-fold, and for F200G increased by 4.8-fold, relative to native PNP (Table 1 and Figure 6A).

^{13}C NMR Detection of Hydrolysis under High Enzyme Concentrations

To analyze the PNP reaction for slow hydrolysis, 0.9 mM $[1'-^{13}\text{C}]$ inosine, 0.2 mM (monomers) of PNPs, and increased




Figure 5. ^{13}C NMR (I) and ^1H NMR (II) Spectra of $[1'-^{13}\text{C}]$ Inosine with Human PNP in the Presence of Phosphate

$[1'-^{13}\text{C}]$ Inosine (5.3 mM) was incubated (A) alone or with (B) 6 μM native human PNP, (C) 14 μM Y88G, (D) 33 μM F159G, (E) 151 μM H257G, or (F) 264 μM F200G. All spectra were acquired in the presence of 50 mM KH_2PO_4 , pH 7.4, at 25°C in 10% D_2O and after $\sim 2 \times 10^7$ enzymatic turnovers (20–24 h). [1] Ino, inosine; [2] Hx, hypoxanthine; [4] R-1-P, α -D-ribose 1-phosphate; [5] N3-Ino, N3-isoinosine.

example for F159G). Attempts to perform similar solvolysis experiments with aqueous mixtures of methanol (up to 20% v/v) failed to yield any detectable methyl riboside by ^{13}C or ^1H NMR, even after prolonged incubations (>48 hr).

DISCUSSION

Inosine Hydrolysis by Human PNP

Without phosphate, solvent H_2O is expected to fill the relatively large phosphate binding site of PNP. From its position in the phosphate site, water acts as an inefficient nucleophile to capture the ribocation. In contrast, bound phosphate is known to be highly polarized and activated at the catalytic site (Deng, et al., 2004). Phosphate also participates in ribocation formation and geometric stabilization by forming an ion pair with the developing ribocation and H-bonds to the ribosyl 2'- and 3'-hydroxyls (Figure 2). Our results suggest that when these interactions are missing, the ribosyl group might not be highly constrained in the catalytic site, and once it forms, it can rebound to form inosine (Figure 1, path a), react with water (path c), or react with N3 of hypoxanthine to form N3-isoinosine (path d; see below). Interestingly, Richard and coworkers have also characterized the partitioning

of an enzyme-generated oxocarbenium ion among reactant reformation, capture by various nucleophiles and capture by solvent water (Richard, et al., 1996).

Human PNP hydrolyzes inosine 2.7-fold more slowly than the bovine enzyme (Kline and Schramm, 1992, 1995). The hydrolytic reaction rate of $7.5 \times 10^{-4} \text{ s}^{-1}$ for native HsPNP is approximately 5×10^{-6} of the on-enzyme chemical turnover rate of the phosphorylytic reaction (154 s^{-1}) (Saen-Oon, et al., 2008a). Bound phosphate therefore contributes a factor of approximately 10^5 to TS formation.

Ribocation Reactivity

Transition state lifetimes of nonenzymatic chemical reactions have been shown to be $\sim 10^{-13} \text{ s}$, as determined with the use of femtosecond time-resolved spectroscopy (Baumert, et al., 2001). Postulates that enzymes preferentially stabilize their transition states suggest that their lifetimes may be substantially longer. Kinetic isotope effect studies indicate that the TS of human PNP is a fully dissociated ribocation with complete leaving-group departure ($\text{C}1'-\text{N}9$ distance = 3 Å) and a similar

acquisition times were used. These conditions enable ^{13}C -NMR detection of ribose resulting from the hydrolysis of a single equivalent of inosine, even if subsequent turnovers are hindered by rate-limiting hypoxanthine release, as previously found for bovine PNP (Kline and Schramm, 1995). The ^{13}C NMR spectra with $[1'-^{13}\text{C}]$ inosine under these conditions showed the appearance of D-ribose [3] (β -D-ribofuranose, δ 94.22) (Figure 6B). After 20 hr, H257G had converted all of the 0.9 mM inosine to D-ribose [3] (β - and α -D-ribofuranose) (Figure 6B, spectrum E). Although inosine hydrolysis by H257G is 1.6-fold slower than native PNP, it formed hypoxanthine and ribose without any detectable isomerization to N3-inosine. Native human, bovine, and the other three catalytic-site glycine mutants of PNP showed incomplete hydrolysis of $[1'-^{13}\text{C}]$ inosine [1] and accumulation of D-ribose [3] (Figure 6B). N3-Isoinosine [5] was also readily detected with native (human or bovine) PNPs and to a lesser extent with F200G (Figure 6B). ^{13}C NMR spectra of similar experiments for Y88G, F159G and F200G recorded after 48 hr showed significant accumulations of ribose and N3-isoinosine at approximately equal concentrations (Figure S2, as an




Figure 6. Hydrolysis Reactions under High Enzyme Concentrations

(A) Single-turnover hydrolysis reaction kinetics, plotting the concentration of hypoxanthine (isolated on a C18 analytical column by reverse-phase HPLC) formed during the hydrolysis reaction of native (●), Y88G (●), F159G (●), H257G (●), and F200G human PNP (●) as a function of mixing time prior to acid quenching. Data were fit to Equation (1).

(B) ^{13}C NMR spectra of the PNP hydrolysis reactions. $[1'-^{13}\text{C}]$ inosine (900 μM) with (A) native bovine PNP (BtPNP), (B) native human PNP (HsPNP), (C) Y88G, (D) F159G, (E) H257G, and (F) F200G. All spectra were acquired after 20 hr at 25°C with 200 μM PNP in 10% D_2O . [1] Ino, inosine; [3], D-ribose; [5] N3-Ino, N3-isoinosine.

distance to the phosphate nucleophile (i.e., an $\text{S}_{\text{N}}1$ mechanism) (Lewandowicz and Schramm, 2004; Schramm, 2005). The TS lifetime for human PNP has been estimated by computational transition path sampling to be ~ 10 fs (10^{-14} s) with reaction coordinate motion occurring over ~ 70 fs (Saen-Oon, et al., 2008b). On-enzyme chemistry occurs over a much longer time-scale (154 s $^{-1}$, once every 6×10^{-3} s), reflecting the necessity for multiple geometric alignments between the protein and reactants to permit formation of the TS (Saen-Oon, et al., 2008a, 2008b). Because the lifetime of the ribocation is between 10^{-14} (the computed TS lifetime) and 6×10^{-3} s, we designed an experimental test to determine if the ribocation is stabilized (near 6×10^{-3} s) or has a lifetime near the time-scale of a single bond vibration lifetime (near 10^{-14} s). The lifetime of the ribocation can be estimated by catalytic-site mutations that are designed to permit water access to the reactive ribocation.

Undetectable inosine hydrolysis in the presence of phosphate establishes steric or temporal exclusion for water capture of the ribocation. Transition state analysis (Lewandowicz and Schramm, 2004) and X-ray structures of PNP complexed with

TS analogs (Figure 2) reveal a $> 3 \text{ \AA}$ separation between the bound phosphate and C-1', a distance that is expected to permit the diffusional approach of water at assay temperatures. Based on the X-ray structures, phosphate is poised more closely to the ribocation than structurally observed catalytic site waters. If the ribocation is an intermediate and has a lifetime permitting the diffusion of solvent or catalytic site waters to the ribocation, then some ribose formation would be observed. The lack of hydrolysis is consistent with a ribocation lifetime that is too short to allow diffusion of a water molecule to within reactive distance of the ribocation C-1'.

Times for water diffusion are a function of distance and environment. The shortest trajectory from solvent water to C1' of the ribocation is approximately 10 \AA by way of a wide channel located at the interface of the two adjacent subunits, surrounded by H257, F159', F200, and Y88. Solvent water must diffuse around the ribosyl ring, a path that is partially obstructed by phosphate and enzyme residues. An alternative consideration is water bound within the catalytic site, as is presumed to exist in the absence of phosphate. X-ray structures of phosphate-bound PNPs reveal the existence of a structural water within 4.4 \AA of C-1' (Figure 2B). Diffusion of approximately 3.0 \AA would permit formation of ribose. The upper rate of water diffusion is approximately 1 \AA ps^{-1} (based on the water diffusion coefficient of $3.01 \times 10^{-5} \text{ cm}^2 \cdot \text{s}^{-1}$ at 25°C [Wang, 1951]), which sets lower limits of 10 and 3 ps, respectively, for the time of diffusion of solvent water and the catalytic-site water. The lower limit on the rate of diffusion, however, is not defined for the specific environment in the catalytic site of PNP. If we assume it is slowed by a factor of five, we can, with this assumption, set upper limits of 50 and 15 ps, respectively, for these water molecules. The lifetimes of carbohydrate oxocarbenium ions in water are estimated to be ~ 1 ps (Amyes and Jencks, 1989). Based on our upper estimates of diffusion time, we can propose that the ribocation formed at the TS of PNP has a lifetime less than the 15 to 50 ps required for diffusional capture by catalytic-site or solvent water.

Formation of N3-Isoinosine

A control for the ribocation capture experiment is the rate of inosine solvolysis in the absence of phosphate. Removal of phosphate permits solvent access to the catalytic site and the unequivocal formation of ribose. Thus, phosphate-free PNP forms the ribocation, and without phosphate, most is captured by water to give ribose. In addition to the hydrolytic reaction, a novel product, N3-isoinosine, was also formed. The mechanism proposed for N3-isoinosine formation involves ribocation formation in a catalytic site where mutations weaken the geometric alignment of the purine base and/or the ribocation (Figure 2). Without tight geometric constraints on the ribocation and displaced base, nucleophilic rebound can occur to both N9 (the normal reaction) and, by misalignment, to N3, causing isomerization. Phosphate is directly involved in this alignment by its H-bond contacts with the 2'- and 3'-hydroxyl groups of ribose (Figure 2). In the presence of phosphate, the alignment of N9-hypoxanthine with the ribocation fails less than once in a million transfers through the catalytic site, as evidenced by the failure to detect any N3-isomer in multiple-turnover NMR studies. In the absence of phosphate, the purine leaving-group interactions assist in ribocation formation, but the poor reactivity of the disordered waters in the phosphate-binding site allows the ribocation to wobble on its nucleophilic rebound.

The slow formation of N3-isoinosine in the absence of phosphate is consistent with the full formation of the ribocation. With water molecules in the phosphate site, ribocation capture occurs with N9 (to reform inosine), N3 to form isoinosine, and solvent to form ribose. In the presence of phosphate, an alternative mechanism for isomerization involving N3 attack upon the ribose 1-phosphate product cannot be excluded, but more complex mechanisms depart from known features of the physiological PNP mechanism.

N3-Isoinosine is a byproduct in the chemical synthesis of inosine (Lehikoinen et al., 1989; Ryu et al., 2004; Wolfenden et al., 1966) and is an alternative substrate for PNPs (Bzowska et al., 1996; Lehikoinen et al., 1989), but it has never been reported as a product in any enzymatic reactions. Nucleoside deoxyribosyltransferases have been demonstrated to form 2'-deoxynucleosides with N3- and N7-fused purine derivatives (Huang et al., 1983; Steenkamp, 1991); however, unlike with PNP, these isomers are likely formed via the covalent deoxyribosylated enzyme intermediate that is known to form during its function with natural substrates (Smar et al., 1991).

The lifetime of PNP intermediates can be compared with a nonenzymatic isomerization reaction for cyclopropane stereomutation (Hrovat et al., 1997). The mechanism of isomerization proceeds via C–C bond homolysis to generate a trimethylene biradical, which rotates about the linearized bonds before closing. Their calculations estimated a biradical lifetime of 140 fs, within the range of the 10–50,000 fs lifetime for the PNP-generated ribocation estimated in our computational (Saen-Oon et al., 2008b) and solvolysis experiments, respectively.

Inosine Hydrolysis and Isomerization by Catalytic-Site Mutants of Human PNP

Four glycine mutants of the residues that are proposed to shield the catalytic site from solvent were prepared to create a water leak into the catalytic site and thereby introduce a hydrolytic

branch point in the reaction mechanism (Figure 1, path c). The hydrolytic rate is a measure of the water access and reactivity as well as the lifetime of the highly reactive ribocation. The mutants are reduced in k_{cat} by up to 44-fold relative to the native enzyme. All enzymes were capable of inosine hydrolysis with rates similar to or faster than native PNP.

The formation of N3-isoinosine in the absence of phosphate is consistent with the formation of a ribocation TS that reacts with hypoxanthine at N9 (to form inosine) or N3 (yielding N3-isoinosine) or with water (yielding ribose). Unlike the native enzyme, isomerization to N3-isoinosine was not observed with any of the mutants under the conditions of the hydrolysis experiments. In the presence of phosphate, the PNP species with relatively high k_{cat}/K_m values (i.e., native, Y88G and F159G; Table S1) efficiently transfer the ribocation to phosphate without any detectable side reactions. Less efficient PNP mutants (i.e., H257G and F200G) occasionally misdirect the electrophile, resulting in formation of N3-isoinosine. Reaction with N3 is rare—only 1 of 40,000 (F200G) or 80,000 (H257G) catalytic cycles generates N3-inosine, based on its rate of formation. H257 anchors the 5'-hydroxyl group of the ribosyl moiety, and its removal provides the ribosyl cation sufficient positional flexibility to permit slow ribocation capture by N3. The crystal structure of human PNP complexed with a TS-analog inhibitor (Immuticillin-H) and phosphate indicates that the phenyl-group side chain of residue F200 is involved in a herringbone-type interaction with the purine base (Figure 2) (de Azevedo et al., 2003; Ghanem et al., 2009). In F200G, the positional stability of the purine base is compromised. In this mutant, the ribocation contacts are unchanged from native PNP, but weakened interactions to hypoxanthine permit both N9 and N3 to react with the ribocation.

SIGNIFICANCE

Native human PNP and catalytic-site glycine mutants slowly hydrolyze inosine when phosphate is absent but not in the presence of phosphate. These mutants were designed to permit solvent leaks; however, the lifetime of the ribocation is too short to permit solvent capture of the ribocation in competition with phosphate. In the absence of phosphate, human PNP catalyzes the capture of the ribocation by N3 of the purine group, resulting in isomerization of the substrate. The TS of human PNP involves a fully dissociated ribo-oxacarbenium ion (i.e., S_N1 character). The chemical properties of a ribocation are consistent with the isomerization at N3, in which hypoxanthine departure is required to permit $N9 \rightarrow N3$ isomerization. The chemical reactivity of PNP and its catalytic site mutants provide experimental evidence for a ribocation lifetime too short to permit efficient solvent capture of the ribocation. Amino acids that stabilize both the ribocation (e.g., H257) and hypoxanthine (e.g., F200) are required for fidelity of product formation. Atomic freedom of motion around the TS results in the formation of N3-isoinosine during phosphorolysis by these mutants.

EXPERIMENTAL PROCEDURES

Steady-State Enzyme Assays

Activity assays for native and mutant PNPs with inosine as a substrate were carried out by monitoring the conversion of hypoxanthine to uric acid

($\epsilon_{293} = 12.9 \text{ mM}^{-1} \text{ cm}^{-1}$) (Kim et al., 1968) in a coupled assay containing 60 milliunits xanthine oxidase and variable concentrations of inosine in 50 mM KH_2PO_4 (pH 7.4) at 25°C (Lewandowicz et al., 2003). Substrate concentration (inosine or [$1'$ - ^{13}C]inosine) was determined spectrophotometrically using the published millimolar extinction coefficient of 12.3 at 248.5 nm, pH 7 (Dawson et al., 1986).

[$1'$ - ^{13}C]Inosine

[$1'$ - ^{13}C]Adenosine (Omicron Biochemicals, Inc.) was converted to [$1'$ - ^{13}C]inosine by treatment with calf spleen adenosine deaminase (Sigma) (Pfrogner, 1967a, 1967b), with spectrophotometric monitoring at 267 nm ($\Delta\epsilon_{267} = 6.5 \text{ mM}^{-1} \text{ cm}^{-1}$) (Schramm and Baker, 1985) at pH 7. [$1'$ - ^{13}C]inosine was purified by reverse-phase HPLC (5% methanol in 50 mM triethylammonium acetate [pH 5.0] at 1 ml min^{-1}) using an analytical C18 column (Waters Deltapak, 15 μm , 300 Å, 300 × 3.9 mm).

Steady-State NMR Detection for Hydrolysis, Phosphorolysis, and Isomerization

The ^{13}C and ^1H NMR spectra of reaction mixtures with PNP and labeled inosine were obtained by mixing 6 μM native human PNP with 5.3 mM [$1'$ - ^{13}C]inosine in 20 mM Tris-HCl (pH 7.4) and 10% D_2O , in the presence or the absence of 50 mM KH_2PO_4 (pH 7.4) for $\sim 2 \times 10^7$ enzymatic turnovers (~ 20 to 24 hr). Native human PNP has a chemical turnover of 154 s^{-1} and steady-state rate of 44 s^{-1} , limited by the rate of hypoxanthine release (Ghanem et al., 2008a). During 24 hr, the entire inosine sample passes through the catalytic site approximately 15,000 times (according to chemical turnovers) in this freely reversible reaction. Thus, a hydrolytic reaction rate of once in every 300,000 phosphorolytic reactions ($5 \times 10^{-4} \text{ s}^{-1}$) would cause 5% of the inosine to accumulate as ribose and would be readily detected by NMR. For mutant PNPs, enzyme was added at concentrations that yield the same steady-state phosphorolytic rate as native PNP (e.g., $[\text{H257G}] = [\text{Native PNP}] \times k_{\text{cat, native}} \div k_{\text{cat, H257G}} = 6 \mu\text{M} \times 44 \text{ s}^{-1} \div 1.75 \text{ s}^{-1} = 150 \mu\text{M}$). Standard references included free [$1'$ - ^{13}C]inosine and [$1'$ - ^{13}C]ribose obtained through the irreversible arsenolysis of [$1'$ - ^{13}C]inosine in the presence of PNP and 50 mM Na_2HAsO_4 (pH 7.4). All spectra were acquired using a Bruker 300 MHz spectrometer with 5 mm BBI probe at 298 K. ^1H -decoupled ^{13}C NMR spectra were collected with 256 scans of 64 K points ($\sim 15 \text{ min/spectrum}$), a recycle delay of 2 s and a carbon sweep width of 315 ppm with the carrier set to 126 ppm. ^1H NMR spectra were collected with 64 scans of 16 K points, a recycle delay of 1.3 s and a proton sweep width of 14 ppm with the carrier set to 4.7 ppm. Spectra were processed with an exponential window function of 1 Hz. Spectra were assigned as described previously (Dorman and Roberts, 1970; Teixeira et al., 1997; Tindall et al., 1972).

^{13}C NMR Detection for Hydrolysis under High Enzyme Concentrations

^{13}C NMR spectra of reaction mixtures with high molar ratios of PNP to inosine were obtained by mixing 200 μM (monomers) of PNP with 900 μM of [$1'$ - ^{13}C]inosine in 20 mM Tris-HCl (pH 7.4) and 10% D_2O , for ~ 20 to 24 hr. ^1H -decoupled ^{13}C NMR spectra were collected with 1024 scans of 64 K points ($\sim 58 \text{ min}$), a recycle delay of 2 s and a carbon sweep width of 315 ppm with the carrier set to 126 ppm using a Bruker 300 MHz spectrometer with 5 mm BBI probe at 298 K. See Supplemental Data for calculation of k_{iso} .

Single-Turnover Hydrolysis

Equal amounts of inosine and enzyme (80 μM final) were incubated at 25°C for 30 s and up to 72 hr, followed by quenching with HCl (1 N final) and removal of the denatured protein by centrifugation. Hypoxanthine (retention time 4.3 min) and inosine (retention time 7.3 min) were quantitated by UV detection at 249 nm using reverse-phase HPLC on a Waters Deltapak C18 column (15 μm , 300 Å, 300 × 3.9 mm), equilibrated with 5% methanol in 50 mM triethylammonium acetate (pH 5.0) at 1 ml min^{-1} . See Supplemental Data for calculation of k_{hyd} .

Characterization of N3-Isoinosine

N3-Isoinosine was isolated from incubations of inosine with human PNP in the absence of phosphate by reverse-phase HPLC using an analytical C18 column (Phenomenex Luna 5 μm , 100 Å, 150 × 4.6 mm), equilibrated with an ion-pair

buffer (100 mM KH_2PO_4 , 8 mM Bu_4NHSO_4 [pH 6.0]) at 1 ml min^{-1} , with monitoring by a Waters 996 Photodiode Array Detector. Under these conditions, hypoxanthine, inosine, and N3-isoinosine eluted with retention times of 3.2, 8.07, and 8.8 min, respectively (Figure S1). The N3-isomer peak was collected and reduced to 400 μl in a Speedvac concentrator, and this sample was desalted by application to a second C18 column (Waters Deltapak 15 μm , 300 Å, 300 × 3.9 mm), equilibrated with water at 1.5 ml min^{-1} . N3-Isoinosine was collected, concentrated to dryness in a Speedvac concentrator, redissolved in water, and analyzed by ESI-MS and by UV/V is spectroscopy at pH 1, 7, and 11.

SUPPLEMENTAL DATA

Supplemental Data include three tables, two figures, and Supplemental Experimental Procedures and can be found with this article online at [http://www.cell.com/chemistry-biology/supplemental/S1074-5521\(09\)00251-8](http://www.cell.com/chemistry-biology/supplemental/S1074-5521(09)00251-8).

ACKNOWLEDGMENTS

We acknowledge the US National Institutes of Health grants GM41916 and GM068036 for funding.

Received: April 30, 2009

Revised: June 26, 2009

Accepted: July 23, 2009

Published: September 24, 2009

REFERENCES

- Amyes, T.L., and Jencks, W.P. (1989). Lifetimes of oxocarbenium ions in aqueous solution from common ion inhibition of the solvolysis of alpha-azido ethers by added azide ion. *J. Am. Chem. Soc.* *111*, 7888–7900.
- Baumert, T., Frohnmeyer, T., Kiefer, B., Niklaus, P., Strehle, M., Gerber, G., and Zewail, A.H. (2001). Femtosecond transition state dynamics of cis-stilbene. *Appl. Phys. B* *72*, 105–108.
- BioCryst (2009). BioCryst research pipeline (<http://www.biocryst.com/pipeline.htm>). BioCryst Pharmaceuticals, Inc.
- Bzowska, A., Kulikowska, E., Poopko, N.E., and Shugar, D. (1996). Kinetics of phosphorolysis of 3-(beta-D-ribofuranosyl)adenine and 3-(beta-D-ribofuranosyl)hypoxanthine, non-conventional substrates of purine-nucleoside phosphorylase. *Eur. J. Biochem.* *239*, 229–234.
- Chenon, M.T., Pugmire, R.J., Grant, D.M., Panzica, R.P., and Townsend, L.B. (1975a). Carbon-13 magnetic resonance. XXV. A basic set of parameters for the investigation of tautomerism in purines. Established from carbon-13 magnetic resonance studies using certain purines and pyrrolo[2,3-d]pyrimidines. *J. Am. Chem. Soc.* *97*, 4627–4636.
- Chenon, M.T., Pugmire, R.J., Grant, D.M., Panzica, R.P., and Townsend, L.B. (1975b). Carbon-13 magnetic resonance. XXVI. A quantitative determination of the tautomeric populations of certain purines. *J. Am. Chem. Soc.* *97*, 4636–4642.
- Dawson, R.M.C., Elliott, D., Elliott, W.H., and Jones, K.M. (1986). Spectral data and pKa values for purines, pyrimidines, nucleosides, and nucleotides. In *Data for Biochemical Research*, Third Edition, R.M.C. Dawson, ed. (New York: Oxford Univ. Press), pp. 103–114.
- de Azevedo, W.F., Jr., Canduri, F., dos Santos, D.M., Silva, R.G., de Oliveira, J.S., de Carvalho, L.P., Basso, L.A., Mendes, M.A., Palma, M.S., et al. (2003). Crystal structure of human purine nucleoside phosphorylase at 2.3 Å resolution. *Biochem. Biophys. Res. Commun.* *308*, 545–552.
- Deng, H., Lewandowicz, A., Schramm, V.L., and Callender, R. (2004). Activating the phosphate nucleophile at the catalytic site of purine nucleoside phosphorylase: a vibrational spectroscopic study. *J. Am. Chem. Soc.* *126*, 9516–9517.
- Dorman, D.E., and Roberts, J.D. (1970). Nuclear magnetic resonance spectroscopy: ^{13}C spectra of some common nucleotides. *Proc. Natl. Acad. Sci. USA* *65*, 19–26.

- Ebrahimi, M., Rossi, P., Rogers, C., and Harbison, G.S. (2001). Dependence of ^{13}C NMR chemical shifts on conformations of rna nucleosides and nucleotides. *J. Magn. Reson.* *150*, 1–9.
- Fedorov, A., Shi, W., Kicska, G., Fedorov, E., Tyler, P.C., Furneaux, R.H., Hanson, J.C., Gainsford, G.J., Laresse, J.Z., et al. (2001). Transition state structure of purine nucleoside phosphorylase and principles of atomic motion in enzymatic catalysis. *Biochemistry* *40*, 853–860.
- Ghanem, M., Li, L., Wing, C., and Schramm, V.L. (2008a). Altered thermodynamics from remote mutations altering human toward bovine purine nucleoside phosphorylase. *Biochemistry* *47*, 2559–2564.
- Ghanem, M., Saen-oon, S., Zhadin, N., Wing, C., Cahill, S.M., Schwartz, S.D., Callender, R., and Schramm, V.L. (2008b). Tryptophan-free human PNP reveals catalytic site interactions. *Biochemistry* *47*, 3202–3215.
- Ghanem, M., Zhadin, N., Callender, R., and Schramm, V.L. (2009). Loop-tryptophan human PNP reveals submillisecond protein dynamics. *Biochemistry* *48*, 3568–3668.
- Hrovat, D.A., Fang, S., Borden, W.T., and Carpenter, B.K. (1997). Investigation of cyclopropane stereomutation by quasiclassical trajectories on an analytical potential energy surface. *J. Am. Chem. Soc.* *119*, 5253–5254.
- Huang, M.C., Montgomery, J.A., Thorpe, M.C., Stewart, E.L., Secrist, J.A., 3rd, and Blakley, R.L. (1983). Formation of 3-(2'-deoxyribofuranosyl) and 9-(2'-deoxyribofuranosyl) nucleosides of 8-substituted purines by nucleoside deoxyribosyltransferase. *Arch. Biochem. Biophys.* *222*, 133–144.
- Kenneth, N., Drew, J.Z., Bondo, G., Bose, B., and Serianni, A.S. (1998). ^{13}C -labeled aldopentoses: detection and quantitation of cyclic and acyclic forms by heteronuclear 1D and 2D NMR spectroscopy. *Carbohydr. Res.* *307*, 199–209.
- Kim, B.K., Cha, S., and Parks, R.E., Jr. (1968). Purine nucleoside phosphorylase from human erythrocytes. II. Kinetic analysis and substrate-binding studies. *J. Biol. Chem.* *243*, 1771–1776.
- Kline, P.C., and Schramm, V.L. (1992). Purine nucleoside phosphorylase. Inosine hydrolysis, tight binding of the hypoxanthine intermediate, and third-site reactivity. *Biochemistry* *31*, 5964–5973.
- Kline, P.C., and Schramm, V.L. (1993). Purine nucleoside phosphorylase. Catalytic mechanism and transition-state analysis of the arsenolysis reaction. *Biochemistry* *32*, 13212–13219.
- Kline, P.C., and Schramm, V.L. (1995). Pre-steady-state transition-state analysis of the hydrolytic reaction catalyzed by purine nucleoside phosphorylase. *Biochemistry* *34*, 1153–1162.
- Koellner, G., Luic, M., Shugar, D., Saenger, W., and Bzowska, A. (1997). Crystal structure of calf spleen purine nucleoside phosphorylase in a complex with hypoxanthine at 2.15 Å resolution. *J. Mol. Biol.* *265*, 202–216.
- Lehikoinen, P.K., Sinnott, M.L., and Krenitsky, T.A. (1989). Investigation of α -deuterium kinetic isotope effects on the purine nucleoside phosphorylase reaction by the equilibrium-perturbation technique. *Biochem. J.* *257*, 355–359.
- Lewandowicz, A., and Schramm, V.L. (2004). Transition state analysis for human and *Plasmodium falciparum* purine nucleoside phosphorylases. *Biochemistry* *43*, 1458–1468.
- Lewandowicz, A., Shi, W., Evans, G.B., Tyler, P.C., Furneaux, R.H., Basso, L.A., Santos, D.S., Almo, S.C., and Schramm, V.L. (2003). Over-the-barrier transition state analogues and crystal structure with *Mycobacterium tuberculosis* purine nucleoside phosphorylase. *Biochemistry* *42*, 6057–6066.
- Montgomery, J.A., and Thomas, H.J. (1969). Ribosyl derivatives of hypoxanthine. *J. Org. Chem.* *34*, 2646–2650.
- Murkin, A.S., Birck, M.R., Rinaldo-Matthis, A., Shi, W., Taylor, E.A., Almo, S.C., and Schramm, V.L. (2007). Neighboring group participation in the transition state of human purine nucleoside phosphorylase. *Biochemistry* *46*, 5038–5049.
- Nunez, S., Wing, C., Antoniou, D., Schramm, V.L., and Schwartz, S.D. (2006). Insight into catalytically relevant correlated motions in human purine nucleoside phosphorylase. *J. Phys. Chem. A* *110*, 463–472.
- Pfrogner, N. (1967a). Adenosine deaminase from calf spleen. I. Purification. *Arch. Biochem. Biophys.* *119*, 141–146.
- Pfrogner, N. (1967b). Adenosine deaminase from calf spleen. II. Chemical and enzymological properties. *Arch. Biochem. Biophys.* *119*, 147–154.
- Richard, J.P., Huber, R.E., Heo, C., Amyes, T.L., and Lin, S. (1996). Structure-reactivity relationships for beta-galactosidase (*Escherichia coli*, lac Z). 4. Mechanism for reaction of nucleophiles with the galactosyl-enzyme intermediates of E461G and E461Q beta-galactosidases. *Biochemistry* *35*, 12387–12401.
- Rinaldo-Matthis, A., Murkin, A.S., Ramagopal, U.A., Clinch, K., Mee, S.P., Evans, G.B., Tyler, P.C., Furneaux, R.H., Almo, S.C., et al. (2008). L-Enantiomers of transition state analogue inhibitors bound to human purine nucleoside phosphorylase. *J. Am. Chem. Soc.* *130*, 842–844.
- Ryu, K.-S., Kim, C., Park, C., and Choi, B.-S. (2004). NMR analysis of enzyme-catalyzed and free-equilibrium mutarotation kinetics of monosaccharides. *J. Am. Chem. Soc.* *126*, 9180–9181.
- Saen-oon, S., Ghanem, M., Schramm, V.L., and Schwartz, S.D. (2008a). Remote mutations and active site dynamics correlate with catalytic properties of purine nucleoside phosphorylase. *Biophys. J.* *94*, 4078–4088.
- Saen-oon, S., Quaytman-Machleder, S., Schramm, V.L., and Schwartz, S.D. (2008b). Atomic detail of chemical transformation at the transition state of an enzymatic reaction. *Proc. Natl. Acad. Sci. USA* *105*, 16543–16548.
- Schramm, V.L. (2005). Enzymatic transition states: thermodynamics, dynamics and analogue design. *Arch. Biochem. Biophys.* *433*, 13–26.
- Schramm, V.L., and Baker, D.C. (1985). Spontaneous epimerization of (S)-deoxycoformycin and interaction of (R)-deoxycoformycin, (S)-deoxycoformycin, and 8-ketodeoxycoformycin with adenosine deaminase. *Biochemistry* *24*, 641–646.
- Shi, W., Ting, L.M., Kicska, G.A., Lewandowicz, A., Tyler, P.C., Evans, G.B., Furneaux, R.H., Kim, K., Almo, S.C., et al. (2004). *Plasmodium falciparum* purine nucleoside phosphorylase: crystal structures, Immucillin inhibitors, and dual catalytic function. *J. Biol. Chem.* *279*, 18103–18106.
- Smar, M., Short, S.A., and Wolfenden, R. (1991). Lyase activity of nucleoside 2-deoxyribosyltransferase: transient generation of ribal and its use in the synthesis of 2'-deoxynucleosides. *Biochemistry* *30*, 7908–7912.
- Steenkamp, D.J. (1991). The purine-2-deoxyribonucleosidase from *Crithidia luciliae*. Purification and trans-N-deoxyribosylase activity. *Eur. J. Biochem.* *197*, 431–439.
- Stoekler, J.D., Agarwal, R.P., Agarwal, K.C., and Parks, R.E., Jr. (1978). Purine nucleoside phosphorylase from human erythrocytes. *Methods Enzymol.* *51*, 530–538.
- Stoekler, J.D., Cambor, C., and Parks, R.E., Jr. (1980). Human erythrocytic purine nucleoside phosphorylase: reaction with sugar-modified nucleoside substrates. *Biochemistry* *19*, 102–107.
- Teixeira, M., Santana, L., and Uriarte, E. (1997). Assignment of the ^{13}C NMR spectra of some adenine, hypoxanthine and guanine carbonucleosides. *Magn. Reson. Chem.* *35*, 806–807.
- Tindall, C.G., Jr., Robins, R.K., Tolman, R.L., and Hutzenlaub, W. (1972). Directed glycosylation of 8-bromo-adenine. Synthesis and reactions of 8-substituted 3-glycosyladenine derivatives. *J. Org. Chem.* *37*, 3985–3989.
- Wang, J.H. (1951). Self-diffusion and structure of liquid water. II. Measurement of self-diffusion of liquid water with O18 as tracer. *J. Am. Chem. Soc.* *73*, 4181–4183.
- Wolfenden, R., Sharpless, T.K., Ragade, I.S., and Leonard, N.J. (1966). Enzymatic and chemical deamination of 3-(β -D-ribofuranosyl)adenine. *J. Am. Chem. Soc.* *88*, 185–186.