

Commentary

The structure of the O-linked carbohydrate chain of bovine seminal plasma protein PDC-109 revised by $^1\text{H-NMR}$ spectroscopy

A correction

Gerrit J. Gerwig^a, Juan J. Calvete^{b,*}, Edda Töpfer-Petersen^b, Johannes F.G. Vliegthart^a

^a*Bijvoet Center for Biomolecular Research, Department of Bio-Organic Chemistry, Utrecht University, Utrecht, The Netherlands*

^b*Institut für Reproduktionsmedizin, Tierärztliche Hochschule Hannover, Bünteweg 15, 30559 Hannover-Kirchrode, Germany*

Received 16 April 1996

PDC-109, a 13-kDa protein [1], is the major secretory product of the seminal vesicles of the bull (*Bos taurus*) and constitutes the most abundant protein of seminal plasma (16–25 mg/ml) [2]. It possesses heparin- and phosphorylcholine-binding activities [3] and coats the spermatozoa surface at ejaculation [4], enhancing their fertilizing capability [5]. PDC-109 has a single O-glycosylated residue (threonine 11) [4]. Carbohydrate composition, mass spectrometric, and lectin mapping analyses suggested the Neu5Ac(α 2-6)-Gal(β 1-3)-GalNAc structure for the carbohydrate chain of bovine PDC-109 [4]. Here, we report the revision of the oligosaccharide structure of PDC-109 after $^1\text{H-NMR}$ spectroscopic analysis.

PDC-109 was isolated from the seminal plasma of Holstein bulls by chromatography on a DEAE-Sephadex A25 equilibrated with 10 mM Tris/HCl, 1 M NaCl, 5 mM EDTA, 0.025% NaN₃, pH 7.4. PDC-109 was eluted with column buffer containing 10 mM *o*-phosphorylcholine. O-linked carbohydrate chains were released from PDC-109 (200 ml) by alkaline borohydride treatment (β -elimination) [6] with 50 ml of 0.1 M NaOH, 1 M NaBH₄ for 48 h at 37°C in the dark and under a nitrogen atmosphere. After centrifugation (3000 rpm, 20 min), the supernatant was acidified to pH 6 with 4 M acetic acid at 0°C, applied to a 11 \times 2.5 cm Dowex 50X8 (H⁺ form) column, washed with \sim 200 ml cold water, and the eluate lyophilized. Boric acid was removed by co-evaporation with methanol. The remaining material was fractionated on a BioGel P2 column (46 \times 1.6 cm, BioRad) eluting with 5 mM NH₄HCO₃. Fractions, monitored at 206 nm, were tested for hexose content by the orcinol/H₂SO₄ method. The major carbohydrate-containing fraction was purified by gel filtration chromatography on a 46 \times 1 cm BioGel P4 column (BioRad).

Carbohydrate-containing fractions were repeatedly treated with D₂O (99.96 atom% D, Isotec Inc.) with intermediate lyophilization. High-resolution 500-MHz $^1\text{H-NMR}$ spectra were recorded with a Bruker AMX-500 spectrometer at a probe temperature of 300 K. Acetone (δ 2.225 ppm) was used as internal standard [7].

Monosaccharide analysis of O-linked carbohydrates released from PDC-109 by β -elimination demonstrated that GalNAc was completely converted into GalNAc-ol (Table 1). The little amount of Gal-ol found (\sim 10% of total Gal) indicated that peeling may have occurred during the β -elimination procedure [8]. Fractionation on BioGel P2 yielded six fractions, denoted P2-I to P2-VI. Fraction P2-V contained free monosaccharide alditols. Fraction P2-IV contained neutral material and $^1\text{H-NMR}$ spectroscopy proved the structure to be Gal(β 1-3)-GalNAc-ol [9]. Fraction P2-I was further purified on BioGel P4, and four subfractions, designated P4-I to P4-IV, were recovered. Fractions P4-I and P4-II did not contain carbohydrate. Fraction P4-IV contained a small amount of free sialic acid. $^1\text{H-NMR}$ spectroscopy of fraction P4-III clearly demonstrated the structure Neu5Ac(α 2-3)-Gal(β 1-3)-GalNAc-ol: the β -Gal residue was identified by the H-1 and H-4 signals at δ 4.547 ($J_{1,2}$ 7.9 Hz) and δ 3.931 ppm, respectively. The Gal H-3 was observed at δ 4.122 ppm. The Gal(β 1-3)-GalNAc-ol core was characterized by the H-2 and H-5 signals of GalNAc-ol at δ 4.389 and 4.188 ppm, respectively. The NAc singlet of GalNAc-ol was observed at δ 2.046 ppm. The structural-reporter groups of Neu5Ac, namely NAc signal at δ 2.034 ppm, H-3a at δ 1.800 ppm, and H-3e at δ 2.773 ppm, indicated an α -Neu5Ac residue linked to C3 of the galactose residue.

Table 1
Monosaccharide analysis of PDC-109 and derived fractions

Monosaccharide	PDC-109		BioGel P2			BioGel P4	
	native	after β -elimination	P2-I	P2-IV	P2-V	P4-III	P4-IV
Gal	1.0	0.9	1.1	1.1	\pm	1.3	—
Gal-ol	—	0.1	0.1	0.1	1.1	—	\pm
GalNAc	0.9	—	—	—	—	—	—
GalNAc-ol	—	1.0	1.0	1.0	1.0	1.0	—
Neu5Ac	1.1	1.1	1.2	\pm	—	1.2	1.0

Values are given in mol saccharide per mol protein. \pm means present, but less than 0.1.

*Corresponding author. Fax: (49) (511) 953 8504.
E-mail: JCalvete@Repro.TiHo-Hannover.De

This study confirms that the major (>80%) carbohydrate structure of PDC-109 is a trisaccharide and, in addition, unambiguously establishes that the anomery of the *N*-acetyl neuraminic acid-galactose linkage is α 2–3 rather than α 2–6.

Acknowledgements: This work was partly financed by grants PB92-0096 from the Dirección General de Investigación Científica y Técnica, Madrid, Spain, BMBF 01KY9503 from the Bundesministerium für Bildung, Forschung und Technologie, Bonn, Germany, and the Netherlands Foundation for Chemical Research (SON/NWO).

References

- [1] Esch, F.S., Ling, N.C., Böhlen, P., Ying, S.Y. and Guillemin, R. (1983) *Biochem. Biophys. Res. Commun.* 113, 861–867.
- [2] Scheit, K.-H., Kemme, M., Aumüller, G., Seitz, J., Hagendorff, G. and Zimmer, M. (1988) *Biosci. Rep.* 8, 569–608.
- [3] Desnoyers, L. and Manjunath, P. (1992) *J. Biol. Chem.* 267, 10149–10155.
- [4] Calvete, J.J., Raida, M., Sanz, L., Wempe, F., Scheit, K.-H., Romero, A. and Töpfer-Petersen, E. (1994) *FEBS Lett.* 350, 203–206.
- [5] Thérien, I., Bleau, G. and Manjunath, P. (1995) *Biol. Reprod.* 52, 1372–1379.
- [6] Damm, J.B.L., Kamerling, J.P., Van Dedem, G.W.K. and Vliegthart, J.F.G. (1987) *Glycoconjugate J.* 4, 129–144.
- [7] Vliegthart, J.F.G., Dorland, L. and Van Halbeek, H. (1983) *Adv. Carbohydr. Chem. Biochem.* 41, 209–374.
- [8] Lloyd, K.O., Kabat, E.A. and Licerio, E. (1968) *Biochemistry* 7, 2976–2990.
- [9] Kamerling, J.P. and Vliegthart, J.F.G. (1992) *Biol. Magn. Reson.* 90, 1–94.