



## Solvent effects in gold-catalysed A<sup>3</sup>-coupling reactions



Gregory A. Price, Alan K. Brisdon\*, Kevin R. Flower\*, Robin G. Pritchard, Peter Quayle\*

School of Chemistry, The University of Manchester, Manchester M13 9PL, UK

### ARTICLE INFO

#### Article history:

Received 28 July 2013

Revised 25 October 2013

Accepted 29 October 2013

Available online 6 November 2013

#### Keywords:

Gold

A<sup>3</sup>-reaction

Trifluoroethanol

Microwave

Solvent effect


### ABSTRACT

Gold-catalysed A<sup>3</sup>-reactions proceed efficiently when conducted in 2,2,2-trifluoroethanol as solvent. The rates of these reactions are accelerated considerably when conducted in a microwave reactor.

© 2013 The Authors. Published by Elsevier Ltd. Open access under [CC BY license](http://creativecommons.org/licenses/by/3.0/).

The development of domino, multi-component and cascade reactions continues to be an area of intense academic interest<sup>1</sup> even though they were first documented over a century ago.<sup>2</sup> A particularly useful example is the 'A<sup>3</sup>-coupling reaction', an atom-economic, single step process for the preparation of synthetically useful propargyl amines<sup>3,4</sup> from aldehydes, alkynes and amines (Scheme 1). This reaction is found to be catalysed by a number of transition metal complexes<sup>5</sup> in a process that does not require the pre-generation of stoichiometric quantities of nucleophilic  $\sigma$ -alkyne organometallic reagents. The use of strong bases (Grignard or organolithium reagents) is therefore obviated in these reactions, a factor which contributes to their greater synthetic potential, flexibility and atom economy.

Concurrent with the development of multi-component reactions there has also been a resurgence of interest in the chemistry of gold, especially as applied to A<sup>3</sup>-coupling reactions.<sup>6</sup> In their seminal Letter<sup>7</sup> Wei and Li noted that the efficiency of gold-catalysed A<sup>3</sup>-reactions was greatly dependent upon the reaction conditions employed: reactions carried out at 100 °C in water for 12 h, using catalyst loadings of 1%, proved to be optimal for the substrates used in these initial investigations. The use of organic solvents (THF, DMF and toluene), rather than water, proceeded in lower conversions and with concomitant formation of undesired by-products.<sup>7</sup>



Scheme 1. Transition metal catalysed A<sup>3</sup> coupling reaction.

Our interest<sup>8a,b</sup> in the chemistry of organogold complexes encouraged us to investigate the use of complexes **1–4** in A<sup>3</sup>-coupling reactions and led us to question whether these complexes could be employed as homogeneous solutions in organic solvents. Complexes **1–4** (Fig. 1) used in this study were initially screened in a standard A<sup>3</sup>-reaction between benzaldehyde (1 mmol) and phenylacetylene (1.5 mmol) using either dibenzylamine or piperidine (1.1 mmol) as the amine partner, in a range of solvents at temperatures in the range between 40 °C and 60 °C.

Initial results indicated that the Au(III) complex **1** was catalytically active and resulted in quantitative conversion to the requisite A<sup>3</sup>-coupled product after 24 h in most of the solvents examined (Table 1, entries 1–4). Complexes **2**, **3** and **4** also promoted the A<sup>3</sup>-reaction, albeit at a much reduced rate, when the reactions were conducted in water (entries 6, 15 and 18; <20% conversion at 40 °C for 24 h). Most notably, exchange of water for chloroform as reaction medium afforded homogeneous reaction mixtures which resulted in a significant improvement in conversion rates, even when the less reactive catalysts **2** and **3** were employed (entries 9 and 16). Further studies clearly indicated that this was a solvent, rather than a temperature effect, as attempted coupling at 100 °C in water failed to force the reaction to go to completion (entries 7 and 8).

\* Corresponding authors. Tel.: +44 (0)1612754619; fax: +44 (0)1612754598.

E-mail address: [peter.quayle@manchester.ac.uk](mailto:peter.quayle@manchester.ac.uk) (P. Quayle).

**Table 1**  
Solvent and catalyst screening experiments for A<sup>3</sup> reactions

Entry	Catalysts	Solvent/T	Amine	Reaction time (h)	Conversion <sup>a,b</sup> , %
1	<b>1</b>	H <sub>2</sub> O/40 °C	HNBn <sub>2</sub>	24	97 (85)
2	<b>1</b>	CHCl <sub>3</sub> /60 °C	HNBn <sub>2</sub>	24	100
3	<b>1</b>	PhCF <sub>3</sub> /60 °C	HNBn <sub>2</sub>	24	90
4	<b>1</b>	CF <sub>3</sub> CH <sub>2</sub> OH/40 °C	HNBn <sub>2</sub>	24	96 (84)
5	<b>1</b>	MeOH/40 °C	HNBn <sub>2</sub>	24	51
6	<b>2</b>	H <sub>2</sub> O/40 °C	Piperidine	24	18
7	<b>2</b>	H <sub>2</sub> O/60 °C	Piperidine	24	33
8	<b>2</b>	H <sub>2</sub> O/100 °C	Piperidine	24	75
9	<b>2</b>	CHCl <sub>3</sub> /60 °C	Piperidine	24	100
10	<b>2</b>	PhCF <sub>3</sub> /60 °C	HNBn <sub>2</sub>	24	15
11	<b>2</b>	CF <sub>3</sub> CH <sub>2</sub> OH/60 °C	Piperidine	24	100
12	<b>2</b>	CF <sub>3</sub> CH <sub>2</sub> OH/60 °C	HNBn <sub>2</sub>	24	100
13	<b>2</b>	CF <sub>3</sub> CH <sub>2</sub> OH/40 °C	HNBn <sub>2</sub>	24	100
14	<b>2</b>	MeOH/40 °C	HNBn <sub>2</sub>	24	54
15	<b>3</b>	H <sub>2</sub> O/40 °C	HNBn <sub>2</sub>	24	9
16	<b>3</b>	CHCl <sub>3</sub> /60 °C	HNBn <sub>2</sub>	24	65
17	<b>3</b>	CF <sub>3</sub> CH <sub>2</sub> OH/60 °C	HNBn <sub>2</sub>	24	88
18	<b>4</b>	H <sub>2</sub> O/40 °C	HNBn <sub>2</sub>	72	10
19	<b>4</b>	CHCl <sub>3</sub> /60 °C	HNBn <sub>2</sub>	24	22
20	<b>4</b>	CHCl <sub>3</sub> /60 °C	HNBn <sub>2</sub>	72	31
21	<b>4</b>	CHCl <sub>3</sub> /60 °C	HNBn <sub>2</sub>	168	62
22	<b>4</b>	CF <sub>3</sub> CH <sub>2</sub> OH/40 °C	HNBn <sub>2</sub>	24	96
23	<b>4</b>	MeOH/40 °C	HNBn <sub>2</sub>	24	39
24	No catalyst	H <sub>2</sub> O/40 °C	HNBn <sub>2</sub>	24	0
25	No catalyst	CHCl <sub>3</sub> /60 °C	HNBn <sub>2</sub>	24	0
26	No catalyst	CF <sub>3</sub> CH <sub>2</sub> OH/40 °C	HNBn <sub>2</sub>	24	0
27	No catalyst	PhCF <sub>3</sub> /60 °C	HNBn <sub>2</sub>	24	0

Reaction conditions: 1 mmol benzaldehyde, 1.1 mmol amine, 1.5 mmol phenylacetylene, 1 mol % catalyst, 2 mL solvent, 24 h.

<sup>a</sup> Conversion determined by <sup>1</sup>H NMR analysis of the crude reaction mixtures based on benzaldehyde conversion.

<sup>b</sup> Isolated yield.

It is conceivable that the increased rate observed for complex **2** when the reactions were conducted in chloroform was simply due to an improved solubility of the complex in this solvent. A similar rate enhancement was also observed when the sterically encumbered complex **4** was employed, although the degree of conversion plateaued at 62% even after extended reaction times (entries 19–21). The use of methanol<sup>9</sup> or benzotrifluoride (PhCF<sub>3</sub>; a nontoxic alternative<sup>10</sup> to CH<sub>2</sub>Cl<sub>2</sub> and CHCl<sub>3</sub>) instead of chloroform proved to be deleterious to conversion rates. The fluorinated alcohol, 2,2,2-trifluoroethanol, (TFE) exhibits a number of unique properties ( $pK_a = 12.4$ ;  $E^N_T = 0.898$ ;  $\alpha = 1.51$ ;  $\beta = 0$ ;  $N = -2.78$ ;  $Y = 3.82$ ), and proved to have a pronounced effect on conversion rates of A<sup>3</sup>-reactions (entries 11–13, 17 and 22).

For the most active catalyst **1**, substitution of water by TFE had no appreciable effect upon the overall rate of conversion or isolated yield of A<sup>3</sup>-products (H<sub>2</sub>O and TFE afford isolated yields of 85% and 84%; entries 1 and 4); these results which should be compared to the relatively poor outcome obtained using methanol as solvent (entry 5). A quite different result was encountered with the less active catalysts **3** and **4**. In the case of the least active catalyst **4**, near quantitative conversion to the desired A<sup>3</sup>-product was achieved when using TFE as solvent after 24 h at 40 °C (entry 22). This outcome should be compared to the poor conversions in either water (10% after 72 h at 40 °C; entry 18) or chloroform (62% conversion after 168 h at 60 °C; entry 21). A similar improvement in conversion, when compared to water as solvent, was also observed with **3** in combination with TFE as reaction solvent (entry 17). It should be noted that A<sup>3</sup>-coupling reactions did not proceed when conducted in the absence of gold in TFE (entries 24–27).

While there are numerous examples of microwave-assisted copper catalysed A<sup>3</sup>-reactions,<sup>12</sup> application of this technique to homogeneous, gold-catalysed A<sup>3</sup>-reactions is, to our knowledge, without literature precedent.<sup>13</sup> The effect of microwave irradiation on gold-catalysed A<sup>3</sup>-reactions is tabulated above (Table 2).

**Table 2**  
Screening experiments for microwave reactions

Entry	Catalysts	Solvent	Conversion <sup>a</sup> (%)
1	<b>1</b>	H <sub>2</sub> O	89
2	<b>1</b>	CHCl <sub>3</sub>	43
3	<b>1</b>	PhCF <sub>3</sub>	33
4	<b>1</b>	Perfluorodecalin	76
5	<b>1</b>	CF <sub>3</sub> CH <sub>2</sub> OH	86
6	<b>1</b>	MeOH	50
7	<b>1</b>	PEG-400	61
8	<b>2</b>	H <sub>2</sub> O	10
9	<b>2</b>	CHCl <sub>3</sub>	10
10	<b>2</b>	PhCF <sub>3</sub>	11
11	<b>2</b>	Perfluorodecalin	30
12	<b>2</b>	MeOH	40
13	<b>2</b>	PEG-400	27
14	<b>2</b>	CF <sub>3</sub> CH <sub>2</sub> OH	77

Reaction conditions: 1 mmol benzaldehyde, 1.1 mmol dibenzylamine, 1.5 mmol phenylacetylene, 1 mol % catalyst, 1 mL solvent, 100 °C,  $\mu$ W, 30 min.

<sup>a</sup> Conversion determined by <sup>1</sup>H NMR analysis of the crude reaction mixtures based on benzaldehyde conversion.

Highest conversions using **1** as catalyst were obtained when either H<sub>2</sub>O (entry 1) or TFE (entry 5) were used as solvents. The use of either *cis/trans*-perfluorodecalin<sup>14</sup> or PEG-400 (an environmentally benign solvent<sup>15</sup>) also proceeded with good efficiencies. This observation may facilitate the development of 'green', A<sup>3</sup>-coupling, reactions where the propargylamine product can be readily separated from the fluorosolvent and also enable recycling of a fluorosolvent catalyst.<sup>16</sup> The effectiveness of **2**, which generally has a lower catalytic activity than **1** in A<sup>3</sup>-reactions, under the conditions of microwave activation, was also briefly investigated. Here the use of either water, chloroform or benzotrifluoride as solvent resulted in low conversion rates (entries 8–10) whereas reactions carried out in *cis/trans*-perfluorodecalin, methanol or PEG-400 resulted in

a slight enhancement in conversion rates (entries 11–13). Most startling however was the observation that, once again, the rate of  $A^3$ -reactions are substantially accelerated when conducted in TFE as solvent (entry 14).

From this study, and in contrast to an earlier Letter,<sup>7</sup> it is clear that the use of solvents, other than water, are compatible with efficient Au-catalysed  $A^3$ -reactions. In particular we note that TFE has a beneficial effect upon both thermal and microwave-promoted gold-catalysed  $A^3$ -reactions, such that these reactions can be driven to completion with short reaction times and without loss in yields. The TFE-modification does not require the use anaerobic conditions, proceeds efficiently in reagent grade solvents, and enables facile product isolation (TFE has a bp of 77 °C).





Figure 1. Gold complexes utilised in this investigation.



Thermal:  $CF_3CH_2OH$ , 40 °C, 24 h, 1 mol% **1**;  
 $\mu W$ :  $CF_3CH_2OH$ , 30 min, 100 °C, 1 mol% **1**.

Figure 2. Representative  $A^3$ -reactions conducted in TFE.

Table 3  
 Selected properties of solvents<sup>10,11</sup>

Solvent	$E^N_T$ (polarity)	HBD	Dipole moment (D)
Perfluorodecalin	–	0	0
$PhCF_3$	0.241	–	2.9
$CH_2Cl_2$	0.309	0.13	1.57
$CHCl_3$	0.259	0.1	1.2
PEG-400	0.575	0.31	4.3
MeOH	0.762	0.93	1.7
TFE	0.898	1.51	2.03
$H_2O$	1.0	1.17	1.87

We have also applied our standard reaction conditions to other  $A^3$ -coupling reactions (Fig. 2). We find that either thermal or microwave activation proves to be effective in the promotion of these reactions in TFE and that product isolation using this solvent system proves to be trivial.<sup>17</sup>

## Conclusions

Gold complexes **1–4** were found to be effective catalysts for the  $A^3$ -coupling reaction. The use of TFE as solvent results in improved rates, an effect which can be further augmented by the application of microwave irradiation. We surmise that, in the case of TFE in particular, the observed solvent effects are due to a favourable combination of physical properties as listed in Table 3. TFE is a highly polar, H-bonding, nonnucleophilic solvent which presumably favours both initial iminium ion formation and subsequent capture with an acetylenic gold  $\sigma$ -complex.<sup>18</sup> Studies are now underway for the development of recyclable, fluorine-tagged catalysts for use in  $A^3$ -reactions.

## Acknowledgments

G.A.P. thanks the EPSRC for a research studentship. We thank Alfa Aesar/JJM for kind donation of sodium tetrachloroaurate.

## Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.tetlet.2013.10.141>.

## References and notes

- (a) Peshkov, V. A.; Pereshivko, O. P.; Van der Eycken, E. V. *Chem. Soc. Rev.* **2012**, *41*, 3790–3807; (b) Brauch, S.; Van Berkela, S. S.; Westermann, B. *Chem. Soc. Rev.* **2013**, *42*, 4948–4962; (c) Gupta, A. K.; Singh, N.; Singh, K. N. *Curr. Org. Chem.* **2013**, *17*, 474–490; (d) Liu, Y.; Wang, H.; Wan, J. *Asian J. Org. Chem.* **2013**, *2*, 374–386; (e) Green, N. J.; Sherburn, M. S. *Aust. J. Chem.* **2013**, *66*, 267–283; (f) Pellissier, H. *Chem. Rev.* **2013**, *113*, 442–524; (g) Marson, C. M. *Chem. Soc. Rev.* **2012**, *41*, 7712–7722.
- (a) Strecker, A. *Justus Liebigs Annal. Chem.* **1850**, *75*, 27–45; (b) Mannich, C.; Krosche, W. *Arch. Pharm.* **1912**, *250*, 647–667; (c) Robinson, R. J. *Chem. Soc.* **1917**, *111*, 762–768.
- Jiang, B.; Xu, M. *Angew. Chem., Int. Ed.* **2004**, *43*, 2543–2546.
- (a) Wright, J. L.; Gregory, T. F.; Kesten, S. R.; Boxer, P. A.; Serpa, K. A.; Meltzer, L. T.; Wise, L. D.; Espitia, S. A.; Konkoy, C. S.; Whittemore, E. R.; Woodward, R. M. *J. Med. Chem.* **2000**, *43*, 3408–3419; (b) Kung, K. K.-Y.; Li, G. L.; Zou, L.; Chong, H.-C.; Leung, Y.-C.; Wong, K.-H.; Lo, V. K.-Y.; Che, C.-M.; Wong, M.-K. *Org. Biomol. Chem.* **2012**, *10*, 925–930; (c) Uhlig, N.; Li, C. J. *Org. Lett.* **2012**, *14*, 3000–3003; (d) Renhack, A.; Jumpertz, T.; Ness, J.; Baches, S.; Pietrzik, C. U.; Weggen, S.; Bulic, B. *Bioorg. Med. Chem.* **2012**, *20*, 6523–6532.
- (a) Zani, L.; Bolm, C. *Chem. Commun.* **2006**, 4263–4275; (b) Zhou, X.; Lu, Y.; Zhai, L.-L.; Zhao, Y.; Liu, Q.; Sun, W.-Y. *RSC Advances* **2013**, *3*, 1732; (c) Yao, X.; Li, C.-J. *Org. Lett.* **2005**, *7*, 4395–4398; (d) Wei, C.; Li, Z.; Li, C.-J. *Org. Lett.* **2003**, *5*, 4473–4475; (e) Li, C.-J.; Wei, C. *Chem. Commun.* **2002**, 268–269; (f) Koradin, C.; Polborn, K.; Knochel, P. *Angew. Chem., Int. Ed.* **2002**, *41*, 2535–2538; (g) Bieber, L. W.; da Silva, M. F. *Tetrahedron Lett.* **2004**, *45*, 8281–8283; (h) Wie, C.; Li, C.-J. *J. Am. Chem. Soc.* **2002**, *124*, 5638–5639; (i) Li, P.; Zhang, Y.; Wang, L. *Chem. Eur. J.* **2009**, *15*, 2045–2049; (j) Chen, W.-W.; Nguyen, R. V.; Li, C.-J. *Tetrahedron Lett.* **2009**, *50*, 2895–2898; (k) Fischer, C.; Carreira, E. M. *Org. Lett.* **2001**, *3*, 4319–4321; (l) Pin-Hua, L.; Lei, W. *Chin. J. Chem.* **2005**, 1076–1080.
- (a) Schmidbaur, H.; Schier, A. *Arab. J. Sci. Eng.* **2012**, *37*, 1187–1225; (b) De Haro, T.; Nevado, C. *Synlett* **2011**, 2530–2539; (c) Oña-Burgos, P.; Fernández, I.; Rocas, L.; Fernández, L. T.; García-Granda, S.; Ortiz, F. L. *Organometallics* **2009**, *28*, 1739–1747; (d) Kidwai, M.; Bansal, V.; Kumar, A.; Mozumdar, S. *Green Chem.* **2007**, *9*, 742–745; (e) Graf, T. A.; Anderson, T. K.; Bowden, N. B. *Adv. Synth. Catal.* **2011**, *353*, 1033–1038; (f) Elie, B. T.; Levine, C.; Ubarretxena-Belandia, I.; Varela-Ramírez, A.; Aguilera, R. J.; Ovalle, R.; Contel, M. *Eur. J. Inorg. Chem.* **2009**, 3421–3430; (g) Zhang, X.; Corma, A. *Angew. Chem., Int. Ed.* **2008**, *47*, 4358–4361; (h) Lili, L.; Xin, Z.; Jinsen, G.; Chunming, X. *Green Chem.* **2012**, *14*, 1710–1720; (i) Lo, V. K.-Y.; Liu, Y.; Wong, M.-K.; Che, C.-M. *Org. Lett.* **2006**, *8*, 1529–1532; (j) Villaverde, G.; Corma, A.; Iglesias, M.; Sánchez, F. *ACS Catal.* **2012**, *2*, 399–406.
- Wie, C.; Li, C.-J. *J. Am. Chem. Soc.* **2003**, *125*, 9584–9585.
- (a) Price, G. A.; Flower, K. R.; Pritchard, R. G.; Brisdon, A. K.; Quayle, P. *Dalton Trans.* **2011**, *40*, 11696–11697; (b) Price, G. A.; Brisdon, A. K.; Randall, S.; Whittaker, D. M.; Pritchard, R. G.; Muryn, C. A.; Flower, K. R.; Quayle, P. *Dalton Trans.* **2013** (submitted).

9. Chen, M.-T.; Landers, B.; Navarro, O. *Org. Biomol. Chem.* **2012**, *10*, 2206–2208.
10. (a) Ogawa, A.; Curran, D. P. *J. Org. Chem.* **1997**, *62*, 450–451; (b) Maul, J. J.; Ostrowski, P. J.; Ublacker, G. A.; Linclau, B.; Curran, D. P. *Top. Curr. Chem.* **1999**, *206*, 79–105.
11. (a) Shuklov, I.; Dubrovina, N.; Börner, A. *Synthesis* **2007**, 2925–2943; (b) Brégué, J.-P.; Bonnet-Delpon, D.; Crousse, B. *Synlett* **2004**, 18–29.
12. (a) Shi, L.; Tu, Y.; Wang, M.; Zhang, F.; Fan, C. *Org. Lett.* **2004**, *6*, 1001–1003; (b) Ju, Y.; Li, C.-J.; Varma, R. *QSAR Comb. Sci.* **2004**, *23*, 891–894; (c) Kabalka, G.; Wang, L.; Pagni, R. *Synlett* **2001**, 676–678; (d) Bariwal, J. B.; Ermolat'ev, D. S.; Van der Eycken, E. V. *Chem. Eur. J.* **2010**, *16*, 3281–3284.
13. For a MACOS  $A^3$ -reaction see: Shore, G.; Yoo, W.-J.; Organ, M. G. *Chem. Eur. J.* **2010**, *16*, 126–133.
14. For the use of *cis/trans*-perfluorodecalin see: (a) Hong, M.; Cai, C.; Yi, W.-B. *J. Fluorine Chem.* **2010**, *131*, 111–114; (b) Shi, M.; Cui, S.-C. *Chem. Commun.* **2002**, 994–995; For use of perfluoromethyldecalin in microwave-assisted reactions see: (c) Benali, O.; Deal, M.; Farrant, E.; Tapolczay, D.; Wheeler, R. *Org. Process Res. Dev.* **2008**, *12*, 1007–1011.
15. For multicomponent reactions in PEG: (a) Gu, Y. *Green Chem.* **2012**, *14*, 2091–2128; Cu-catalysed  $A^3$ -reactions in PEG: (b) Zhang, Q.; Chen, J.-X.; Gao, W.-X.; Ding, J.-C. *Appl. Organometal. Chem.* **2010**, *24*, 809–812.
16. (a) Hong, M.; Cai, C.; Yi, W. B. *Chin. Chem. Lett.* **2011**, *22*, 322–325; (b) Vincent, J.-M. *Top. Curr. Chem.* **2012**, *308*, 153–174.
17. Full experimental details can be found in the Supplementary data.
18. For similar solvent effects in related condensation–addition reactions of aldehydes see: (a) Heydari, A.; Khaksar, S.; Tajbaksh, M. *Tetrahedron Lett.* **2009**, *50*, 77–80; (b) Hosseinzadeh, R.; Alinezhad, H.; Ghahari, S.; Heydari, A. *Synthesis* **2011**, 490–496; (c) Khaksar, S.; Alipour, M. *Monatsh Chem.* **2013**, *144*, 395–398.