

# NAADP receptors are present and functional in the heart

Judit Bak<sup>\*†</sup>, Richard A. Billington<sup>\*</sup>, Gyoergy Timar<sup>†</sup>, Alice C. Dutton<sup>\*</sup>  
and Armando A. Genazzani<sup>\*</sup>

**Alongside the well-studied inositol 1,4,5 trisphosphate and ryanodine receptors, evidence is gathering that a new intracellular release mechanism, gated by the pyridine nucleotide nicotinic acid adenine dinucleotide phosphate (NAADP), is present in numerous organisms, ranging from plant to mammalian cells (reviewed in [1]). Most cells have been shown to express at least two Ca<sup>2+</sup>-release mechanisms controlled by different messengers, and this can lead to redundancy, convergence, or divergence of responses. One exception appears to be muscle and heart contractile tissues. Here, it is thought that the dominant intracellular channel is the ryanodine receptor, while IP<sub>3</sub> receptors are poorly expressed and their role appears to be negligible. We now report that NAADP receptors are functional and abundant in cardiac microsomes. NAADP binds specifically and with high affinity (130 pM and 4 nM) to two sites on cardiac microsomes and releases Ca<sup>2+</sup> with an apparent EC<sub>50</sub> of 323 ± 14 nM. Furthermore, binding experiments show that this receptor displays both positive and negative cooperativity, a peculiarity unique among intracellular Ca<sup>2+</sup> channels. Therefore, we show that the heart possesses multiple mechanisms to increase the complexity of Ca<sup>2+</sup> signaling and that NAADP may be integral in the functioning of this organ.**

Addresses: <sup>\*</sup>Department of Pharmacology, CB2 1QJ Cambridge, United Kingdom. <sup>†</sup>Department of Medical Biochemistry, Semmelweis University, Budapest, Hungary.

Correspondence: Armando A. Genazzani  
E-mail: aag25@cam.ac.uk

Received: 16 March 2001  
Revised: 17 April 2001  
Accepted: 3 May 2001

Published: 26 June 2001

**Current Biology** 2001, 11:987–990

0960-9822/01/\$ – see front matter  
© 2001 Elsevier Science Ltd. All rights reserved.


## Results and discussion

First, we investigated whether NAADP could induce Ca<sup>2+</sup> release from adult rabbit heart microsomes loaded passively with <sup>45</sup>Ca<sup>2+</sup> (see Materials and methods in the Supplementary material available with this article online) and

compared it to cADPR-induced Ca<sup>2+</sup> release. In this assay, both NAADP (1 μM) and cADPR (1 μM) induced a fast Ca<sup>2+</sup> efflux from the microsomes, which differed significantly from control microsomes (Figure 1a). In the first 3 seconds after its addition, NAADP released 8.3 ± 0.31 nmol Ca<sup>2+</sup>/mg protein, while cADPR released 9.9 ± 0.23 nmol Ca<sup>2+</sup>/mg protein, compared to the 5.2 ± 0.32 nmol Ca<sup>2+</sup>/mg protein released by the control release solution (Figure 1a). It is thought that the latter release occurs in part due to Ca<sup>2+</sup>-induced Ca<sup>2+</sup> release (CICR), since an optimal concentration of this ion is present in the release buffer (1 μM). To evaluate the efficacy of release by pyridine nucleotides, we compared Ca<sup>2+</sup> release to that induced by caffeine, an established agonist of the ryanodine receptor. Caffeine (5 mM) released 10.9 ± 0.88 nmol Ca<sup>2+</sup>/mg protein in 3 s (44.3% of ionomycin release, n = 3). In the same set of experiments, cADPR released 9.9 ± 0.32 nmol Ca<sup>2+</sup>/mg protein (91% of caffeine release, n = 3), and NAADP released 8.6 ± 0.22 nmol Ca<sup>2+</sup>/mg protein (80% of caffeine release, n = 3). Similar to previous data in brain microsomes [2], NAADP appeared insensitive to the extravesicular Ca<sup>2+</sup> concentrations, while controls, as expected, appeared to be the most sensitive (10 nM Ca<sup>2+</sup> in the buffer induced 63% ± 7.2% of 1 μM Ca<sup>2+</sup>). When Ca<sup>2+</sup> was buffered to 10 nM, cADPR released 9.2 ± 0.65 nmol Ca<sup>2+</sup>/mg protein (143% of the control), while NAADP released 10.12 ± 0.93 nmol/mg (158% of the control; data not shown) after 3 s. This would suggest that NAADP-mediated Ca<sup>2+</sup> release is robust and could play a modulatory role toward other release mechanisms. This possibility is further substantiated by the data from other model systems, where NAADP-induced Ca<sup>2+</sup> release is capable of recruiting other release mechanisms [3–5]. To further discriminate between the effect of NAADP and any artifacts of the technique, we loaded microsomes in an active manner with 1 mM ATP in the medium. Ca<sup>2+</sup> ATPases rapidly induced Ca<sup>2+</sup> uptake in microsomes (half-maximal uptake occurred between 15 and 20 s, and maximal uptake occurred after 30–40 s). After 2 min of incubation with ATP, stores were fully loaded, and release was initiated. In this scenario, NAADP (2 μM) and cADPR (2 μM) released similar amounts of Ca<sup>2+</sup> (5.2 ± 0.56 and 6.4 ± 0.56 nmol/mg protein after 5 s, respectively; Figure 1b). Total NAADP and cADPR release was 32%–38% and 40%–43% of ionomycin-releasable Ca<sup>2+</sup>, respectively (calculated after 20 s of the addition of the agonist in two different cardiac preparations). It has been previously suggested that ATP and cADPR might compete for the same site on the ryanodine receptor complex [6], but in our experimental conditions, we could detect cADPR release in the presence of 1 mM ATP at room temperature. When a concentration-response curve

**Figure 1**

(a)  $^{45}\text{Ca}^{2+}$  release from passively loaded cardiac microsomes induced by CICR (solid line, filled squares), NAADP (1  $\mu\text{M}$ ; broken line, filled circles), and cADPR (1  $\mu\text{M}$ ; broken line, open circles). (b) Active  $^{45}\text{Ca}^{2+}$  uptake into cardiac microsomes. Release by NAADP (2  $\mu\text{M}$ ; filled circles) and cADPR (2  $\mu\text{M}$ ; open circles). (c) The concentration dependence of NAADP- (filled circles) and cADPR- (open circles) induced  $\text{Ca}^{2+}$  release. (d) A comparison of the effects of various pharmacological agents on NAADP- (1  $\mu\text{M}$ ; open bars) and cADPR- (1  $\mu\text{M}$ ; filled bars) induced release. Pharmacological agents include verapamil, 100  $\mu\text{M}$ ; diltiazem, 100  $\mu\text{M}$ ; ryanodine, 100  $\mu\text{M}$ ; ruthenium red, 5  $\mu\text{M}$ ; and 8-NH<sub>2</sub>-cADPR, 10  $\mu\text{M}$ . Values are mean  $\pm$  SEM of 3–9 determinations.




was performed after passive loading and the release at 3 s was used as a measure of the total response induced by NAADP, it was found that NAADP releases  $\text{Ca}^{2+}$  with an  $\text{EC}_{50}$  of  $323 \pm 14$  nM and a Hill slope of  $2.32 \pm 0.21$  (Figure 1c). cADPR appeared to be slightly more potent, with an  $\text{EC}_{50}$  of  $199 \pm 6.1$  nM and a Hill slope of  $2.2 \pm 0.11$ . NAADP has been shown to release  $\text{Ca}^{2+}$  via a distinct channel in all systems examined thus far, although interplay between  $\text{Ca}^{2+}$ -release systems has been reported. Similarly, ruthenium red and 8-NH<sub>2</sub>-cADPR, antagonists of the ryanodine receptor-mediated mechanism, had no effect on NAADP-induced  $\text{Ca}^{2+}$  release, while these compounds were capable of partially inhibiting cADPR-mediated responses (Figure 1d). High concentrations of verapamil and diltiazem (100  $\mu\text{M}$ ) selectively, but only partially, inhibited NAADP responses versus cADPR responses, consistent with data from earlier reports in sea urchin eggs [7] (Figure 1d). Although these antagonists did not completely abolish NAADP-induced release, it did not seem appropriate to increase their concentrations, since 100  $\mu\text{M}$  is approximately two orders of magnitude higher than the concentration required to selectively block L-type  $\text{Ca}^{2+}$  channels and already influences non-specifically numerous channels. Furthermore, the coaddition of heparin and 8-NH<sub>2</sub>-cADPR did not affect NAADP-induced  $\text{Ca}^{2+}$  release (data not shown).

If NAADP, as suggested by our data, releases  $\text{Ca}^{2+}$  by

a separate mechanism from that of ryanodine and IP<sub>3</sub> receptors, then we would expect to find a specific binding site in cardiac microsomes. Indeed, [<sup>32</sup>P]NAADP bound avidly to its receptor (see Supplementary material for a [<sup>32</sup>P]NAADP production and binding protocol) with an apparent  $K_d$  of  $4.2 \pm 0.71$  nM (Figure 2a). Furthermore, the Hill slope was close to 2 ( $2.0 \pm 0.89$ ), mimicking closely the cooperativity observed for  $\text{Ca}^{2+}$  release. The discrepancy between the  $\text{EC}_{50}$  of  $\text{Ca}^{2+}$  release and the  $K_d$  is consistent with sea urchin egg data [1], has been shown for other ligands (e.g., IP<sub>3</sub> [8]), and might be attributable to the different kinetics of release versus binding [8]. Our data also suggests that there is an additional specific displacement when more than 1  $\mu\text{M}$  cold ligand was used (Figure 2a), which could represent an NADP<sup>+</sup> binding protein. When adding low concentrations of cold ligand to the 150 pM [<sup>32</sup>P]NAADP, we observe an increase in bound molecules (up to 150% of the control value, Figure 2a). This suggests a positive cooperativity similar to the one observed for some plasma membrane receptors (e.g., insulin). When the data obtained was linearized by Scatchard analysis (data not shown), we detected the presence of at least three binding sites. Alternatively, this could be interpreted as the presence of a single class of binding sites whose affinity changes with receptor occupancy, i.e., negative cooperativity (see below). Alongside a low-affinity site (identified to be the same as the one mentioned above), another two sites were identifiable. They showed

Figure 2



(a) The displacement of  $[^{32}\text{P}]\text{NAADP}$  binding by NAADP (solid line, filled circles) and NADP (broken line, open circles; purified once as described in Supplementary material). (b) The association and dissociation kinetics of  $[^{32}\text{P}]\text{NAADP}$  binding. Dissociation was performed by either 100-fold dilution (open squares) or the addition of excess cold ligand (open circles). Values are mean  $\pm$  SEM of 6–9 determinations.

affinity constants of approximately 130 pM and 4 nM, with  $B_{\text{max}}$  values of 25 and 507 fmol/mg, respectively. Recently, a similar assay has been used to investigate NAADP binding in rat brain [9]. Although, in this report, binding appeared specific, the  $K_{\text{d}}$  was found to be considerably higher ( $\sim 200$  nM), and the Hill slope differed considerably ( $\sim 1$ ). It, therefore, appears that, similar to  $\text{IP}_3$  and ryanodine receptors, different NAADP receptor subtypes may be expressed in different tissues.

Not surprisingly, cADPR had no influence on  $[^{32}\text{P}]\text{NAADP}$  binding. The possibility that this binding site was not specific to NAADP was further tested by challenging  $[^{32}\text{P}]\text{NAADP}$  binding with high concentra-

Table 1

Displacement of  $[^{32}\text{P}]\text{NAADP}$  binding by structural analogs and  $\text{Ca}^{2+}$  mobilizing agents.

Ligand	% Bound
Total	100 $\pm$ 1.23
10 $\mu\text{M}$ NAADP	9.22 $\pm$ 0.71
10 $\mu\text{M}$ NAD	128 $\pm$ 17.74
10 $\mu\text{M}$ NAAD	125 $\pm$ 15.2
10 $\mu\text{M}$ cADPR	113 $\pm$ 7.70
10 $\mu\text{M}$ ryanodine	107 $\pm$ 17.3
100 nM ADPRP	96.7 $\pm$ 5.27
10 $\mu\text{M}$ ADPRP	16.3 $\pm$ 2.16

Values are  $\pm$  SEM of 3–6 determinations.

tions of structurally similar compounds (Table 1).  $\text{NAD}^+$  (10  $\mu\text{M}$ ) lacked affinity for the binding site, as did NAAD, suggesting that, as in sea urchin eggs, the phosphate group is critical for receptor-ligand recognition [10]. On the other hand,  $\text{NADP}^+$  and ADP ribose phosphate (ADPRP) showed some affinity for the receptor.  $\text{NADP}^+$  had an affinity of  $206 \pm 24.5$  nM (Hill slope of  $1.97 \pm 0.12$ ; Figure 2a), while ADPRP had no effect at a concentration of 100 nM but displaced  $60\% \pm 3.9\%$  of the bound  $[^{32}\text{P}]\text{NAADP}$  at 1  $\mu\text{M}$ . The  $\text{NADP}^+$  effect is similar to that observed in sea urchin eggs [11, 12], where it displays a 500-fold lower affinity than NAADP, and part of this effect has been attributed to the contamination present in the commercial sources of  $\text{NADP}^+$ , which is extremely resistant to purification [13]. In support of this,  $\text{NADP}^+$  (20  $\mu\text{M}$ ) bought from commercial sources (Sigma) released 34% of the  $\text{Ca}^{2+}$  released by NAADP (1  $\mu\text{M}$ ), suggesting that NAADP is present in the sample. However, ADPRP seems to be recognized differently by the mammalian receptor compared to that in sea urchin, where the  $K_{\text{d}}$  appears to be greater than 10  $\mu\text{M}$  (R.A.B. and A.A.G., unpublished data). Nonetheless, this data supports the notion that NAADP binding is specific, that the phosphate group is crucial for recognition, and that the pyridine moiety dictates affinity. We, therefore, suggest that it is likely that the binding site in the heart is not identical to that present in sea urchin eggs.

We then investigated the association and dissociation kinetics of  $[^{32}\text{P}]\text{NAADP}$  binding. NAADP associated rapidly to its receptor on ice, with a  $t_{1/2}$  of 0.6 min (Figure 2b). Once again, this is in stark contrast to brain, where association appeared to be  $\sim 10$ -fold slower [9]. To measure dissociation, we allowed full association and then added an excess amount of cold NAADP. Unlike in sea urchin,  $[^{32}\text{P}]\text{NAADP}$  was displaced rapidly, with  $\sim 56.0 \pm 6.5\%$  of specific counts displaced after 15 s of addition of the displacing ligand. To further analyze this, we allowed full association for 20 min and diluted the microsomes 100-fold. This procedure, by reducing the concentration of free ligand in the medium, should induce a similar dissociation to that shown with the addition of cold ligand. Al-

most no dissociation was measured during the time course of the experiment. To our knowledge, the only other proteins that display similar characteristics are plasma membrane peptide receptors (e.g., insulin, GDNF), and this is the first report of an intracellular mechanism displaying such a property. In the case of insulin receptors, the presence of negative cooperativity, whereby the ligand binding site appears to be able to switch between a high-affinity, slow-dissociating state to a low-affinity, fast-dissociating state as their occupancy increases, has been postulated [14–16]. It is highly likely that a similar situation occurs with NAADP. Our data parallels functional observations in pancreatic and submandibular acinar cells [4, 5, 17], where 100  $\mu\text{M}$  NAADP does not elicit responses, but 50 nM does. Nonetheless, whether and how these two observations are mechanistically linked is unclear at the present time.

From our data it appears that NAADP signaling could play a role in cardiac  $\text{Ca}^{2+}$  homeostasis. This possibility is strengthened by the recent finding that the enzyme responsible for the production of NAADP is present on the intracellular membranes of the myocardium [18]. Furthermore, a preliminary report suggests that NAADP enhances spark occurrence in permeabilized cardiomyocytes [19]. This contribution, together with a recent report that  $\text{IP}_3$  receptors might play a role in the atria in excitation-contraction coupling [20], suggests that the heart possesses multiple release mechanisms. Whether NAADP-sensitive signaling represents a “backup” system for ryanodine receptor-mediated signaling, whether it represents a signal that is decoded differently (allowing the myocyte to discriminate between a signal to contract and a signal to perform a different task), or whether it coordinates ryanodine-receptor signaling (as proposed recently in pancreatic acinar cells, starfish, and sea urchins) remains to be established.

#### Supplementary material

The Materials and methods section, including information on microsome preparation, radioligand binding, and  $\text{Ca}^{2+}$ -release assays, can be found with the Supplementary material, which is available at <http://images.cellpress.com/supmat/supmatin.htm>.

#### Acknowledgements

This work was funded by the Biotechnology and Biological Sciences Research Council (David Phillips Fellowship, A.A.G.), the Department of Pharmacology (Departmental studentship, R.A.B.), and the Hungarian Research Funds (J.B., OTKA D-25594, ETT 414).

#### References

1. Lee HC: **Physiological functions of cyclic ADP-ribose and NAADP as calcium messengers.** *Annu Rev Pharmacol Toxicol* 2001, **41**:317-345.
2. Bak J, White P, Timar G, Missiaen, RL, Genazzani, AA, Galione, A: **Nicotinic acid adenine dinucleotide phosphate triggers  $\text{Ca}^{2+}$  release from brain microsomes.** *Curr Biol* 1999, **9**:751-754.
3. Santella L, Kyojuka K, Genazzani AA, De Riso L, Carafoli E: **Nicotinic acid adenine dinucleotide phosphate-induced  $\text{Ca}^{2+}$  release. Interactions among distinct  $\text{Ca}^{2+}$  mobilizing mechanisms in starfish oocytes.** *J Biol Chem* 2000, **275**:8301-8306.
4. Cancela JM, Churchill GC, Galione A: **Coordination of agonist-induced  $\text{Ca}^{2+}$ -signalling patterns by NAADP in pancreatic acinar cells.** *Nature* 1999, **398**:74-76.
5. Cancela JM, Gerasimenko OV, Gerasimenko JV, Tepikin AV, Petersen OH: **Two different but converging messenger pathways to intracellular  $\text{Ca}^{2+}$  release: the roles of nicotinic acid adenine dinucleotide phosphate, cyclic ADP-ribose and inositol trisphosphate.** *EMBO J* 2000, **19**:2549-2557.
6. Sitsapesan R, McGarry SJ, Williams AJ: **Cyclic ADP-ribose competes with ATP for the adenine nucleotide binding site on the cardiac ryanodine receptor  $\text{Ca}^{2+}$ -release channel.** *Circ Res* 1994, **75**:596-600.
7. Genazzani AA, Mezna M, Dickey DM, Michelangeli F, Walseth TF, Galione A: **Pharmacological properties of the  $\text{Ca}^{2+}$ -release mechanism sensitive to NAADP in the sea urchin egg.** *Br J Pharmacol* 1997, **121**:1489-1495.
8. Ribeiro-do-Valle RM, Poitras M, Boulay G, Guillemette G: **The important discrepancy between the apparent affinity observed in  $\text{Ca}^{2+}$  mobilization studies and the  $K_d$  measured in binding studies is a consequence of the quantal process by which inositol 1,4,5-trisphosphate releases  $\text{Ca}^{2+}$  from bovine adrenal cortex microsomes.** *Cell Calcium* 1994 **15**:79-88.
9. Patel S, Churchill GC, Sharp T, Galione A: **Widespread distribution of binding sites for the novel  $\text{Ca}^{2+}$ -mobilizing messenger, nicotinic acid adenine dinucleotide phosphate, in the brain.** *J Biol Chem* 2000, **275**:36495-36497.
10. Lee HC, Aarhus R: **Structural determinants of nicotinic acid adenine dinucleotide phosphate important for its calcium-mobilizing activity.** *J Biol Chem* 1997, **272**:20378-20383.
11. Patel, S., G.C. Churchill, A. Galione: **Unique kinetics of nicotinic acid-adenine dinucleotide phosphate (NAADP) binding enhance the sensitivity of NAADP receptors for their ligand.** *Biochem J* 2000, **352 Pt 3**:725-729.
12. Billington RA, Genazzani AA: **Characterization of NAADP<sup>+</sup> binding in sea urchin eggs.** *Biochem Biophys Res Commun* 2000, **276**:112-116.
13. Dickey DM, Aarhus R, Walseth TF, Lee HC: **Thio-NADP is not an antagonist of NAADP.** *Cell Biochem Biophys* 1998, **28**:63-73.
14. Cik M, Masure S, Lesage AS, Van Der Linden I, Van Gompel P, Pangalos MN, et al.: **Binding of GDNF and neurturin to human GDNF family receptor alpha 1 and 2. Influence of cRET and cooperative interactions.** *J Biol Chem* 2000, **275**:27505-27512.
15. DeMeyts P, Bainco AR, Roth J: **Site-site interactions among insulin receptors. Characterization of the negative cooperativity.** *J Biol Chem* 1976, **251**:1877-1888.
16. Marsh JW, Westley J, Steiner DF: **Insulin-receptor interactions. Presence of a positive cooperative effect.** *J Biol Chem* 1984, **259**:6641-6649.
17. Harmer AR, Gallacher DV, Smith PM: **Role of  $\text{Ins}(1,4,5)\text{P}_3$ , cADP-ribose and nicotinic acid-adenine dinucleotide phosphate in  $\text{Ca}^{2+}$  signalling in mouse submandibular acinar cells.** *Biochem J* 2001, **353**:555-560.
18. Chidambaram N, Wong ET, Chang CF: **Differential oligomerization of membrane-bound CD38/ADP-ribosyl cyclase in porcine heart microsomes.** *Biochem Mol Biol Int* 1998, **44**:1225-1233.
19. Rakovic S, Churchill GC, Patel S, Webb R, Galione A, Terrar DA: **Synthesis, degradation, binding and actions of NAADP in cardiac tissue.** *Biophys J* 2001, **80**:458a.
20. Lipp P, Laine M, Tovey SC, Burrell KM, Berridge MJ, Li W, et al.: **Functional  $\text{InsP}_3$  receptors that may modulate excitation-contraction coupling in the heart.** *Curr Biol* 2000, **10**:939-942.