

# The role of phospholipase A<sub>2</sub> in lipid peroxidation-induced fall of membrane potential of rat liver mitochondria

V.G. Gogvadze, N.N. Brustovetsky and A.A. Zhukova

*Institute of Biological Physics, USSR Academy of Sciences, Pushchino 142292. USSR*

Received 13 March 1990

Cumene hydroperoxide (230  $\mu$ M)-induced fall of the membrane potential takes place only in Ca<sup>2+</sup>-loaded mitochondria. Inhibitor of phospholipase A<sub>2</sub> *p*-bromphenacyl bromide prevents uncoupling of mitochondria, having no effect on the accumulation of lipid peroxidation products.

Mitochondria; Lipid peroxidation; Calcium; Phospholipase A<sub>2</sub>

## 1. INTRODUCTION

It is well established that activation of lipid peroxidation regardless of the nature of prooxidant (Fe<sup>2+</sup>-ascorbate, organic hydroperoxides) causes the time-dependent fall of  $\Delta\phi$  and efflux of Ca<sup>2+</sup> and other cations from mitochondria [1,2]. In the absence of prooxidant the same changes occur as a result of so called 'massive' Ca<sup>2+</sup> loading in the presence of P<sub>i</sub>, and are potentiated by activation of mitochondrial Ca<sup>2+</sup>-dependent phospholipase A<sub>2</sub> and accumulation of free fatty acids and lysophospholipids in the mitochondrial membrane [3,4]. It should be mentioned that phospholipase A<sub>2</sub> activity also increases when mitochondria are peroxidized by iron and ascorbate [5]. So it is of interest to clarify the role of phospholipase A<sub>2</sub> in prooxidant induced deenergization of mitochondria. In the present paper we show that inhibition of phospholipase A<sub>2</sub> by *p*-bromphenacyl bromide prevents uncoupling of mitochondria, having no effect on the accumulation of lipid peroxidation products. The result obtained suggests that prooxidant-induced uncoupling of mitochondria is mediated by phospholipase A<sub>2</sub> activation.

## 2. MATERIALS AND METHODS

Rat liver mitochondria were prepared by the conventional procedure in 0.3 M sucrose containing 0.2 mM EGTA and 5 mM Tris-HCl, pH 7.5. EGTA was omitted from the final washing solution and

*Correspondence address:* V.G. Gogvadze, Institute of Biological Physics, USSR Academy of Sciences, Pushchino 142292, USSR

*Abbreviations:* CuOOH, cumene hydroperoxide; TPP<sup>+</sup>, tetraphenylphosphonium; MDA, malonic dialdehyde; TBA, 2-thiobarbituric acid;  $\Delta\phi$ , mitochondrial inner membrane potential; *p*-BrPhBr, *p*-bromphenacyl bromide; CCCP, carbonyl cyanide *m*-chlorophenylhydrazone

sedimented mitochondria were suspended in the same solution at 60–70 mg protein/ml. The standard incubation medium contained 0.1 M sucrose, 0.1 M KCl, 2 mM KH<sub>2</sub>PO<sub>4</sub>, 2  $\mu$ M rotenone, 2  $\mu$ M TPP<sup>+</sup> chloride, 5 mM potassium succinate and 5 mM Tris-HCl.  $\Delta\phi$  changes were evaluated by TPP<sup>+</sup> distribution between the incubation medium and the mitochondrial matrix with a TPP<sup>+</sup>-selective electrode [6]. The changes in Ca<sup>2+</sup> and O<sub>2</sub> concentrations in the incubation medium were recorded using a Ca<sup>2+</sup>-selective electrode (Orion 93-90, USA) and Clark-type electrode, respectively. Lipid peroxidation was estimated by accumulation of MDA using the TBA-test [7]. Protein concentration was determined by Lowry's method [8]. All incubations were carried out at 24°C.

## 3. RESULTS AND DISCUSSION

Fig. 1 shows that addition of CuOOH to Ca<sup>2+</sup> loaded mitochondria induced the fall of  $\Delta\phi$ . This fall was associated with rapid accumulation of lipid peroxidation products. It should be mentioned, that in the absence of prooxidant mitochondria can retain  $\Delta\phi$  for a long time (data not shown).

CuOOH-induced deenergization of mitochondria was found to be Ca<sup>2+</sup>-dependent, since addition of EGTA in the incubation medium prevented deenergization without any effect on lipid peroxidation. Taking into account the ability of Ca<sup>2+</sup> to activate mitochondrial phospholipase A<sub>2</sub>, it can be supposed that prooxidant-induced uncoupling of mitochondria is mediated by activation of enzyme. To clarify the role of phospholipase A<sub>2</sub>, we inhibited enzyme by addition of 10  $\mu$ M *p*-BrPhBr prior to CuOOH. It has been shown [9] that higher concentrations also inhibited lipid peroxidation.

As can be seen from Fig. 2, *p*-BrPhBr completely prevented the drop of  $\Delta\phi$  and Ca<sup>2+</sup> release from mitochondria (until uncoupler CCCP was added) without any decrease in MDA production.

Inhibition of phospholipase A<sub>2</sub> abolished prooxidant-induced disturbance of oxidative phosphory-




Fig. 1. Effect of EGTA on the  $CuOOH$ -induced fall of  $\Delta\phi$  and accumulation of lipid peroxidation products. Mitochondria (MCH), 1 mg/ml;  $Ca^{2+}$ , 60  $\mu M$ ;  $CuOOH$ , 230  $\mu M$ ; EGTA (curve 2, black symbols), 0.2 mM.

lation. As seen from Table I  $CuOOH$  activated State 4 respiration and reduced respiratory control coefficient in  $Ca^{2+}$ -loaded mitochondria. Addition of  $p$ -BrPhBr partially restored respiratory control and prevented  $CuOOH$ -induced acceleration of State 4 respiration.

A number of studies have demonstrated an association between lipid peroxidation and phospholipase  $A_2$




Fig. 2. Effect of  $p$ -BrPhBr on the  $CuOOH$ -induced fall of  $\Delta\phi$ ,  $Ca^{2+}$  efflux and accumulation of lipid peroxidation products. MCH, 1 mg/ml;  $Ca^{2+}$ , 60  $\mu M$ ;  $CuOOH$ , 230  $\mu M$ ;  $p$ -BrPhBr (curve 2, black symbols), 10  $\mu M$ .

activity [5,9,10]. It has been shown that activity of enzyme is dependent on the physico-chemical state of the membrane lipid phase [11]; lipid peroxidation induces alteration of membrane lipids [12], enhancing the

Table I  
Effect of  $p$ -BrPhBr on  $CuOOH$ -induced alteration of mitochondrial respiration

Additions	Rate of respiration nmol $O_2$ /min mg protein		Respiratory control coefficient
	$V_3$	$V_4$	
$Ca^{2+}$	53.6 ± 2.8	15.2 ± 1.4	3.5 ± 0.3
$Ca^{2+} + p$ -BrPhBr	55.1 ± 2.2	16.8 ± 1.2	3.3 ± 0.2
$Ca^{2+} + CuOOH$	59.2 ± 1.8	30.4 ± 2.4*	1.9 ± 0.2*
$Ca^{2+} + CuOOH + p$ -BrPhBr	58.2 ± 2.5	19.2 ± 1.4	2.7 ± 0.3

Mitochondria, 1 mg/ml;  $Ca^{2+}$ , 60  $\mu M$ ;  $CuOOH$ , 230  $\mu M$ ;  $p$ -BrPhBr, 10  $\mu M$ ; ADP, 300  $\mu M$ . Results are mean ± SE.

\* Statistically significant difference from other values,  $P < 0.05$

susceptibility of phospholipids to phospholipase A<sub>2</sub> attack [13].

Since addition of *p*-BrPhBr in incubation medium keeps mitochondria intact in spite of TBA-active product accumulation, it can be concluded that activation of phospholipase A<sub>2</sub> plays a key role in the prooxidant-induced fall of  $\Delta\phi$  and Ca<sup>2+</sup> efflux from mitochondria. The process of peroxidation appears to facilitate phospholipid hydrolysis, while Ca<sup>2+</sup> is necessary for phospholipase A<sub>2</sub> activation.

## REFERENCES

- [1] Marshansky, V.N., Novgorodov, S.A. and Yaguzhinsky, L.S. (1983) FEBS Lett. 158, 27-31.
- [2] Masini, A., Trenti, T., Ceccarelli-Stanzani, D. and Ventura, E. (1985) Biochim. Biophys. Acta 810, 27-32.
- [3] Pfeiffer, D.R., Schmid, P.C., Beatrice, M.C. and Schmid, H.H.O. (1979) J. Biol. Chem. 254, 11485-11494.
- [4] Palmer, J.W. and Pfeiffer, D.R. (1981) J. Biol. Chem. 256, 6742-6750.
- [5] Yasuda, M. and Fujita, T. (1977) Jpn. J. Pharmacol. 27, 429-435.
- [6] Kamo, N., Miratsugu, M., Hongoh, R. and Kobatake, Y. (1979) J. Membr. Biol. 49, 105-121.
- [7] Ohkawa, H., Onishi, N. and Yagi, K. (1979) Anal. Biochem. 95, 351-358.
- [8] Lowry, O., Rosebrough, N.J., Farr, A.L. and Randall, R.J. (1951) J. Biol. Chem. 193, 265-275.
- [9] Borowitz, S.M. and Montgomery, C. (1989) Biochem. Biophys. Res. Commun. 158, 1021-1028.
- [10] Ungemach, F.R. (1987) Chem. Phys. Lipids. 45, 171-205.
- [11] Verheij, H.M., Slootboom, A.J. and De Haas, G.H. (1981) Rev. Physiol. Biochem. Pharmacol. 91, 91-203.
- [12] Demopoulos, H.B., Flamm, E.S., Pietronigro, D.D. and Seligman, M. (1980) Acta Physiol. Scand. 492, 91-119.
- [13] Sevanian, A., Wratten, M.L., McLeod, L.L. and Kim, E. (1988) Biochim. Biophys. Acta 961, 316-327.