

Contents lists available at [ScienceDirect](http://ScienceDirect.com)

# Physiology & Behavior

journal homepage: [www.elsevier.com/locate/phb](http://www.elsevier.com/locate/phb)

## Salivary oxytocin mediates the association between emotional maltreatment and responses to emotional infant faces



Ritu Bhandari <sup>a,b</sup>, Marian J. Bakermans-Kranenburg <sup>a,b</sup>, Rixt van der Veen <sup>a,b</sup>, Christine E. Parsons <sup>c</sup>, Katherine S. Young <sup>c</sup>, Karen M. Grewen <sup>d</sup>, Alan Stein <sup>c</sup>, Morten L. Kringelbach <sup>c</sup>, Marinus H. van IJzendoorn <sup>a,b,\*</sup>

<sup>a</sup> Centre for Child and Family Studies, Leiden University, Leiden, Netherlands

<sup>b</sup> Leiden Institute for Brain and Cognition, Leiden University, Leiden, Netherlands

<sup>c</sup> University of Oxford, Oxford, United Kingdom

<sup>d</sup> Department of Psychiatry, University of North Carolina, Chapel Hill, NC, USA

### HIGHLIGHTS

- Emotional maltreatment is positively associated with oxytocin
- OXTR rs53576 does not moderate relation between emotional maltreatment and oxytocin
- Salivary oxytocin is associated with positive evaluation of happy expressions
- Maltreatment indirectly influences responses to happy faces by modulating oxytocin

### ARTICLE INFO

#### Article history:

Received 3 July 2013

Received in revised form 27 February 2014

Accepted 14 April 2014

Available online 24 April 2014

#### Keywords:

Oxytocin

Emotional maltreatment

Healthy females

Infant faces

Face processing

Emotion processing

### ABSTRACT

Childhood emotional maltreatment has been associated with a higher risk for maltreating one's own offspring. In the current study, we explored a possible role of oxytocin in mediating the association between childhood emotional maltreatment and participants' interpretation of infant facial expressions. Oxytocin levels were measured in 102 female participants using saliva samples. They rated the mood of thirteen infants with happy, sad and neutral facial expressions. Emotional maltreatment indirectly influenced responses to happy infant faces by modulating oxytocin levels: higher self-reported emotional maltreatment was related to higher levels of salivary oxytocin which were in turn related to a more positive evaluation of happy infant expressions, but not to the evaluation of sad infant expressions. Oxytocin receptor polymorphism rs53576 did not moderate the relation between maltreatment experiences and salivary oxytocin levels. Early emotional maltreatment might indirectly affect emotional information processing by altering the oxytonergic system.

© 2014 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/3.0/>).

### 1. Introduction

Childhood emotional maltreatment has a worldwide prevalence of 27% [1], and it has enduring effects on neural, physiological and behavioral development [2–4]. Emotional maltreatment has been associated with increased vulnerability to mental health problems such as depression and anxiety, and antisocial behavior [5,6]. In parenting, it might also lead to negative appraisal of infant signals [7] and a higher risk for maltreating one's own offspring [8–11]. In the current study, we explored a possible role of oxytocin in mediating the association

between childhood emotional maltreatment and participants' evaluation of infants' mood as derived from infants' facial emotional expressions.

Oxytocin is a neuro-peptide associated with trust, empathy and emotion recognition [12,13]. It is well known for its anxiolytic effects and its effects on prosocial behaviors such as in-group favoritism [14] and sensitive parenting [15–17]. Oxytocin levels can be affected by stressful early life experiences [18,19]. However, the direction of effects is unclear. One study demonstrated a negative relation between early life stress and oxytocin levels, with more stress leading to lower levels of oxytocin [18] whereas other studies showed a positive relation [19]. The diverging outcomes might be (partly) explained by technical differences between these studies, as the oxytocin levels were measured in different body fluids, e.g. in the cerebrospinal fluid in the study of Heim et al. [18] and in plasma in the study of Pierrehumbert et al. [19]. In the present study, we assessed oxytocin levels in saliva. Salivary

\* Corresponding author at: Centre for Child and Family Studies, Leiden University, P.O. Box 9555, 2300 RB Leiden, the Netherlands. Tel.: +31 71 5273435; fax: +31 71 5273945. E-mail address: [vanijzen@fsw.leidenuniv.nl](mailto:vanijzen@fsw.leidenuniv.nl) (M.H. van IJzendoorn).

oxytocin levels have been found to be correlated to plasma oxytocin levels (although modestly; for a review see [20]) and associated with parent and child's social engagement, affect synchrony and positive communication sequences [21]. We examined the relation between early life emotional maltreatment experiences and salivary oxytocin levels. Moreover, we tested a possible moderating role of oxytocin receptor polymorphism (OXTR) *rs53576* in the relation between emotional maltreatment and oxytocin levels [22]. In previous studies OXTR *rs53576* has been suggested to be a moderator of the effects of intranasal oxytocin administration on the preferences for infant faces as compared to adult faces. Intranasal oxytocin increased the preference for infant faces, but this effect was present only for individuals with GG genotype [22].

Using intranasal administration, oxytocin has been mostly investigated for its role in facial emotion recognition (for meta analysis see [12]). Although most of these studies show increased recognition of happy faces [23], some also report increased recognition of other emotions such as anger and fear [24]. To our knowledge, no studies have examined whether individual differences in endogenous levels of oxytocin are associated with differences in emotion recognition. Moreover, none of the previous studies assessed the association of oxytocin with recognition of emotional infant faces as opposed to adult faces. From an evolutionary perspective, oxytonergic functioning is strongly implicated in parent–infant bonding. For example, oxytocin selectively increases the preference for infant faces as compared to adult faces [16,22]. In the present study we used images of sad and happy infants to test whether salivary oxytocin levels were related to the evaluation of infant mood as derived from infants' facial emotional expressions.

Emotional maltreatment has been associated with processing biases for some negative emotional stimuli such as anger and sadness, yielding heightened responsivity and reactivity [25–28]. Impaired emotion recognition might result in more problematic interpersonal relationships [29] and more insensitive behavior towards offspring. We therefore examined the involvement of the oxytonergic system in the underlying mechanism responsible for the processing bias of emotional information in individuals with emotional maltreatment experiences.

To summarize, we hypothesized that childhood experiences of emotional maltreatment would be associated with oxytocin levels. However, given the contradictory findings from previous studies, we were not able to predict whether more maltreatment experiences would lead to higher or to lower oxytocin levels. Furthermore, we explored the potential role of OXTR *rs53576* in moderating the relation between emotional maltreatment and oxytocin levels. Moreover, in line with previous studies with intranasal administration, we hypothesized that for women with higher levels of oxytocin, infant emotional expressions would be more salient than for women with lower levels of oxytocin. Furthermore, we tested whether oxytocin mediated any association between childhood maltreatment experiences and mood ratings as derived from infants' facial expressions.

## 2. Methods

### 2.1. Participants and procedure

Three hundred and fifty three undergraduate students from the Leiden University departments of Education and Child studies and Psychology completed online questionnaires about their childhood experiences of abuse and neglect. Of these, 102 healthy female students (age  $M = 19.86$  years,  $SD = 1.42$ ) were randomly invited to participate in the present study. All were nulliparous, a majority (85%) of the participants reported being in the luteal phase of their menstrual cycle (third or fourth week following menstruation) and 74% of the participants reported using oral contraceptives. Exclusion criteria were use of steroidal or any other interfering medications such as analgesics and anti-inflammatory drugs. All participants were non-smokers and reported not having used any recreational drugs in the six months before the

experiment. Participants reported not having any current or past neurological or psychological disorder(s). Two participants were excluded from the analysis due to technical errors in the computer session. Seven participants were not included in the final analysis because of missing values of salivary oxytocin levels. The study was approved by the Leiden University Medical Center ethics committee.

Participants were invited to the laboratory for a computer session. The sessions started at 0900, 1200 or 1500 h. Participants provided saliva samples that were used to determine oxytocin levels. Next, they rated the emotion of images of faces of thirteen infants with happy, sad and neutral facial expressions. Participants also rated several infant faces following this task, which is described elsewhere [30].

### 2.2. Task and measures

#### 2.2.1. Stimuli

The stimuli consisted of grayscale pictures of thirteen infants, each of which was shown with neutral, happy and sad expressions. Infant images were obtained from a standardized database (for detailed description of the stimuli see [31]). Faces were all forward facing, with direct eye gaze, matched for size (300 × 300 pixels) and luminosity. Participants saw a facial image at the center of a 15.3 inch monitor with a visual analog scale immediately to the right. The participants were instructed to rate the mood of the presented picture from *very positive* to *very negative*. The scale ranged from 4 to –4.

Reliability analyses of the mood ratings of the neutral, happy and sad infant faces showed that the internal consistency for the happy and sad faces was high, Cronbach's  $\alpha$  (happy) = .81; Cronbach's  $\alpha$  (sad) = .84. However, the internal consistency for neutral faces' mood rating was low (Cronbach's  $\alpha$  = .49); responses to neutral facial expressions were therefore not analyzed. Two scales, one for the mood rating of the happy facial expression and one for the mood rating of the sad facial expression, were created by averaging the ratings for the thirteen infants. The scale for sad facial expression was reversed so that a higher value would represent a more negative rating.

#### 2.2.2. Salivary oxytocin

The saliva samples were immediately stored at –20 °C until batch assay. The samples were assayed using the standard procedures described in detail elsewhere, using the commercially available Enzyme Immuno Assay (EIA) kit (ADI-900-153, Enzo Life Science, Plymouth Meeting, PA) [32–34]. An extraction step was performed to concentrate the sample, increase precision and reduce matrix interference. A strata-X 33  $\mu$ m polymeric reversed phase SPE sorbent was equilibrated in a 96-well plate containing 60 mg sorbent per well, Phenomenex, Torrance CA, by adding 1 ml MeOH followed by 1 ml of water. Next, 0.8 ml of saliva was acidified with 0.4 ml of 1.5% trifluoroacetic acid (TFA) and centrifuged at 6000  $\times$ g for 20 min at 4 °C. The supernatant was loaded onto the pre-treated strata-X plate. The wells were slowly washed with 1.5 ml of 0.1% TFA, and then the peptide was eluted with 1 ml of 80% acetonitrile. The eluant was collected in a polystyrene tube and evaporated to dryness under a N<sub>2</sub> stream. The residue was reconstituted in 250  $\mu$ l of assay buffer. Oxytocin extraction efficiency was 94%, as determined by spiking with a known amount of hormone and extracting this known amount along with the samples. Oxytocin levels in extracted saliva were then quantified using the oxytocin EIA, in which the endogenous oxytocin hormone competes with exogenously added alkaline phosphatase linked oxytocin, for binding sites on oxytocin antibody. After overnight incubation at 4 °C, excess reagents were washed away and the bound oxytocin phosphatase was incubated with a substrate. After 1 h this enzyme reaction, which generates a yellow color, was stopped and the optical density (OD) was measured on a Sunrise plate reader (Tecan, Research Triangle Park, NC) at 405 nm. The intensity of the color is inversely proportional to the concentration of endogenous oxytocin. The hormone content (in pg/ml) was determined by plotting the OD of each sample

against a standard curve. One individual had extremely high oxytocin levels (>3 standard deviations above the mean). The oxytocin level values for this individual were substituted with the next highest score (Winsorized; [35]).

### 2.2.3. Genotyping

Another sample of saliva was collected using specialized DNA collection kits (Oragene DNA OG-500, DNA genotech) and was used to determine oxytocin receptor (OXTR) polymorphism, rs53576. Samples were analyzed at BaseClear laboratories, Leiden, Netherlands. DNA was extracted from the samples using the Chemagic buccal swab kit on a Chemagen Module I workstation (Chemagen Biopolymer Technologie AG, Baesweiler, Germany). DNA concentrations were measured using the Quant-iT DNA Assay kit (Invitrogen, Breda, Netherlands). The average yield was 4 µg of genomic DNA per sample. DNA was amplified in a polymerase chain reaction (PCR) using forward primer (5'-GCCACCA TGCTCTCCACATC-3') and a reverse primer (5'-GCTGGACTCAGGAGGA ATAGGGAC-3'). PCR reactions contained between 10 and 100 ng genomic DNA template and 10 pmol of forward and reverse primers. PCR was carried out in the presence of 5% dimethyl sulfoxide with 0.3 U of BioThermAB polymerase (GeneCraft, Munster, Germany) in a total volume of 30 µl using the following cycling conditions: an initial denaturation step of 3 min at 95 °C, followed by 40 cycles of 30 s at 95 °C, 30 s at 60 °C, 1 min at 72 °C and a final extension step of 3 min at 72 °C. To determine the A/G polymorphism, PCR fragments were sequenced using the forward primer and dye terminator chemistry (BigDye v3.1, Applied Biosystems).

As might be expected in this predominantly Caucasian sample (>90% Caucasian), the number of AA carriers was small [36], AA = 6, AG = 36, and GG = 51. The genotype distribution was in the Hardy-Weinberg equilibrium,  $\chi^2(1, n = 93) = 0.01, p = 0.92$ . AA and AG genotypes were combined in the analyses.

### 2.2.4. Childhood Trauma Questionnaire

Participants indicated their experiences of abuse and neglect until the age of 16 using the Childhood Trauma Questionnaire Short Form (CTQ-SF; [37]). Twenty-eight items covered experiences of physical abuse, emotional abuse, sexual abuse, physical neglect, or emotional neglect. Each item (e.g. "During my childhood I felt hated by family") was rated on a 5-point scale ranging from *never true* to *very often true*. An emotional maltreatment scale was created by averaging the items tapping into the 'emotional abuse' and 'emotional neglect' dimensions. The internal consistency of this scale was high (Cronbach's  $\alpha = .91$ ). The mean scale score was  $M = 1.49$  ( $SD = 0.48$ , range = 1–3.60). The distribution of scores was positively skewed, with all but one cases having experienced none to moderate maltreatment (range 1–3.10) and only one case (score 3.60) having extreme emotional maltreatment experiences (based on the classification suggested by the developers of CTQ). Log transformation was performed to obtain a normal distribution. The scores on the items of physical abuse and neglect and sexual abuse had a low variance and skewed distribution. These scales were not used in the current study.

### 2.3. Statistical analyses

Bivariate associations between the variables were computed. Participants' use of contraceptives and the phase of menstrual cycle were recorded but as they were not associated with any of the independent or dependent variables (Table 1), they were not co-varied in the analyses. Two mediation analyses were performed using a macro package for SPSS, available on-line (<http://www.afhayes.com/spss-sas-and-mplus-macros-and-code.html>) developed by Preacher and Hayes [38]. The mood ratings of the happy expression and sad expression were the dependent variables. Log transformed emotional maltreatment was entered as the independent variable and salivary oxytocin level as the mediator. We also tested whether the relation between emotional maltreatment and oxytocin levels was moderated by OXTR polymorphism. Centered predictors (emotional maltreatment and OXTR rs53576 genotype) were entered first in a multiple regression and the interaction term of the polymorphism with emotional maltreatment was added in the next step.

## 3. Results

### 3.1. Bivariate associations

Table 1 presents the bivariate correlations between the variables. The mood ratings of happy infant faces and sad infant faces were correlated ( $r = .53, p < .001$ ). Participants who rated the mood of the happy faces as more positive rated the mood of sad faces as more negative. Childhood experience of emotional maltreatment was correlated with salivary oxytocin levels ( $r = .25, p = .02$ ). Participants who experienced more emotional maltreatment had elevated levels of oxytocin. Oxytocin levels were associated with the mood ratings of happy infant faces ( $r = .33, p = .001$ ). Participants with higher oxytocin levels rated the mood of the happy faces as more positive than participants with lower oxytocin levels. The association between oxytocin level and mood rating of the sad faces was not significant ( $r = .16, p = .14$ ). OXTR polymorphism was not associated with the mood ratings (all  $p > .05$ ).

### 3.2. Multivariate analysis


Results of the mediation analysis with the mood rating of the happy infants as the dependent variable showed that the indirect path from emotional maltreatment through oxytocin level to mood rating was significant ( $b = 0.33, SE = 0.15, CI = .08, .69$ ). Emotional maltreatment was positively related to oxytocin level ( $b = 2.75, SE = 1.12, p = .02$ ), and oxytocin levels were positively related to the mood ratings of happy infants ( $b = 0.12, SE = 0.03, p < .001$ ). The direct path between emotional maltreatment and infant mood rating was not significant ( $b = -0.09, SE = 0.39, p = .82$ ). The strength of this path increased somewhat after controlling for oxytocin levels, but it did not reach significance ( $b = -0.12, SE = 0.38, p = .27$ ). This shows that the effects of emotional maltreatment on infant mood rating were indirect and emotional maltreatment appears to exert its influence by

**Table 1**

Descriptive statistics and bivariate correlations between the predictors and ratings of infant facial expressions.

	Range	Mean (SD)	1	2	3	4	5
1. Oxytocin receptor rs53576	GG/AA;AG	55% (GG)		0.00	0.05	−0.01	−0.01
2. Oxytocin levels (pg/ml)	0.90 to 7.00	2.65 (1.32)			0.25*	0.33***	0.16
3. Emotional maltreatment (Log)	0.00 to 0.56	0.16 (0.12)				−0.02	0.15
4. Mood rating (happy expression)	1.53 to 3.93	2.63 (.45)					.53***
5. Mood rating (sad expression)	1.30 to 3.70	2.48 (.46)					

Note. \*  $p < .05$ ; \*\*  $p < .01$ ; \*\*\*  $p < .001$ .



**Fig. 1.** Indirect effect of childhood emotional maltreatment on the mood rating of happy infant faces is mediated by oxytocin. The overall indirect effect is significant,  $b = 0.33$ ,  $SE = .15$ ,  $CI = .08, .69$ . Note. \*  $p < .05$ , \*\*\*  $p < .001$ .  $\beta$ -values are presented in the figure.

modulating oxytocin levels (see Fig. 1 in which  $\beta$  values are presented). For the mood ratings of the infant faces with sad expressions, the indirect path from emotional maltreatment through oxytocin level was not significant ( $b = 0.12$ ,  $SE = 0.11$ ,  $CI = -.02, .42$ ). The direct path between emotional maltreatment and negative mood rating was also not significant ( $b = 0.57$ ,  $SE = 0.39$ ,  $p = .15$ ).

We performed a hierarchical multiple regression analysis to examine whether the relation between emotional maltreatment and oxytocin levels was moderated by the OXTR genotype. The main effect of the OXTR genotype (GG vs. AG/AA) on oxytocin levels was not significant ( $b = -0.02$ ,  $SE = 0.14$ ,  $p = .90$ ). The interaction between emotional maltreatment and OXTR genotype was not significant either ( $b = -0.45$ ,  $SE = 1.21$ ,  $p = .71$ ), showing that the relation between emotional maltreatment and oxytocin level was not moderated by OXTR *rs53576*.

#### 4. Discussion

More emotional maltreatment experiences predicted higher salivary oxytocin levels. Moreover, as expected, higher salivary oxytocin levels compared to lower oxytocin levels were associated with more positive ratings of infants' mood when happy infant faces were shown, indicating better recognition of happy infant faces. Our results suggest that having been emotionally maltreated may indirectly influence responses to emotional infant faces by modulating oxytocin levels.

Elevated oxytocin levels in participants with more emotional maltreatment experiences are in line with a study on sexually abused females (with mixed gender control subjects) showing higher plasma oxytocin levels in individuals with more maltreatment experiences [19]. A recent study with older children also showed higher urinary oxytocin levels in girls who experienced physical abuse as compared to physically abused boys and non-maltreated age-matched controls [39]. Our finding of an association between maltreatment and salivary oxytocin is however in contrast with a small pilot study ( $N = 22$ ) showing decreased cerebrospinal oxytocin levels in females (of mixed ethnicity) with a history of maltreatment, in particular emotional abuse [18]. Our study differs from the above mentioned studies in several respects. The source of oxytocin levels varies across the studies (concentrations in cerebrospinal fluid, plasma, urine or saliva), as did the type of abuse (sexual versus emotional maltreatment), whereas variations in gender and ethnic differences may also play a role. Moreover, our study included a non-clinical sample of female university students. It is not clear whether the previous studies used participants with clinical or non-clinical post traumatic symptoms of abuse.

Experiments with intranasally administered oxytocin have shown that recognition of various emotions at low intensities is enhanced after oxytocin administration [12,23,24]. In the present correlational study participants rated the mood of emotional infant faces. Higher levels of oxytocin appeared to be associated with more positive perceptions of infant mood when viewing happy infant expressions, which is generally consistent with experimental evidence of enhanced emotion recognition after intranasal administration of oxytocin. Of course, salivary oxytocin levels might not be direct reflections of oxytocin levels in brain regions relevant for empathic concern, an issue that also

remains to be solved in experimental oxytocin studies on humans [40], but see [41] for evidence in rodents).

Our results might provide some insight into the mechanisms by which oxytocin leads to more positive caretaking behaviors [16]. Oxytocin has been proposed to facilitate interpersonal trust by increasing sensitivity to happiness cues of adult faces [23]. Similarly, more positive appraisals of infant cues might represent an increased reward value of infant stimuli for the caretaker [42], thereby facilitating more sensitive parenting responses towards the infants.

Previous studies with adult faces have proposed various mechanisms explaining the effects of oxytocin on enhanced emotion recognition, such as modulation of eye gaze [24,43,44], decreased effortful processing of affective information [45], wider pupil dilatation [46], or decreased amygdala activation [47,48]. Oxytocin not only reduces amygdala activation but also enhances functional connectivity between the amygdala and the orbitofrontal cortex, the anterior cingulate, the hippocampus, the precuneus, the supramarginal gyri, and the middle temporal gyrus [42]. Increased functional connectivity between the amygdala and these brain regions involved in emotion regulation may reduce negative emotional arousal while enhancing the incentive salience of the infant laughter [42,49]. Similar enhanced neural connectivity might be present or elicited more easily in those individuals who have higher naturally produced oxytocin levels when happy infant faces are processed.

Emotional maltreatment has been associated with difficulty in discriminating emotional expressions [26]. In the present study, we found evidence for an indirect effect of emotional maltreatment on the perception of emotional infant faces via elevated oxytocin levels: more emotional maltreatment was associated with higher levels of salivary oxytocin which were related to a more positive assessment of happy infant expression. Various animal and human studies have shown increased oxytocin levels in response to stressors [50,51]. Higher oxytocin levels have been found in females with more attachment anxiety [51,52]. The anxiolytic role of oxytocin [53,54] might explain why emotionally maltreated individuals 'need' more of it to cope with stressful events and situations which may contribute to resilience in some individuals [55,56].

We did not find evidence for a role of OXTR *rs53576* in influencing salivary oxytocin levels, in this relatively small sample. The polymorphism did not moderate the relation between maltreatment experiences and salivary oxytocin levels either. To our knowledge, only one study examined the association between OXTR polymorphism *rs2254298* and plasma oxytocin levels, showing that the presence of the minor A allele was associated with higher oxytocin levels [57]. A recent meta-analysis however failed to find a significant association between the two most popular OXTR polymorphisms (*rs53576* and *rs2254298*) and various social and behavioral outcomes [36]. The current study supports the general conclusion of the meta-analysis that the functional role of OXTR *rs53576* has not yet been established.

There were a number of limitations in this study. Firstly, our sample consisted of nulliparous female university students who did not report many severe past or current post-traumatic symptoms of emotional maltreatment. Generalizing our findings to experienced mothers, to males, or to individuals with clinical levels of post traumatic symptoms because of emotional maltreatment is not possible based on this sample. Oxytocin may follow an inverted U-shape relation (as shown between plasma oxytocin and trust, [58]) resulting in lower oxytocin levels with severe maltreatment experiences. Such a relation with maltreatment, if present, might show a different pattern of results compared to the findings presented here. Nevertheless clinical levels of post traumatic symptoms might not be needed to trigger depressive or neglectful parenting responses in individuals with child emotional maltreatment experiences, and the current results might thus be applicable to a more general, non-clinical population. Secondly, oxytocin levels were assessed in saliva, which might not reflect the actual working oxytocin levels in the brain. Studies using intranasal oxytocin

administration however provide some evidence for the association between central oxytocin levels and salivary oxytocin levels [40,59], and for the association between intranasally elevated levels of oxytocin (also reflected in salivary levels; [34]) and altered resting state brain activity [49]. Moreover, differences in studies due to different assay methods (for e.g. extraction methods for reducing interference and radio- versus enzyme-immunoassay techniques) as well as the controversies regarding the differences and similarities between oxytocin levels measured in various body fluids such as plasma, cerebrospinal fluid and urine should be taken into account while interpreting these findings and should be addressed in future studies [20]. Finally, the moderating effect of social support on the relation between emotional maltreatment and oxytocin levels might be important and should be examined in future work. Social support has been shown to be associated with oxytocin levels as well as with sensitivity in infant caretaking [32, 60,61], and is also an important factor in post-traumatic coping [56].

To summarize, our findings suggest that experiences of childhood emotional maltreatment may alter salivary oxytocin levels, which in turn are related to more positive perceptions of infant stimuli with positive emotion. Through this indirect effect, early emotional maltreatment might influence the processing of emotional information, although the causal relationship implied in this interpretation would need to be the subject of further investigation.

### Role of funding source

MJB-K and MHvIJ were supported by research awards from the Netherlands Organization for Scientific Research (MHvIJ: NWO SPINOZA prize; MJBK: VICI grant no. 453-09-003). MLK was supported by funding from the TrygFonden Charitable Foundation and KSY was supported by a Medical Research Council Studentship. The funding source had no input in the design and conduct of this study or in any other aspect of this work.

### Conflict of interest

None of the participating institutions and authors reported biomedical financial interests or potential conflicts of interest regarding this study.

### Acknowledgments

We thank Lisanne Marijs and Renée Smulders for their contributions to data-collection.

### References

- [1] Stoltenborgh M, Bakermans-Kranenburg MJ, Alink LRA, Van IJzendoorn MH. The universality of childhood emotional abuse: a meta-analysis of worldwide prevalence. *J Aggress Maltreat Trauma* 2012;21:870–90.
- [2] Spinhoven P, Elzinga BM, Hovens JGFM, Roelofs K, Zitman FG, van Oppen P, et al. The specificity of childhood adversities and negative life events across the life span to anxiety and depressive disorders. *J Affect Disord* 2010;126:103–12.
- [3] Van Harmelen A-L, de Jong PJ, Glaschouwer KA, Spinhoven P, Penninx BWJH, Elzinga BM. Child abuse and negative explicit and automatic self-associations: the cognitive scars of emotional maltreatment. *Behav Res Ther* 2010;48:486–94.
- [4] Van Harmelen A-L, Van Tol M-J, Demenescu LR, van der Wee NJA, Veltman DJ, Aleman A, et al. Enhanced amygdala reactivity to emotional faces in adults reporting childhood emotional maltreatment. *Soc Cogn Affect Neurosci* 2013;8:362–9.
- [5] Egeland B, Yates T, Appleyard K, van Dulmen M. The long-term consequences of maltreatment in the early years: a developmental pathway model to antisocial behavior. *Child Serv* 2002;5:249–60.
- [6] Gibb BE, Chelminski I, Zimmerman M. Childhood emotional, physical, and sexual abuse, and diagnoses of depressive and anxiety disorders in adult psychiatric outpatients. *Depression Anxiety* 2007;24:256–63.
- [7] Crouch JL, Skowronski JJ, Milner JS, Harris B. Parental responses to infant crying: the influence of child physical abuse risk and hostile priming. *Child Abuse Negl* 2008;32:702–10.
- [8] Berlin LJ, Appleyard K, Dodge KA. Intergenerational continuity in child maltreatment: mediating mechanisms and implications for prevention. *Child Dev* 2011;82:162–76.
- [9] Belsky J, Vondra J. Lessons from child abuse: The determinants of parenting. In: Carlson DCV, editor. *Child maltreatment: theory and research on the causes and consequences of child abuse and neglect*. New York, NY, US: Cambridge University Press; 1989. p. 153–202.
- [10] Van IJzendoorn MH. Intergenerational transmission of parenting: a review of studies in nonclinical populations. *Dev Rev* 1992;12:76–99.
- [11] Dixon L, Browne K, Hamilton-Giachritsis C. Risk factors of parents abused as children: a mediational analysis of the intergenerational continuity of child maltreatment (Part I). *J Child Psychol Psychiatry* 2005;46:47–57.
- [12] Van IJzendoorn MH, Bakermans-Kranenburg MJ. A sniff of trust: meta-analysis of the effects of intranasal oxytocin administration on face recognition, trust to in-group, and trust to out-group. *Psychoneuroendocrinology* 2012;37:438–43.
- [13] MacDonald K, MacDonald TM. The peptide that binds: a systematic review of oxytocin and its prosocial effects in humans. *Harv Rev Psychiatry* 2010;18:1–21.
- [14] De Dreu CKW, Greer LL, Handgraaf MJJ, Shalvi S, Van Kleef GA, Baas M, et al. The neuropeptide oxytocin regulates parochial altruism in intergroup conflict among humans. *Science* 2010;328:1408–11.
- [15] Feldman R. Oxytocin and social affiliation in humans. *Horm Behav* 2012;61:380–91.
- [16] Feldman R, Weller A, Zagoory-Sharon O, Levine A. Evidence for a neuroendocrinological foundation of human affiliation: plasma oxytocin levels across pregnancy and the postpartum period predict mother–infant bonding. *Psychol Sci* 2007;18:965–70.
- [17] Naber F, Poslowsky I, Van IJzendoorn MH, Engeland H, Bakermans-Kranenburg MJ. Brief report: oxytocin enhances paternal sensitivity to a child with autism: a double-blind within-subject experiment with intranasally administered oxytocin. *J Autism Dev Disord* 2013;43:224–9.
- [18] Heim C, Young LJ, Newport DJ, Mletzko T, Miller AH, Nemeroff CB. Lower CSF oxytocin concentrations in women with a history of childhood abuse. *Mol Psychiatry* 2008;14:954–8.
- [19] Pierrehumbert B, Torrisi R, Lauffer D, Halfon O, Ansermet F, Beck Popovic M. Oxytocin response to an experimental psychosocial challenge in adults exposed to traumatic experiences during childhood or adolescence. *Neuroscience* 2010;166:168–77.
- [20] McCullough ME, Churchland PS, Mendez AJ. Problems with measuring peripheral oxytocin: can the data on oxytocin and human behavior be trusted? *Neurosci Biobehav Rev* 2013;37:1485–92.
- [21] Feldman R, Gordon I, Zagoory-Sharon O. Maternal and paternal plasma, salivary, and urinary oxytocin and parent–infant synchrony: considering stress and affiliation components of human bonding. *Dev Sci* 2011;14:752–61.
- [22] Marsh A, Yu H, Pine D, Gorodetsky E, Goldman D, Blair RJR. The influence of oxytocin administration on responses to infant faces and potential moderation by OXTR genotype. *Psychopharmacology* 2012;224:469–76.
- [23] Marsh A, Yu H, Pine D, Blair RJR. Oxytocin improves specific recognition of positive facial expressions. *Psychopharmacology* 2010;209:225–32.
- [24] Lischke A, Berger C, Prehn K, Heinrichs M, Herpertz SC, Domes G. Intranasal oxytocin enhances emotion recognition from dynamic facial expressions and leaves eye-gaze unaffected. *Psychoneuroendocrinology* 2012;37:475–81.
- [25] Pollak SD, Cicchetti D, Hornung K, Reed A. Recognizing emotion in faces: developmental effects of child abuse and neglect. *Dev Psychol* 2000;36:679–88.
- [26] Pollak SD, Cicchetti D, Klorman R, Brumaghim JT. Cognitive brain event-related potentials and emotion processing in maltreated children. *Child Dev* 1997;68:773–87.
- [27] Masten CL, Guyer AE, Hodgson HB, McClure EB, Charney DS, Ernst M, et al. Recognition of facial emotions among maltreated children with high rates of post-traumatic stress disorder. *Child Abuse Negl* 2008;32:139–53.
- [28] Curtis WJ, Cicchetti D. Affective facial expression processing in young children who have experienced maltreatment during the first year of life: an event-related potential study. *Dev Psychopathol* 2011;23:373–95.
- [29] Crawford E, Wright MOD. The impact of childhood psychological maltreatment on interpersonal schemas and subsequent experiences of relationship aggression. *J Emot Abuse* 2007;7:93–116.
- [30] Parsons CE, Young KS, Bhandari R, van IJzendoorn MH, Bakermans-Kranenburg MJ, Stein A. The bonnie baby: experimentally manipulated temperament affects perceived cuteness and motivation to view infant faces. *Dev Sci* 2014;17:257–69.
- [31] Kringelbach ML, Lehtonen A, Squire S, Harvey AG, Craske MG, Holliday IE, et al. A specific and rapid neural signature for parental instinct. *PLoS One* 2008;3:e1664.
- [32] Holt-Lunstad J, Birmingham WA, Light KC. Influence of a “warm touch” support enhancement intervention among married couples on ambulatory blood pressure, oxytocin, alpha amylase, and cortisol. *Psychosom Med* 2008;70:976–85.
- [33] Grewen KM, Davenport RE, Light KC. An investigation of plasma and salivary oxytocin responses in breast- and formula-feeding mothers of infants. *Psychophysiology* 2010;47:625–32.
- [34] Van IJzendoorn MH, Bhandari R, Van der Veen R, Grewen K, Bakermans-Kranenburg MJ. Elevated salivary levels of oxytocin persist more than seven hours after intranasal administration. *Front Neurosci* 2012;6.
- [35] Tabachnick BG, Fidell LS. *Using multivariate statistics*. 5th ed. Allyn & Bacon, Inc; 2006.
- [36] Bakermans-Kranenburg MJ, van IJzendoorn MH. A sociability gene? Meta-analysis of oxytocin receptor (OXTR) genotype effects in humans. *Psychiatr genet* 2014;24:45–51.
- [37] Bernstein DP, Stein JA, Newcomb MD, Walker E, Pogge D, Ahluvalia T, et al. Development and validation of a brief screening version of the Childhood Trauma Questionnaire. *Child Abuse Negl* 2003;27:169–90.
- [38] Preacher K, Hayes A. Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. *Behav Res Methods* 2008;40:879–91.
- [39] Seltzer LJ, Ziegler T, Connolly MJ, Proskoski AR, Pollak SD. Stress-induced elevation of oxytocin in maltreated children: evolution, neurodevelopment, and social behavior. *Child Dev* 2014;85:501–12.

- [40] Guastella AJ, MacLeod C. A critical review of the influence of oxytocin nasal spray on social cognition in humans: evidence and future directions. *Horm Behav* 2012;61:410–8.
- [41] Neumann ID, Maloumy R, Beiderbeck DI, Lukas M, Landgraf R. Increased brain and plasma oxytocin after nasal and peripheral administration in rats and mice. *Psychoneuroendocrinology* 2013;38:1985–93.
- [42] Riem MME, Van IJzendoorn MH, Tops M, Boksem MAS, Rombouts SAR, Bakermans-Kranenburg MJ. No laughing matter: intranasal oxytocin administration changes functional brain connectivity during exposure to infant laughter. *Neuropsychopharmacology* 2012;37:1257–66.
- [43] Domes G, Steiner A, Porges SW, Heinrichs M. Oxytocin differentially modulates eye gaze to naturalistic social signals of happiness and anger. *Psychoneuroendocrinology* 2012;38:1198–202.
- [44] Guastella AJ, Mitchell PB, Dadds MR. Oxytocin increases gaze to the eye region of human faces. *Biol Psychiatry* 2008;63:3–5.
- [45] Ellenbogen MA, Linnen A-M, Grumet R, Cardoso C, Joobar R. The acute effects of intranasal oxytocin on automatic and effortful attentional shifting to emotional faces. *Psychophysiology* 2012;49:128–37.
- [46] Prehn K, Kazzer P, Lischke A, Heinrichs M, Herpertz SC, Domes G. Effects of intranasal oxytocin on pupil dilation indicate increased salience of socioaffective stimuli. *Psychophysiology* 2013;50:528–37.
- [47] Petrovic P, Kalisch R, Singer T, Dolan RJ. Oxytocin attenuates affective evaluations of conditioned faces and amygdala activity. *J Neurosci* 2008;28:6607–15.
- [48] Domes G, Heinrichs M, Gläscher J, Büchel C, Braus DF, Herpertz SC. Oxytocin attenuates amygdala responses to emotional faces regardless of valence. *Biol Psychiatry* 2007;62:1187–90.
- [49] Riem MME, Van IJzendoorn MH, Tops M, Boksem MAS, Rombouts SAR, Bakermans-Kranenburg MJ. Oxytocin effects on complex brain networks are moderated by experiences of maternal love withdrawal. *Eur Neuropsychopharmacol* 2013;23:1288–95.
- [50] Tops M, Van Peer JM, Korf J, Wijers AA, Tucker DM. Anxiety, cortisol, and attachment predict plasma oxytocin. *Psychophysiology* 2007;44:444–9.
- [51] Taylor SE, Saphire-Bernstein S, Seeman TE. Are plasma oxytocin in women and plasma vasopressin in men biomarkers of distressed pair-bond relationships? *Psychol Sci* 2010;21:3–7.
- [52] Weisman O, Zagoory-Sharon O, Schneiderman I, Gordon I, Feldman R. Plasma oxytocin distributions in a large cohort of women and men and their gender-specific associations with anxiety. *Psychoneuroendocrinology* 2013;38:694–701.
- [53] Neumann ID, Landgraf R. Balance of brain oxytocin and vasopressin: implications for anxiety, depression, and social behaviors. *Trends Neurosci* 2012;35:649–59.
- [54] Ditzen B, Schaer M, Gabriel B, Bodenmann G, Ehlert U, Heinrichs M. Intranasal oxytocin increases positive communication and reduces cortisol levels during couple conflict. *Biol Psychiatry* 2009;65:728–31.
- [55] DuMont KA, Widom CS, Czaja SJ. Predictors of resilience in abused and neglected children grown-up: the role of individual and neighborhood characteristics. *Child Abuse Negl* 2007;31:255–74.
- [56] Feder A, Nestler EJ, Charney DS. Psychobiology and molecular genetics of resilience. *Nat Rev Neurosci* 2009;10:446–57.
- [57] Feldman R, Zagoory-Sharon O, Weisman O, Schneiderman I, Gordon I, Maoz R, et al. Sensitive parenting is associated with plasma oxytocin and polymorphisms in the OXTR and CD38 genes. *Biol Psychiatry* 2012;72:175–81.
- [58] Zhong S, Monakhov M, Mok HP, Tong T, Lai PS, Chew SH, et al. U-shaped relation between plasma oxytocin levels and behavior in the trust game. *PLoS One* 2012;7:e51095.
- [59] Weisman O, Zagoory-Sharon O, Feldman R. Intranasal oxytocin administration is reflected in human saliva. *Psychoneuroendocrinology* 2012;37:1582–6.
- [60] Quittner AL, Glueckauf RL, Jackson DN. Chronic parenting stress: moderating versus mediating effects of social support. *J Pers Soc Psychol* 1990;59:1266–78.
- [61] Grewen KM, Girdler SS, Amico J, Light KC. Effects of partner support on resting oxytocin, cortisol, norepinephrine, and blood pressure before and after warm partner contact. *Psychosom Med* 2005;67:531–8.