

The MicroRNA-Regulated SBP-Box Transcription Factor SPL3 Is a Direct Upstream Activator of *LEAFY*, *FRUITFULL*, and *APETALA1*

Ayako Yamaguchi,¹ Miin-Feng Wu,¹ Li Yang,¹ Gang Wu,¹ R. Scott Poethig,¹ and Doris Wagner^{1,*}

¹Department of Biology, University of Pennsylvania, Philadelphia, PA 19104-6018, USA

*Correspondence: wagnerdo@sas.upenn.edu

DOI 10.1016/j.devcel.2009.06.007

SUMMARY

When to form flowers is a developmental decision that profoundly impacts the fitness of flowering plants. In *Arabidopsis* this decision is ultimately controlled by the induction and subsequent activity of the transcription factors *LEAFY* (*LFY*), *FRUITFULL* (*FUL*), and *APETALA1* (*AP1*). Despite their central importance, our current understanding of the regulation of *LFY*, *FUL*, and *AP1* expression is still incomplete. We show here that all three genes are directly activated by the microRNA-targeted transcription factor SQUAMOSA PROMOTER BINDING PROTEIN-LIKE 3 (*SPL3*). Our findings suggest that *SPL3* acts together with other microRNA-regulated SPL transcription factors to control the timing of flower formation. Moreover, the identified SPL activity defines a distinct pathway in control of this vital developmental decision.

INTRODUCTION

Flowering plants are sessile organisms that use environmental as well as endogenous cues to optimize the progression through developmental phases. During each developmental phase, distinct structures form from the primordia at the flanks of the shoot apical meristem (Araki, 2001; Poethig, 2003; Steeves and Sussex, 1989), which differ in morphology and function. Developmental phases and the transitions between them have been well studied in the reference plant *Arabidopsis thaliana*. In this species, three main phase transitions can be distinguished: the vegetative phase change, the reproductive or floral transition, and the meristem identity (MI) transition. After germination, the first primordia formed give rise to juvenile leaves. Following the vegetative phase change, adult leaves are formed instead, which differ from juvenile leaves in many traits (Poethig, 2003). Production of these leaves, collectively referred to as rosette leaves, ceases following the reproductive transition, which is regulated by several independent flowering time pathways (reviewed in Kobayashi and Weigel, 2007; Turck et al., 2008). After this transition, the primary inflorescence grows upward (bolts) and primordia give rise to secondary inflorescence branches subtended by cauline leaves (Araki, 2001; Steeves and Sussex, 1989). Finally, the MI transition controls the onset of the last

developmental phase, in which the primordia give rise to the reproductive structures, the flowers (Blazquez et al., 2006).

Proper timing of the MI transition is important for fitness of flowering plants (Roux et al., 2006). This transition is controlled by two types of positive regulators: the plant-specific transcription factor *LFY* and members of the *AP1/FUL* (formerly *AGL8*) clade of MADS-box transcription factors (Benlloch et al., 2007; Litt and Irish, 2003). The role of these regulators is conserved in different plant species (Benlloch et al., 2007). Studies in *Arabidopsis* have shown that *LFY*, *FUL*, and *AP1* expression increases just prior to the MI transition, with *LFY* and *FUL* activated very early and *AP1* upregulation occurring later (for example see Hempel et al., 1997). The expression of the three MI genes overlaps at the sites where flowers are specified, the inflorescence meristem and associated primordia (reviewed in Kobayashi and Weigel, 2007; Turck et al., 2008). *LFY* is considered a master regulator of the MI transition since loss of *LFY* function causes the most dramatic delay in flower formation (Weigel et al., 1992). *ful* single mutants exhibit a significant delay in the MI transition (Ferrandiz et al., 2000; Melzer et al., 2008). *FUL* acts together with other MADS-box transcription factors including *AP1* to regulate this process (Ferrandiz et al., 2000). *ap1* single mutants also cause a delay of the MI transition, and *AP1* plays an important role in this process both downstream of and together with *LFY* (Bowman et al., 1993). Finally, elevated levels of either *LFY*, *AP1*, or *FUL* cause a precocious MI transition, suggesting that upregulation of any one of these regulators is sufficient to trigger this developmental switch (Ferrandiz et al., 2000; Mandel and Yanofsky, 1995; Weigel and Nilsson, 1995). Proper control of the *LFY*, *FUL*, and *AP1* accumulation is therefore key for the correct timing of the onset of flower formation.

Despite their importance in regulation of a vital developmental transition, our understanding of the regulation of *LFY*, *FUL*, and *AP1* expression is incomplete. Recently, microRNA-regulated SQUAMOSA PROMOTER BINDING PROTEIN (SBP)-box transcription factors have been implicated in regulation of multiple developmental transitions in *Arabidopsis* and other plant species (Cardon et al., 1997; Chuck et al., 2007; Gandikota et al., 2007; Schwarz et al., 2008; Wu and Poethig, 2006; Xie et al., 2006). Consistent with their role in developmental timing, transcript accumulation of several members of this family strongly increases during development (Cardon et al., 1999; Schmid et al., 2003; Wu and Poethig, 2006). Concomitantly, levels of the microRNA miR156 that specifically targets these genes decreases (Schwab et al., 2005; Wu and Poethig, 2006). How the SBP transcription factors exert their roles in developmental




Figure 1. SPL3 Activates Meristem Identity Gene Expression

(A) Schematic representation of the known interactions involved in the meristem identity (MI) switch (see text for details). Flowering time regulators are indicated in orange, MI regulators in pink. Solid arrows denote direct interactions; dashed arrows denote direct or indirect interactions.

(B–D) qRT-PCR analysis of flowering time gene (B) or MI gene expression (C and D) in 35S:SPL3Δ (SPL3OX) compared to wild-type plants. Plants were grown in short-day (SD) for 4 and 10 days (B and C) or in long-day (LD) for 7 and 9 days (D).

(E) Temporal expression of *SPL3*, *LFY*, *FUL*, and *AP1* in wild-type plants grown in LD. The ratio of the expression at each time point over the final expression level is graphed.

(B–E) Values are mean \pm SEM. Black solid horizontal line, 2-fold increase; black dotted line, 2-fold decrease.

(F) Confocal image of pSPL3:GFP-SPL3 (top view of inflorescence meristem) and in situ hybridization of *LFY*, *FUL*, and *AP1* (longitudinal section through inflorescence meristem). Arrowheads point to the incipient flower primordium; numbers denote stages of flower development. Bar: 50 μ m.

transitions is an area of intense investigation. We show here that one well-studied member of this family, SPL3, directly activates *LFY*, *FUL*, and *AP1* expression in *Arabidopsis*, identifying a direct molecular link between SPL transcription factors and one of the developmental transitions they regulate. We further show that SPL3, together with additional microRNA-regulated SPL transcription factors, is required for *LFY*, *FUL*, and *AP1* upregulation during the MI transition, implicating SPL transcription factors as a central node in the regulatory network that controls flower formation.

RESULTS

SPL3 Upregulates *LFY*, *FUL*, and *AP1*

Mutations in the microRNA-regulated transcription factor SPL3 cause no visible phenotype, most likely because of functional

redundancy with other related SPL proteins in *Arabidopsis* (Wang et al., 2008; Wu and Poethig, 2006). To examine the role of SPL3 in the MI transition, we constitutively expressed the *SPL3* mRNA without the microRNA target site in the 3'UTR (35S:SPL3Δ; Wu and Poethig, 2006) using a gain-of-function approach. 35S:SPL3Δ plants exhibited a precocious reproductive transition (fewer rosette leaves formed) and MI transition (fewer secondary inflorescences and cauline leaves formed) compared to the wild-type in both long-day and short-day conditions (see Table S1 available online).

The rapid MI transition in 35S:SPL3Δ could be due to upregulation of upstream regulators of *LFY*, *FUL*, or *AP1* (Figure 1A). To test this possibility, we examined expression of known activators of the MI genes. We scored two time points (day 4 and day 10) in short-day growth conditions to assess the temporal upregulation of genes in 35S:SPL3Δ compared to the wild-type. Both time points precede the MI transition, which occurs at day 25 in short day (data not shown). The MADS box transcription factors and flowering time regulators SUPPRESSOR OF OVEREXPRESSION OF CO1 (SOC1) and AGAMOUS-LIKE 24 (AGL24) are direct upstream transcriptional activators of *LFY* (Lee et al., 2008; Liu et al., 2008). Neither *AGL24* nor *SOC1* expression was strongly elevated in short-day grown 35S:SPL3Δ seedlings (Figure 1B). We note that *AGL24* levels are reduced in 35S:SPL3Δ; the reason for this reduction is currently not clear. The plant hormone gibberellin induces *LFY* expression through GAMYB transcription factors in short day (Achard et al., 2004; Blazquez and Weigel, 2000). Expression of *MYB33*, which encodes one of the GAMYB transcription factors known to bind the *LFY* promoter in vitro (Gocal et al., 2001), was unaltered in 35S:SPL3Δ (Figure S1). FLOWERING LOCUS T (FT) is a known photoperiod pathway flowering time regulator that acts upstream of *AP1* and *FUL* (Abe et al., 2005; Teper-Bamnolker and Samach, 2005; Wigge et al., 2005). *FT* expression was not much altered in 35S:SPL3Δ compared to the wild-type (Figure 1B). Expression of *FD*, which encodes a bZIP transcription factor that acts together with FT upstream of *AP1* and *FUL*, was slightly increased (1.8-fold) at the later time point, in 10-day-old short-day-grown 35S:SPL3Δ plants. Hence, expression of the flowering time regulators was not strongly upregulated in 35S:SPL3Δ with the possible exception of *FD*.

By contrast, when we examined expression of *LFY*, *FUL*, and *AP1* in the same conditions, we observed a strong increase in *LFY* levels (2.6-fold at day 4 and 4.4-fold at day 10) and *FUL* expression (greater than 25-fold at both time points) in 35S:SPL3Δ compared to wild-type seedlings (Figure 1C). *AP1* upregulation was only observed at day 10 (8-fold increase, Figure 1C). The combined data suggest that SPL3 activates *LFY*, *FUL*, and *AP1*. To further test whether *LFY*, *FUL*, and *AP1* were also upregulated by SPL3 in different growth conditions, we examined their expression in inductive photoperiod (continuous light) in 35S:SPL3Δ compared to the wild-type (Figure 1D). Again, the two time points chosen (day 7 and day 9) precede the MI transition, which occurs at day 11 under these growth conditions (Figure 1E, first upregulation of *AP1*). We observed a very similar increase in *LFY* and *FUL* expression in inductive photoperiod to that observed in short-day conditions (compare Figure 1C to Figure 1D). In addition, *AP1* levels were strongly increased at both time points tested (Figure 1D). Thus SPL3 may regulate the MI transition via upregulation of *LFY*, *FUL*, and *AP1*.

Table 1. Effect of Loss of Function of Meristem Identity Genes on Phenotype of 35S:SPL3Δ

Genotype	Number of Rosette Leaves	Number of Cauline Leaves	Number of Secondary Inflorescences
Wild-type (Col)	12.9 ± 0.2 (30)	3.2 ± 0.1 (30)	3.2 ± 0.1 (30)
35S:SPL3Δ	5.6 ± 0.2 (30)	1.6 ± 0.1 (30)	1.6 ± 0.1 (30)
35S:SPL3Δ <i>lfy-1</i>	5.5 ± 0.2 (11)	11.6 ± 0.9 (11) ^a	13.4 ± 0.1 (11)
<i>lfy-1</i>	13.5 ± 0.4 (12)	10.9 ± 0.5 (16) ^a	21.3 ± 0.6 (16)
Wild-type (Col)	12.5 ± 0.3 (12) ^a	2.7 ± 0.1 (12)	2.7 ± 0.1 (12)
35S:SPL3Δ	5.5 ± 0.2 (42)	1.5 ± 0.1 (42)	1.5 ± 0.1 (42)
35S:SPL3Δ <i>ful-2</i>	9.6 ± 0.4 (12)	3.5 ± 0.2 (12) ^a	3.5 ± 0.2 (12) ^a
<i>ful-2</i>	11.8 ± 0.3 (11) ^a	3.7 ± 0.2 (11) ^a	3.7 ± 0.2 (11) ^a
Wild-type (Col)	12.6 ± 0.5 (7)	3.4 ± 0.3 (7)	3.4 ± 0.3 (7)
35S:SPL3Δ	5.7 ± 0.2 (30)	1.4 ± 0.1 (30) ^b	1.4 ± 0.1 (30)
35S:SPL3Δ <i>ap1-10</i>	5.4 ± 0.2 (19)	1.8 ± 0.3 (19) ^b	8.4 ± 1.6 (9) ^c 42.1% ^d
<i>ap1-10</i>	10.9 ± 0.5 (12)	3.2 ± 0.1 (12)	5.8 ± 1.0 (5) ^c 41.7% ^d

Shown is the mean ± SEM (n). Plants were grown on soil at 22°C in long-day conditions (LDs; 16 hr light/8 hr dark, ~110 μmol m² s⁻¹).

^aNo statistically significant difference (Student's t test, p > 0.4) was detected between the marked genotypes.

^bStatistically significant difference (Student's t test, p < 0.05) was detected between the marked genotypes.

^cNumber of secondary inflorescences in plants that ceased secondary inflorescence and branched flower production prior to node 20.

^dPercent of plants that formed more than 20 secondary inflorescences and branched flowers.

If SPL3 induces *LFY*, *FUL*, and *AP1*, we expect temporal and spatial overlap in the expression of these regulators. In long-day-grown wild-type seedlings, *SPL3*, *LFY*, and *FUL* levels showed very similar temporal upregulation, increasing gradually from day 7 onward up to the MI transition at day 11 (first accumulation of *AP1*; Figure 1E). The expression of the three genes increased at a similar rate after the MI transition and peaked during late reproductive development (day 13 to 15; Figure 1E). By contrast, *AP1* levels started to increase later (at day 9) and increased at a slow rate up to the MI transition. A similar delay in *AP1* induction has been observed in short-day-grown plants in response to photoinduction (Hempel et al., 1997; Schmid et al., 2003). After the MI transition, *AP1* expression increased very rapidly until day 13, likely because of the elaboration of the first flower primordia, which strongly express *AP1* (Figure 1E; Hempel et al., 1997). The temporal expression of *SPL3* and *FUL*, *LFY*, and *AP1* is thus consistent with a role for SPL3 in the regulation of the MI genes.

Previous mRNA in situ hybridization data suggest that *SPL3* and *FUL*, *LFY*, and *AP1* are expressed in the inflorescence meristem and associated young flower primordia (Cardon et al., 1997; Hempel et al., 1997; Mandel and Yanofsky, 1995; Weigel et al., 1992). To test for spatial expression overlap between *SPL3* and *FUL*, *LFY*, and *AP1*, we generated a translational fusion protein of *SPL3* with GFP (pSPL3:GFP-SPL3). This construct contains the microRNA binding site in the 3'UTR of *SPL3*. Two independent transgenic pSPL3:GFP-SPL3 lines showed GFP fluorescence in adult leaves and in the inflorescence meristem when grown in long-day conditions and did not exhibit early phase transitions (Figures 1F and 3C; data not shown), suggesting that the transgene recapitulates endogenous *SPL3* expression and activity. When we compared *SPL3* protein accumulation and mRNA accumulation of *FUL*, *LFY*, and *AP1* in young inflorescences (Figure 1F), we observed coexpression at the sites of the MI transition in the inflorescence:

in incipient flower primordia (*SPL3*, *FUL*, and *LFY*) and very young flowers (*SPL3*, *AP1*, *LFY*, and, to a lesser extent, *FUL*; Figure 1F). The observed spatial expression patterns support a role for *SPL3* upstream of *FUL*, *LFY*, and *AP1* in flower formation.

LFY, AP1, and FUL Act Genetically Downstream of SPL3

Since 35S:SPL3Δ plants exhibit a precocious MI transition and elevated *LFY*, *FUL*, and *AP1* levels, this raises the question whether the MI regulators are required for the precocious MI transition of 35S:SPL3Δ plants. We first examined the requirement for *LFY* by crossing 35S:SPL3Δ to the *lfy-1* null mutant (Schultz and Haughn, 1991). *lfy-1* was epistatic to 35S:SPL3Δ with regard to cauline leaf formation (Table 1). 35S:SPL3Δ *lfy-1* plants formed the same number of cauline leaves as *lfy-1*, despite the strong reduction in cauline leaf number observed in 35S:SPL3Δ compared to the wild-type (Table 1). In addition, a strong increase was observed in the number of secondary inflorescences in 35S:SPL3Δ *lfy-1* compared to 35S:SPL3Δ (Table 1, Figure 2A). By contrast, *lfy-1* had no effect on the number of rosette leaves formed in 35S:SPL3Δ (Table 2). 35S:SPL3Δ plants exhibit a precocious vegetative phase change (Wu and Poethig, 2006), which was not altered in the absence of *LFY* (Table S2). Thus, consistent with its known role (Blazquez et al., 1997; Weigel et al., 1992), *LFY* is specifically required for the precocious MI transition observed in 35S:SPL3Δ.

35S:SPL3Δ *ful-2* plants had MI phenotypes identical to those of strong loss-of-function *ful-2* single mutants (Ferrandiz et al., 2000), both with respect to the number of secondary inflorescences and cauline leaves formed (Table 1, Figure 2B). Thus, *ful-2* is epistatic to 35S:SPL3Δ in regulation of the MI transition. *ful-2* also partly suppressed the precocious reproductive transition observed in 35S:SPL3Δ (Table 1), suggesting that upregulation of *FUL* expression contributes to the early flowering of 35S:SPL3Δ. These findings are consistent with the previously




Figure 2. The Precocious Meristem Identity Phenotype of 35S:SPL3 Δ Requires a Functional Copy of LFY, FUL, and AP1

Phenotype of 35S:SPL3 Δ double mutants with *lfy-1* (A), *ful-2* (B), and *ap1-10* (C).

Top: side view of same-age plants. Note the increased number of secondary inflorescences in the double mutants (left) compared to 35S:SPL3 Δ (right).

Bottom: top view of shoot apex close-ups from 5 cm tall primary inflorescences. Note the leafy appearance of 35S:SPL3 Δ *lfy-1* and 35S:SPL3 Δ *ap1-10* due to the presence of additional cauline leaves. Plants were grown in LD. Upper panel bar, 1 cm; lower panel bar, 1 mm.

described strong and subtle roles of FUL in regulation of the MI and the reproductive transition, respectively (Ferrandiz et al., 2000; Melzer et al., 2008). In addition, *ful-2* partially suppressed the precocious vegetative phase transition of 35S:SPL3 Δ (Table S2), suggesting a potential novel role for FUL in vegetative phase change. Hence FUL may act downstream of SPL3 in all three developmental transitions.

AP1 is also required for the precocious MI transition in 35S:SPL3 Δ (Table 1, Figure 2C). Like the strong *ap1-10* mutant (Schultz and Haughn, 1993), 35S:SPL3 Δ *ap1-10* plants often continued to produce secondary inflorescences or branched flowers after the 20th node, and the remaining plants showed a strong increase in the number of secondary inflorescences formed (8.4 in 35S:SPL3 Δ *ap1-10* compared to 5.8 in *ap1-10* plants; Table 1). In addition, a slight increase in the number of cauline leaves was observed in 35S:SPL3 Δ *ap1-10* (Table 1).

As expected based on the role of AP1 as an MI regulator (Bowman et al., 1993), *ap1-10* had no effect on the vegetative transition (Table S2) or the reproductive transition (Table 1; Cardon et al., 1997) of 35S:SPL3 Δ . Taken together, these analyses provide strong support for the hypothesis that SPL3 acts upstream of LFY, AP1, and FUL in the same genetic pathway to control flower formation.

SPL3 Directly Regulates Expression of MI Genes

Our combined findings suggest a possible direct role of SPL3 in the upregulation of LFY, FUL, and AP1. Therefore, we computationally identified candidate SPL transcription factor binding sites in the LFY, FUL, and AP1 loci (Figure 3D) using a matrix-based approach (<http://www.athamap.de/index.php>; Galuschka et al., 2007). To test whether SPL3 directly regulates MI gene expression by binding to any of these *cis* elements, we probed

Table 2. Phenotype of 35S:FT-GFP Alone and in Combination with 35S:miR156a or 35S:SPL3 Δ

Genotype	Number of Rosette Leaves	Number of Cauline Leaves	Number of Secondary Inflorescences
Wild-type (Col)	11.9 \pm 0.3 (12)	2.9 \pm 0.1 (12)	2.9 \pm 0.1 (12)
35S:FT-GFP	4.2 \pm 0.2 (16)	2.0 \pm 0.1 (16)	2.1 \pm 0.1 (16)
35S:FT-GFP \times 35S:miR156a	3.9 \pm 0.2 (24)	2.2 \pm 0.1 (24)	1.8 \pm 0.1 (24)
35S:miR156a \times Col	39.1 \pm 1.9 (13)	ND	ND
35S:FT-GFP \times Col	3.3 \pm 0.1 (57) ^a	1.3 \pm 0.1 (57)	1.3 \pm 0.1 (57) ^a
35S:FT-GFP \times 35S:SPL3 Δ	2.0 \pm 0 (29) ^a	1.4 \pm 0.1 (29)	1.0 \pm 0.1 (29) ^a
35S:SPL3 Δ \times Col	5.1 \pm 0.2 (15)	1.5 \pm 0.1 (15)	1.5 \pm 0.1 (15)

Shown is the mean \pm SEM (n). ND denotes not determined. Plants were grown on soil at 22°C in LDs.

^aStatistically significant difference (Student's t test, $p < 0.005$) was detected between marked genotypes.




Figure 3. SPL3 Binds Regulatory Regions of Meristem Identity Genes In Vivo

(A) Early flowering phenotype of LD-grown *SPL3*-overexpressing plants compared to wild-type. The asterisk marks the primary inflorescence bolt. Bar, 1 cm.

(B) Fluorescence image of 35S:*SPL3*Δ-GFP in leaf one (B) and p*SPL3*:GFP-*SPL3* in leaf four (C). Inset in (B): close-up confocal microscopy image. Bar, 1 mm.

(D) Schematic of the *LFY*, *FUL*, and *AP1* loci. Pale blue and green boxes represent untranslated regions and exons, respectively. Asterisks indicate computationally identified SPL consensus motifs. Pink horizontal lines: fragments amplified in by qPCR after ChIP. Blue boxes: regulatory regions in the *LFY* promoter previously identified (Blazquez and Weigel, 2000).

(E and F) qPCR of anti-GFP ChIP in 35S:*SPL3*Δ-GFP (E), p*SPL3*:GFP-*SPL3*, and wild-type (F). Plants were grown in SD for 10 days (E) or LD for 7 days (F). Immunoprecipitated DNA enrichment is presented as percent input DNA. Shown is the mean \pm SEM.

SPL3 occupancy on genomic DNA in vivo by chromatin immunoprecipitation (ChIP). Toward this end, we generated GFP-tagged 35S:*SPL3*Δ (35S:*SPL3*Δ-GFP and 35S:*GFP-SPL3*Δ). Both transgenes recapitulated the phenotype of 35S:*SPL3*Δ (Table S1, Figure 3A; data not shown). The tagged SPL3 protein was constitutively expressed (Figure 3B; data not shown) and localized to the nucleus (inset in Figure 3B). Using two independent transgenic lines and two different polyclonal anti-GFP antibodies, the same genomic regions showed selective SPL3 occupancy in 35S:*SPL3*Δ-GFP plants (Figure S2), suggesting that the assay accurately detects SPL3 binding.

No candidate SPL binding sites were identified in the *LFY* promoter, but consensus binding sites were present in the first exon/intron as well as in the second intron of *LFY* (region 3 and region 4, respectively; Figure 3D). Using ChIP-qPCR, we detected strong and selective occupancy of SPL3 at region 3 of the *LFY* locus, which contains three consensus SPL binding motifs (Figure 3E). By contrast, the predicted SPL binding sites in the second intron of *LFY* (region 4) were not bound (Figure 3E). The close proximity of the three predicted SPL binding sites in region 3 of the *LFY* locus (less than 200 bp apart) is below the resolution limit of ChIP-qPCR since average size of the sonicated DNA is 500 bp. Thus, we were unable to determine which of the three consensus binding sites in region 3 are occupied by SPL3. We also tested for *SPL3*Δ-GFP occupancy at *LFY* promoter regions previously shown to be important for photoperiodic induction or gibberellin-dependent induction (regions 1 and 2 in the diagram in Figure 3D; Blazquez and Weigel, 2000). No strong SPL3 binding was observed for either region (Figure 3E). Hence SPL3 binds regulatory regions outside of the *LFY* promoter in vivo.

ChIP-qPCR. SPL3 did not bind to the distal *FUL* promoter (region 1; Figure 3E). By contrast, strong SPL3 binding was observed in region 2, which contains a cluster of promoter proximal candidate SPL binding sites (Figure 3E, Figure S2; Lannenpaa et al., 2004). In addition, we observed strong SPL3 occupancy in region 3 in the first *FUL* intron (Figure 3E). Again, the ChIP resolution does not allow us to determine which of the predicted SPL binding sites in region 2 or 3 of *FUL* are occupied by SPL3. Region 4 of *FUL*, which contains a putative SPL binding site in the fifth intron, was not bound by SPL3. Hence SPL3 binds regulatory regions in the *FUL* locus.

AP1 is a direct target of *LFY* (Busch et al., 1999; Wagner et al., 1999; William et al., 2004); therefore, the upregulation of *AP1* observed in 35S:*SPL3*Δ may be an indirect consequence of the elevated *LFY* levels in these plants. However, prior in vitro studies in both *Antirrhinum* and *Arabidopsis* instead point to a direct role of SPL3 in *AP1* activation (Klein et al., 1996; Cardon et al., 1997). To examine these possibilities, we tested for *SPL3*Δ-GFP occupancy at *AP1*-regulatory regions using ChIP. A total of five regions were tested (Figure 3D), including the SPL3 binding site previously identified by electrophoretic mobility shift assays in the *AP1* promoter (region 4; Cardon et al., 1997). We detected strong in vivo SPL3 binding to regions 2, 3, and 4 of the *AP1* promoter (Figure 3E, Figure S2), suggesting that SPL3 occupies multiple binding sites in the *AP1* intergenic region. A more distal predicted SPL3 binding motif in the intergenic region upstream of the *AP1* transcription start site (region 1) as well as a predicted SPL binding site at the end of the first intron of *AP1* (region 5; Figure 3D) were not occupied by *SPL3*Δ-GFP (Figure 3E). Hence SPL3 can bind *AP1* regulatory regions in vivo.

These experiments suggest that SPL3 may directly upregulate the expression of *LFY*, *FUL*, and *AP1* to cause a precocious MI transition in 35S:SPL3 Δ , strongly implicating SPL3 in control of this vital phase transition. To further test this hypothesis, we examined whether SPL3 also binds to regulatory regions of MI genes in wild-type plants expressing pSPL3:GFP-SPL3 (described above). ChIP-qPCR in two independent pSPL3:GFP-SPL3 transgenic lines yielded very similar results (data not shown), suggesting that this construct likely accurately reflects endogenous SPL3 occupancy on the genomic DNA. Representative ChIP-qPCR results for 7-day-old long-day-grown pSPL3:GFP-SPL3 seedlings are shown in Figure 3F, with the same regions tested for occupancy as in Figure 3E. A total of six regions were strongly and selectively occupied by SPL3 in pSPL3:GFP-SPL3 (Figure 3F), including region 3 of the *LFY* locus; regions 2 and 3 in the *FUL* locus; and regions 2, 3, and 4 of the *AP1* locus. The regions occupied by SPL3 in pSPL3:GFP-SPL3 are identical to those bound by SPL3 in 35S:SPL3 Δ -GFP lines. We noted some quantitative differences in relative binding strength in 35S:SPL3 Δ -GFP compared to pSPL3:GFP-SPL3. These differences may be due to the different SPL3 protein levels in the two lines. Amplification of SPL3-occupied regions after ChIP of nontransgenic wild-type plants yielded no detectable signal under identical conditions (Figure 3F), indicating that the observed enrichment is highly specific. Since SPL3 binds to regulatory regions in the *LFY*, *FUL*, and *AP1* loci in seedlings that exhibit wild-type growth and development, our combined data indicate that *LFY*, *FUL*, and *AP1* are direct *in vivo* targets of SPL3.

SPL3 Acts Redundantly with Other SPL Proteins to Control the MI Transition

Although SPL3 regulates expression of MI genes and directly binds their regulatory regions *in vivo* under physiological conditions, *sp3* single mutants do not exhibit any morphological defects, likely due to functional redundancy with other SPL proteins (Wang et al., 2008; Wu and Poethig, 2006). SPL3 is a member of a family of 16 SBP-box transcription factors in *Arabidopsis*, 11 of which (including SPL3) are regulated post-transcriptionally by the miR156 family of microRNAs (Gandikota et al., 2007; Schwab et al., 2005; Wu and Poethig, 2006). SPL3 has two close paralogs: SPL4 and SPL5; the three genes are grouped together into one SBP subfamily (Guo et al., 2008). Members of this SBP subfamily differ from the remaining seven miR156 targets in two important ways: the genes and cDNAs for *SPL3*, *SPL4*, and *SPL5* are much smaller than the other *SPL* genes and encode primarily the DNA binding domain. In addition, the miR156 recognition motif is located in the 3'UTR and not in the coding region in these three *SPLs* (Gandikota et al., 2007; Guo et al., 2008; Schmid et al., 2003; Wu and Poethig, 2006). Because of these observations, we hypothesized that *SPL4* and *SPL5* are most likely to have overlapping roles with *SPL3*.

To test for a role of *SPL4* and *SPL5* in MI gene expression, we examined expression of *LFY*, *FUL*, and *AP1* in plants overexpressing *SPL4* or *SPL5* without the miR156 target site in the 3'UTR (35S:SPL4 Δ or 35S:SPL5 Δ ; Wu and Poethig, 2006). 35S:SPL4 Δ and 35S:SPL5 Δ plants exhibit similar-strength phenotypes as 35S:SPL3 Δ (Table S1; Wu and Poethig, 2006)

and had similar transgene expression levels (*SPL4* Δ expression was slightly lower than that of *SPL3* Δ and *SPL5* Δ ; Figure S3A). We observed a similar increase in *LFY* and *FUL* levels in 10-day-old 35S:SPL4 Δ and 35S:SPL5 Δ plants grown in short-day conditions as we had for 35S:SPL3 Δ (compare Figure 1C and Figure 4A). *AP1* levels were only increased in 35S:SPL5 Δ and not 35S:SPL4 Δ plants, perhaps owing to the slightly lower transgene levels in the latter. Consistent with this, expression of all three MI genes was increased in 35S:SPL4 Δ and 35S:SPL5 Δ plants grown in inductive photoperiod (Figure 4B). Since *SPL3* levels are not elevated in 35S:SPL4 Δ and 35S:SPL5 Δ (Figure S3B), our results suggest that *SPL4* and *SPL5* act redundantly with *SPL3* in upregulation of *LFY*, *FUL*, and *AP1* expression. However, in contrast to *SPL3* overexpression, which does not result in strong upregulation of any of the flowering time genes tested (Figure 1E), overexpression of *SPL4* or *SPL5* resulted in increased expression of several flowering time genes (Figure S4A). Hence although *SPL3*, *SPL4*, and *SPL5* are very similar to one another based on primary sequence and regulate MI gene expression, they do not have identical roles and activities, even when constitutively expressed.

It is difficult to test whether these SPL proteins are necessary for *LFY*, *FUL*, and *AP1* regulation since *sp3* mutants have no morphological defect and no knockout alleles are available for *SPL4* and *SPL5* (Wang et al., 2008; Wu and Poethig, 2006). We therefore instead used transgenic plants overexpressing miR156, a microRNA that targets these and additional *SPL* genes. The *Arabidopsis* genome encodes for six expressed miR156 genes, which are conserved in different plant species (Xie et al., 2005). Constitutive overexpression of each of these microRNAs causes a delay in the reproductive transition (Schwab et al., 2005; Schwarz et al., 2008; Wu and Poethig, 2006), a phenotype opposite to that observed in plants with elevated levels of *SPL3*, *SPL4*, or *SPL5* (Table 1; Wu and Poethig, 2006). Furthermore, miR156b overexpression reduces the expression level of all *SPL* genes with a miR156 recognition motif (Schwab et al., 2005). Indeed, we detected a greater than 2-fold reduction of *SPL3*, *SPL4*, and *SPL5* (Figure S3C) in a transgenic line overexpressing miR156a (Wu and Poethig, 2006).

We assayed expression of the MI genes in 10-day-old 35S:miR156a compared to wild-type seedlings grown in inductive photoperiod. Compared to the wild-type, *LFY*, *FUL*, and *AP1* expression was markedly reduced in 35S:miR156a seedlings (Figure 4C). This reduction was specific, because expression of other genes regulating developmental transitions such as *FT* and *AGL24* was not reduced in 35S:miR156a (Figure S4B). These results strongly suggest that miR156-regulated SBP transcription factors, including *SPL3*, *SPL4*, and *SPL5*, are required for upregulation of *LFY*, *FUL*, and *AP1* in *Arabidopsis*.

We next wanted to test whether, consistent with the observed molecular phenotype, 35S:miR156a plants exhibit a delayed MI transition. Unfortunately, we were unable to score the MI phenotype of these plants. As previously described, 35S:miR156a plants were late flowering in long-day conditions (Wu and Poethig, 2006). In addition, 35S:miR156a plants exhibit severely reduced apical dominance (Figure S5; Chuck et al., 2007; Schwarz et al., 2008; Xie et al., 2005). This phenotype, while potentially highly interesting as discussed below, disrupts the




Figure 4. SPL4 and SPL5 Upregulate Meristem Identity Genes

qRT-PCR analysis of meristem identity gene expression in 35S:SPL4Δ and 35S:SPL5Δ compared to wild-type plants (A and B) and in 35S:miR156a compared to wild-type plants (C). Plants were grown in SD for 10 days (A) and in LD for 9 days (B) or 10 days (C). Shown is the mean ± SEM relative to the wild-type. Black solid horizontal line, 2-fold increase; black dotted line, 2-fold decrease.

normal progression of developmental transitions. Hence it was impossible to assess the timing of the MI transition in 35S:miR156a plants.

Interactions between FT and SPL3

We have identified SPL3 as a direct upstream regulator of *LFY*, *FUL*, and *AP1*. Interestingly, the flowering time regulator FT also upregulates these three MI genes. Together with the bZIP transcription factor FD, FT directly activates *AP1* (Abe et al., 2005; Wigge et al., 2005) and FT/FD either directly or indirectly activate *FUL* (Teper-Bamnolker and Samach, 2005). FT is thought to indirectly activate *LFY* via upregulation of *SOC1* (Searle et al., 2006). This raises the possibility that FT may upregulate *FUL* or *LFY* via *SPL3* induction. To test whether FT is sufficient to induce *SPL3* expression, we generated transgenic plants overexpressing FT (35S:FT-GFP), which flowered precociously and had a more rapid MI transition (fewer cauline leaves and secondary inflorescences

formed than in the wild-type) consistent with previous reports for FT overexpression (Table 2; Kobayashi and Weigel, 2007). No elevated *SPL3* expression was observed in 4-day-old 35S:FT-GFP plants grown in inductive photoperiod. However, *SPL3* levels were increased more than 2-fold at day 6 in this line (Figure 5A). Since none of the six expressed miR156 precursors showed a reduced expression in 6-day-old 35S:FT-GFP compared to wild-type plants (data not shown), the observed increase in *SPL3* expression is not likely to be mediated by miR156. Together with previous findings (Schmid et al., 2003), the data suggest that FT acts at least in part upstream of *SPL3*. Interestingly, *LFY*, *FUL*, *AP1*, as well as *SOC1* levels were already elevated by day 4 in 35S:FT-GFP (Figure 5A). Since upregulation of *FUL* and *LFY* precedes that of *SPL3*, FT is not likely to activate *FUL* and *LFY* via *SPL3* induction.

To further test the regulatory interaction between FT and *SPL3*, we examined the timing of developmental transitions in 35S:FT-GFP plants that had strongly reduced *SPL* levels. F1 progeny of a cross between 35S:FT-GFP and 35S:miR156a still exhibited a precocious reproductive and MI transition compared to the wild-type (Table 2). Overall, the phenotype of the F1 progeny was additive with plants forming slightly more rosette leaves, cauline leaves, and secondary inflorescences than the F1 progeny of the control cross (35S:FT-GFP × wild-type; Table 2). As expected, the F1 progeny of 35S:miR156a crossed to wild-type flowered much later than wild-type, more like the 35S:miR156a parental line (Table 2 and data not shown). The phenotype of 35S:FT-GFP 35S:miR156a plants is shown in Figure 5B (top). In a parallel approach, we examined developmental transitions in the F1 progeny of the cross of 35S:SPL3Δ to 35S:FT-GFP. These plants displayed a more precocious reproductive and MI transition than either parental line (Figure 5B, bottom). Indeed, 35S:FT-GFP 35S:SPL3Δ plants exhibited a strong significant decrease ($p < 0.0001$) in the number of rosette leaves and a subtle but statistically significant ($p < 0.001$) decrease in the number of secondary inflorescences formed compared to the control cross (35S:FT-GFP × wild-type; Table 2). Because 35S:miR156a is not epistatic to 35S:FT-GFP with respect to the reproductive and MI transition, while 35S:SPL3Δ enhances 35S:FT-GFP, our genetic data suggest that *SPL3* acts at least in part in a pathway parallel to the FT pathway to regulate developmental timing in *Arabidopsis*.

DISCUSSION

Proper timing of the onset of reproduction is of central importance for fitness across species (Roux et al., 2006). Recently, the SBP-box family of transcription factors has been implicated in regulating this process (Cardon et al., 1997; Schwarz et al., 2008; Wu and Poethig, 2006). Here, we show that the SBP-box transcription factor *SPL3* directly activates the three key MI genes *LFY*, *FUL*, and *AP1* during the MI transition (Figure 5C). *SPL3* and related proteins form an independent regulatory pathway in the network that controls the onset of reproduction in *Arabidopsis*. These findings provide a major advance in our understanding of the regulation of this important life history trait. Because of the molecular and functional conservation of the MI regulators (Benlloch et al., 2007; Litt and Irish, 2003; Maizel et al., 2005) and the *SPL* proteins (Guo et al., 2008; Xie et al., 2006), the




Figure 5. Regulatory Interactions between Flowering Time and Meristem Identity Regulators

(A) qRT-PCR analysis of the known FT targets *LFY*, *FUL*, *AP1*, and *SOC1* as well as *SPL3* in 35S:FT-GFP plants grown in LD. Plants were harvested at day 4 and day 6. Shown is the expression mean \pm SEM relative to the wild-type. Black solid horizontal line, 2-fold increase.

(B) Phenotypes of F1 progeny of 35S:FT-GFP \times 35S:miR156a (top) and 35S:SPL3 Δ \times 35S:FT-GFP (bottom). Star, secondary inflorescences; arrow, flower subtended by cauline leaf; asterisk, primary inflorescence. Bar, 1 cm.

(C) Role of SPL3 in meristem identity transition. Solid arrows denote direct interactions; dashed arrows denote direct or indirect interactions. SPL3 directly regulates *LFY*, *FUL*, and *AP1* transcription (red arrows) and acts in a pathway that is partly in parallel with the FT pathway (red dashed arrow).

identified regulatory interactions may also play a role in the timing of flower formation in other plant species.

SPL3, a Direct Upstream Activator of *LFY*, *FUL*, and *AP1*

The timing of the MI transition is exquisitely sensitive to *LFY* levels (Blazquez et al., 1997), and precise spatial and temporal *LFY* upregulation is likely also important for additional life history traits such as species-specific inflorescence architecture (Prusinkiewicz et al., 2007) and irreversibility of the switch to flower formation (Parcy et al., 2002). Thus far, very few direct upstream activators of *LFY* are known (Gocal et al., 2001; Lee et al., 2008; Liu et al., 2008), and loss- or gain-of-function mutations in these cause only modest MI phenotypes, suggesting that additional types of transcriptional regulators are required for *LFY* upregulation. Here, we identify the SBP-box containing transcription factors as additional *LFY* activators.

The *LFY* promoter, which consists of the entire 2.3 kb intergenic region upstream of *LFY* (Blazquez et al., 1997), does not contain any consensus SPL binding motifs (this study; Cardon et al., 1997). In agreement with this, *SPL3* overexpressing plants do not upregulate expression of a reporter gene whose expression is driven by the *LFY* promoter (Figure S6). Hence regulatory elements outside of the *LFY* promoter are required for upregulation by *SPL3*. Indeed, we show that *SPL3* strongly binds in vivo to a region in the first exon/intron of *LFY* that contains three consensus SPL binding sites. *LFY* introns are known to be critical

for proper expression of *LFY* in monocots (Bomblied and Doebley, 2005; Prasad et al., 2003; Rao et al., 2008), and recent genome-wide transcription factor binding studies support the idea that important *cis* elements may also reside in exons in *Arabidopsis* (C.M. Winter and D. Wagner, unpublished; Oh et al., 2009).

Our findings combined with previous studies suggest that *LFY* accumulation is regulated by a surprisingly complex array of partly redundantly acting transcription factor families and regulatory regions (Blazquez and Weigel, 2000; Lee et al., 2008; Liu et al., 2008). This complexity may form the basis for species-specific differences in induction of flower formation and of inflorescence architecture.

The MADS-box transcription factor *FUL* plays a role in both the reproductive transition and the MI transition together with other MADS-box transcription factors (Ferrandiz et al., 2000; Teper-Bamnolker and Samach, 2005). More recently, *FUL* and *SOC1* were shown to control the annual growth habit of *Arabidopsis* (Melzer et al., 2008). Hence precise regulation of *FUL* accumulation is of critical importance for the life strategy of flowering plants. Thus far, no direct upstream activator of *FUL* is known, although there is evidence that *FUL* acts downstream of the photoperiod pathway regulators FT and FD (Abe et al., 2005; Teper-Bamnolker and Samach, 2005; Wigge et al., 2005). Here, we identify *SPL3* as a direct upstream activator of *FUL*. In addition, we show that *FUL* is not only required for the precocious MI transition but also for the precocious vegetative phase change and reproductive transition observed in 35S:SPL3 Δ plants. Moreover, we observed very strong activation of *FUL* by *SPL3* and that occupancy of *SPL3* at *FUL* regulatory regions was highly robust. Taken together, these findings suggest that *FUL* is an important target of SPL transcription factors in control of developmental timing at multiple stages in the plant life cycle.

Interestingly, in 35S:miR156a plants, axillary inflorescences bolt prior to the outgrowth of the primary inflorescence and plants have a multiple rosette phenotype (Figure S5 and data not shown; Schwab et al., 2005; Xie et al., 2006). This phenotype is reminiscent of the perennial growth habit observed in plants with simultaneous downregulation of *FUL* and *SOC1* (Melzer et al., 2008). We show that 35S:miR156a plants have reduced levels of *FUL* and *SOC1* (Figure 4C and Figure S4B). One possibility suggested by our results that warrants further investigation is that SPL transcription factors may play a role in the regulation of annual versus perennial growth habit in different species of flowering plants via modulation of *FUL* and *SOC1* expression.

We show that *AP1* is a direct in vivo target of *SPL3*, with strong occupancy of *SPL3* at multiple regions in the *AP1* promoter. In contrast to the upregulation of *LFY* and *FUL* by *SPL3*, *AP1* upregulation is only observed late in noninductive photoperiod, suggesting that *SPL3* is not sufficient for *AP1* activation. A delay of *AP1* induction compared to *FUL* and *LFY* upregulation is also observed in wild-type plants in many different growth conditions (this study; Hempel et al., 1997; Schmid et al., 2003) and may be important for proper timing of the commitment to floral fate. FT and FD together directly upregulate *AP1* (Abe et al., 2005; Wigge et al., 2005), and we detect a subtle increase in *FD* levels in 35S:SPL3 Δ that coincides with *AP1* upregulation. While we cannot rule out that other factors contribute to the delay in *AP1* accumulation, our results suggest that FT/FD may be required

for *AP1* induction by SPL3. This conclusion is supported by the finding that *AP1* is more strongly upregulated in 35S:SPL3 Δ plants grown in inductive photoperiod when FT is present.

The Position of SPL3 in the Regulatory Network Controlling the Onset of Reproduction

The MI transition is the final step in reproductive development and leads to formation of the first flowers. Recent investigations determined that flowering time regulators directly activate MI genes (Figure 5C; Abe et al., 2005; Lee et al., 2008; Liu et al., 2008; Wigge et al., 2005). Since SPL3 and related proteins are important activators of the three main MI genes, this raises the question of how these proteins fit in the known regulatory network of flowering time and MI genes. SPL3 and paralogous genes are strongly upregulated in response to inductive photoperiod in an FT-dependent fashion (Schmid et al., 2003). Moreover, we show that plants overexpressing *FT* have elevated SPL3 levels. On the other hand, gain of SPL3 and FT function are additive, and reduction of SPL3 function is not epistatic to gain of FT function, suggesting that SPL3 and FT act in parallel pathways, which converge on upregulation of MI genes (Figure 5C). Similarly, FT and LFY act in parallel pathways that converge on a common target gene (Abe et al., 2005; Wigge et al., 2005). These regulatory interactions likely allow exquisite fine-tuning of important developmental decisions through integration of multiple environmental and endogenous inputs.

Our findings also point to the presence of a second SPL3 activation pathway (X in Figure 5C). Since we did not detect a decrease in miR156 gene expression in 35S:FT-GFP plants with elevated SPL3 levels, it is tempting to speculate that this second pathway may involve posttranscriptional regulation of SPL3 levels by this family of microRNAs. Future investigation will be needed to test this hypothesis.

EXPERIMENTAL PROCEDURES

Plant Materials and Growth Conditions

Accession numbers of all genes used in this study are listed in Table S3. Columbia (Col) was used as wild-type. *lfy-1*, *ful-2*, and *ap1-10* mutants have been described (Ferrandiz et al., 2000; Schultz and Haughn, 1991, 1993). 35S:SPL3 Δ , 35S:SPL4 Δ , and 35S:SPL5 Δ as well as 35S:miR156a have been described (Wu and Poethig, 2006). For phenotype analysis, plants were grown on soil at 22°C in long-day (16 hr light/8 hr dark) conditions using white fluorescent lights (~110 $\mu\text{mol m}^{-2} \text{s}^{-1}$) or in short-day (10 hr light/14 hr dark) conditions with a 3:1 mixture of white fluorescent and GrowLux lights (~120 $\mu\text{mol m}^{-2} \text{s}^{-1}$). For expression and chromatin immunoprecipitation analysis, plants were grown on half-strength Murashige and Skoog (MS) medium under long-day (continuous light) conditions with white fluorescent lights (~90 $\mu\text{mol m}^{-2} \text{s}^{-1}$) or short-day (10 hr light/14 hr dark) as described above. Seeds were stratified at 4°C for 2 to 4 days and then transferred to 22°C at day 0.

Plasmid Construction and Transgenic Plants

To construct the 35S:SPL3 Δ -GFP and 35S:FT-GFP lines, the coding region of SPL3 and FT was amplified by PCR with primer sets containing the attB sequence for Gateway-based cloning and recombined into pDONR221 using BP reaction (Invitrogen, USA). Primers used are listed in Table S4. After sequencing, each coding sequence was recombined into pGWB5 (Nakagawa et al., 2007) using LR reactions (Invitrogen, USA). The resulting binary vectors were introduced into *Agrobacterium* strain GV3101::pMP90 and transformed into Col plants. Transgenic plants flowering earlier than wild-type were further analyzed.

To construct pSPL3:GFP-SPL3, a 2.9 kb genomic fragment upstream of the ATG and a 0.9 kb fragment downstream of the ATG of SPL3 were cloned before and after eGFP, respectively, in the binary vector pCAMBIA3300 (CAMBIA, Australia) and transformed into Col plants. Primers used are listed in Table S4.

qRT-PCR and In Situ Hybridization

RNA extraction and reverse transcription was essentially as described (William et al., 2004), except treatment with RNase-free DNase (QIAGEN, USA) was included and the Superscript III kit (Invitrogen, USA) was used. Real-time PCR reactions were performed using Power SYBR Green PCR master mix (Applied Biosystems). The relative amount of a given mRNA was determined based on the threshold cycle number required for amplification compared to the standard curve and then normalized by the expression values of the eukaryotic translation initiation factor 4A-1 (*EIF4A*(At1g54270)) in each sample. Fold change was calculated by dividing the normalized value of the experimental genotype by that of the wild-type. The mean and standard error were determined using one to two biological replicates with three technical replicates each. Primers used are listed in Table S5.

In situ hybridization was performed as in Long and Barton (1998) using probes previously described (Hempel et al., 1997).

Chromatin Immunoprecipitation

300 mg of 10-day-old seedlings grown in short-day conditions (35S:SPL3 Δ -GFP) or 7-day-old seedlings grown in long-day conditions (pSPL3:GFP-SPL3) were employed for ChIP following the procedure previously described (Kwon et al., 2005; William et al., 2004). Anti-GFP antibodies A6455 (Invitrogen) and ab290 (Abcam) were used. Real-time PCR was performed using Power SYBR Green PCR master mix (Applied Biosystems). To estimate SPL3 occupancy on genomic DNA, we computed the ratio of ChIP over input DNA (% Input) by comparing the reaction threshold cycle for each the ChIP sample to a dilution series of the corresponding input sample. Primers used are summarized in Table S6.

Fluorescence Microscopy

GFP fluorescence was visualized using a fluorescent microscope (Olympus, MVX10) or a confocal microscope (Leica, LCS SL).

SUPPLEMENTAL DATA

Supplemental Data include six figures and six tables and can be found with this article online at [http://www.cell.com/developmental-cell/supplemental/S1534-5807\(09\)/](http://www.cell.com/developmental-cell/supplemental/S1534-5807(09)/).

ACKNOWLEDGMENTS

We thank Xuemei Chen, Kim Gallagher, Sarah Liljegren, and John Wagner, as well as Wagner lab members for comments on this manuscript, Koji Koizumi for help with fluorescence microscopy, Gary Ditta and Marty Yanofsky for the *ful-2* allele and the *FUL* probe, and Jia-Wei Wang and Detlef Weigel for exchange of unpublished information. This work was supported by NSF IBN grant 0516622 to D.W., NIH R01 GM051893 to R.S.P., and a JSPS fellowship to A.Y.

Received: February 7, 2009

Revised: May 12, 2009

Accepted: June 1, 2009

Published: August 17, 2009

REFERENCES

- Abe, M., Kobayashi, Y., Yamamoto, S., Daimon, Y., Yamaguchi, A., Ikeda, Y., Ichinoki, H., Notaguchi, M., Goto, K., and Araki, T. (2005). FD, a bZIP protein mediating signals from the floral pathway integrator FT at the shoot apex. *Science* 309, 1052–1056.
- Achard, P., Herr, A., Baulcombe, D.C., and Harberd, N.P. (2004). Modulation of floral development by a gibberellin-regulated microRNA. *Development* 131, 3357–3365.

- Araki, T. (2001). Transition from vegetative to reproductive phase. *Curr. Opin. Plant Biol.* 4, 63–68.
- Benlloch, R., Berbel, A., Serrano-Mislata, A., and Madueno, F. (2007). Floral initiation and inflorescence architecture: a comparative view. *Ann. Bot. (Lond.)* 100, 659–676.
- Blazquez, M.A., and Weigel, D. (2000). Integration of floral inductive signals in *Arabidopsis*. *Nature* 404, 889–892.
- Blazquez, M.A., Soowal, L.N., Lee, I., and Weigel, D. (1997). LEAFY expression and flower initiation in *Arabidopsis*. *Development* 124, 3835–3844.
- Blazquez, M.A., Ferrandiz, C., Madueno, F., and Parcy, F. (2006). How floral meristems are built. *Plant Mol. Biol.* 60, 855–870.
- Bombliès, K., and Doebley, J.F. (2005). Molecular evolution of FLORICAULA/LEAFY orthologs in the Andropogoneae (Poaceae). *Mol. Biol. Evol.* 22, 1082–1094.
- Bowman, J.L., Alvarez, J., Weigel, D., Meyerowitz, E.M., and Smyth, D.R. (1993). Control of flower development in *Arabidopsis thaliana* by *Apetala1* and interacting genes. *Development* 119, 721–743.
- Busch, M.A., Bombliès, K., and Weigel, D. (1999). Activation of a floral homeotic gene in *Arabidopsis*. *Science* 285, 585–587.
- Cardon, G.H., Hohmann, S., Nettekheim, K., Saedler, H., and Huijser, P. (1997). Functional analysis of the *Arabidopsis thaliana* SBP-box gene SPL3: a novel gene involved in the floral transition. *Plant J.* 12, 367–377.
- Cardon, G., Hohmann, S., Klein, J., Nettekheim, K., Saedler, H., and Huijser, P. (1999). Molecular characterisation of the *Arabidopsis* SBP-box genes. *Gene* 237, 91–104.
- Chuck, G., Cigan, A.M., Saetern, K., and Hake, S. (2007). The heterochronic maize mutant *Comgrass1* results from overexpression of a tandem microRNA. *Nat. Genet.* 39, 544–549.
- Ferrandiz, C., Gu, Q., Martienssen, R., and Yanofsky, M.F. (2000). Redundant regulation of meristem identity and plant architecture by FRUITFULL, APETALA1 and CAULIFLOWER. *Development* 127, 725–734.
- Galuschka, C., Schindler, M., Bulow, L., and Hehl, R. (2007). AthaMap web tools for the analysis and identification of co-regulated genes. *Nucleic Acids Res.* 35, D857–D862.
- Gandikota, M., Birkenbihl, R.P., Hohmann, S., Cardon, G.H., Saedler, H., and Huijser, P. (2007). The miRNA156/157 recognition element in the 3' UTR of the *Arabidopsis* SBP box gene SPL3 prevents early flowering by translational inhibition in seedlings. *Plant J.* 49, 683–693.
- Gocal, G.F., Sheldon, C.C., Gubler, F., Moritz, T., Bagnall, D.J., MacMillan, C.P., Li, S.F., Parish, R.W., Dennis, E.S., Weigel, D., et al. (2001). GAMYB-like genes, flowering, and gibberellin signaling in *Arabidopsis*. *Plant Physiol.* 127, 1682–1693.
- Guo, A.Y., Zhu, Q.H., Gu, X., Ge, S., Yang, J., and Luo, J. (2008). Genome-wide identification and evolutionary analysis of the plant specific SBP-box transcription factor family. *Gene* 418, 1–8.
- Hempel, F.D., Weigel, D., Mandel, M.A., Ditta, G., Zambryski, P.C., Feldman, L.J., and Yanofsky, M.F. (1997). Floral determination and expression of floral regulatory genes in *Arabidopsis*. *Development* 124, 3845–3853.
- Klein, J., Saedler, H., and Huijser, P. (1996). A new family of DNA binding proteins includes putative transcriptional regulators of the *Antirrhinum majus* floral meristem identity gene SQUAMOSA. *Mol. Gen. Genet.* 250, 7–16.
- Kobayashi, Y., and Weigel, D. (2007). Move on up, it's time for change—mobile signals controlling photoperiod-dependent flowering. *Genes Dev.* 21, 2371–2384.
- Kwon, C.S., Chen, C., and Wagner, D. (2005). WUSCHEL is a primary target for transcriptional regulation by SPLAYED in dynamic control of stem cell fate in *Arabidopsis*. *Genes Dev.* 19, 992–1003.
- Lannenpää, M., Janonen, I., Holttä-Vuori, M., Gardemeister, M., Porali, I., and Sopanen, T. (2004). A new SBP-box gene BpSPL1 in silver birch (*Betula pendula*). *Physiol. Plant.* 120, 491–500.
- Lee, J., Oh, M., Park, H., and Lee, I. (2008). SOC1 translocated to the nucleus by interaction with AGL24 directly regulates LEAFY. *Plant J.* 55, 832–843.
- Litt, A., and Irish, V.F. (2003). Duplication and diversification in the APETALA1/FRUITFULL floral homeotic gene lineage: implications for the evolution of floral development. *Genetics* 165, 821–833.
- Liu, C., Chen, H., Er, H.L., Soo, H.M., Kumar, P.P., Han, J.H., Liou, Y.C., and Yu, H. (2008). Direct interaction of AGL24 and SOC1 integrates flowering signals in *Arabidopsis*. *Development* 135, 1481–1491.
- Long, J.A., and Barton, M.K. (1998). The development of apical embryonic pattern in *Arabidopsis*. *Development* 125, 3027–3035.
- Maizel, A., Busch, M.A., Tanahashi, T., Perkovic, J., Kato, M., Hasebe, M., and Weigel, D. (2005). The floral regulator LEAFY evolves by substitutions in the DNA binding domain. *Science* 308, 260–263.
- Mandel, M.A., and Yanofsky, M.F. (1995). A gene triggering flower formation in *Arabidopsis*. *Nature* 377, 522–524.
- Meizer, S., Lens, F., Gennen, J., Vanneste, S., Rohde, A., and Beeckman, T. (2008). Flowering-time genes modulate meristem determinacy and growth form in *Arabidopsis thaliana*. *Nat. Genet.* 40, 1489–1492.
- Nakagawa, T., Suzuki, T., Murata, S., Nakamura, S., Hino, T., Maeo, K., Tabata, R., Kawai, T., Tanaka, K., Niwa, Y., et al. (2007). Improved Gateway binary vectors: high-performance vectors for creation of fusion constructs in transgenic analysis of plants. *Biosci. Biotechnol. Biochem.* 71, 2095–2100.
- Oh, E., Kang, H., Yamaguchi, S., Park, J., Lee, D., Kamiya, Y., and Choi, G. (2009). Genome-wide analysis of genes targeted by PHYTOCHROME INTERACTING FACTOR 3-LIKE5 during seed germination in *Arabidopsis*. *Plant Cell* 21, 403–419.
- Parcy, F., Bombliès, K., and Weigel, D. (2002). Interaction of LEAFY, AGAMOUS and TERMINAL FLOWER1 in maintaining floral meristem identity in *Arabidopsis*. *Development* 129, 2519–2527.
- Poethig, R.S. (2003). Phase change and the regulation of developmental timing in plants. *Science* 301, 334–336.
- Prasad, K., Kushalappa, K., and Vijayraghavan, U. (2003). Mechanism underlying regulated expression of RFL, a conserved transcription factor, in the developing rice inflorescence. *Mech. Dev.* 120, 491–502.
- Prusinkiewicz, P., Erasmus, Y., Lane, B., Harder, L.D., and Coen, E. (2007). Evolution and development of inflorescence architectures. *Science* 316, 1452–1456.
- Rao, N.N., Prasad, K., Kumar, P.R., and Vijayraghavan, U. (2008). Distinct regulatory role for RFL, the rice LFY homolog, in determining flowering time and plant architecture. *Proc. Natl. Acad. Sci. USA* 105, 3646–3651.
- Roux, F., Touzet, P., Cuguen, J., and Le Corre, V. (2006). How to be early flowering: an evolutionary perspective. *Trends Plant Sci.* 11, 375–381.
- Schmid, M., Uhlenhaut, N.H., Godard, F., Demar, M., Bressan, R., Weigel, D., and Lohmann, J.U. (2003). Dissection of floral induction pathways using global expression analysis. *Development* 130, 6001–6012.
- Schultz, E.A., and Haughn, G.W. (1991). LEAFY, a homeotic gene that regulates inflorescence development in *Arabidopsis*. *Plant Cell* 3, 771–781.
- Schultz, E.A., and Haughn, G.W. (1993). Genetic analysis of the floral induction process (FLIP) in *Arabidopsis*. *Development* 119, 745–765.
- Schwab, R., Palatnik, J.F., Rießer, M., Schommer, C., Schmid, M., and Weigel, D. (2005). Specific effects of microRNAs on the plant transcriptome. *Dev. Cell* 8, 517–527.
- Schwarz, S., Grande, A.V., Bujdoso, N., Saedler, H., and Huijser, P. (2008). The microRNA regulated SBP-box genes SPL9 and SPL15 control shoot maturation in *Arabidopsis*. *Plant Mol. Biol.* 67, 183–195.
- Searle, I., He, Y., Turck, F., Vincent, C., Fornara, F., Krober, S., Amasino, R.A., and Coupland, G. (2006). The transcription factor FLC confers a flowering response to vernalization by repressing meristem competence and systemic signaling in *Arabidopsis*. *Genes Dev.* 20, 898–912.
- Steeves, T.A., and Sussex, I.M. (1989). *Patterns in Plant Development* (Cambridge, UK: Cambridge University Press).
- Teper-Bamnolker, P., and Samach, A. (2005). The flowering integrator FT regulates SEPALLATA3 and FRUITFULL accumulation in *Arabidopsis* leaves. *Plant Cell* 17, 2661–2675.

- Turck, F., Fornara, F., and Coupland, G. (2008). Regulation and identity of florigen: FLOWERING LOCUS T moves center stage. *Annu. Rev. Plant Biol.* 59, 573–594.
- Wagner, D., Sablowski, R.W., and Meyerowitz, E.M. (1999). Transcriptional activation of APETALA1 by LEAFY. *Science* 285, 582–584.
- Wang, J.W., Schwab, R., Czech, B., Mica, E., and Weigel, D. (2008). Dual effects of miR156-targeted SPL genes and CYP78A5/KLUH on plastochron length and organ size in *Arabidopsis thaliana*. *Plant Cell* 20, 1231–1243.
- Weigel, D., and Nilsson, O. (1995). A developmental switch sufficient for flower initiation in diverse plants. *Nature* 377, 495–500.
- Weigel, D., Alvarez, J., Smyth, D.R., Yanofsky, M.F., and Meyerowitz, E.M. (1992). LEAFY controls floral meristem identity in *Arabidopsis*. *Cell* 69, 843–859.
- Wigge, P.A., Kim, M.C., Jaeger, K.E., Busch, W., Schmid, M., Lohmann, J.U., and Weigel, D. (2005). Integration of spatial and temporal information during floral induction in *Arabidopsis*. *Science* 309, 1056–1059.
- William, D.A., Su, Y., Smith, M.R., Lu, M., Baldwin, D.A., and Wagner, D. (2004). Genomic identification of direct target genes of LEAFY. *Proc. Natl. Acad. Sci. USA* 101, 1775–1780.
- Wu, G., and Poethig, R.S. (2006). Temporal regulation of shoot development in *Arabidopsis thaliana* by miR156 and its target SPL3. *Development* 133, 3539–3547.
- Xie, K., Wu, C., and Xiong, L. (2006). Genomic organization, differential expression, and interaction of SQUAMOSA promoter-binding-like transcription factors and microRNA156 in rice. *Plant Physiol.* 142, 280–293.
- Xie, Z., Allen, E., Fahlgren, N., Calamar, A., Givan, S.A., and Carrington, J.C. (2005). Expression of *Arabidopsis* MIRNA genes. *Plant Physiol.* 138, 2145–2154.