



ELSEVIER

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

Journal of Combinatorial Theory, Series A 110 (2005) 31–41

Journal of
Combinatorial
Theory

Series A

www.elsevier.com/locate/jcta

A family of bijections between G -parking functions and spanning trees

Denis Chebikin, Pavlo Pylyavskyy

Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA 02139, USA

Received 30 July 2003

Available online 9 December 2004

Abstract

For a directed graph G on vertices $\{0, 1, \dots, n\}$, a G -parking function is an n -tuple (b_1, \dots, b_n) of non-negative integers such that, for every non-empty subset $U \subseteq \{1, \dots, n\}$, there exists a vertex $j \in U$ for which there are more than b_j edges going from j to $G - U$. We construct a family of bijective maps between the set \mathcal{P}_G of G -parking functions and the set \mathcal{T}_G of spanning trees of G rooted at 0, thus providing a combinatorial proof of $|\mathcal{P}_G| = |\mathcal{T}_G|$.

© 2004 Elsevier Inc. All rights reserved.

Keywords: Parking functions; Spanning trees

1. Introduction

The classical parking functions are defined in the following way. There are n drivers, labeled $1, \dots, n$, and n parking spots, $0, \dots, n - 1$, arranged linearly in this order. Each driver i has a favorite parking spot b_i . Drivers enter the parking area in the order in which they are labeled. Each driver proceeds to his favorite spot and parks there if it is free, or parks at the next available spot otherwise. The sequence (b_1, \dots, b_n) is called a *parking function* if every driver parks successfully by this rule. The most notable result about parking functions is a bijective correspondence between such functions and trees on $n + 1$ labeled vertices. The number of such trees is $(n + 1)^{n-1}$ by Cayley's theorem. For more on parking functions, see for example [9].

Postnikov and Shapiro [8] suggested the following generalization of parking functions. Let G be a directed graph on $n + 1$ vertices indexed by integers from 0 to n . A G -parking

E-mail addresses: chebikin@mit.edu (D. Chebikin), pasha@mit.edu (P. Pylyavskyy).

unction is a sequence (b_1, \dots, b_n) of non-negative integers that satisfies the following condition: for each subset $U \subseteq \{1, 2, \dots, n\}$ of vertices of G , there exists a vertex $j \in U$ such that the number of edges from j to vertices outside of U is greater than b_j . For the complete graph $G = K_{n+1}$, these are the classical parking functions (we view K_{n+1} as the digraph with exactly one edge (i, j) for all $i \neq j$).

A *spanning tree* of G rooted at m is a subgraph of G such that, for each $i \in \{0, 1, \dots, n\}$, there is a unique path from i to m along the edges of the spanning tree. Note that these are the spanning trees of the graph in the usual sense with each edge oriented towards m . The number of such trees is given by the matrix-tree theorem; see [9]. In [8], it is shown that the number of spanning trees of G rooted at 0 is equal to the number of G -parking functions for any digraph G .

An equivalent fact was originally discovered by Dhar [2], who studied the sandpile model. The so-called recurrent states of the sandpile model are in one-to-one correspondence with G -parking functions for certain graphs G , including all symmetric graphs. A bijection between recurrent states and spanning trees for symmetric graphs G is mentioned in [6], and a class of bijections is constructed in [1]. The sandpile model was also studied by Gabrielov in [5]. This paper also contains an extensive list of references on the topic.

In this paper, we present a family of bijections between G -parking functions and rooted spanning trees of G . Given a spanning tree T of G , we establish a total order on the vertices of T satisfying two conditions, and each such order gives rise to a bijection in the family. In [3] Francon used a similar concept, which he called selection procedures, to construct a family of bijections between parking functions and rooted trees in the classical case $G = K_{n+1}$. Thus our result provides a generalization of Francon's construction.

2. A family of bijections

Let G be a directed graph on vertices $\{0, \dots, n\}$. We allow G to have multiple edges but not loops. To distinguish between multiple edges of G , we fix an order on the set of edges going from i to j for all $i \neq j$.

A *subtree* of G rooted at m is a subgraph T of G containing m such that for every vertex i of T , there is a unique path in T from i to m . A subtree is called a *spanning tree* if it contains all vertices of G .

Let \mathbb{T}_G be the set of subtrees of G rooted at 0, and let \mathcal{T}_G be the set of spanning trees of G rooted at 0. Unless stated otherwise, all spanning trees in this paper are assumed to be rooted at 0. Let \mathcal{P}_G be the set of G -parking functions. In this section we give a bijection between \mathcal{T}_G and \mathcal{P}_G .

For every $T \in \mathbb{T}_G$, let $\pi(T)$ be a total order on the vertices of T , and write $i <_{\pi(T)} j$ to denote that i is smaller than j in this order. We call the set $\Pi(G) = \{\pi(T) \mid T \in \mathbb{T}_G\}$ a *proper set of tree orders* if the following conditions hold for all $T \in \mathbb{T}_G$:

- (1) if (j, i) is an edge of T , then $i <_{\pi(T)} j$;
- (2) if t is a subtree of T rooted at 0, then the order $\pi(t)$ is consistent with $\pi(T)$; in other words, $i <_{\pi(t)} j$ if and only if $i <_{\pi(T)} j$ for $i, j \in t$.

We give several examples of proper sets of tree orders in Section 3.

For $T \in \mathbb{T}_G$ and a vertex j of G , the order $\pi(T)$ induces the order on the edges going from j to vertices of T in which (j, i) is smaller than (j, i') whenever $i <_{\pi(T)} i'$ and which is consistent with the previously fixed order on multiple edges. We write $e <_{\pi(T)} e'$ to denote that e is smaller than e' in this order.

Given a proper set of tree orders $\Pi(G)$, define the map $\Theta_{\Pi,G} : \mathcal{T}_G \rightarrow \mathcal{P}_G$ as follows. For $T \in \mathcal{T}_G$ and a vertex $j \in \{1, \dots, n\}$, let e_j be the edge of T going out of j . Set $\Theta_{\Pi,G}(T) = (b_1, \dots, b_n)$, where b_j is the number of edges e going out of j such that $e <_{\pi(T)} e_j$. For the rest of the section, we write Θ instead of $\Theta_{\Pi,G}$.

Theorem 2.1. *The map Θ is a bijection between \mathcal{T}_G and \mathcal{P}_G .*

Proof. We begin by checking that $\Theta(T)$ is a G -parking function.

Lemma 2.2. $\Theta(T) \in \mathcal{P}_G$ for $T \in \mathcal{T}_G$.

Proof. For a subset $U \subseteq \{1, \dots, n\}$, let j be the smallest vertex of U in the order $\pi(T)$. Let $e_j = (j, i)$ be the edge of T coming out of j . Then $i <_{\pi(T)} j$, so $i \notin U$ by choice of j . For each of the b_j edges $e = (j, i')$ such that $e <_{\pi(T)} e_j$, we have $i' \leq_{\pi(T)} i <_{\pi(T)} j$, so $i' \notin U$. Thus there are at least $b_j + 1$ edges going from j to vertices outside of U . \square

Next, we define the inverse map $\Phi_{\Pi,G} : \mathcal{P}_G \rightarrow \mathcal{T}_G$. Given $P = (b_1, \dots, b_n) \in \mathcal{P}_G$, we construct the corresponding tree $\Phi_{\Pi,G}(P)$ one edge at a time. Initially, let t_0 be the subtree of G consisting of the vertex 0 alone, and put $p_0 = 0$. For $1 \leq m \leq n$, we choose the vertex p_m and construct the subtree t_m rooted at 0 inductively as follows. Let U_m be the set of vertices not in t_{m-1} , and let V_m be the set of vertices $j \in U_m$ such that the number of edges from j to t_{m-1} is at least $b_j + 1$. Note that $|V_m| \geq 1$ by definition of a G -parking function. For each $j \in V_m$, let e_j be the edge from j to t_{m-1} such that exactly b_j edges e from j to t_{m-1} satisfy $e <_{\pi(t_{m-1})} e_j$. Let t be the tree obtained by adjoining each vertex $j \in V_m$ to t_{m-1} by means of the edge e_j . Set p_m to be the smallest vertex of V_m in the order $\pi(t)$, and set t_m to be the tree obtained by adjoining p_m to t_{m-1} by means of the edge e_{p_m} . Obviously, t_m is a subtree of G . In the end, set $\Phi_{\Pi,G}(P) = T = t_n$. For the rest of the section, we write Φ instead of $\Phi_{\Pi,G}$.

An example of constructing $\Phi(P)$ is shown in Fig. 1. Let G be the graph shown in the figure, and let $P = (0, 1, 0, 1)$. Let Π be the tree order in which vertex i is smaller than vertex j if i is closer to the root than j , or else if i and j are equidistant to the root, and $i < j$. Initially, $U_1 = \{1, 2, 3, 4\}$ and $V_1 = \{1\}$, so vertex 1 is attached to the root to produce the subtree t_1 . Then we have $V_2 = \{3, 4\}$ with $e_3 = (3, 1)$ and $e_4 = (4, 1)$. Adjoining vertices 3 and 4 to t_1 by means of e_3 and e_4 places vertices 3 and 4 the same distance away from the root, making $p_2 = 3$, so vertex 3 is attached by means of the edge $e_3 = (3, 1)$ to produce t_2 . At the next step, we have $V_3 = \{2, 4\}$ with $e_2 = (2, 3)$ and $e_4 = (4, 1)$. Adjoining vertices 2 and 4 to t_2 by means of e_2 and e_4 makes vertex 4 closer to the root than vertex 2, so we select vertex 4 and attach it to vertex 1 to form t_3 . Finally, we attach vertex 2 to 3 to form $t_4 = \Phi(P)$.

Lemma 2.3. *In the above construction, $p_0 <_{\pi(T)} \dots <_{\pi(T)} p_n$.*




Fig. 1. An example of constructing $\Phi(P)$.

Proof. Since $\Pi(G)$ is a proper set of tree orders, it follows that the root 0 is the smallest vertex of T in the order $\pi(T)$. Hence $p_0 <_{\pi(T)} p_1$. Suppose that $p_0 <_{\pi(T)} \dots <_{\pi(T)} p_m$ for some $1 \leq m \leq n - 1$. We show that $p_m <_{\pi(T)} p_{m+1}$. We consider the following two cases.

Case 1: $p_{m+1} \notin V_m$. Then the number of edges from p_{m+1} to t_{m-1} is at most $b_{p_{m+1}}$. Since $p_{m+1} \in V_{m+1}$, the number of edges from p_{m+1} to t_m is at least $b_{p_{m+1}} + 1$. It follows that there is at least one edge (p_{m+1}, p_m) in G and that p_{m+1} is adjoined to t_m by means of such an edge. Thus $p_m <_{\pi(T)} p_{m+1}$ because $\Pi(G)$ is a proper set of tree orders.

Case 2: $p_{m+1} \in V_m$. Let $e_{p_{m+1}}$ be the edge from p_{m+1} to t_{m-1} such that exactly $b_{p_{m+1}}$ edges e from p_{m+1} to t_{m-1} satisfy $e <_{\pi(t_{m-1})} e_{p_{m+1}}$. Since p_m is the largest vertex of t_m in the order $\pi(T)$ and hence in the order $\pi(t_m)$, and $e_{p_{m+1}}$ goes from p_{m+1} to $t_{m-1} = t_m - p_m$, it follows that $e <_{\pi(t_{m-1})} e_{p_{m+1}}$ if and only if $e <_{\pi(t_m)} e_{p_{m+1}}$ because the order $\pi(t_{m-1})$ is consistent with $\pi(t_m)$. Therefore, exactly $b_{p_{m+1}}$ edges e from p_{m+1} to t_m satisfy $e <_{\pi(t_m)} e_{p_{m+1}}$, hence p_{m+1} is adjoined to t_m by means of the edge $e_{p_{m+1}}$.

Let e_{p_m} be the edge of T coming out of p_m , and let t be the tree in the construction of T obtained by adjoining the vertices of V_m to t_{m-1} . Let t' be the tree obtained from t_{m-1} by adjoining the vertices p_m and p_{m+1} by means of the edges e_{p_m} and $e_{p_{m+1}}$. Then t' is a subtree of both t and T . By choice of p_m , we have $p_m <_{\pi(t)} p_{m+1}$, so $p_m <_{\pi(t')} p_{m+1}$ and $p_m <_{\pi(T)} p_{m+1}$ because the order $\pi(t')$ is consistent with both $\pi(t)$ and $\pi(T)$. \square

We now check that Θ and Φ are inverses of each other.

Lemma 2.4. $\Theta(\Phi(P)) = P$ for $P \in \mathcal{P}_G$.

Proof. Put $P = (b_1, \dots, b_n)$ and $T = \Phi(P)$. Consider the process of constructing T . For $j \in \{1, \dots, n\}$, we have $j = p_m$ for some $1 \leq m \leq n$. Let e_j be the edge of T coming out of j . The edge e_j goes from j to t_{m-1} . Since the set of vertices of t_{m-1} is $\{p_0, \dots, p_{m-1}\}$, it follows from Lemma 2.3 that if an edge e coming out of j satisfies $e <_{\pi(T)} e_j$, then e goes from j to t_{m-1} . Thus, $e <_{\pi(T)} e_j$ if and only if $e <_{\pi(t_{m-1})} e_j$ because the order $\pi(t_{m-1})$ is consistent with $\pi(T)$. By construction of T , the number of edges e satisfying $e <_{\pi(t_{m-1})} e_j$

is b_j , hence the number of edges e satisfying $e <_{\pi(T)} e_j$ is also b_j . We conclude that $\Theta(T) = P$. \square

Lemma 2.5. $\Phi(\Theta(T')) = T'$ for $T' \in \mathcal{T}_G$.

Proof. Put $P = \Theta(T') = (b_1, \dots, b_n)$. Consider the process of constructing $T = \Phi(P)$. We show by induction that for $0 \leq m \leq n$, the tree t_m is a subtree of T' and that p_0, \dots, p_m are the smallest $m + 1$ vertices in the order $\pi(T')$. Since the root 0 is the smallest vertex in $\pi(T')$, the assertion is true for $m = 0$.

Now, suppose that t_{m-1} is a subtree of T' and that p_0, \dots, p_{m-1} are the smallest m vertices in the order $\pi(T')$. Let k be the $(m + 1)$ th smallest vertex in the order $\pi(T')$, and let $e'_k = (k, i)$ be the edge coming out of k in T' . Then $i <_{\pi(T')} k$, so $i \in \{p_0, \dots, p_{m-1}\}$ and $i \in t_{m-1}$. Hence if an edge e coming out of k satisfies $e <_{\pi(T')} e'_k$, then e goes from k to t_{m-1} . There are b_k edges e satisfying $e <_{\pi(T')} e'_k$. These b_k edges together with the edge e'_k give $b_k + 1$ edges going from k to t_{m-1} . It follows that $k \in V_m$.

As before, for every $j \in V_m$, let e_j be the edge from j to t_{m-1} such that exactly b_j edges e from j to t_{m-1} satisfy $e <_{\pi(t_{m-1})} e_j$. Since the vertices of t_{m-1} are the smallest m vertices in the order $\pi(T')$, it follows that if an edge e coming out of j satisfies $e <_{\pi(T')} e_j$, then e goes from j to t_{m-1} . Thus, $e <_{\pi(T')} e_j$ if and only if $e <_{\pi(t_{m-1})} e_j$ because the order $\pi(t_{m-1})$ is consistent with $\pi(T')$. There are b_j edges e satisfying $e <_{\pi(t_{m-1})} e_j$, hence there are b_j edges e satisfying $e <_{\pi(T')} e_j$. It follows from the choice of b_j that e_j is an edge of T' . Therefore, the tree t obtained by adjoining the vertices $j \in V_m$ by means of the edges e_j is a subtree of T' . Consequently, the smallest vertex p_m of V_m in the order $\pi(t)$ is the smallest vertex of V_m in the order $\pi(T')$. Since k is the smallest vertex of U_m in the order $\pi(T')$ and $k \in V_m \subseteq U_m$, it follows that $p_m = k$. The induction step is complete.

Finally, we obtain $T' = t_n = T$. \square

Theorem 2.1 follows from Lemmas 2.4 and 2.5. \square

3. Examples

In this section, we give examples of proper sets of tree orders and the resulting bijections between \mathcal{T}_G and \mathcal{P}_G from the family of bijections defined in Section 2.

We begin by introducing the *breadth-first search order* $\pi_{bf}(T)$ on the vertices of a tree $T \in \mathbb{T}_G$. For a vertex $i \in T$, we define the *height* $h_T(i)$ of i in T to be the number of edges in the unique path from i to the root 0. We set $i <_{\pi_{bf}(T)} j$, or $i <_{bf} j$, if $h_T(i) < h_T(j)$ or else if $h_T(i) = h_T(j)$ and $i < j$. It is easy to check that $\pi_{bf}(T)$ is a total order on the vertices of T and that $\Pi_{bf}(G) = \{\pi_{bf}(T) \mid T \in \mathbb{T}_G\}$ is a proper set of tree orders.

The *depth-first search order* $\pi_{df}(T)$ on the vertices of a tree $T \in \mathbb{T}_G$ is defined as follows. For a vertex $i \in T$, let $T(i)$ denote the branch of T rooted at i . In other words, $T(i)$ consists of all vertices k of T such that the unique path from k to 0 in T contains i . If (i, ℓ) is an edge of T , then we set $\ell <_{\pi_{df}(T)} i$. Furthermore, if (j, ℓ) is an edge of T such that




Fig. 2. An example of constructing $\Theta_{bf,G}(T)$, $\Theta_{df,G}(T)$, and $\Theta_{va,G}(T)$.

$i < j$, then we set $i' <_{\pi_{df}(T)} j'$ for $i' \in T(i)$ and $j' \in T(j)$. We use the symbol $<_{df}$ with the same meaning as $<_{\pi_{df}(T)}$. It is not hard to see that $\Pi_{df}(G) = \{\pi_{df}(T) \mid T \in \mathbb{T}_G\}$ is a proper set of tree orders.

Our third example is the *vertex-adding order* $\pi_{va}(T)$ on the vertices of $T \in \mathbb{T}_G$. Construct the sequence $p_0, \dots, p_{|T|-1}$ inductively as follows. Set $p_0 = 0$, and, for $1 \leq m \leq |T| - 1$, let p_m be the smallest vertex j in $G - \{p_0, \dots, p_{m-1}\}$ such that there is an edge in T from j to $\{p_0, \dots, p_{m-1}\}$. Note that the sequence $p_0, \dots, p_{|T|-1}$ contains each vertex of T exactly once. Put $p_0 <_{\pi_{va}(T)} \dots <_{\pi_{va}(T)} p_{|T|-1}$, and let the symbol $<_{va}$ have the same meaning as $<_{\pi_{va}(T)}$. Clearly, $<_{va}$ is a total order on the vertices of T . Also, $\Pi_{va}(G) = \{\pi_{va}(T) \mid T \in \mathbb{T}_G\}$ is a proper set of tree orders. Indeed, if t is a subtree of T , then adding or not adding a vertex of $T - t$ to $\{p_0, \dots, p_{m-1}\}$ does not affect the order in which the vertices of t are added.

Let $\Theta_{bf,G}$, $\Theta_{df,G}$, and $\Theta_{va,G}$ be the maps $\Theta_{\Pi_{bf,G}}$, $\Theta_{\Pi_{df,G}}$, and $\Theta_{\Pi_{va,G}}$ constructed in Section 2. Fig. 2 shows a sample graph G and a spanning tree $T \in \mathbb{T}_G$. To compute $\Theta_{bf,G}(T)$, $\Theta_{df,G}(T)$, and $\Theta_{va,G}(T)$, we first determine the orders $\pi_{bf}(T)$, $\pi_{df}(T)$, and $\pi_{va}(T)$. We have $h_T(0) = 0$, $h_T(2) = h_T(6) = 1$, $h_T(3) = h_T(4) = 2$, and $h_T(1) = h_T(5) = 3$, so

$$0 <_{bf} 2 <_{bf} 6 <_{bf} 3 <_{bf} 4 <_{bf} 1 <_{bf} 5.$$

Next, we determine $\pi_{df}(T)$. Applying the depth-first search rule with $\ell = 0$, we get $0 <_{df} \{1, 2, 3, 4, 5\} <_{df} 6$ because $T(2)$ contains vertices 1, 2, 3, 4, and 5, and $T(6)$ contains a single vertex 6. Taking $\ell = 2$ we get $0 <_{df} 2 <_{df} \{3, 5\} <_{df} \{1, 4\} <_{df} 6$. Finally, taking $\ell = 3$ and $\ell = 4$ we get

$$0 <_{df} 2 <_{df} 3 <_{df} 5 <_{df} 4 <_{df} 1 <_{df} 6.$$

Also, the order $\pi_{va}(T)$ is the following:

$$0 <_{va} 2 <_{va} 3 <_{va} 4 <_{va} 1 <_{va} 5 <_{va} 6.$$

The edge coming out of vertex 1 in T is $(1, 4)$. The relation $e <_{bf} (1, 4)$ is satisfied for $e = (1, 0), (1, 2), (1, 3), (1, 6)$, so the first component of $\Theta_{bf,G}(T)$ is 4. Similarly, $e <_{df} (1, 4)$ holds for $e = (1, 0), (1, 2), (1, 3), (1, 5)$, so the first component of $\Theta_{df,G}(T)$ is 4. The relation $e <_{va} (1, 4)$ holds for $e = (1, 0), (1, 2), (1, 3)$, so the first component of $\Theta_{va,G}(T)$ is 3. The other components are computed in the same way. Fig. 2 shows the values of $\Theta_{bf,G}(T)$, $\Theta_{df,G}(T)$, and $\Theta_{va,G}(T)$.

Note that for $G = K_{n+1}$, the presented construction yields a family of bijections between the classical parking functions and trees on $n + 1$ labeled vertices. This family includes some of the well-known bijections. For example, using the vertex-adding tree order results in the following simple correspondence defined in terms of drivers and parking spots: given a parking function (b_1, \dots, b_n) , the corresponding tree is obtained by introducing the edge (i, j) whenever driver j ended up parking in spot $b_i - 1$, and the edge $(i, 0)$ whenever $b_i = 0$.

Another bijection involving labeled Dyck paths as an intermediate object, communicated to us by A. Postnikov, results if the right-to-left depth first search tree order is used (this order is the same as the depth first search order described above except that larger numbers are given priority among the children of the same vertex). Given a parking function P , we write numbers 1 through n in the $n \times n$ square so that all numbers j such that $b_j = i$ appear in the i th row in increasing order, and the numbers in a lower row appear to the left of the numbers in a higher row. Such an arrangement defines a Dyck path from the lower-left corner to the upper-right corner of the square, with horizontal steps labeled with integers between 1 and n ; see Fig. 3. To get the spanning tree T corresponding to P , start from the upper-right corner of the square and proceed to the lower-left corner along the Dyck path, keeping track of the current vertex, initially set to be 0. At each horizontal step labeled i , connect the vertex i to the current vertex, and at each vertical step, replace the current vertex with its successor in the right-to-left depth first search order on the tree constructed so far. It is not hard to show that the obtained tree T is precisely $\Phi(P)$ for the right-to-left depth first search tree order.

The bijection obtained using the breadth first search tree order is discussed in Section 5 in connection with the sandpile model.

4. More proper sets of tree orders

We now present a method for constructing proper sets of tree orders. Let $\langle \sigma_1, \dots, \sigma_\ell \rangle$ denote the path consisting of the edges $(\sigma_\ell, \sigma_{\ell-1}), (\sigma_{\ell-1}, \sigma_{\ell-2}), \dots, (\sigma_1, 0)$. Also, let $\langle \rangle$ denote the path consisting of the vertex 0 alone. Define \mathbb{A}_G to be the set of paths $\langle \sigma_1, \dots, \sigma_\ell \rangle$ in G such that $\sigma_1, \dots, \sigma_\ell$ are distinct vertices of $G - \{0\}$, where $\ell \geq 0$. Let $<$ be a partial order on \mathbb{A}_G satisfying the following conditions:

- (i) if $A \cap A' \in \mathbb{A}_G$ for some $A, A' \in \mathbb{A}_G$, then A and A' are comparable;
- (ii) $\langle \sigma_1, \dots, \sigma_{\ell'} \rangle < \langle \sigma_1, \dots, \sigma_{\ell'}, \dots, \sigma_\ell \rangle$ for $\ell' < \ell$.




Fig. 3. A bijection between trees and parking functions involving labeled Dyck paths.

For a tree $T \in \mathbb{T}_G$ and a vertex $i \in T$, let $A_T(i) \in \mathbb{A}_G$ be the unique path in T from i to 0. Introduce the order $\pi_{\prec}(T)$ on the vertices of T in which $i <_{\pi_{\prec}(T)} j$ whenever $A_T(i) < A_T(j)$. Put $\Pi_{\prec}(G) = \{\pi_{\prec}(T) \mid T \in \mathbb{T}_G\}$.

Proposition 4.1. $\Pi_{\prec}(G)$ is a proper set of tree orders.

Proof. Let $T \in \mathbb{T}_G$, and let i and j be vertices of $T - \{0\}$. Since $A_T(i)$ and $A_T(j)$ are the unique paths in T from i and j to 0, it follows that $A_T(i) \cap A_T(j) \in \mathbb{A}_G$. Therefore, $\pi_{\prec}(T)$ is a total order on the vertices of T , by property (i) of \prec .

If (j, i) is an edge of T , then $A_T(i) = \langle \sigma_1, \dots, \sigma_\ell, i \rangle$ and $A_T(j) = \langle \sigma_1, \dots, \sigma_\ell, i, j \rangle$, so $A_T(i) < A_T(j)$, by property (ii) of \prec , so $i <_{\pi_{\prec}(T)} j$.

If t is a subtree of T , then $A_t(i) = A_T(i)$ for all vertices $i \in t$, so the order $\pi_{\prec}(t)$ is consistent with the order $\pi_{\prec}(T)$.

The proposition follows. \square

The orders $\pi_{bf}(T)$, $\pi_{df}(T)$, and $\pi_{va}(T)$ described in Section 3 can be obtained as $\pi_{\prec}(T)$ via an appropriate choice of \prec . Setting \prec to be the lexicographic order on the paths $\langle \sigma_1, \dots, \sigma_\ell \rangle$ viewed as sequences of integers yields the order $\pi_{df}(T)$. To obtain $\pi_{bf}(T)$, set $\langle \sigma_1, \dots, \sigma_\ell \rangle < \langle \sigma'_1, \dots, \sigma'_{\ell'} \rangle$ if $\ell < \ell'$, or else if $\ell = \ell'$ and $\sigma_\ell < \sigma'_{\ell'}$. Finally, setting \prec




Fig. 4.

to be the order in which $A \prec A'$ whenever $A \cap A' \in \mathbb{A}_G$ and the largest vertex of $A \setminus A'$ is smaller than the largest vertex of $A' \setminus A$, yields the order $\pi_{va}(T)$.

We can obtain other proper sets of tree orders from partial orders \prec on \mathbb{A}_G satisfying the conditions above. For example, we can set $A = \langle \sigma_1, \dots, \sigma_\ell \rangle \prec A' = \langle \sigma'_1, \dots, \sigma'_{\ell'} \rangle$ if $\ell < \ell'$, or else if $\ell = \ell'$ and the increasing rearrangement of A is smaller than that of A' in the lexicographic order. Another example is setting $A \prec A'$ if $\sum \sigma_k < \sum \sigma'_k$, or else if $\sum \sigma_k = \sum \sigma'_k$ and $\sigma_\ell < \sigma'_{\ell'}$.

Similar examples of partial orders on \mathbb{A}_G yielding proper sets of tree orders can be obtained by using an arbitrary numbering of the edges of G instead of vertex labels.

It is worth noting that not all proper sets of tree orders are induced by a partial order on \mathbb{A}_G satisfying the above conditions. Consider the following simple example. Let G be the graph shown in Fig. 4. Let $e_{1,2}$ be the two edges of G going from vertex 3 to 1, and let $f_{1,2}$ be the two edges going from vertex 4 to 2. For $1 \leq i, j \leq 2$, let T_{ij} be the spanning tree of G containing edges e_i and f_j . Let $\Pi = \{\pi(T_{ij}) \mid 1 \leq i, j \leq 2\}$ be the proper set of tree orders defined as follows:

$$0 <_{\pi(T_{ij})} 1 <_{\pi(T_{ij})} 2 <_{\pi(T_{ij})} 3 <_{\pi(T_{ij})} 4$$

for $i \neq j$, and

$$0 <_{\pi(T_{ii})} 1 <_{\pi(T_{ii})} 2 <_{\pi(T_{ii})} 4 <_{\pi(T_{ii})} 3.$$

Let A_{e_i} (resp., A_{f_i}) be the unique path in \mathbb{A}_G from vertex 3 (resp., 4) to the root 0 containing the edge e_i (resp., f_i). Then in order for Π to be induced by some partial order \prec on \mathbb{A}_G , we must have $A_{e_1} \prec A_{f_2}$ so that relation $3 <_{\pi(T_{12})} 4$ holds. Similarly, to achieve relations $4 <_{\pi(T_{22})} 3$, $3 <_{\pi(T_{21})} 4$, and $4 <_{\pi(T_{11})} 3$, we must have $A_{f_2} \prec A_{e_2}$, $A_{e_2} \prec A_{f_1}$, and $A_{f_1} \prec A_{e_1}$. We obtain a contradiction $A_{e_1} \prec A_{e_1}$, hence Π is not induced by a partial order on \mathbb{A}_G .

5. G-parking functions and the sandpile model

In [1], Cori and Le Borgne construct a family of bijections between the rooted spanning trees of a digraph G and the recurrent states of the sandpile model defined on G . It was

shown by Gabrielov [4] that if for all vertices of G except the root, the out-degree is greater than or equal to the in-degree, then recurrent states coincide with the so-called *allowed configurations* of the model, which correspond to G -parking functions: if d_i is the out-degree of vertex i , then (u_1, \dots, u_n) is an allowed configuration if and only if $(d_1 - u_1, \dots, d_n - u_n)$ is a G -parking function. In particular, this observation is valid for symmetric graphs, in which the number of edges from i to j is equal to the number of edges from j to i for all $i \neq j$; such graphs can be naturally viewed as undirected graphs. Thus for these graphs the result of Cori and Le Borgne provides a bijective correspondence between rooted spanning trees of G and G -parking functions. For the rest of the section, we assume that G is a symmetric graph.

The construction described in [1] begins by fixing an arbitrary order on the edges of G . Given a spanning tree T of G , an edge e in $G - T$ is called *externally active* with respect to T if in the unique cycle of $T + e$, the edge e is the smallest in the chosen order. A key property of the obtained bijection is that the sum of the values of a recurrent state is equal to the number of externally active edges with respect to the corresponding spanning tree. It follows that in the resulting bijection between G -parking functions and spanning trees, G -parking functions with the same sum of values are mapped to spanning trees with the same number of externally active edges.

To show that the bijections presented in this paper are substantially different from the ones in [1], consider the case $G = K_{n+1}$, and let P be the path obtained as follows: start at the root vertex 0, and then append the remaining vertices one by one, so that at each step the appended edge is the smallest, in the chosen edge order, among all edges that can possibly be appended. There are no externally active edges with respect to P since every edge (i, j) not in P , where j is closer to the root in P than i , is greater than the edge (i', j) , where i' is the vertex appended after j in the construction of P , by choice of i' . On the other hand, if a path P' does not include the smallest edge in the chosen edge order, then this edge is externally active with respect to P' . Hence there is a different number of externally active edges with respect to P and P' . However, every bijection $\Theta_{\Pi, G}$ maps both P and P' to permutations of $(0, \dots, n - 1)$, so the sum of values of the corresponding G -parking functions is the same. Hence for $G = K_{n+1}$, none of the bijections $\Theta_{\Pi, G}$ coincides with a bijection from the family constructed in [1].

Dhar defined the *burning algorithm* for determining whether a given configuration is allowed; see [7]. In our setting this task corresponds to the question whether a function $P : \{1, \dots, n\} \rightarrow \mathbb{N}$ is a G -parking function, and an equivalent formulation of Dhar's burning algorithm is the following. We mark vertices of the graph, starting with the root 0. At each iteration of the algorithm, we mark all vertices v that have more marked neighbors than the value of the function at v . If in the end all vertices are marked, then we have a G -parking function, as it is not hard to see directly from definition. Conversely, for every G -parking function, this algorithm marks all vertices.

We claim that our bijection corresponding to the breadth first search order π_{bf} is a natural generalization of Dhar's algorithm. Given a parking function $P = (b_1, \dots, b_n)$, perform the construction of $T = \Phi_{\Pi, G}(P)$ as described in Section 3. We know that T contains all vertices if and only if we started with a G -parking function. Let us group the vertices of T by height, setting W_i to be the set of vertices of T of height i .

Proposition 5.1. W_i is exactly the set of vertices marked at the i th step of the burning algorithm.

Proof. For $i = 0$ the claim is true because the root 0 is marked at the 0th step. We prove the claim by induction. Suppose that for $k < i$, the vertices in W_k are marked at the k th step of the Dhar's algorithm. Let $e_j = (j, w_j)$ be the edge going out of j in T . Each vertex $j \in W_i$ has more than b_j edges going to vertices not larger than w_j in π_{bf} order. All vertices not larger than w_j are in $\cup_{k < i} W_k$ since $w_j \in W_{i-1}$. Therefore, all vertices in W_i are marked at the i th step of Dhar's algorithm. On the other hand, every vertex marked at the i th step of the algorithm in our is to be attached in T to a vertex from $\cup_{k < i} W_k$ since we add vertices to T in the order π_{bf} . Thus each such vertex is in W_i . Hence, Dhar's burning algorithm is realized by our bijection for the breadth first search tree order. \square

Acknowledgments

The authors thank Prof. Alexander Postnikov for suggesting this problem and for many helpful discussions. We are also grateful to Prof. Igor Pak and Prof. Richard Stanley for feedback and advice. Finally, we thank the anonymous referees for supplying useful references and questions related to our results.

References

- [1] R. Cori, Y. Le Borgne, The sand-pile model and Tutte polynomials, *Adv. Appl. Math.* 30 (2003) 44–52.
- [2] D. Dhar, Self-organised critical state of the sandpile automaton models, *Phys. Rev. Lett.* 64 (14) (1990) 1613–1616.
- [3] J. Francon, Acyclic and parking functions, *J. Combin. Theory Ser. A* 18 (1975) 27–35.
- [4] A. Gabrielov, Abelian avalanches and Tutte polynomials, *Physica A* 195 (1993) 253–274.
- [5] A. Gabrielov, Asymmetric abelian avalanches and sandpiles, preprint 93-65, MSI, Cornell University, 1993.
- [6] E.V. Ivashkevich, V.B. Priezzhev, Introduction to the sandpile model, *Physica A* 254 (1998) 97–116.
- [7] R. Meester, F. Redig, D. Znamenski, The Abelian sandpile; a mathematical introduction, *Markov Process. Related Fields* 7 (2001) 509–523.
- [8] A. Postnikov, B. Shapiro, Trees, parking functions, syzygies, and deformations of monomial ideals, *Trans. Amer. Math. Soc.* 356 (2004) 3109–3142.
- [9] R.P. Stanley, *Enumerative Combinatorics*, vol. 2, Cambridge Studies in Advanced Mathematics, vol. 62, Cambridge University Press, Cambridge, 1999.