brought to you by CORE

JOURNAL OF FOOD AND DRUG ANALYSIS 23 (2015) 447-452



Available online at www.sciencedirect.com

## **ScienceDirect**

journal homepage: www.jfda-online.com



### **Original Article**

## Simultaneous determination of some common food dyes in commercial products by digital image analysis



### Mohammad-Hossein Sorouraddin<sup>a,\*</sup>, Masoud Saadati<sup>a,b</sup>, Fariba Mirabi<sup>a</sup>

<sup>a</sup> Department of Analytical Chemistry, Faculty of Chemistry, University of Tabriz, Tabriz, Iran <sup>b</sup> Department of Science, Ardabil Allameh Tabatabaei Campus, Farhangian University, Iran

#### ARTICLE INFO

Article history: Received 20 May 2014 Received in revised form 21 October 2014 Accepted 22 October 2014 Available online 3 January 2015

Keywords: flatbed scanner food additive food color image analysis simultaneous determination

#### ABSTRACT

A simple and relatively fast image-analysis method using digital images, obtained with a flatbed scanner, has been described. The method was used for the simultaneous determination of four common food dyes, namely, carmoisine, brilliant blue, sunset yellow, and quinoline yellow, in binary mixtures in commercial products without a need for any prior separation steps. The results obtained were validated against a standard high-performance liquid chromatography method and a good agreement was obtained. The parameters affecting the experimental results were optimized. Under the optimal conditions, the method provided acceptable linear ranges (20–250 mg/L) with correlation coefficients higher than 0.998, suitable precision (relative standard deviation  $\leq$  4.5%), and limits of detection between 4.82 and 8.05 mg/L.

Copyright © 2014, Food and Drug Administration, Taiwan. Published by Elsevier Taiwan LLC. Open access under CC BY-NC-ND license.

#### 1. Introduction

Food additives are commonly used in processed foodstuff to improve appearance, flavor, taste, color, texture, nutritive value, and preservation. Color is the first quality attribute of food evaluated by consumers, and is therefore an important component of food quality relevant to market acceptance. Although increasing evidence in recent years indicates that the abuse of dyes may cause adverse effects, many kinds of dyes are still widely used due to their low price, high effectiveness, and excellent stability. Therefore, determination of food dyes is important to quality control and control the amount of use permitted [1]. Numerous methods such as sensitive spectrophotometry [2], differential spectrophotometry [3], derivative spectrophotometry [4], spectrophotometry using chemometrics [5,6], colorimetry using a homemade double-beam photocolorimeter [7], reflectometry using a homemade reflectometer [8], electrochemical methods [9,10], capillary electrophoresis [11], ion-pair high-performance liquid chromatography (HPLC) [12], electrokinetic chromatography [13], supercritical fluid extraction–capillary liquid chromatography (LC) [14], thin-layer chromatography [15], HPLC–mass spectrophotometry (MS) [16], solid-phase

<sup>\*</sup> Corresponding author. Department of Analytical Chemistry, Faculty of Chemistry, University of Tabriz, Tabriz, Iran. E-mail address: soruraddin@tabrizu.ac.ir (M.-H. Sorouraddin). http://dx.doi.org/10.1016/j.jfda.2014.10.007

<sup>1021-9498/</sup>Copyright © 2014, Food and Drug Administration, Taiwan. Published by Elsevier Taiwan LLC. Open access under CC BY-NC-ND license.

extraction—HPLC [17], HPLC—diode-array detector detection [18,19], LC—MS [20], LC—hybrid linear analysis presented by Goicoechea and Olivieri [21], and gas chromatography—MS [22] have been developed for determination or resolving mixtures of colorants in foodstuff.

Image analysis is the extraction of meaningful information from images, especially from digital images by digital-imageprocessing techniques. Several techniques have been used for obtaining digital images for different purposes. These techniques include some types of spectroscopy and obtaining images using digital color cameras and scanners [23,24].

The aim of this work is to propose a simple and cheap image-analysis method using flatbed scanners for simultaneous determination of some food colors [quinoline yellow (QY), sunset yellow (SY), carmoisine (CA), and brilliant blue (BL)] in commercial food products. The flatbed scanners in comparison with other imaging devices are relatively inexpensive with the ability to digitize images into a stored array of pixels within a computer. Therefore, the proposed method proved to be simple, relatively fast, and economical for the analysis of the target analytes.

#### 2. Methods

#### 2.1. Chemicals

Dyes of purity higher than 95% were purchased from Grodab Chemie GmbH (Hamburg, Germany). The dyes used were QY, SY, CA, and BL. All other chemicals were obtained from Merck (Germany) and were of the analytical reagent grade. Standard solutions (1000 mg/L) of dyes were prepared using pure water.

#### 2.2. Apparatus and software

The experiments were carried out using a flatbed scanner (LiDE20; Canon, China). All HPLC measurements of colors were carried out using a liquid chromatograph (Waters, Milford, MA, USA), which was equipped with the following: Waters 515 pump, Waters 717plus autosampler, and Waters 2487 dual-wavelength absorbance detector (double channel). A filter paper (ALBET ashless), a thin-chromatography Whatman paper (1 Chr Whatman paper, Thickness, 0.18 mm, UK), and a thick Whatman paper (No. 43; thickness, 0.23 mm, UK) were used as spotting surfaces. Photoshop CS5 (Adobe Systems) and MATLAB 7.0 (MathWorks, Natick, MA, USA) were used for investigating the color intensities from the saved images and calculations, respectively.

#### 2.3. Preparation of the real sample solutions

The proposed method was applied for determination of QY, SY, CA, and BL dyes present as individual components or in binary mixtures within commercial products. The paper chromatography method was used for identification of food dyes in commercial products. In this method, the chromatogram is developed by applying aqueous samples and standard dye solutions to a normal-phase paper. The solutions were developed in the mobile phase of sodium citrate (2.4% w/ v):pure ethanol (1:1). Retention factor values, that is, the ratio of distance traveled by the dye to the distance traveled by the solvent, were estimated for the samples and the standard solutions. Various solid jelly powders (Farmand; Iran) and traditional chocolate (Morvarid, Minoo, Iran) samples were purchased from local supermarkets (Tabriz, Iran). An unnamed shoddy orange soft-drink sample was supplied by the Quality Control Laboratory of the Food and Drug Department of Tabriz University of Medical Sciences. Approximately 6 g of color-coated chocolates (shells) were transferred into a flask. The colored shells were then dissolved in distilled water, centrifuged, and diluted with an equal volume of 3M CH<sub>3</sub>COOH solution. Approximately 200 mg (per 10 mL of sample) of white commercial wool yarn was washed with detergent, rinsed with distilled water, dried, and then added to the aforementioned diluted sample, and the mixture was heated (60 minutes at 60°C or 10 minutes at 90°C). The colored yarn was then taken out, and washed with plenty of distilled water. The dyes were recovered by mixing the yarns with 10 mL of  $NH_3$  (2M) and heating for 10 minutes at 90°C.

#### 2.4. Analytical procedures

Different concentrations of dye samples were prepared by appropriate dilutions of stock solutions (1000 mg/L) with deionized water (Ghazi Company, Tabriz, Iran). To prepare the color spots of the dyes, the prepared solutions (20  $\mu$ L) were spotted on Whatman No. 43 paper, left for a while (40 minutes), and placed on the glass plate of the flatbed scanner. The spots on the paper were separated by Microsoft Paint (XP) software. The selected images were saved in the bitmap format composed in the three channels of the RGB model using Photoshop. The magnitude of each component (red, green, and blue) was determined, and the color value was calculated by subtracting the suitable RGB channel intensities of samples from those of the white sheet. This procedure was performed using MATLAB.

#### 2.5. High-performance liquid chromatography

The aforementioned method was validated by running a parallel HPLC test for the commercial food products. Measurements were made at 460 nm for CA, SY, and QY and at 640 nm for BL using  $NaH_2PO_4/Na_2HPO_4$  buffer solution (pH = 6.0); acetonitrile (65:35 v/v) was used as the mobile phase with a flow rate of 0.9 mL/minute at room temperature. The retention times were 2.91, 3.06, and 3.51 minutes at the first channel (460 nm) for QY, SY, and CA, and 3.47 minutes at the second channel (640 nm) for BL. No efficient interaction between BL and CA was observed as they had similar retention times.

#### 3. Results and discussion

## 3.1. The RGB color model and selection of the suitable RGB component

A digital image is intrinsically a multivariate system, which is a collection of data stored in pixels, each usually highly correlated to its neighbors [25]. The numerical information of each pixel can be decomposed into three channels that correspond to red, green, and blue, which are added in various ways to reproduce a broad array of colors. This is known as the *RGB model* [26]. The abbreviation *RGB* is the initials of the three primary colors, red, green, and blue. Thus, a color in the *RGB* model is described by indicating how much of each component (red, green, or blue) is included (each component can vary from zero to a defined maximum value). Fig. 1 shows the linear *RGB* values corresponding to each wavelength. As shown in the figure, there are *R*, *G*, and *B* values on this plot for every wavelength.

The scanner optically scans the images and converts them to digital images (each point is expressed as an RGB triplet, the components of which vary from 0 to 255). These three parameters are the color intensities of the red (R), green (G), and blue (B) channels [27].

The absorption spectra of the dyes (at pH 6 and 11; recorded and represented in Fig. 2) indicate that the dyes were active in the ultraviolet and visible regions. Within the 400-700-nm range, the spectra of QY, CA, and SY are highly overlapped, whereas within the same range, a lower spectral overlapping was observed between BL and other dyes. Furthermore, a negligible spectral shift was observed, when the solution was made alkaline. Such a high overlapping (particularly between QY, SY, and CA) limits the analysis of these dyes in food products by conventional methods, whereas the proposed method was successfully applied for determining some dyes in binary mixtures. Considering the wavelengths of the maximum absorbance of the dyes (Fig. 2) and RGB values for each wavelength (Fig. 1), the suitable RGB component was selected for determination of any food color as an individual component or as a binary mixture (Table 1).

#### 3.2. Optimization of parameters

#### 3.2.1. Selection of suitable surface for dyes spotting

To study the effect of spotted surface on the analytical results, black surfaces (paper and fabric) were selected at the first step, because the RGB values are zero for black backgrounds. However, the color spots were not detectable on the scanned images with black backgrounds. Thus, white surfaces (paper and fabric) were investigated next. Because the RGB factors on the various white fabrics were not constant at 255 in different points of each surface, paper surfaces consisting of filter paper, paperboard, and two types of Whatman paper (1 Chr and No. 43) were tested for further studies. Because of the weak absorbance strength of the paperboard, the best spotted surface was determined by comparison of repeatability of



Fig. 1 – RGB values for the pure spectral lines.



Fig. 2 – Absorption spectra of brilliant blue (1, 2), carmoisine (3, 4), sunset yellow (5, 6), and quinoline yellow (7, 8) at pH 6 and 11, respectively. Concentrations of all dyes are 10 g/L.

analytical signals with the other three surfaces. Based on the data presented in Table 2, the Whatman paper no. 43 (i.e., the thick one) was selected as the optimized spotted surface.

#### 3.2.2. Optimization of spotted sample volume

The optimum spotted sample volume was selected based on its effect on repeatability of results and determination efficiency. The influence of the spotted sample volume was studied in the 5–50  $\mu$ L range. The results obtained showed that by increasing the volume of sample from 5 to 20  $\mu$ L, analytical signals increase and then remain constant until 50  $\mu$ L. With increases in volume, the repeatability also improved. Based on these results, 20  $\mu$ L was selected as the suitable volume of spotted sample for scanning.

mixtures.		
Analyte	Food color with	Suitable component of RGB
Brilliant blue	_	R
Brilliant blue	Sunset yellow	R
Brilliant blue	Carmoisine	R
Brilliant blue	Quinoline yellow	R
Carmoisine	-	G
Carmoisine	Brilliant blue	G
Carmoisine	Quinoline yellow	G
Carmoisine	Sunset yellow	G
Sunset yellow	-	В
Sunset yellow	Brilliant blue	В
Sunset yellow	Carmoisine	В
Quinoline yellow	_	В
Quinoline yellow	Brilliant blue	В
Quinoline yellow	Carmoisine	В

# Table 1 – Suitable components of RGB for determination of food dyes as individual components and in binary mixtures.

Table 2 $-$ Statistical data for selection of spotted surface.					
Relative standard deviation of signals (C = 50 mg/L, n = 6)	Analyte	Kind of paper			
9.3	Brilliant blue	Filter paper			
8.9	Carmoisine				
7.7	Quinoline yellow				
9.1	Sunset yellow				
14.2	Brilliant blue	Whatman paper			
11.8	Carmoisine	(1 Chr)			
15.2	Quinoline yellow				
14.3	Sunset yellow				
2.6	Brilliant blue	Whatman paper			
2.1	Carmoisine	(No. 43)			
2.8	Quinoline yellow				
1.9	Sunset yellow				

Table 3 - Quantitative features of the method for th	е
selected dyes.	

Analyte	LR (mg/L)	R <sup>2</sup>	LOD (mg/L)	LOQ (mg/L)		
Brilliant blue	20-250	0.9996	4.82	16.06		
Carmoisine	30-180	0.9982	7.23	24.11		
Quinoline yellow	30-250	0.9991	8.05	26.84		
Sunset yellow	30-250	0.9989	7.70	25.68		
$LOD = limit$ of detection (S/N = 3); $LOQ = limit$ of quantification (S/N = 10); $LR = linear$ range; $R^2 =$ square of correlation coefficient.						

#### 3.2.3. Optimization of drying time of spots

In this study, *drying time* was defined as the time interval between spotting the analyte on paper and scanning it. To obtain constant signal for each spot, the time of its drying was studied between 0 and 70 minutes. The results obtained show that by increasing drying time, analytical signals fluctuate until the  $40^{\rm th}$  minute, after which they remain constant. Therefore, the optimal drying time was selected as 40 minutes.

#### 3.2.4. Effect of scanner dots per inch

Dots per inch (DPI) is a measure of spatial printing or video dot density, in particular the number of individual dots that can

Table 5 – Tolerable concentration ratios for determination of 30 mg/L of each dye with the proposed method (relative error  $\pm$  5%). The analytes and interfering dyes are shown in horizontal and vertical columns, respectively.

	BL	CA	QY	SY
BL	_	50	60	60
CA	30	—	60	50
QY	60	60	-	—
SY	30	50	—	—
BL = brill SY = sunse	iant blue; CA et yellow.	A = carmoisine;	QY = quinoline	yellow;

be placed in a line within the span of 2.54 cm (1 in.). By increasing the DPI of scanner, the quality of digital image and the time required for scanning can be increased. The effect of DPI was studied in the range of 75–300, and it was found that resolution of images had no effect on the obtained responses and on their reproducibility. Therefore, the resolution of scanner was adjusted at 75 DPI (the minimum value in scanners) in this study.

#### 3.2.5. Effect of surrounding light

To investigate the effect of light on the obtained signals, two papers were spotted under the same conditions. For drying these spots, one of the papers was exposed to surrounding light and the other one was placed in a darkroom. It was concluded that the drying condition does not affect the analytical signals. Thus, other experiments were carried out under normal laboratory conditions (i.e., no need for a darkroom).

#### 3.3. Quantitative features of the method

The analytical performance of the proposed method was validated by considering the dynamic linear range, square of correlation coefficient, limit of detection (LOD), and limit of quantification (LOQ) for the analytes (summarized in Table 3). In general, the LOD is taken as the lowest concentration of an analyte in a sample that can be detected, but not necessarily

Table 4 – Analytical performances of some similar methods reported in the literature for determination of food dyes.							
Parameter	Proposed method	Altınöz and Toptan <mark>[4]</mark> .	Al-Degs and El-Sheikh [5].	Al-Degs [21].	Alves et al [29].	Huang et al [30].	Berzas et al <mark>[31]</mark> .
Method	Image analysis	Spec	Spec	Spec and HPLC	HPLC	CE	Spec
Data processing	NN	Derivative spectrophotometry	PARAFAC and BLLS/RBL	HLA	NN	NN	PLS/PCR
Dynamic range (mg/L)	20-250	1.0–52	1.0-180	0.5–16	1-100	1-500	0.8–20
Number of dyes studied	4	2	3	3	5	8	3
RSD (%)	0.79-4.46	1.50-3.73	<5	2.2-3.2	<3.87	2.04-4.96	0.4-1.15
LOD (mg/L)	4.82-8.05	0.002-0.05	0.22-0.54	0.04-0.3	0.01-0.1	<0.5	0.003-0.06
LOQ (mg/L)	16.06-26.84	NA	NA	NA	0.03-0.5	NA	NA

BLLS/RBL = bilinear least squares/residual bilinearization; CE = capillary electrophoresis; HLA = hybrid linear analysis; HPLC = high-performance liquid chromatography; LOD = limit of detection; LOQ = limit of quantification; NA = not available; NN = not needed; PARAFAC = parallel factor analysis; PLS/PCR = partial least squares/principal component regression; RSD = relative standard deviation; Spec = spectrophotometry.

Table 6 – Results obtained for determination of dyes in food samples with the proposed method and HPLC (mg/L).								
		Image analysis			HPLC			
	BL	CA	SY	QY	BL	CA	SY	QY
Chocolate—green	ND	-	-	$34 \pm 3$	$4 \pm 1$	-	-	34 ± 2
Chocolate—yellow	—	—	—	36 ± 3	_	—	—	37 ± 3
Chocolate—red	—	50 ± 3	—	—	—	47 ± 2	—	—
Chocolate—orange	—	—	35 ± 3	—	—	—	33 ± 3	—
Soft drink—orange	—	ND	57 ± 2	—	—	$4 \pm 1$	56 ± 3	—
Jelly powder—banana	—	—	ND	$44 \pm 2$	—	—	5 ± 1	46 ± 2
Jelly powder—blue raspberry	$40 \pm 1$	—	—	—	$41 \pm 3$	—	—	—
Jelly powder—watermelon		$48 \pm 1$	30 ± 2	—	—	47 ± 3	32 ± 2	—
BL = brilliant blue; CA = carmoisine; HPLC = high-performance liquid chromatography; ND = not detected; QY = quinoline yellow; SY = sunset yellow.								

quantified, under the stated conditions of the test. The LOQ is the lowest concentration of an analyte in a sample that can be determined with acceptable precision and accuracy under the stated conditions of the test [28].

Despite some limitations, the proposed model offers some advantages over other similar methods, such as simplicity (Table 4). The results indicated that wide linear ranges and good linearity ( $R^2 > 0.99$ ) were achievable for the selected food colors. Acceptable detection and quantification limits in the range of 4.82–8.05 and 16.06–26.84 mg/L, respectively, were obtained by image analysis. Repeatability of the method was assessed by its application on the six similar standard solutions at three different concentrations (40, 90, and 180 mg/L). The relative standard deviations were obtained in the 0.79–4.46% range.

#### 3.4. Study of the interference

The interference of a food dye over other dyes was investigated by analyzing solutions containing two dyes. The results obtained with tolerable maximum error of 5% in the reflectance reading are presented in Table 5. As it can be seen, the interference of a dye over another increases with increasing the molar absorptivity of the interfering dye and spectral overlapping of the dyes. Because of the similarity of the RGB factors and the signals obtained for QY and SY in the software used, the examination of the interference of these dyes on each other was not possible.

#### 3.5. Determination of food dyes in real samples

The BL, CA, SY, and QY dyes in various samples (solid jelly powders, traditional chocolate, and untitled shoddy orange soft drink) were analyzed. The solutions of food samples were prepared as described in the "Preparation of the Real Sample Solutions" section. The dyes in each sample were then identified by paper chromatography. These samples were also subjected to an HPLC method. The results obtained for each of the two methods were then compared (Table 6). As it can be seen, the results obtained from the proposed method was in a good agreement with those obtained for the HPLC. There was also no significant difference between the results obtained by each of the two methods (t test).

#### 4. Conclusion

The proposed analytical approach is a cheap analytical method for obtaining digital images and determination of food colors by a scanner. The optimized technique in conjunction with flatbed scanner was considered to be a facile, economical procedure, and nearly fast for analyzing some food colors in commercial samples. The method has been successfully used for the analysis of some dyes (BL, CA, QY, and SY) as model analytes. The method does not require any preliminary sample-preparation step. The methodology showed acceptable linear ranges for the analytes, with correlation coefficients higher than 0.998, and suitable precision. In this digital imaging method, the costs of equipment and software required are low, and the experimental setup and operating conditions are simple.

#### **Conflicts of interest**

All contributing authors declare no conflicts of interest.

#### REFERENCES

- Wu D, Sun D-W. Colour measurements by computer vision for food quality control—a review. Trends Food Sci Tech 2014;29:5–20.
- [2] Sahraei R, Farmany A, Mortazavi SS. A nanosilver-based spectrophotometry method for sensitive determination of tartrazine in food samples. Food Chem 2013;138:1239–42.
- [3] Komissarchik S, Nyanikova G. Test systems and a method for express detection of synthetic food dyes in drinks. LWT Food Sci Technol 2014;58:315–20.
- [4] Altınöz S, Toptan S. Simultaneous determination of indigotin and Ponceau-4R in food samples by using Vierordt's method, ratio spectra first order derivative and derivative UV spectrophotometry. J Food Compos Anal 2003;16:517–30.
- [5] Al-Degs YS, El-Sheikh AH. Spectrophotometric determination of food dyes in soft drinks by second order multivariate calibration of the absorbance spectra-pH data matrices. Dyes Pigm 2013;97:330–9.
- [6] Schenone AV, Culzoni MJ, Marsili NR, Goicoechea HC. Determination of tartrazine in beverage samples by stoppedflow analysis and three-way multivariate calibration of non-

linear kinetic-spectrophotometric data. Food Chem 2013;138:1928–35.

- [7] Sorouraddin M-H, Rostami A, Saadati M. A simple and portable multi-colour light emitting diode based photocolourimeter for the analysis of mixtures of five common food dyes. Food Chem 2011;127:308–13.
- [8] Sorouraddin M-H, Saadati M. A simple fabrication of portable diffuse reflectometer for simultaneous analysis of common food dyes. Sens Actuators B Chem 2010;145:367–72.
- [9] Yilmaz UT, Ergun F, Yilmaz H. Determination of the food dye carmine in milk and candy products by differential pulse polarography. J Food Drug Anal 2014;22:329–35.
- [10] Gan T, Sun J, Meng W, Song L, Zhang Y. Electrochemical sensor based on graphene and mesoporous TiO<sub>2</sub> for the simultaneous determination of trace colourants in food. Food Chem 2013;141:3731–7.
- [11] Dossi N, Toniolo R, Pizzariello A, Susmel S, Perennes F, Bontempelli G. A capillary electrophoresis microsystem for the rapid in-channel amperometric detection of synthetic dyes in food. J Electroanal Chem 2007;601:1–7.
- [12] de Andrade FI, Florindo Guedes MI, Pinto Vieira ÍG, Pereira Mendes FN, Salmito Rodrigues PA, Costa Maia CS, Marques Ávila MM, de Matos Ribeiro L. Determination of synthetic food dyes in commercial soft drinks by TLC and ion-pair HPLC. Food Chem 2014;157:193–8.
- [13] Huang HY, Chuang CL, Chiu CW, Chung MC. Determination of food colorants by microemulsion electrokinetic chromatography. Electrophoresis 2005;26:867–77.
- [14] Ávila M, Zougagh M, Escarpa A, Ríos Á. Determination of Sudan dyes in food samples using supercritical fluid extraction–capillary liquid chromatography. J Supercrit Fluids 2011;55:977–82.
- [15] Vlajković J, Andrić F, Ristivojević P, Radoičić A, Tešić Ž, Milojković-Opsenica D. Development and validation of a TLC method for the analysis of synthetic food-stuff dyes. J Liq Chromatogr Relat Technol 2013;36:2476–88.
- [16] Zou T, He P, Yasen A, Li Z. Determination of seven synthetic dyes in animal feeds and meat by high performance liquid chromatography with diode array and tandem mass detectors. Food Chem 2013;138:1742–8.
- [17] Long C, Mai Z, Yang Y, Zhu B, Xu X, Lu L, Zou X. Synthesis and characterization of a novel molecularly imprinted polymer for simultaneous extraction and determination of water-soluble and fat-soluble synthetic colorants in chilli products by solid phase extraction and high performance liquid chromatography. J Chromatogr A 2009;1216:8379–85.
- [18] Shen Y, Zhang X, Prinyawiwatkul W, Xu Z. Simultaneous determination of red and yellow artificial food colourants and carotenoid pigments in food products. Food Chem 2014;157:553–8.

- [19] Bonan S, Fedrizzi G, Menotta S, Elisabetta C. Simultaneous determination of synthetic dyes in foodstuffs and beverages by high-performance liquid chromatography coupled with diode-array detector. Dyes Pigm 2013;99:36–40.
- [20] Murty MRVS, Chary NS, Prabhakar S, Prasada Raju N, Vairamani M. Simultaneous quantitative determination of Sudan dyes using liquid chromatography–atmospheric pressure photoionization–tandem mass spectrometry. Food Chem 2009;115:1556–62.
- [21] Al-Degs YS. Determination of three dyes in commercial soft drinks using HLA/GO and liquid chromatography. Food Chem 2009;117:485–90.
- [22] He L, Su Y, Fang B, Shen X, Zeng Z, Liu Y. Determination of Sudan dye residues in eggs by liquid chromatography and gas chromatography–mass spectrometry. Anal Chim Acta 2007;594:139–46.
- [23] Fashandi H, Amirshahi SH, Amani Tehran M, Gorji Kandi S. Evaluation of scanner capability for measuring the color of fabrics with different textures in different setups. Fiber Polym 2010;11:767–74.
- [24] Soponar F, Catalin Mot A, Sarbu C. Quantitative determination of some food dyes using digital processing of images obtained by thin-layer chromatography. J Chromatogr A 2008;1188:295–300.
- [25] Facco P, Bezzo F, Barolo M, Mukherjee R, Romagnoli JA. Monitoring roughness and edge shape on semiconductors through multiresolution and multivariate image analysis. AlChE J 2009;55:1147–60.
- [26] Geladi PLM, Grahn HF, Burger JE. Multivariate images, hyperspectral imaging: background and equipment. In: Grahn HF, Geladi P, editors. Techniques and applications of hyperspectral image analysis. Chichester, UK: John Wiley & Sons, Ltd; 2007. p. 1–15.
- [27] Giorgianni EJ, Madden TE. Digital color management: encoding solutions. Boston, MA: Addison-Wesley Longman Publishing Co; 1998.
- [28] Shrivastava A, Gupta VB. Methods for the determination of limit of detection and limit of quantitation of the analytical methods. Chron Young Sci 2011;2:21–5.
- [29] Alves SP, Brum DM, Branco de Andradea ÉC, Netto ADP. Determination of synthetic dyes in selected foodstuffs by high performance liquid chromatography with UV-DAD detection. Food Chem 2008;107:489–96.
- [30] Huang HY, Shih YC, Chen YC. Determining eight colorants in milk beverages by capillary electrophoresis. J Chromatogr A 2002;959:317–25.
- [31] Berzas JJ, Flores JR, Villaseñor Llerena MJ, Fariñas NR. Spectrophotometric resolution of ternary mixtures of tartrazine, patent blue V and indigo carmine in commercial products. Anal Chim Acta 1999;391:353–64.